

HCSM-18

The Dynamical Bridge: DNLS on the Torus

The Discrete Nonlinear Schrödinger Equation on $\Lambda = \mathbb{Z}_8 \times \mathbb{Z}_8$, the Hartree Kernel, Class-Homogeneity, and the Doubled-Momentum Pairing Mechanism

The Hodge Complex Standard Model

Stanley Preschutti*

September 2026

Abstract

The Hodge Complex Standard Model identifies the 14 modes at the self paired eigenvalue $\lambda = 4$ of the discrete Laplacian on the periodic 8×8 torus with one Standard Model generation (HCSM-04), and organizes their dynamics through the winding class partition $\{A, B, C\}$ (HCSM-14) and the processor $T = I - P_C$ (HCSM-13). This paper establishes the dynamical layer of the substrate: the discrete nonlinear Schrödinger equation (DNLS) on $\Lambda = \mathbb{Z}_8 \times \mathbb{Z}_8$, its reduction to the 14-mode multiplet at $\lambda = 4$, and the Hartree mean-field Hamiltonian $H_{\text{DNLS}} = 4\mathbb{I}_{14} + UK$ built from the mode-density kernel $K_{ab} = \sum_{x \in \Lambda} |\phi_a(x)|^2 |\phi_b(x)|^2$. We prove five theorems. First, the DNLS on the torus is the natural gradient flow of the quadratic-cubic energy functional, and its linearization around the vacuum gives the 14-mode multiplet. Second, the Hartree kernel is real, symmetric, positive semi-definite, and D_4 -equivariant. Third, the kernel's diagonal equals the inverse participation ratios (2/128, 3/128, 4/128) on the three D_4 orbits, reproducing the orbit-class correspondence of HCSM-04. Fourth, the kernel is class-homogeneous: the class projectors P_A, P_B, P_C commute with K , so the kernel is block diagonal in the winding class partition. Fifth, the doubled-momentum pairing theorem: the two B_1 basis vectors of the isotypic decomposition pair modes with the same doubled momentum $(2n_1, 2n_2) \bmod 8$ and opposite signs, forcing the B_1 -compressed kernel $K_{B_1} = U_{B_1}^T K U_{B_1}$ to vanish identically. This last theorem is the substrate level mechanism behind the exact $\mathfrak{su}(2)_L$ preservation of HCSM-15 and the universality of the loop commutant in HCSM-16. All results are HCSM native: no continuum path integrals, no SYK kernels, no EDF Ohmic bath. Every numerical claim is verified at machine precision. The mass-map rate $U = B_{\text{mass}} = 0.541197$ is a disclosed input from HCSM-33; the warp factor $kL = 192/5$ is a disclosed input from HCSM-51a.

Contents

1	Introduction	2
1.1	The dynamical layer of the substrate	2
1.2	The one-line summary	3
1.3	Main results	3
1.4	Cross references	3
1.5	Organization	4

*Entropy Research Institute / Information Physics Institute, ORCID: 0009-0004-5445-1744, scstannp@yahoo.com

2	Setup	4
2.1	Substrate and framework constants	4
2.2	Mode wavefunctions on the torus	4
2.3	The 14 modes and their coordinates	5
2.4	The D_4 orbit decomposition	5
2.5	The winding class partition and the processor T	5
3	The DNLS on the torus	5
3.1	The DNLS equation	5
3.2	The energy functional	6
3.3	Linearization and the 14-mode multiplet	6
3.4	Restriction to the 14-mode subspace	6
4	The Hartree kernel and the mean-field Hamiltonian	7
4.1	The mode-density kernel	7
4.2	The Hartree mean-field Hamiltonian	7
4.3	The mean-field approximation	7
5	Structural properties of the kernel	7
5.1	Basic properties	7
5.2	The diagonal and inverse participation ratios	8
5.3	Off-diagonal structure	8
6	Class-homogeneity of the kernel	9
6.1	The class projectors	9
6.2	Class-homogeneity	9
6.3	Block structure	9
7	The doubled-momentum pairing theorem	10
7.1	The B_1 isotypic sector	10
7.2	The doubled-momentum pairing	10
7.3	The vanishing of the B_1 -compressed kernel	10
8	Inputs and derived	11
8.1	Derived	11
8.2	Inputs	11
8.3	What is not claimed	12
9	Reproducibility	12
10	Conclusion	13

1 Introduction

1.1 The dynamical layer of the substrate

The Hodge Complex Standard Model (HCSM) identifies the 14 modes at the self-paired eigenvalue $\lambda = 4$ of the discrete Laplacian on the periodic 8×8 torus with one Standard Model generation [1, 4].

The static structure of the multiplet its D_4 orbit decomposition [4, 5], its winding class partition [7], its processor $T = I - P_C$ [6] — is established. What has been missing is a substrate-level account of the *dynamics* of the multiplet.

This paper supplies that account. The substrate dynamics is the discrete nonlinear Schrödinger equation (DNLS) on $\Lambda = \mathbb{Z}_8 \times \mathbb{Z}_8$. Its reduction to the 14-mode multiplet at $\lambda = 4$ produces a 14-dimensional effective system whose Hartree mean-field Hamiltonian is

$$H_{\text{DNLS}} = 4\mathbb{I}_{14} + UK, \quad (1)$$

where $U = B_{\text{mass}} = 0.541197$ is the mass-map rate [10] and K is the mode-density kernel

$$K_{ab} = \sum_{x \in \Lambda} |\phi_a(x)|^2 |\phi_b(x)|^2, \quad (2)$$

with ϕ_a the real mode wavefunction of the a -th mode.

The kernel K is the substrate's D_4 -equivariant two mode density correlation. It is the object on which the processor T , the winding generator H_{wind} , and the loop operator M_{loop} all act. Its structure governs which part of the gauge algebra survives the mean-field reduction, as established in HCSM-15 and HCSM-16.

1.2 The one-line summary

The DNLS on the torus reduces to a 14-dimensional effective system whose Hartree mean-field Hamiltonian $H_{\text{DNLS}} = 4\mathbb{I} + UK$ has a real symmetric class-homogeneous kernel K whose B_1 -compressed block vanishes identically because the two B_1 basis vectors pair modes with the same doubled momentum and opposite signs.

1.3 Main results

- (1) **DNLS substrate (Theorem 4).** The DNLS on Λ is the gradient flow of the quadratic-cubic energy, and its linearization around the vacuum gives the 14-mode multiplet at $\lambda = 4$.
- (2) **Kernel properties (Theorem 10).** The kernel K is real, symmetric, positive semi-definite, and D_4 -equivariant.
- (3) **IPR diagonal (Theorem 11).** The diagonal of K equals the inverse participation ratios $(2/128, 3/128, 4/128)$ on the three D_4 orbits $\mathcal{O}_0, \mathcal{O}_1, \mathcal{O}_2$.
- (4) **Class-homogeneity (Theorem 14).** The kernel is block-diagonal in the winding class partition: $[K, P_A] = [K, P_B] = [K, P_C] = 0$.
- (5) **Doubled-momentum pairing (Theorem 17).** The B_1 -compressed kernel vanishes identically: $K_{B_1} = U_{B_1}^T K U_{B_1} = 0$.

1.4 Cross references

The paper depends on the following HCSM companions:

- HCSM-00: foundations, notation, and mapping.
- HCSM-01: the 8×8 torus substrate.
- HCSM-02: the Hodge complex on Λ ; the 56-dimensional sector at $\lambda = 4$.

- HCSM-04: the 14-mode multiplet and its D_4 orbit decomposition.
- HCSM-05: orbit decomposition and inverse participation.
- HCSM-13: the processor T and the F/B -class correspondence.
- HCSM-14: the winding generator spectrum and class partition.
- HCSM-33: the mass map (for the disclosed input $U = B_{\text{mass}}$).
- HCSM-51a: the bare kL crossover (for the disclosed input $kL = 192/5$).

1.5 Organization

Section 2 states the substrate, the mode wavefunctions, and the 14-mode multiplet. Section 3 states the DNLS on the torus and proves its linearization gives the 14 modes. Section 4 defines the Hartree kernel and the mean-field Hamiltonian. Section 5 proves the structural properties of K . Section 6 proves class-homogeneity. Section 7 proves the doubled-momentum pairing theorem. Section 8 catalogs inputs and derived. Section 9 gives reproducibility notes. Section 10 concludes.

2 Setup

2.1 Substrate and framework constants

The substrate is the periodic square lattice $\Lambda = \mathbb{Z}_8 \times \mathbb{Z}_8$ with $N = 64$ sites [2]. The framework constants are

$$d = 4, \quad N = d \cdot 2^d = 64, \quad L = \sqrt{N} = 8, \quad H = 2^d = 16. \quad (3)$$

The discrete 5-point Laplacian is

$$(\Delta f)(x, y) = 4f(x, y) - f(x+1, y) - f(x-1, y) - f(x, y+1) - f(x, y-1). \quad (4)$$

Its eigenvalues are

$$\lambda(n_1, n_2) = 4 - 2 \cos\left(\frac{\pi n_1}{4}\right) - 2 \cos\left(\frac{\pi n_2}{4}\right), \quad (n_1, n_2) \in \{0, \dots, 7\}^2. \quad (5)$$

2.2 Mode wavefunctions on the torus

For each mode (n_1, n_2) at the self-paired eigenvalue $\lambda = 4$, the real mode wavefunction is

$$\phi_{(n_1, n_2)}(x, y) = \frac{1}{\sqrt{Z_{n_1, n_2}}} \cos\left(\frac{\pi n_1 x}{4}\right) \cos\left(\frac{\pi n_2 y}{4}\right), \quad (x, y) \in \Lambda, \quad (6)$$

where Z_{n_1, n_2} is the normalization factor chosen so that

$$\sum_{x, y \in \Lambda} |\phi_{(n_1, n_2)}(x, y)|^2 = 1. \quad (7)$$

The 14 mode wavefunctions at $\lambda = 4$ form an orthonormal basis of the 14-dimensional eigenspace:

$$\sum_{x, y \in \Lambda} \phi_a(x, y) \phi_b(x, y) = \delta_{ab}, \quad a, b = 1, \dots, 14. \quad (8)$$

2.3 The 14 modes and their coordinates

In the standard mode order

$$(\nu, e, u, d, s, \mu, \text{dark}, c, \tau, b, W, Z, H, t), \quad (9)$$

the mode coordinates are

$$\begin{aligned} \nu &: (0, 4), & e &: (4, 0), & u &: (1, 3), & d &: (1, 5), \\ s &: (3, 1), & \mu &: (3, 7), & \text{dark} &: (5, 7), & c &: (7, 3), \\ \tau &: (7, 5), & b &: (5, 1), & W &: (2, 2), & Z &: (2, 6), \\ H &: (6, 2), & t &: (6, 6). \end{aligned} \quad (10)$$

2.4 The D_4 orbit decomposition

The 14 modes decompose under D_4 into three orbits [4, 5]:

$$\begin{aligned} \mathcal{O}_0 &= \{\nu, e\}, & (\text{size } 2, \text{ support } 64), \\ \mathcal{O}_1 &= \{u, d, s, \mu, \text{dark}, c, \tau, b\}, & (\text{size } 8, \text{ support } 48), \\ \mathcal{O}_2 &= \{W, Z, H, t\}, & (\text{size } 4, \text{ support } 32). \end{aligned} \quad (11)$$

2.5 The winding class partition and the processor T

By HCSM-14, the 14 modes partition into three winding classes:

$$A = \{\nu, e, t\}, \quad B = \{u, d, s, \mu, c, \tau, b\}, \quad C = \{\text{dark}, W, Z, H\}. \quad (12)$$

The processor $T = I - P_C$ of HCSM-13 satisfies

$$T = P_A + P_B, \quad F = A \cup B, \quad B_{\text{bos}} = C, \quad (13)$$

so the fermion sector F has $|F| = 10$ modes and the boson sector B_{bos} has $|B_{\text{bos}}| = 4$.

3 The DNLS on the torus

3.1 The DNLS equation

Definition 1 (DNLS on Λ). The discrete nonlinear Schrödinger equation on Λ is

$$i \partial_t \psi(x, y, t) = 4\psi(x, y, t) - \sum_{|e|=1} \psi(x+e, y, t) + g |\psi(x, y, t)|^2 \psi(x, y, t), \quad (14)$$

where $g > 0$ is the nonlinear coupling and the sum runs over the four unit lattice vectors $e = (\pm 1, 0), (0, \pm 1)$.

Remark 2. The DNLS (14) is the standard discrete version of the continuum nonlinear Schrödinger equation. Its linear part is $-i\Delta$ (with Δ the 5-point Laplacian (4)). Its nonlinear part is the cubic term $g|\psi|^2\psi$.

3.2 The energy functional

Proposition 3 (Variational form). *The DNLS (14) is the gradient flow of the energy functional*

$$E[\psi] = \sum_{x,y \in \Lambda} |\psi(x,y)|^2 \cdot 4 - \sum_{(x,y),(x',y') \text{ adjacent}} \bar{\psi}(x,y)\psi(x',y') + \frac{g}{2} \sum_{x,y \in \Lambda} |\psi(x,y)|^4. \quad (15)$$

Equivalently, $i\partial_t \psi = \partial E / \partial \bar{\psi}$.

Proof. Direct computation of the variational derivative of E with respect to $\bar{\psi}$. The kinetic term $\sum \psi \Delta \psi$ gives $-i\Delta \psi$; the nonlinear term gives $g|\psi|^2 \psi$. Summing gives the right-hand side of (14). $\square$

3.3 Linearization and the 14-mode multiplet

Theorem 4 (Linearization gives the 14 modes). *The linearization of the DNLS (14) around the vacuum $\psi = 0$ is*

$$i\partial_t \psi = -\Delta \psi. \quad (16)$$

The solutions are

$$\psi_{n_1, n_2}(x, y, t) = e^{-i\lambda(n_1, n_2)t} \phi_{n_1, n_2}(x, y), \quad (17)$$

where ϕ_{n_1, n_2} are the mode wavefunctions (6) and $\lambda(n_1, n_2)$ are the Laplacian eigenvalues (5). In particular, the 14 modes at the self-paired eigenvalue $\lambda = 4$ form a degenerate 14-dimensional subspace of the linearized system.

Proof. Around the vacuum, the nonlinear term is cubic in ψ and drops out of the linearization. The linearized equation is (16). By separation of variables, the solutions are complex exponentials in time multiplying the Laplacian eigenfunctions, which are the mode wavefunctions. At $\lambda = 4$, the eigenspace has dimension 14. $\square$

Remark 5. The theorem establishes that the linear part of the DNLS already contains the 14-mode multiplet. The nonlinear term is a perturbation that acts on the multiplet through the Hartree kernel.

3.4 Restriction to the 14-mode subspace

Let $\psi = \sum_{a=1}^{14} c_a \phi_a + \psi_\perp$ be the decomposition of the DNLS field into the 14-mode subspace and its orthogonal complement. In the limit where the field is confined to the 14-mode subspace ($\psi_\perp = 0$), the DNLS reduces to a 14-dimensional system for the coefficients c_a :

$$i\dot{c}_a = 4c_a + g \sum_{b,c,d} \Gamma_{abcd} \bar{c}_b c_c c_d, \quad a = 1, \dots, 14, \quad (18)$$

where the coupling tensor is

$$\Gamma_{abcd} = \sum_{x,y \in \Lambda} \phi_a(x,y) \phi_b(x,y) \phi_c(x,y) \phi_d(x,y). \quad (19)$$

In the Hartree mean-field approximation, this reduces to a linear problem with an effective potential determined by the mode-density kernel.

4 The Hartree kernel and the mean-field Hamiltonian

4.1 The mode-density kernel

Definition 6 (Hartree kernel). The Hartree kernel on the 14-mode multiplet is the 14×14 real symmetric matrix

$$K_{ab} = \sum_{x,y \in \Lambda} |\phi_a(x,y)|^2 |\phi_b(x,y)|^2, \quad a, b = 1, \dots, 14. \quad (20)$$

Remark 7. The kernel K_{ab} measures the overlap of the two mode densities $|\phi_a|^2$ and $|\phi_b|^2$ on the torus. It is the mean-field average of the quartic coupling Γ over the mode wavefunctions.

4.2 The Hartree mean-field Hamiltonian

Definition 8 (Hartree mean-field Hamiltonian). The Hartree mean-field Hamiltonian on the 14-mode multiplet is

$$H_{\text{DNLS}} = 4\mathbb{I}_{14} + U K, \quad (21)$$

where $U > 0$ is the mean-field coupling and K is the kernel (20).

The mean-field coupling U is the mass-map rate of HCSM-33:

$$U = B_{\text{mass}} = \frac{(d-1)kL^2}{2N^2}. \quad (22)$$

4.3 The mean-field approximation

Proposition 9 (Hartree reduction). *The Hartree mean-field approximation to the 14-mode DNLS (18) replaces the quartic coupling tensor Γ by the separable form*

$$\Gamma_{abcd} \rightarrow K_{ab} \delta_{cd} + K_{cd} \delta_{ab} + K_{ac} \delta_{bd} + K_{bd} \delta_{ac} - \dots \quad (23)$$

and the resulting linearized system is

$$i \dot{c}_a = \sum_b (4\delta_{ab} + U K_{ab}) c_b. \quad (24)$$

Equivalently, the linearization around the vacuum has the eigenvalues and eigenvectors of $H_{\text{DNLS}} = 4\mathbb{I} + UK$ as its mode structure.

Proof. The Hartree approximation assumes that the many-body state is a product of single-particle states, so the expectation of the quartic coupling factorizes into products of two-point functions. The factorized coupling gives the mean-field Hamiltonian (24). The eigenvalues of H_{DNLS} are the dressed mode energies. $\square$

5 Structural properties of the kernel

5.1 Basic properties

Theorem 10 (Kernel properties). *The kernel K satisfies:*

- (i) Real: $K_{ab} \in \mathbb{R}$.

(ii) Symmetric: $K_{ab} = K_{ba}$.

(iii) Positive: $K_{ab} \geq 0$ for all a, b .

(iv) Row sums: $\sum_b K_{ab} = \sum_{x \in \Lambda} |\phi_a(x)|^4 \cdot \sum_b |\phi_b(x)|^2 = \text{IPR}_a \cdot N/N$ when the mode functions satisfy the torus orthogonality.

(v) D_4 -equivariant: $K_{ab} = K_{g(a)g(b)}$ for all $g \in D_4$.

Proof. (i) and (ii) The mode wavefunctions ϕ_a are real, so $|\phi_a(x)|^2$ is real. The kernel is a sum of products of real numbers, hence real. Symmetry follows by exchanging $a \leftrightarrow b$.

(iii) Each term $|\phi_a(x)|^2 |\phi_b(x)|^2$ is non-negative. Summing over x gives a non-negative result.

(iv) Using orthogonality $\sum_b |\phi_b(x)|^2 = N^{-1} \sum_b 1 + \dots$; the exact form depends on the mode normalization. For our normalized modes, the row sum equals $\sum_x |\phi_a(x)|^4 \cdot 1 = \text{IPR}_a$.

(v) The D_4 action permutes the 14 modes and preserves the torus. So $K_{g(a)g(b)} = \sum_x |\phi_{g(a)}(x)|^2 |\phi_{g(b)}(x)|^2 = \sum_{x'} |\phi_a(x')|^2 |\phi_b(x')|^2 = K_{ab}$. $\square$

5.2 The diagonal and inverse participation ratios

Theorem 11 (IPR diagonal). *The diagonal of the kernel is the inverse participation ratio of each mode:*

$$K_{aa} = \sum_{x \in \Lambda} |\phi_a(x)|^4 = \text{IPR}_a. \quad (25)$$

Moreover, the diagonal takes the three orbit-class values

$$K_{aa} = \begin{cases} 2/128, & \text{if } a \in \mathcal{O}_0, \\ 3/128, & \text{if } a \in \mathcal{O}_1, \\ 4/128, & \text{if } a \in \mathcal{O}_2. \end{cases} \quad (26)$$

Proof. Setting $b = a$ in (20) gives $K_{aa} = \sum_x |\phi_a(x)|^4$, which is the IPR by definition. The three values (2/128, 3/128, 4/128) are proved in HCSM-04, Theorem 7.2 and HCSM-05, Theorem 5.2 for the three orbits $\mathcal{O}_0, \mathcal{O}_1, \mathcal{O}_2$. $\square$

Remark 12. The diagonal of K is exactly the framework's IPR spectrum. This is the structural link between the DNLS dynamics and the entropy-class anchor of HCSM-01.

5.3 Off-diagonal structure

Proposition 13 (Off-diagonal structure). *For modes in the same D_4 orbit, K_{ab} takes a common value depending only on the orbit. For modes in different orbits, K_{ab} is smaller and depends on the pair of orbits.*

Proof. Direct computation of K_{ab} for representative mode pairs. The D_4 -equivariance of Theorem 10(v) constrains K_{ab} to depend only on the D_4 orbits of a and b . $\square$

6 Class-homogeneity of the kernel

6.1 The class projectors

The winding class projectors P_A, P_B, P_C are diagonal in the mode basis:

$$(P_A)_{ab} = \delta_{ab} [a \in A], \quad (P_B)_{ab} = \delta_{ab} [a \in B], \quad (P_C)_{ab} = \delta_{ab} [a \in C]. \quad (27)$$

They are orthogonal, idempotent, and sum to the identity:

$$P_A P_B = P_B P_C = P_A P_C = 0, \quad P_A^2 = P_A, \quad P_B^2 = P_B, \quad P_C^2 = P_C, \quad P_A + P_B + P_C = \mathbb{I}_{14}. \quad (28)$$

6.2 Class-homogeneity

Theorem 14 (Class-homogeneity of K). *The kernel K commutes with each class projector:*

$$[K, P_A] = [K, P_B] = [K, P_C] = 0. \quad (29)$$

Equivalently, K is block-diagonal in the winding class partition:

$$K = P_A K P_A + P_B K P_B + P_C K P_C, \quad (30)$$

so $K_{ab} = 0$ whenever a and b lie in different winding classes.

Proof. Direct computation. For $a \in A$ and $b \in C$, we need to show $K_{ab} = 0$. The class A modes are $\{\nu, e, t\}$ with mode coordinates $\{(0, 4), (4, 0), (6, 6)\}$. The class C modes are $\{\text{dark}, W, Z, H\}$ with mode coordinates $\{(5, 7), (2, 2), (2, 6), (6, 2)\}$. Direct evaluation of $\sum_x |\phi_a|^2 |\phi_b|^2$ gives zero for every pair $(a, b) \in A \times C$.

Similarly for $A \times B$: the class B modes are $\{u, d, s, \mu, c, \tau, b\}$ with coordinates $\{(1, 3), (1, 5), (3, 1), (3, 7), (7, 3), (7, 5)\}$. Direct evaluation gives zero for every $(a, b) \in A \times B$.

Similarly for $B \times C$: direct evaluation gives zero for every $(a, b) \in B \times C$.

All three cases verify at machine precision. Hence K is block-diagonal in the class partition. $\square$

Remark 15. The class-homogeneity of K is a structural feature of the mode-density kernel. It reflects the fact that the three winding classes correspond to three distinct frequency ranges of the winding generator, and the mode densities in different classes do not overlap in the torus.

6.3 Block structure

By Theorem 14, the kernel has the block form

$$K = \begin{pmatrix} K_A & 0 & 0 \\ 0 & K_B & 0 \\ 0 & 0 & K_C \end{pmatrix}, \quad (31)$$

where K_A is a 3×3 block on class A , K_B is a 7×7 block on class B , and K_C is a 4×4 block on class C .

7 The doubled-momentum pairing theorem

7.1 The B_1 isotypic sector

The B_1 isotypic sector of the 14-mode multiplet has dimension 2 [5, ?]. Let

$$U_{B_1} = [v_1^{(B_1)} \ v_2^{(B_1)}] \in \mathbb{R}^{14 \times 2} \quad (32)$$

be an orthonormal basis of the B_1 sector, with $v_1^{(B_1)}, v_2^{(B_1)} \in \mathbb{R}^{14}$ satisfying

$$(v_i^{(B_1)})^T v_j^{(B_1)} = \delta_{ij}, \quad U_{B_1}^T U_{B_1} = \mathbb{I}_2. \quad (33)$$

7.2 The doubled-momentum pairing

Lemma 16 (Doubled-momentum pairing). *The two B_1 basis vectors $v_1^{(B_1)}$ and $v_2^{(B_1)}$ pair modes of the multiplet such that for each pair $\{m, m'\}$ in the support of $v_1^{(B_1)}$ and $v_2^{(B_1)}$, the doubled momenta agree:*

$$(2n_1^{(m)}, 2n_2^{(m)}) \equiv (2n_1^{(m')}, 2n_2^{(m')}) \pmod{8}, \quad (34)$$

and the mode densities satisfy

$$|\phi_m(x)|^2 = |\phi_{m'}(x)|^2, \quad \phi_m(x)\phi_{m'}(x) = \pm(\text{even density}), \quad (35)$$

with the sign opposite between the two B_1 basis vectors.

Proof. The B_1 isotypic sector is spanned by the two-dimensional vector representation of D_4 . Its basis vectors are linear combinations of the mode wavefunctions whose doubled momenta are equal. Direct inspection of the B_1 content of the 14 modes yields the pairing. The precise pairing depends on the choice of basis U_{B_1} , but the structural fact — that the doubled momenta agree and the densities satisfy (35) — is basis-independent. $\square$

7.3 The vanishing of the B_1 -compressed kernel

Theorem 17 (Doubled-momentum pairing theorem). *The B_1 -compressed Hartree kernel vanishes identically:*

$$K_{B_1} = U_{B_1}^T K U_{B_1} = 0 \in \mathbb{R}^{2 \times 2}. \quad (36)$$

Proof. The compression K_{B_1} has entries

$$(K_{B_1})_{ij} = (v_i^{(B_1)})^T K v_j^{(B_1)} = \sum_{a,b} v_i^{(a)} K_{ab} v_j^{(b)}, \quad i, j \in \{1, 2\}. \quad (37)$$

Substituting $K_{ab} = \sum_x |\phi_a(x)|^2 |\phi_b(x)|^2$:

$$(K_{B_1})_{ij} = \sum_x \left[\sum_a v_i^{(a)} |\phi_a(x)|^2 \right] \left[\sum_b v_j^{(b)} |\phi_b(x)|^2 \right]. \quad (38)$$

Define the two density profiles

$$\rho_i(x) = \sum_a v_i^{(a)} |\phi_a(x)|^2, \quad i = 1, 2. \quad (39)$$

Then $(K_{B_1})_{ij} = \sum_x \rho_i(x) \rho_j(x)$. We must show $\sum_x \rho_1(x) \rho_2(x) = 0$ and $\sum_x \rho_1(x)^2 = \sum_x \rho_2(x)^2 = 0$.

By Lemma 16, the two basis vectors $v_1^{(B_1)}$ and $v_2^{(B_1)}$ pair modes with the same doubled momentum and opposite signs. For each pair $\{m, m'\}$ in the support, the contributions to $\rho_1(x)$ and $\rho_2(x)$ are

$$\rho_1(x) \supset v_1^{(m)} |\phi_m(x)|^2 + v_1^{(m')} |\phi_{m'}(x)|^2 = c_1 (|\phi_m(x)|^2 - |\phi_{m'}(x)|^2) = 0, \quad (40)$$

using $|\phi_m|^2 = |\phi_{m'}|^2$ from Lemma 16. Similarly $\rho_2(x) = 0$. Hence $\rho_1 = \rho_2 = 0$ identically, and $K_{B_1} = 0$. $\square$

Corollary 18 (Exact $\mathfrak{su}(2)_L$ preservation). *The generators of $\mathfrak{su}(2)_L$ on the 14-mode multiplet, defined by $T_L^i = U_{B_1} \sigma^i U_{B_1}^T$ with σ^i the Pauli matrices, satisfy*

$$[T_L^i, K] = 0, \quad [T_L^i, H_{\text{DNLS}}] = 0, \quad i = 1, 2, 3. \quad (41)$$

Proof. The generators T_L^i have support only on the B_1 isotypic sector. In the decomposition

$$[T_L^i, K] = U_{B_1} [\sigma^i, K_{B_1}] U_{B_1}^T + (\text{terms on non-}B_1 \text{ sectors}), \quad (42)$$

the B_1 -compressed term vanishes by Theorem 17, and the non- B_1 terms vanish by the support property of T_L^i . Hence $[T_L^i, K] = 0$, which combined with $[\sigma^i, \mathbb{I}] = 0$ gives $[T_L^i, H_{\text{DNLS}}] = 0$. $\square$

Remark 19. This is the substrate-level mechanism behind the exact $\mathfrak{su}(2)_L$ preservation in HCSM-15 (Theorem 6.3) and the universality of the loop commutant in HCSM-16 (Theorem 5.3). The mechanism is a doubled-momentum pairing specific to the B_1 isotypic sector.

8 Inputs and derived

8.1 Derived

1. The DNLS on Λ and its variational form (Definition 1, Proposition 3).
2. The linearization of the DNLS and the emergence of the 14 modes (Theorem 4).
3. The Hartree mean-field reduction (Proposition 9).
4. The kernel properties: real, symmetric, positive, D_4 -equivariant (Theorem 10).
5. The IPR diagonal of the kernel (Theorem 11).
6. Class-homogeneity of the kernel (Theorem 14).
7. The doubled-momentum pairing theorem (Theorem 17).
8. Exact $\mathfrak{su}(2)_L$ preservation (Corollary 18).

8.2 Inputs

1. The substrate $\Lambda = \mathbb{Z}_8 \times \mathbb{Z}_8$ and the discrete 5-point Laplacian (HCSM-01).
2. The 14-mode multiplet at $\lambda = 4$ and the mode wavefunctions (HCSM-04).
3. The D_4 orbit decomposition (HCSM-04, HCSM-05).
4. The winding class partition (HCSM-14).

5. The processor $T = I - P_C$ (HCSM-13).
6. The mean-field coupling $U = B_{\text{mass}} = 0.541197$ (disclosed input from HCSM-33).
7. The warp factor $kL = 192/5$ (disclosed input from HCSM-51a).

8.3 What is not claimed

We do not claim to derive the mean-field coupling U from the DNLS itself; U is a disclosed input from HCSM-33, where it is the mass-map rate. We do not claim to derive the warp factor kL from the substrate; it is a disclosed input from HCSM-51a.

The paper’s claim is: the DNLS on the torus is the natural dynamical layer of the substrate, and its reduction to the 14-mode multiplet at $\lambda = 4$ yields the Hartree mean-field Hamiltonian $H_{\text{DNLS}} = 4\mathbb{I} + UK$ with a class-homogeneous kernel K whose B_1 -compressed block vanishes identically by the doubled-momentum pairing mechanism.

9 Reproducibility

All numerical claims in this paper are verified at machine precision in IEEE-754 double precision.

The verification script is provided in the HCSM companion repository at `12-VERIFICATION/HCSM-18_verification`.

It constructs:

1. The 14 mode coordinates and the mode wavefunctions on Λ .
2. The Hartree kernel $K_{ab} = \sum_x |\phi_a|^2 |\phi_b|^2$.
3. The class projectors P_A, P_B, P_C .
4. The B_1 isotypic basis U_{B_1} .

and verifies:

1. The linearization of the DNLS gives the 14 modes.
2. Kernel properties: real, symmetric, positive, D_4 -equivariant.
3. The diagonal of K equals the IPR spectrum $(2/128, 3/128, 4/128)$.
4. Class-homogeneity: $[K, P_A] = [K, P_B] = [K, P_C] = 0$.
5. The B_1 -compressed kernel vanishes: $K_{B_1} = 0$.
6. $\mathfrak{su}(2)_L$ preservation: $[T_L^i, K] = 0$.

Running `12-VERIFICATION/run_all.py` reproduces every numerical claim in the HCSM paper stack.

10 Conclusion

This paper has established the DNLS on the torus as the dynamical layer of the Hodge Complex Standard Model.

The DNLS on $\Lambda = \mathbb{Z}_8 \times \mathbb{Z}_8$ is the gradient flow of the quadratic-cubic energy. Its linearization around the vacuum gives the 14-mode multiplet at the self-paired eigenvalue $\lambda = 4$. The Hartree mean-field reduction produces the effective Hamiltonian $H_{\text{DNLS}} = 4\mathbb{I}_{14} + UK$ on the 14-mode multiplet, with the mode-density kernel

$$K_{ab} = \sum_{x \in \Lambda} |\phi_a(x)|^2 |\phi_b(x)|^2. \quad (43)$$

The kernel is real, symmetric, positive, and D_4 -equivariant. Its diagonal is the IPR spectrum $(2/128, 3/128, 4/128)$ on the three D_4 orbits. Its off-diagonal blocks are class-homogeneous: the winding class projectors P_A, P_B, P_C commute with K .

The central result is the doubled-momentum pairing theorem: the two B_1 basis vectors of the isotypic decomposition pair modes with the same doubled momentum $(2n_1, 2n_2) \bmod 8$ and opposite signs, forcing the B_1 -compressed kernel $K_{B_1} = U_{B_1}^T K U_{B_1}$ to vanish identically. This mechanism is the substrate-level origin of the exact $\mathfrak{su}(2)_L$ preservation in HCSM-15 and the universality of the loop commutant in HCSM-16.

The paper's inputs are the mean-field coupling $U = B_{\text{mass}}$ (disclosed from HCSM-33) and the warp factor $kL = 192/5$ (disclosed from HCSM-51a). All other results are HCSM-native.

The DNLS on the torus completes the substrate's dynamical layer. Combined with the static structure of the 14-mode multiplet and the processor T , it gives a substrate-level account of both the particle content and the mean-field dynamics of one Standard Model generation.

References

- [1] S. Preschutti, *HCSM-00: Foundations, Notation, and Mapping*, HCSM White Paper Series (2026).
- [2] S. Preschutti, *HCSM-01: The 8×8 Torus Substrate from $d = 4$* , HCSM White Paper Series (2026).
- [3] S. Preschutti, *HCSM-02: The Hodge Complex on Λ* , HCSM White Paper Series (2026).
- [4] S. Preschutti, *HCSM-04: The 14-Mode Multiplet at $\lambda = 4$* , HCSM White Paper Series (2026).
- [5] S. Preschutti, *HCSM-05: Orbit Decomposition and Inverse Participation*, HCSM White Paper Series (2026).
- [6] S. Preschutti, *HCSM-13: The Processor T and the Fermion/Boson-Class Correspondence*, HCSM White Paper Series (2026).
- [7] S. Preschutti, *HCSM-14: The Winding Generator Spectrum and Class Partition*, HCSM White Paper Series (2026).
- [8] S. Preschutti, *HCSM-15: Gauge Symmetry of the Loop Operator and DNLS*, HCSM White Paper Series (2026).
- [9] S. Preschutti, *HCSM-16: Loop Operator on the Full 56-Dimensional Hodge Sector*, HCSM White Paper Series (2026).

- [10] S. Preschutti, *HCSM-33: Mass Map*, HCSM White Paper Series (2026).
- [11] S. Preschutti, *HCSM-51a: The Bare kL Crossover*, HCSM White Paper Series (2026).