
HCSM-22.a

The Zero State Selection Principle

The Entropy Well Variational Principle,
the Landauer Minimum Information Route,
and the Vacuum as the Unique Maximizer of $S(s)/S_0 = 1 - \gamma\|M^T s\|^2$

Stanley Preschutti

Entropia Research Institute / Information Physics Institute

ORCID: 0009-0004-5445-1744

scstamp@yahoo.com

September 2026

Sub-paper of HCSM-22 (The Zero State)

Abstract

HCSM-22 constructs the zero state s as the unique nullvector of the 14×19 measurement matrix M of vacuum-native observables, but explicitly does not select s : the zero state is stated there as “a structural fact about the multiplet and the observable set, not a selection rule.” This sub-paper supplies the missing selection rule. We prove that the vacuum is the state of the 14-mode multiplet at the self-paired eigenvalue $\lambda = 4$ that minimizes the total squared expectation value of the 19 observables, i.e. $s = \arg \min_{\|s\|=1} \|M^T s\|^2$. Two independent derivations are given, both from the substrate. First, the *Landauer route*: the processor T of HCSM-13 operates at the deterministic Landauer bound (HCSM-21, Thm. 12.1), the vacuum is the state of minimum information, and minimum information is exactly $M^T s = 0$. Second, the *entropy-well route*: identifying the dimensional deviation $\epsilon = D - 4$ of HCSM-54 with the observable magnitude $\|M^T s\|$, the vacuum entropy well becomes $S(s)/S_0 = 1 - \gamma\|M^T s\|^2 + O(\|M^T s\|^4)$, which is maximized exactly when $M^T s = 0$. The two routes are dual descriptions of the same physical fact: the vacuum is the state from which the machine extracts zero information. The uniqueness of the maximizer is established from $\text{rank}(M) = 13$ and $\text{nullity}(M^T) = 1$ (HCSM-22, Thm. 4). Item 2.1 of the HCSM derivation list (Zero State Selection Principle) is closed. All numerical claims are verified at machine precision in IEEE-754 double precision.

Derivation status summary.

- **DERIVED**: the selection principle $s = \arg \min \|M^T s\|^2$; the Landauer route; the entropy-well route; the equivalence of the two routes; the uniqueness of the maximizer; the inherited structural properties of s (rs-oddness, exact ratios, weight fractions, closed forms in $\mathbb{Q}(\sqrt{2})$).
- **DISCLOSED INPUTS**: the 19 vacuum-native observables (HCSM-22, Table 1); the processor $T = I - P_C$ (HCSM-13); the stability coefficient $\gamma = c_2/I(2)$ (HCSM-54); the Landauer bound (HCSM-21).
- **OPEN**: the exact closed-form reconstruction of the 8 non-verifiable columns of M (charge, three pressure components, P_{flip} , v_x , v_y , $|v|$); the identification of the bridge constant in $\epsilon = c\|M^T s\|$ beyond the normalization choice $c = 1$.

Contents

1	Introduction	3
1.1	The zero state problem	3
1.2	The one-line summary	3
1.3	Main results	3
1.4	Cross references	4
1.5	Organization	4
2	Setup	4
2.1	The substrate and the 14-mode multiplet	4
2.2	The 19 vacuum-native observables	5
2.3	The measurement matrix	5
2.4	The entropy well and the stability coefficient	6
2.5	The processor and the Landauer channel	6
3	The Selection Principle	6
4	The Landauer Route	7
5	The Entropy Well Route	7
6	Equivalence of the Two Routes	8
7	Uniqueness of the Maximizer	9
8	Inherited Structure	9
9	Derivation Status	9
9.1	Derived in this paper	9
9.2	Disclosed inputs	10
9.3	Open items	10
10	Falsification Tests	10
11	Reproducibility	10
12	Conclusion	11

1 Introduction

1.1 The zero state problem

The Hodge Complex Standard Model (HCSM) identifies the 14 modes at the self paired eigenvalue $\lambda = 4$ of the discrete 5-point Laplacian on the periodic 8×8 torus $\Lambda = \mathbb{Z}_8 \times \mathbb{Z}_8$ with one Standard Model generation [1, 2, ?]. HCSM-22 [6] constructs the zero state: the unique configuration s of the 14-mode multiplet consistent with the simultaneous vanishing of 19 vacuum native observables. The 19 observables form a 14×19 real measurement matrix M , whose transpose has rank 13 and nullity 1. The unique nullvector s (up to scale) is the zero state.

HCSM-22 establishes the following structural properties of s :

- (i) s is odd under the chiral reflection rs : $s(rs(m)) = -s(m)$ for all 14 modes (Thm. 8).
- (ii) The two rs -fixed modes $W = (2, 2)$ and $t = (6, 6)$ have exactly zero amplitude (Cor. 9).
- (iii) The six independent amplitudes $a_1, \dots, a_6$ satisfy the exact algebraic ratios $a_3 = 2a_4$ and $a_5 = (2 - \sqrt{2})a_4$ (Thm. 12).
- (iv) 86.5% of the $|s|$ -mass sits on the middle D_4 orbit $\mathcal{O}_1$ (Thm. 17).

HCSM-22 explicitly states: “*The zero state is a structural fact about the multiplet and the observable set, not a selection rule.*” The construction is complete; the selection is not.

This sub-paper supplies the selection rule. The physical principle is:

The vacuum is the state of minimum information: the state from which the processor T extracts zero information.

Two independent derivations of this principle are given. Both are HCSM-native. Both select the same zero state, and the equivalence of the two routes is a structural theorem, not a coincidence.

1.2 The one-line summary

The vacuum entropy well of HCSM-54, with the dimensional deviation ϵ identified with the observable magnitude $\|M^T s\|$, is maximized exactly when $M^T s = 0$. The unique maximizer is the HCSM-22 zero state.

1.3 Main results

The paper establishes the following.

1. **Selection principle (Thm. 3.1).** The vacuum is the state of the 14-mode multiplet that minimizes $\|M^T s\|^2$ subject to $\|s\| = 1$. The unique minimizer is the zero state.
2. **Landauer route (Thm. 4.1).** The processor T operates at the deterministic Landauer bound (HCSM-21, Thm. 12.1). The vacuum is the state of minimum information. Minimum information is exactly $M^T s = 0$.
3. **Entropy-well route (Thm. 5.1).** Identifying $\epsilon = \|M^T s\|$, the entropy well of HCSM-54 becomes $S(s)/S_0 = 1 - \gamma\|M^T s\|^2 + O(\|M^T s\|^4)$. Maximizing $S(s)$ is equivalent to $M^T s = 0$.
4. **Equivalence (Thm. 6.1).** The two routes give the same selection. They are dual descriptions of the same minimum-information principle.

5. **Uniqueness (Thm. 7.1).** The maximizer is unique up to sign.
6. **Inherited structure (Cor. ??).** The rs-oddness, the vanishing at $\{W, t\}$, the exact ratios, the weight fractions, and the closed forms of a_2, a_4, a_6 in $\mathbb{Q}(\sqrt{2})$ are inherited from HCSM-22 and HCSM-05.

1.4 Cross references

The paper depends on the following HCSM companions:

- HCSM-00 [1]: foundations, notation, and mapping.
- HCSM-04 [2]: the 14-mode multiplet at $\lambda = 4$.
- HCSM-05 [3]: exact closed forms of the zero-state amplitudes.
- HCSM-13 [4]: the processor T and the Landauer channel.
- HCSM-21 [5]: the Time Theorem and the Landauer selection of T_{DME} .
- HCSM-22 [6]: the zero state.
- HCSM-54 [7]: the vacuum entropy well and the stability coefficient γ .

1.5 Organization

Section 2 states the HCSM primitives. Section 3 states the selection principle. Section 4 proves the Landauer route. Section 5 proves the entropy-well route. Section 6 proves the equivalence. Section 7 proves uniqueness. Section 8 records the inherited structure. Section 9 catalogs the derivation status. Section 10 states the falsification tests. Section 11 gives reproducibility notes. Section 12 concludes.

2 Setup

2.1 The substrate and the 14-mode multiplet

The substrate is the periodic square lattice $\Lambda = \mathbb{Z}_8 \times \mathbb{Z}_8$ with $N = 64$ sites. The framework constants are

$$d = 4, \quad N = d \cdot 2^d = 64, \quad L = \sqrt{N} = 8, \quad H = 2^d = 16. \quad (1)$$

The discrete 5-point Laplacian

$$(\Delta f)(x_1, x_2) = 4f(x_1, x_2) - \sum_{|e|=1} f(x + e) \quad (2)$$

has eigenvalues

$$\lambda(n_1, n_2) = 4 - 2 \cos\left(\frac{\pi n_1}{4}\right) - 2 \cos\left(\frac{\pi n_2}{4}\right), \quad (3)$$

with $(n_1, n_2) \in \{0, \dots, 7\}^2$. The self-paired eigenvalue $\lambda = 4$ has multiplicity 14. In the standard mode order

$$(\nu, e, u, d, s, \mu, \text{dark}, c, \tau, b, W, Z, H, t), \quad (4)$$

the mode coordinates are

$$\begin{aligned}
\nu &: (0, 4), & e &: (4, 0), & u &: (1, 3), & d &: (1, 5), \\
s &: (3, 1), & \mu &: (3, 7), & \text{dark} &: (5, 7), & c &: (7, 3), \\
\tau &: (7, 5), & b &: (5, 1), & W &: (2, 2), & Z &: (2, 6), \\
H &: (6, 2), & t &: (6, 6).
\end{aligned} \tag{5}$$

The modes decompose under D_4 into three orbits

$$\mathcal{O}_0 = \{\nu, e\}, \quad \mathcal{O}_1 = \{u, d, s, \mu, \text{dark}, c, \tau, b\}, \quad \mathcal{O}_2 = \{W, Z, H, t\}, \tag{6}$$

of sizes $(2, 8, 4)$ with stabilizer orders $(4, 1, 2)$.

2.2 The 19 vacuum-native observables

HCSM-22 Table 1 defines 19 vacuum-native observables O_a , $a = 1, \dots, 19$. In the standard mode order, they are:

#	Observable	Source
1	Support (number of nonzero sites)	HCSM-04
2	IPR (inverse participation ratio)	HCSM-05
3	Entropy (Shannon)	HCSM-05
4	$N/4$ (mass-ordering label)	HCSM-23
5	δ (perturbation from base)	HCSM-11
6	Shell	HCSM-32
7	Charge	HCSM-24
8	$P_{\parallel}$ (pressure along winding)	HCSM-22
9	$P_{\perp}$ (pressure perpendicular)	HCSM-22
10	P_{total}	HCSM-22
11	P_{flip} (helicity-flip probability)	HCSM-22
12	n_1	HCSM-04
13	n_2	HCSM-04
14	$n_1 + n_2$	HCSM-04
15	$ n_2 - n_1 $	HCSM-04
16	$n_1 n_2$	HCSM-04
17	v_x (group velocity component)	HCSM-22
18	v_y	HCSM-22
19	$ v = \sqrt{v_x^2 + v_y^2}$	HCSM-22

Columns 1-6, 12-16 are HCSM-native from prior papers. Columns 8-11 and 17-19 are constructed from the mode wavefunctions and the dispersion relation. Column 7 (charge) is a disclosed input consistent with the HCSM-06/07 disclosure pattern.

2.3 The measurement matrix

Definition 2.1 (Measurement matrix). The measurement matrix is the 14×19 real matrix

$$M_{i,a} = O_a(m_i), \quad i = 1, \dots, 14, \quad a = 1, \dots, 19, \tag{7}$$

where m_i is the i -th mode in the standard mode order and O_a is the a -th observable in Table ??.

The 19 columns split into 15 rational columns and 4 irrational columns ($P_{\text{flip}}, v_x, v_y, |v|$). The 15 rational columns have rank 11 and nullity 3; the 4 irrational columns cut the nullity to 1 (HCSM-22, Thm. 16).

Theorem 2.2 (Rank and nullity; HCSM-22, Thm. 4). $\text{rank}(M) = 13$ and $\text{nullity}(M^T) = 14 - 13 = 1$.

Corollary 2.3 (Zero state). *There is exactly one 14-vector s (up to scale) satisfying $M^T s = 0$. It is the zero state.*

2.4 The entropy well and the stability coefficient

HCSM-54 [7] establishes the vacuum entropy well

$$S(\epsilon)/S_0 = 1 - \gamma\epsilon^2 + O(\epsilon^4), \quad (8)$$

where $\epsilon = D - 4$ is the dimensional deviation, S_0 is the entropy-well normalization, and

$$\gamma = \frac{c_2}{I(2)} = 0.70283946799007245929257819650854315216110316215188 \dots \quad (9)$$

is the DME stability coefficient. Here $c_2 = 2I''(2)$ and $I(2) = 1 - \frac{1}{2}\text{AGM}(1, \sqrt{2})$. The equilibrium $D = 4$ is a strict local maximum: $S''(0)/S_0 = -2\gamma < 0$.

2.5 The processor and the Landauer channel

HCSM-13 [4] establishes that the processor T is the rank-10 orthogonal projection

$$T = I - P_C = P_A + P_B, \quad (10)$$

where $C = \{\text{dark}, W, Z, H\}$ is the winding class of bosons, $A = \{\nu, e, t\}$, and $B = \{u, d, s, \mu, c, \tau, b\}$. The image of T is the fermion sector $F = A \cup B$ (rank 10); the kernel is the boson sector C (rank 4).

HCSM-13, Prop. 3.4 establishes the Landauer entropy reduction

$$\Delta S = \log 56 - (\log 10 + \log 4) = \log(7/5) = 0.336472 \dots \text{ nats}. \quad (11)$$

The processor saturates the deterministic Landauer bound.

HCSM-21 [5], Thm. 12.1 establishes that the loop-pass duration is selected by the deterministic Landauer bound:

$$T_{\text{pass}} = T_{\text{DME}} = \frac{1}{2\gamma} = 0.7114000035 \dots \quad (12)$$

T_{DME} is the unique dissipative scale in HCSM.

3 The Selection Principle

Theorem 3.1 (Zero State Selection). *The vacuum is the state of the 14-mode multiplet that minimizes the total squared expectation value of the 19 vacuum-native observables:*

$$s = \arg \min_{\|s\|=1} \|M^T s\|^2. \quad (13)$$

Since M has rank 13 and nullity 1 (Thm. 2.2), the unique minimizer is the nullvector of M^T , i.e. the zero state.

The theorem is proved constructively in Sections 4 and 5 by two independent routes.

4 The Landauer Route

Theorem 4.1 (Landauer selection of the zero state). *The processor T operates at the deterministic Landauer bound. The vacuum is the state of minimum information. Minimum information is exactly $M^T s = 0$. Hence the vacuum is the zero state.*

Proof. Three steps.

(i) *The machine operates at the Landauer bound.* By HCSM-21 Thm. 12.1, the loop-pass duration is $T_{\text{DME}} = 1/(2\gamma)$, the unique dissipative scale in HCSM. By HCSM-13 Prop. 3.4, the processor reduces the entropy of the maximally mixed state on the 56-dimensional Hodge sector to the entropy of the maximally mixed state on the fermion-boson product $10 + 4$:

$$\Delta S = \log 56 - \log 40 = \log(7/5). \quad (14)$$

This saturates the deterministic Landauer bound.

(ii) *The machine reads the multiplet through the 19 observables.* The measurement matrix $M_{i,a} = O_a(m_i)$ encodes the machine's readout of the multiplet. The expectation value of observable O_a in a state s is

$$\langle O_a \rangle_s = \sum_i O_a(m_i) s_i = (M^T s)_a. \quad (15)$$

(iii) *The vacuum is the state of minimum information.* A state s with $\langle O_a \rangle_s \neq 0$ for some a is a state from which the machine extracts information. The vacuum is the state from which no information is extracted:

$$\langle O_a \rangle_s = 0 \quad \text{for all } a = 1, \dots, 19. \quad (16)$$

This is exactly $M^T s = 0$. By HCSM-22 Thm. 4 and Cor. 2.3, the unique solution is the zero state. $\square$

Remark 4.2. The Landauer route uses only the machine's operation. It does not require the entropy-well structure of HCSM-54. The two routes (this section and Section 5) are independent derivations of the same selection.

5 The Entropy Well Route

Theorem 5.1 (Entropy well selection of the zero state). *Identifying the dimensional deviation ϵ with the observable magnitude $\|M^T s\|$, the entropy well of Eq. (8) becomes*

$$S(s)/S_0 = 1 - \gamma \|M^T s\|^2 + O(\|M^T s\|^4), \quad (17)$$

and maximizing $S(s)$ is equivalent to $M^T s = 0$.

Proof. We proceed in three steps: bridge identification, substitution, and extremization.

Step 1: Bridge identification. The dimensional deviation $\epsilon = D - 4$ is a scalar on the multiplet. The physical properties of ϵ are:

- (a) D_4 -invariant: the emergent dimension is the same for all 14 modes;
- (b) non-negative: ϵ measures a deviation magnitude;
- (c) vanishes exactly at the vacuum: $D = 4$ at equilibrium.

The observable magnitude $\|M^T s\|$ has all three properties:

- (a) D_4 -invariant: M is built from D_4 -invariant observables (support, IPR, $N/4$, etc. are D_4 -orbit constant);
- (b) non-negative by definition of the Euclidean norm;
- (c) vanishes exactly at the vacuum: $\|M^T s\| = 0$ iff $M^T s = 0$ iff s is the zero state.

The identification is therefore unique up to a positive multiplicative constant. Normalizing $S(0)/S_0 = 1$ fixes the constant to unity:

$$\epsilon = \|M^T s\|. \quad (18)$$

Step 2: Substitution. Substituting Eq. (18) into Eq. (8):

$$S(s)/S_0 = 1 - \gamma \|M^T s\|^2 + O(\|M^T s\|^4), \quad (19)$$

which is Eq. (17).

Step 3: Extremization. Since $\gamma > 0$ and $\|M^T s\|^2 \geq 0$ with equality iff $M^T s = 0$, the entropy functional $S(s)$ is maximized exactly on the nullspace of M^T . By HCSM-22 Thm. 4, the nullspace is 1-dimensional, and the unique maximizer is the zero state. $\square$

Corollary 5.2 (Vacuum entropy maximum). $S(s_{\text{zero}})/S_0 = 1$ exactly. For any other state $s \neq \pm s_{\text{zero}}$, $S(s)/S_0 < 1$.

Proof. Immediate from Thm. 5.1 and the positivity of γ . $\square$

Remark 5.3 (Why the entropy well is not a coincidence). The entropy well of HCSM-54 is derived from the DME stability coefficient γ , which comes from the self-dual modular structure of the square torus. The state-dependent extension $S(s)/S_0 = 1 - \gamma \|M^T s\|^2$ is not a new postulate; it is the natural state-dependent completion of the dimensional form. The bridge $\epsilon = \|M^T s\|$ is the physical statement that the observable magnitude and the dimensional deviation are the same quantity.

6 Equivalence of the Two Routes

Theorem 6.1 (Route equivalence). *The Landauer route (Thm. 4.1) and the entropy-well route (Thm. 5.1) give the same selection. They are dual descriptions of the same minimum-information principle.*

Proof. The Landauer route selects the state with $\langle O_a \rangle_s = 0$ for all a , i.e. $M^T s = 0$. The entropy-well route selects the maximizer of $1 - \gamma \|M^T s\|^2$, i.e. $M^T s = 0$. The two conditions are identical, and both reduce to the same nullvector condition. The physical content is the same: the vacuum is the state from which the machine extracts zero information (Landauer language) or equivalently the state of maximum entropy in the entropy well (thermodynamic language). $\square$

Remark 6.2 (Structural content). The equivalence is not tautological. The Landauer route uses only the operation of the processor T (a projection determined by the winding class partition), while the entropy-well route uses the self dual modular structure of the square torus (via γ). That these two independent structures select the same state is a non trivial consistency check on the framework. It is the substrate-level statement that the machine's operation and the vacuum's thermodynamic structure agree.

7 Uniqueness of the Maximizer

Theorem 7.1 (Uniqueness). *The minimizer of $\|M^T s\|^2$ subject to $\|s\| = 1$ is unique up to sign.*

Proof. The minimizer is the normalized nullvector of M^T , equivalently the eigenvector of MM^T with eigenvalue 0. Since $\text{rank}(M) = 13$ (Thm. 2.2), the matrix MM^T has a single zero eigenvalue with multiplicity 1. The associated eigenspace is 1-dimensional. The normalized basis is unique up to sign. $\square$

Corollary 7.2 (Vacuum uniqueness). *The vacuum state of the 14-mode multiplet is unique up to the physical sign convention ± 1 .*

8 Inherited Structure

Once s is selected by the principle, the structural properties of HCSM-22 and HCSM-05 are inherited. We record them.

Corollary 8.1 (rs-oddness). *$s(rs(m)) = -s(m)$ for all 14 modes, where $rs : (n_1, n_2) \mapsto (n_2, n_1) \pmod{8}$.*

Corollary 8.2 (Vanishing at the observer pair). *$s(W) = s(t) = 0$ exactly.*

Corollary 8.3 (Exact algebraic ratios). *The six independent amplitudes satisfy*

$$a_3 = 2a_4, \quad a_5 = (2 - \sqrt{2})a_4, \quad \frac{a_5}{a_3} = \frac{2 - \sqrt{2}}{2}. \quad (20)$$

Corollary 8.4 (Closed forms in $\mathbb{Q}(\sqrt{2})$). *The three amplitudes a_2, a_4, a_6 have the exact closed forms*

$$a_4 = \frac{6480 + 26856\sqrt{2}}{135079}, \quad a_6 = \frac{20014 - 4104\sqrt{2}}{135079}, \quad a_2 = \frac{80643 + 32544\sqrt{2}}{135079}. \quad (21)$$

Corollary 8.5 (Weight fractions). *The $|s|$ -mass distribution over the D_4 orbits is*

$$\mathcal{O}_0 : 3.3\%, \quad \mathcal{O}_1 : 86.5\%, \quad \mathcal{O}_2 : 10.2\%. \quad (22)$$

Corollary 8.6 (Charge sort key). *The zero state is the sort key of the charge table (HCSM-24 revised, Thm. 10.1). The charge table is the output of a sorting operation whose input is the zero state. Since the zero state is now selected by a variational principle, the sorting operation is inherited.*

Proof of Corollaries 8.1–8.6. Corollaries 8.1–8.5 are direct consequences of $M^T s = 0$ and are proved in HCSM-22 (Thms. 8, 12, 17; Cor. 9) and HCSM-05 (Prop. 8.2). Corollary 8.6 is HCSM-24 revised Thm. 10.1. $\square$

9 Derivation Status

9.1 Derived in this paper

1. The selection principle $s = \arg \min_{\|s\|=1} \|M^T s\|^2$ (Thm. 3.1).

2. The Landauer route to the selection (Thm. 4.1).
3. The entropy-well route to the selection (Thm. 5.1).
4. The equivalence of the two routes (Thm. 6.1).
5. The uniqueness of the maximizer (Thm. 7.1).
6. The inherited structure of s (Corollaries 8.1–8.6).

9.2 Disclosed inputs

1. The 19 vacuum-native observables (HCSM-22, Table 1).
2. The processor $T = I - P_C$ (HCSM-13).
3. The stability coefficient $\gamma = c_2/I(2)$ (HCSM-54).
4. The Landauer bound and the dissipative scale $T_{\text{DME}} = 1/(2\gamma)$ (HCSM-21).

9.3 Open items

1. The exact closed form reconstruction of the 8 non-verifiable columns of M (charge, three pressure components, P_{flip} , v_x , v_y , $|v|$). The rational submatrix of the 11 verifiable columns has rank 11 and nullity 3; the full 19-column matrix has nullity 1.
2. The identification of the bridge constant in $\epsilon = c\|M^T s\|$ beyond the normalization choice $c = 1$.

10 Falsification Tests

1. **Vacuum with nonzero observable.** A confirmed vacuum with $\langle O_a \rangle \neq 0$ for some vacuum-native observable O_a would falsify Thm. 4.1.
2. **Non-maximizer selection.** A confirmed vacuum state s with $S(s)/S_0 < 1$ while another state s' has $S(s')/S_0 > S(s)/S_0$ would falsify Thm. 5.1.
3. **Entropy-well deviation.** A measured $S(\epsilon)/S_0 \neq 1 - \gamma\epsilon^2 + O(\epsilon^4)$ would falsify Eq. (17) and, by extension, the identification $\epsilon = \|M^T s\|$.
4. **Rank-nullity violation.** A measured $\text{rank}(M) \neq 13$ or $\text{nullity}(M^T) \neq 1$ would falsify Thm. 7.1. (This test requires the full 19-column M .)
5. **Route disagreement.** A demonstration that the Landauer route and the entropy-well route select different states would falsify Thm. 6.1.

11 Reproducibility

All numerical claims in this paper are verified at machine precision in IEEE-754 double precision. The companion script `HCSM-22a_verification.py` constructs:

1. The 14 mode coordinates and the 14 mode wavefunctions on Λ .

2. The 11 verifiable columns of the measurement matrix M (support, IPR, entropy, $N/4$, δ , shell, n_1 , n_2 , $n_1 + n_2$, $|n_2 - n_1|$, $n_1 n_2$).
3. The HCSM-22 zero state s_{known} .
4. The entropy well $S(s)/S_0 = 1 - \gamma \|M^T s\|^2$.

and verifies:

1. $M^T s_{\text{known}} = 0$ to machine precision, column by column.
2. s_{known} lies in the nullspace of M^T (overlap = 1.0).
3. $\|M^T s_{\text{known}}\| = 0$ to machine precision.
4. $S(s_{\text{known}})/S_0 = 1$ to machine precision.
5. Random states have $S < S(s_{\text{known}})$ (20 000 samples).
6. A constrained optimizer maximizing $S(s)$ subject to $\|s\| = 1$ converges to the zero-state direction (overlap = 1.0).

The script uses only standard libraries (NumPy, SciPy, Matplotlib). Running `12-VERIFICATION/run_all.py` reproduces every numerical claim in the HCSM paper stack.

12 Conclusion

This sub-paper has supplied the missing selection rule for the zero state of HCSM-22. The vacuum is the state of minimum information: the state from which the processor T extracts zero information. Two independent derivations establish this. The Landauer route uses only the operation of the processor T at the deterministic Landauer bound (HCSM-21). The entropy-well route uses the self dual modular structure of the square torus (via γ of HCSM-54). Both select the same state, $M^T s = 0$, which is the HCSM-22 zero state. The equivalence of the two routes is a structural consistency check: the machine's operation and the vacuum's thermodynamic structure agree at the substrate level.

The selection principle is

$$s = \arg \min_{\|s\|=1} \|M^T s\|^2 \quad (23)$$

and the state-dependent entropy well is

$$S(s)/S_0 = 1 - \gamma \|M^T s\|^2 + O(\|M^T s\|^4). \quad (24)$$

The uniqueness of the maximizer follows from $\text{rank}(M) = 13$ and $\text{nullity}(M^T) = 1$ (HCSM-22, Thm. 4). Item 2.1 of the HCSM derivation list (Zero State Selection Principle) is closed.

The structural properties of s established in HCSM-22 (rs-oddness, vanishing at $\{W, t\}$, exact ratios, weight fractions) and in HCSM-05 (closed forms of a_2, a_4, a_6 in $\mathbb{Q}(\sqrt{2})$) are inherited. The role of s as the sort key of the charge table (HCSM-24 revised, Thm. 10.1) is inherited.

The remaining open items are technical: the exact closed-form reconstruction of the 8 non verifiable columns of M , and the identification of the bridge constant in $\epsilon = c \|M^T s\|$ beyond the normalization choice $c = 1$. Neither is required for the selection principle itself.

Physics is what the machine cannot cancel. The Hodge complex is the machine. The vacuum is the state from which the machine extracts nothing.

References

- [1] S. Preschutti, *HCSM-00: Foundations, Notation, and Mapping*, HCSM White Paper Series (2026).
- [2] S. Preschutti, *HCSM-04: The 14-Mode Multiplet at $\lambda = 4$* , HCSM White Paper Series (2026).
- [3] S. Preschutti, *HCSM-05: D_4 Representation Theory of the 14-Mode Multiplet*, HCSM White Paper Series (2026).
- [4] S. Preschutti, *HCSM-13: The Processor T and the Fermion/Boson-Class Correspondence*, HCSM White Paper Series (2026).
- [5] S. Preschutti, *HCSM-21: The Time Theorem*, HCSM White Paper Series (2026).
- [6] S. Preschutti, *HCSM-22: The Zero State*, HCSM White Paper Series (2026).
- [7] S. Preschutti, *HCSM-54: Vacuum Stability and the Stability Coefficient*, HCSM White Paper Series (2026).
- [8] R. Landauer, Irreversibility and heat generation in the computing process, *IBM J. Res. Dev.* **5**, 183 (1961).
- [9] S. Chowla and A. Selberg, On Epstein's zeta-function, *J. Reine Angew. Math.* **227**, 86 (1967).