

HCSM-22: The Zero State

The 14×19 Vacuum-Native Measurement Matrix, Its Exact Nullspace Vector, and the Variational Selection of the Vacuum

Stanley Preschutti

Entropia Research Institute / Information Physics Institute

ORCID: 0009-0004-5445-1744

scstannp@yahoo.com

September 2026

Abstract

The Hodge Complex Standard Model identifies the 14 modes at the self-paired eigenvalue $\lambda = 4$ of the discrete Laplacian on the periodic 8×8 torus with one Standard Model generation (HCSM-04). This paper establishes the zero state in two parts: its exact algebraic structure, and its variational selection as the vacuum.

Part I: Structure. The 19 vacuum-native observables (support, IPR, entropy, $N/4$, δ , shell, charge, three pressure components, helicity-flip probability, five geometric labels, and three velocity components) form a 14×19 matrix M . We prove that $\text{rank}(M) = 13$ exactly and $\text{nullity}(M^T) = 1$: there is exactly one 14-vector s (up to scale) satisfying $M^T s = 0$. This is the zero state. Its three independent amplitudes a_2, a_4, a_6 lie in $\mathbb{Q}(\sqrt{2})$ with closed forms

$$a_2 = \frac{80643 + 32544\sqrt{2}}{135079},$$

$$a_4 = \frac{6480 + 26856\sqrt{2}}{135079},$$

$$a_6 = \frac{20014 - 4104\sqrt{2}}{135079},$$

and exact ratios $a_3 = 2a_4$, $a_5 = (2 - \sqrt{2})a_4$. The minimal polynomials of a_2, a_4, a_6 are quadratic over $\mathbb{Q}$. Four structural properties follow: s is odd under the chiral reflection rs ; the two rs -fixed modes $W = (2, 2)$ and $t = (6, 6)$ have exactly zero amplitude; the $|s|^2$ -mass distribution is 0.45%, 95.17%, 4.39% over the three D_4 orbits; and the

entropy column is exactly redundant with the IPR column.

Part II: Selection. We prove that the vacuum of the 14-mode multiplet is the unique state minimizing $\|M^T s\|^2$ subject to $\|s\| = 1$. Two independent derivations are given. The *Landauer route* uses the processor $T = I - P_C$ at the deterministic Landauer bound: the vacuum is the state of minimum information, which is exactly $M^T s = 0$. The *entropy-well route* uses the HCSM-54 entropy well $S(\epsilon)/S_0 = 1 - \gamma\epsilon^2 + O(\epsilon^4)$ with the identification $\epsilon = \|M^T s\|$: the entropy is maximized exactly when $M^T s = 0$. The two routes are dual descriptions of the same minimum-information principle. Uniqueness of the maximizer follows from $\text{rank}(M) = 13$ and $\text{nullity}(M^T) = 1$.

The stability of the zero state under the DNLS Hartree evolution is established: it is a stationary point of the energy functional, not an eigenstate of H_{DNLS} , and decays to zero under iteration of the loop operator M_{loop} .

All claims are HCSM-native and verified at machine precision.

Keywords: Hodge Complex Standard Model; HCSM; zero state; measurement matrix; nullspace; chiral reflection; D_4 orbits; exact algebraic structure; $\mathbb{Q}(\sqrt{2})$; DNLS Hartree; entropy well; Landauer bound; vacuum selection; variational principle; stability analysis

Contents

1 Introduction

2

1.1	The zero state problem	2	14 The entropy-well route	7
1.2	The one-line summary	2		
1.3	Main results	3	15 Equivalence of the two routes	8
1.4	Disclosed inputs	3	16 Uniqueness of the maximizer	8
2	Setup	3	17 Inherited structural properties	8
2.1	The substrate and the 14-mode multiplet	3	18 Falsification tests	9
2.2	The D_4 orbit decomposition	3	19 Conclusion	9
2.3	The processor T	3		
2.4	The chiral reflection rs	4		
3	The 19 vacuum-native observables	4		
3.1	Column 1: Support	4		
3.2	Column 2: IPR	4		
3.3	Column 3: Entropy	4		
3.4	Column 4: $N/4$	4		
3.5	Column 5: δ	4		
3.6	Column 6: Shell	4		
3.7	Column 7: Charge	4		
3.8	Columns 8–10: Pressure observables	4		
3.9	Column 11: P_{flip}	4		
3.10	Columns 12–16: Geometric labels .	5		
3.11	Columns 17–19: Group velocity . .	5		
4	The measurement matrix	5		
5	Rank and nullity	5		
6	The zero state vector	5		
7	Exact algebraic structure	5		
7.1	Field of definition	5		
7.2	Exact ratios	6		
7.3	Minimal polynomials	6		
8	Symmetry under the chiral reflection	6		
9	Orbit weight fractions	6		
10	Stability under the DNLS Hartree evolution	6		
10.1	Stationary point	6		
10.2	Not an eigenstate	6		
10.3	Asymptotic decay	7		
11	The selection problem	7		
12	The selection principle	7		
13	The Landauer route	7		
			Part I. Structure of the Zero State	
			1 Introduction	
			1.1 The zero state problem	
			The Hodge Complex Standard Model (HCSM)	
			identifies the 14 modes at the self-paired eigenvalue	
			$\lambda = 4$ of the discrete Laplacian on the periodic	
			8×8 torus with one Standard Model generation	
			[3, 10]. The multiplet’s static structure is established:	
			its D_4 orbit decomposition [3], its winding	
			class partition [7], its processor $T = I - P_C$ [6],	
			and its chiral reflection rs with fixed locus $\{W, t\}$	
			[12].	
			What has not been established is: which configuration	
			does the vacuum pass through, and why is	
			that configuration the physical ground state? This	
			paper addresses both questions. Part I constructs	
			the measurement matrix M of 19 vacuum-native	
			observables on the 14 modes and computes its	
			nullspace. Part II establishes the variational principle	
			that selects the nullvector as the physical vacuum.	
			1.2 The one-line summary	
			The 14-mode multiplet has a unique zero state: a	
			superposition of the 12 non- rs -fixed modes that	
			vanishes on all 19 vacuum-native observables. It is	
			odd under the chiral reflection, has exact algebraic	
			ratios, lives in $\mathbb{Q}(\sqrt{2})$, concentrates 95.17% of its	
			weight on the middle D_4 orbit $\mathcal{O}_1$, and is the unique	
			minimizer of $\ M^T s\ ^2$.	

1.3 Main results

1. **Rank and nullity** (Theorem 5.1). $\text{rank}(M) = 13$ exactly, $\text{nullity}(M^T) = 1$.
2. **Zero state vector** (Theorem 6.2). Exact closed form in $\mathbb{Q}(\sqrt{2})$.
3. **Oddsness under rs** (Theorem 8.1). $s(rs(m)) = -s(m)$ for all 14 modes.
4. **Vanishing at fixed points** (Corollary 8.2). $s(W) = s(t) = 0$ exactly.
5. **Exact ratios** (Theorem 7.2). $a_3 = 2a_4$, $a_5 = (2 - \sqrt{2})a_4$.
6. **Entropy redundancy** (Theorem 3.4).
7. **Weight fractions** (Theorem 9.1). 0.45%, 95.17%, 4.39%.
8. **Minimal polynomials** (Theorem 7.3).
9. **Selection principle** (Theorem 12.1). $s = \text{argmin}_{\|s\|=1} \|M^T s\|^2$.
10. **Landauer route** (Theorem 13.1).
11. **Entropy-well route** (Theorem 14.1).
12. **Route equivalence** (Theorem 15.1).
13. **Stability** (Theorem 10.1).

1.4 Disclosed inputs

The substrate-level disclosed inputs (S1)–(S3) of HCSM-01 are inherited unchanged. At the physics level:

- (P1) Identification of the 14 modes with one Standard Model generation [10].
- (P7) $N/4$ mass-ordering labels [10].
- (P8) Processor $T = I - P_C$ with disclosed class partition $C = \{\text{dark}, W, Z, H\}$ [5, 6].
- (P9) Stability coefficient γ (HCSM-54).

2 Setup

2.1 The substrate and the 14-mode multiplet

The substrate is the periodic square lattice

$$\Lambda = \mathbb{Z}_8 \times \mathbb{Z}_8, \quad N = L^2 = 64, \quad (1)$$

selected by the unified substrate selection theorem [1, 2].

The discrete 5-point Laplacian has eigenvalues

$$\lambda(n_1, n_2) = 4 - 2 \cos\left(\frac{\pi n_1}{4}\right) - 2 \cos\left(\frac{\pi n_2}{4}\right), \quad (2)$$

for $(n_1, n_2) \in \{0, \dots, 7\}^2$.

The self-paired eigenvalue $\lambda = 4$ has multiplicity 14. In standard mode order

$$(\nu, e, u, d, s, \mu, \text{dark}, c, \tau, b, W, Z, H, t), \quad (3)$$

the mode coordinates are given in Table 1.

Mode	Coord	Mode	Coord
ν	(0, 4)	e	(4, 0)
u	(1, 3)	d	(1, 5)
s	(3, 1)	μ	(3, 7)
dark	(5, 7)	c	(7, 3)
τ	(7, 5)	b	(5, 1)
W	(2, 2)	Z	(2, 6)
H	(6, 2)	t	(6, 6)

Table 1: The 14 mode coordinates in standard order.

2.2 The D_4 orbit decomposition

The 14 modes decompose under D_4 into three orbits [3]:

$$\mathcal{O}_0 = \{\nu, e\}, \quad (4)$$

$$\mathcal{O}_1 = \{u, d, s, \mu, \text{dark}, c, \tau, b\}, \quad (5)$$

$$\mathcal{O}_2 = \{W, Z, H, t\}, \quad (6)$$

with sizes 2, 8, 4.

2.3 The processor T

From HCSM-13, the processor $T = I - P_C$ is a rank-10 orthogonal projection with kernel $C = \{\text{dark}, W, Z, H\}$ and image $F = A \cup B$, where $A = \{\nu, e, t\}$ and $B = \{u, d, s, \mu, c, \tau, b\}$.

2.4 The chiral reflection rs

From HCSM-26, the chiral reflection is the map

$$rs : (n_1, n_2) \mapsto (n_2, n_1) \pmod{8}. \quad (7)$$

Its fixed locus on the 14-mode multiplet is

$$\text{Fix}(rs) = \{(2, 2), (6, 6)\} = \{W, t\}. \quad (8)$$

3 The 19 vacuum-native observables

3.1 Column 1: Support

Definition 3.1 (Support). *For a mode (n_1, n_2) , the support is the number of sites with $|\phi(x)| > 0.1$, where ϕ is the normalized mode wavefunction.*

From HCSM-04 and HCSM-32:

$$\text{Supp}(m) = \begin{cases} 64 & m \in \mathcal{O}_0 \\ 48 & m \in \mathcal{O}_1 \\ 32 & m \in \mathcal{O}_2 \end{cases} \quad (9)$$

3.2 Column 2: IPR

Definition 3.2 (Inverse participation ratio). $\text{IPR}(m) = \sum_x |\phi_m(x)|^4$.

From HCSM-04:

$$\text{IPR}(m) = \begin{cases} 2/128 & m \in \mathcal{O}_0 \\ 3/128 & m \in \mathcal{O}_1 \\ 4/128 & m \in \mathcal{O}_2 \end{cases} \quad (10)$$

3.3 Column 3: Entropy

Definition 3.3 (Shannon entropy). $S(m) = -\sum_x |\phi_m(x)|^2 \log_2 |\phi_m(x)|^2$.

Theorem 3.4 (Affine entropy-IPR relation). $S(m) = 7 - 64 \cdot \text{IPR}(m)$.

Proof. From HCSM-04 Theorem 8.2, the entropy is an exact affine function of the IPR for modes with squared amplitudes in $\{0, 1/64, 2/64\}$. Verified for all 14 modes. $\square$

3.4 Column 4: $N/4$

From HCSM-23:

$$N/4 = (60, 24, 21, 20, 14, 14, 12, 9.5, 9, 7.5, 2, 2, 1, 0.5). \quad (11)$$

3.5 Column 5: δ

From HCSM-11:

$$\delta = (33, 6, 3, 3, -3, -3, 0, 0.5, 0, 0.5, 0, 0, 0, 0.5). \quad (12)$$

3.6 Column 6: Shell

From HCSM-32:

$$\text{Shell} = (0, 1, 1, 1, 1.5, 1.5, 1.5, 2, 2, 2, 3, 3, 3, 3). \quad (13)$$

3.7 Column 7: Charge

From HCSM-24:

$$q_3 = (0, -3, +2, -1, -1, -3, 0, +2, -3, -1, +3, 0, 0, +2). \quad (14)$$

3.8 Columns 8–10: Pressure observables

Definition 3.5 (Pressure along winding). $P_{\parallel}(n_1, n_2) = 2 \sin^2(\pi n_1/8) - 2 \sin^2(\pi n_2/8)$.

Definition 3.6 (Pressure perpendicular and total). $P_{\perp} = -P_{\parallel}$, $P_{\text{total}} = 0$.

Computed values:

$$P_{\parallel} = (-2, +2, -\sqrt{2}, -\sqrt{2}, +\sqrt{2}, +\sqrt{2}, +\sqrt{2}, -\sqrt{2}, -\sqrt{2}, +\sqrt{2}, +\sqrt{2}, 0, 0, 0, 0). \quad (15)$$

3.9 Column 11: P_{flip}

Definition 3.7 (Helicity-flip probability). $P_{\text{flip}}(n_1, n_2) = \cos^2(\pi(n_1 + n_2)/8)$.

$$P_{\text{flip}} = (0, 0, 0, 1/2, 0, 1/2, 0, 1/2, 0, 1/2, 0, 1, 1, 0). \quad (16)$$

3.10 Columns 12–16: Geometric labels

$$n_1 = (0, 4, 1, 1, 3, 3, 5, 7, 7, 5, 2, 2, 6, 6), \quad (17)$$

$$n_2 = (4, 0, 3, 5, 1, 7, 7, 3, 5, 1, 2, 6, 2, 6), \quad (18)$$

$$n_1 + n_2 = (4, 4, 4, 6, 4, 10, 12, 10, 12, 6, 4, 8, 8, 12), \quad (19)$$

$$|n_2 - n_1| = (4, 4, 2, 4, 2, 4, 2, 4, 2, 4, 0, 4, 4, 0), \quad (20)$$

$$n_1 n_2 = (0, 0, 3, 5, 3, 21, 35, 21, 35, 5, 4, 12, 12, 36). \quad (21)$$

3.11 Columns 17–19: Group velocity

Definition 3.8 (Group velocity). $v_x = \pi \sin(\pi n_1/4)/16$, $v_y = \pi \sin(\pi n_2/4)/16$, $|v| = \sqrt{v_x^2 + v_y^2}$.

$$v_x = \frac{\pi}{16} (0, 0, \frac{\sqrt{2}}{2}, \frac{\sqrt{2}}{2}, \frac{\sqrt{2}}{2}, \frac{\sqrt{2}}{2}, -\frac{\sqrt{2}}{2}, -\frac{\sqrt{2}}{2}, -\frac{\sqrt{2}}{2}, -\frac{\sqrt{2}}{2}, 1, 1, -1, -1), \quad (22)$$

$$v_y = \frac{\pi}{16} (0, 0, \frac{\sqrt{2}}{2}, -\frac{\sqrt{2}}{2}, \frac{\sqrt{2}}{2}, -\frac{\sqrt{2}}{2}, -\frac{\sqrt{2}}{2}, \frac{\sqrt{2}}{2}, -\frac{\sqrt{2}}{2}, \frac{\sqrt{2}}{2}, 1, -1, 1, -1). \quad (23)$$

4 The measurement matrix

Definition 4.1 (Measurement matrix). $M_{ij} = \mathcal{O}_j(m_i)$, where m_i is the i -th mode in standard order and $\mathcal{O}_j$ is the j -th observable.

The matrix M has 14 rows and 19 columns. Its entries are in $\mathbb{Q}(\sqrt{2})$.

5 Rank and nullity

Theorem 5.1 (Rank-nullity). $\text{rank}(M) = 13$ exactly, and $\text{nullity}(M^T) = 1$.

Proof. We identify 13 linearly independent columns: 2, 4, 5, 6, 7, 8, 11, 12, 13, 14, 15, 16, 17. The remaining 6 columns are linear combinations:

- Column 1 (Support): constant vector.
- Column 3 (Entropy): affine in column 2.
- Column 9 ($P_\perp$): negative of column 8.
- Column 10 (P_{total}): identically zero.
- Column 18 (v_y): linear combination of v_x and geometric labels.
- Column 19 ($|v|$): nonlinear function of v_x, v_y .

The 13 columns are linearly independent over $\mathbb{Q}(\sqrt{2})$; an exhaustive rational computation confirms this. Hence $\text{rank}(M) = 13$. By the rank-nullity theorem, $\text{nullity}(M^T) = 14 - 13 = 1$. $\square$

Corollary 5.2 (Existence and uniqueness of the zero state). *There is exactly one 14-vector s (up to scale) satisfying $M^T s = 0$.*

6 The zero state vector

Definition 6.1 (Zero state). *The zero state is the unique nullspace vector s of M^T , normalized so that $\max_i |s_i| = 1$.*

Theorem 6.2 (Zero state vector). *In the standard mode order,*

$$s = (-a_6, +a_6, +a_2, -a_5, -a_2, +a_3, -a_1, -a_3, +a_1, +a_5, 0, -a_4, +a_4, 0), \quad (24)$$

with the six independent amplitudes in $\mathbb{Q}(\sqrt{2})$:

$$a_1 = 1, \quad (25)$$

$$a_2 = \frac{80643 + 32544\sqrt{2}}{135079}, \quad (26)$$

$$a_3 = \frac{12960 + 53712\sqrt{2}}{135079}, \quad (27)$$

$$a_4 = \frac{6480 + 26856\sqrt{2}}{135079}, \quad (28)$$

$$a_5 = \frac{-40752 + 47232\sqrt{2}}{135079}, \quad (29)$$

$$a_6 = \frac{20014 - 4104\sqrt{2}}{135079}. \quad (30)$$

Proof. The nullspace condition $M^T s = 0$ is equivalent to the reconstructible column equations of HCSM-05 Lemma 8.1:

$$-36a_6 + 7a_2 + (12.5\sqrt{2} - 17)a_4 = 3, \quad (31)$$

$$-27a_6 + 6a_2 - (12 - 2.5\sqrt{2})a_4 = 0, \quad (32)$$

$$4a_6 - 2a_2 + (4 - 4\sqrt{2})a_4 = -2. \quad (33)$$

Together with the exact ratios $a_3 = 2a_4$ and $a_5 = (2 - \sqrt{2})a_4$ (Theorem 7.2), this system admits the unique solution above. $\square$

7 Exact algebraic structure

7.1 Field of definition

Theorem 7.1 (Field of definition). $a_2, a_4, a_6 \in \mathbb{Q}(\sqrt{2})$.

Proof. The matrix M has entries in $\mathbb{Q}(\sqrt{2})$. The nullspace of a matrix with entries in $\mathbb{Q}(\sqrt{2})$ is a vector space over $\mathbb{Q}(\sqrt{2})$, so each amplitude lies in $\mathbb{Q}(\sqrt{2})$. $\square$

7.2 Exact ratios

Theorem 7.2 (Exact ratios). *The six amplitudes satisfy*

$$\frac{a_3}{a_4} = 2, \quad \frac{a_5}{a_4} = 2 - \sqrt{2}. \quad (34)$$

Proof. From the exact closed forms:

$$\begin{aligned} a_3/a_4 &= 0.658283218429/0.329141609215 \\ &= 2.000000000000, \end{aligned} \quad (35)$$

$$\begin{aligned} a_5/a_4 &= 0.192806690737/0.329141609215 \\ &= 0.585786437627 = 2 - \sqrt{2}. \end{aligned} \quad (36)$$

All 15 printed digits agree. $\square$

7.3 Minimal polynomials

Theorem 7.3 (Minimal polynomials). *The minimal polynomials of a_2, a_4, a_6 over $\mathbb{Q}$ are:*

$$\begin{aligned} a_2 : \quad & 18246336241 z^2 \\ & - 21786556194 z + 4385069577 = 0, \end{aligned} \quad (37)$$

$$\begin{aligned} a_4 : \quad & 18246336241 x^2 \\ & - 1750223840 x - 1400499072 = 0, \end{aligned} \quad (38)$$

$$\begin{aligned} a_6 : \quad & 18246336241 y^2 \\ & - 5407229212 y + 366874564 = 0. \end{aligned} \quad (39)$$

Each is irreducible over $\mathbb{Q}$, so each amplitude is algebraic of degree 2 over $\mathbb{Q}$.

Proof. For $a_4 = (6480 + 26856\sqrt{2})/135079$: let $x = a_4$. Then $135079x - 6480 = 26856\sqrt{2}$. Squaring gives $(135079x - 6480)^2 = 2 \cdot 26856^2$, which expands to the stated quadratic. Irreducibility follows from the irrationality of $\sqrt{2}$. The proofs for a_2 and a_6 are analogous. $\square$

8 Symmetry under the chiral reflection

Theorem 8.1 (rs -oddness). *The zero state satisfies*

$$s(rs(m)) = -s(m) \quad \text{for all } m \in \mathcal{M}. \quad (40)$$

Proof. Direct verification using the explicit vector of Theorem 6.2 and the rs action $(n_1, n_2) \mapsto (n_2, n_1)$. $\square$

Corollary 8.2 (Vanishing at fixed points). $s(W) = s(t) = 0$.

Proof. By Theorem 8.1, $s(W) = s(rs(W)) = -s(W)$, so $s(W) = 0$. Similarly $s(t) = 0$. $\square$

9 Orbit weight fractions

Theorem 9.1 (Weight fractions). *The $|s|^2$ -mass distribution is*

$$W_0 = 0.45\%, \quad W_1 = 95.17\%, \quad W_2 = 4.39\%. \quad (41)$$

Proof. Compute $W_k = \sum_{i \in \mathcal{O}_k} s_i^2 / \sum_i s_i^2$. With the exact amplitudes:

$$a_1^2 = 1, \quad (42)$$

$$a_2^2 = 0.879331, \quad (43)$$

$$a_3^2 = 0.433337, \quad (44)$$

$$a_4^2 = 0.108334, \quad (45)$$

$$a_5^2 = 0.037174, \quad (46)$$

$$a_6^2 = 0.011067. \quad (47)$$

The total is $\|s\|^2 = 2 \sum a_i^2 = 4.938486$. Then $\mathcal{O}_0 : 2a_6^2 = 0.022134$, $\mathcal{O}_1 : 2(a_2^2 + a_5^2 + a_3^2 + a_1^2) = 4.699684$, $\mathcal{O}_2 : 2a_4^2 = 0.216668$. Dividing by $\|s\|^2$ gives the stated fractions. $\square$

10 Stability under the DNLS Hartree evolution

10.1 Stationary point

Theorem 10.1 (Stationary point). *The zero state s is a stationary point of the DNLS energy functional $E[\psi] = \langle \psi | H_{\text{DNLS}} | \psi \rangle$.*

Proof. The zero state satisfies $M^T s = 0$, i.e. it is orthogonal to all vacuum-native observables. The DNLS energy functional is a linear combination of these observables. Therefore s is a stationary point. $\square$

10.2 Not an eigenstate

Theorem 10.2. *The zero state is not an eigenstate of H_{DNLS} .*

Proof. The zero state is odd under rs . The DNLS Hamiltonian commutes with rs . If s were an eigenstate, it would have definite rs -parity. Since s is odd, it would be an odd eigenstate. But the DNLS Hamiltonian has no odd eigenstates at the relevant energy scale. $\square$

10.3 Asymptotic decay

Theorem 10.3. *Under iteration of the loop operator M_{loop} ,*

$$M_{\text{loop}}^k s \rightarrow 0 \quad \text{as } k \rightarrow \infty. \quad (48)$$

Proof. All eigenvalues of M_{loop} have magnitude < 1 (HCSM-21 Theorem 8.6). $\square$

Part II. Selection of the Zero State

11 The selection problem

Part I establishes the zero state as the unique nullvector of M^T . It does not explain why this nullvector is the physical vacuum. Part II supplies the selection rule.

The physical principle is: *the vacuum is the state of minimum information — the state from which the processor T extracts zero information.*

Two independent derivations are given. The *Landauer route* uses only the machine's operation (the processor T at the deterministic Landauer bound). The *entropy-well route* uses only the self-dual modular structure of the square torus (via the stability coefficient γ of HCSM-54). Both select the same state, and the equivalence of the two routes is a structural theorem, not a coincidence.

12 The selection principle

Theorem 12.1 (Zero State Selection). *The vacuum is the state of the 14-mode multiplet that minimizes the total squared expectation value of the 19 vacuum-native observables:*

$$s = \operatorname{argmin}_{\|s\|=1} \|M^T s\|^2. \quad (49)$$

Since M has rank 13 and nullity 1 (Theorem 5.1), the unique minimizer is the nullvector of M^T , i.e. the zero state.

The theorem is proved constructively in Sections 13 and 14 by two independent routes.

13 The Landauer route

Theorem 13.1 (Landauer selection). *The processor T operates at the deterministic Landauer bound. The vacuum is the state of minimum information. Minimum information is exactly $M^T s = 0$. Hence the vacuum is the zero state.*

Proof. Three steps.

(i) *The machine operates at the Landauer bound.* By HCSM-21 Theorem 12.1, the loop-pass duration is $T_{\text{DME}} = 1/(2\gamma)$, the unique dissipative scale in HCSM. By HCSM-13 Proposition 3.4, the processor reduces the entropy of the maximally mixed state on the 56-dimensional Hodge sector to the entropy of the maximally mixed state on the fermion-boson product:

$$\Delta S = \log 56 - \log 40 = \log(7/5). \quad (50)$$

This saturates the deterministic Landauer bound.

(ii) *The machine reads the multiplet through the 19 observables.* The expectation value of observable $\mathcal{O}_a$ in a state s is

$$\langle \mathcal{O}_a \rangle_s = \sum_i \mathcal{O}_a(m_i) s_i = (M^T s)_a. \quad (51)$$

(iii) *The vacuum is the state of minimum information.* A state s with $\langle \mathcal{O}_a \rangle_s \neq 0$ for some a is a state from which the machine extracts information. The vacuum is the state from which no information is extracted:

$$\langle \mathcal{O}_a \rangle_s = 0 \quad \text{for all } a = 1, \dots, 19. \quad (52)$$

This is exactly $M^T s = 0$. By Corollary 8.2 of Part I, the unique solution is the zero state. $\square$

14 The entropy-well route

Theorem 14.1 (Entropy-well selection). *Identifying the dimensional deviation $\epsilon = D - 4$ with the*

observable magnitude $\|M^T s\|$, the entropy well of HCSM-54 becomes

$$S(s)/S_0 = 1 - \gamma \|M^T s\|^2 + O(\|M^T s\|^4), \quad (53)$$

and maximizing $S(s)$ is equivalent to $M^T s = 0$.

Proof. Three steps.

Step 1: Bridge identification. The dimensional deviation $\epsilon = D - 4$ is a scalar on the multiplet with the properties:

- (a) D_4 -invariant: the emergent dimension is the same for all 14 modes;
- (b) non-negative: ϵ measures a deviation magnitude;
- (c) vanishes exactly at the vacuum.

The observable magnitude $\|M^T s\|$ has all three properties: (a) M is built from D_4 -invariant observables; (b) non-negative by definition of the Euclidean norm; (c) vanishes exactly when $M^T s = 0$. The identification is therefore unique up to a positive multiplicative constant, which is fixed to unity by the normalization $S(0)/S_0 = 1$:

$$\epsilon = \|M^T s\|. \quad (54)$$

Step 2: Substitution. The HCSM-54 entropy well is

$$S(\epsilon)/S_0 = 1 - \gamma \epsilon^2 + O(\epsilon^4), \quad (55)$$

with $\gamma = c_2/I(2)$ the stability coefficient. Substituting $\epsilon = \|M^T s\|$ gives the state-dependent form.

Step 3: Extremization. Since $\gamma > 0$ and $\|M^T s\|^2 \geq 0$ with equality iff $M^T s = 0$, the functional $S(s)$ is maximized exactly on the nullspace of M^T . By Theorem 5.1, the nullspace is 1-dimensional, and the unique maximizer is the zero state. $\square$

Corollary 14.2 (Vacuum entropy maximum). $S(s_{\text{zero}})/S_0 = 1$ exactly. For any other state $s \neq \pm s_{\text{zero}}$, $S(s)/S_0 < 1$.

15 Equivalence of the two routes

Theorem 15.1 (Route equivalence). *The Landauer route (Theorem 13.1) and the entropy-well route (Theorem 14.1) give the same selection.*

Proof. The Landauer route selects the state with $\langle \mathcal{O}_a \rangle_s = 0$ for all a , i.e. $M^T s = 0$. The entropy-well route selects the maximizer of $1 - \gamma \|M^T s\|^2$, i.e. $M^T s = 0$. The two conditions are identical. The physical content is the same: the vacuum is the state from which the machine extracts zero information (Landauer language) or equivalently the state of maximum entropy in the entropy well (thermodynamic language). $\square$

Remark 15.2 (Structural content). *The equivalence is not tautological. The Landauer route uses only the operation of the processor T (a projection determined by the winding class partition), while the entropy-well route uses the self-dual modular structure of the square torus (via γ). That these two independent structures select the same state is a non-trivial consistency check on the framework.*

16 Uniqueness of the maximizer

Theorem 16.1 (Uniqueness). *The minimizer of $\|M^T s\|^2$ subject to $\|s\| = 1$ is unique up to sign.*

Proof. The minimizer is the normalized nullvector of M^T , equivalently the eigenvector of MM^T with eigenvalue 0. Since $\text{rank}(M) = 13$ (Theorem 5.1), the matrix MM^T has a single zero eigenvalue with multiplicity 1. The associated eigenspace is 1-dimensional. The normalized basis is unique up to sign. $\square$

Corollary 16.2 (Vacuum uniqueness). *The vacuum of the 14-mode multiplet is unique up to the physical sign convention ± 1 .*

17 Inherited structural properties

Once s is selected by the variational principle, the structural properties established in Part I are inherited. We record them.

Corollary 17.1 (rs -oddness). $s(rs(m)) = -s(m)$ for all 14 modes.

Corollary 17.2 (Vanishing at the observer pair). $s(W) = s(t) = 0$ exactly.

Corollary 17.3 (Exact ratios). $a_3 = 2a_4$, $a_5 = (2 - \sqrt{2})a_4$, $a_5/a_3 = (2 - \sqrt{2})/2$.

Corollary 17.4 (Closed forms in $\mathbb{Q}(\sqrt{2})$). a_2, a_4, a_6 have the exact closed forms of Theorem 6.2.

Corollary 17.5 (Weight fractions). The $|s|^2$ -mass distribution over the D_4 orbits is 0.45%, 95.17%, 4.39%.

18 Falsification tests

1. **Vacuum with nonzero observable.** A confirmed vacuum with $\langle \mathcal{O}_a \rangle \neq 0$ for some vacuum-native observable would falsify Theorem 13.1.
2. **Non-maximizer selection.** A confirmed vacuum s with $S(s)/S_0 < 1$ while another state s' has $S(s')/S_0 > S(s)/S_0$ would falsify Theorem 14.1.
3. **Entropy-well deviation.** A measured $S(\epsilon)/S_0 \neq 1 - \gamma\epsilon^2 + O(\epsilon^4)$ would falsify the identification $\epsilon = \|M^T s\|$.
4. **Rank-nullity violation.** A measured $\text{rank}(M) \neq 13$ or $\text{nullity}(M^T) \neq 1$ would falsify Theorem 16.1.
5. **Route disagreement.** A demonstration that the Landauer route and the entropy-well route select different states would falsify Theorem 15.1.

19 Conclusion

This paper has established the zero state of the Hodge Complex Standard Model in complete form.

The 14-mode multiplet at the self-paired eigenvalue $\lambda = 4$ carries 19 vacuum-native observables, forming a 14×19 measurement matrix M with entries in $\mathbb{Q}(\sqrt{2})$. The matrix has rank 13 and nullity 1, so there is a unique nullspace vector s (up to scale), the zero state.

The zero state has the following structural properties:

1. It is odd under the chiral reflection rs : $s(rs(m)) = -s(m)$ for all 14 modes.
2. The two rs -fixed modes $W = (2, 2)$ and $t = (6, 6)$ have exactly zero amplitude.

3. The six independent amplitudes satisfy the exact ratios $a_3/a_4 = 2$ and $a_5/a_4 = 2 - \sqrt{2}$.
4. The amplitudes lie in $\mathbb{Q}(\sqrt{2})$ and are algebraic of degree 2 over $\mathbb{Q}$, with explicit quadratic minimal polynomials.
5. The $|s|^2$ -mass distribution is 0.45% on $\mathcal{O}_0$, 95.17% on $\mathcal{O}_1$, and 4.39% on $\mathcal{O}_2$.
6. The zero state is a stationary point of the DNLS Hartree energy functional, not an eigenstate of H_{DNLS} , and decays to zero under iteration of the loop operator.

The zero state is selected as the physical vacuum by the variational principle

$$s = \underset{\|s\|=1}{\text{argmin}} \|M^T s\|^2. \quad (56)$$

Two independent derivations establish this selection. The Landauer route uses the processor T at the deterministic Landauer bound; the vacuum is the state of minimum information. The entropy-well route uses the HCSM-54 entropy well with the identification $\epsilon = \|M^T s\|$; the entropy is maximized exactly when $M^T s = 0$. The two routes are dual descriptions of the same minimum-information principle. Uniqueness of the maximizer follows from $\text{rank}(M) = 13$ and $\text{nullity}(M^T) = 1$.

The zero state is the physical vacuum of the 14-mode multiplet.

References

- [1] S. Preschutti, HCSM-00: Foundations, Notation, and Mapping, HCSM White Paper Series (2026).
- [2] S. Preschutti, HCSM-01: The 8×8 Torus Substrate from $d = 4$, HCSM White Paper Series (2026).
- [3] S. Preschutti, HCSM-04: The 14-Mode Multiplet at $\lambda = 4$, HCSM White Paper Series (2026).
- [4] S. Preschutti, HCSM-05: D_4 Representation Theory of the 14-Mode Multiplet, HCSM White Paper Series (2026).

- [5] S. Preschutti, HCSM-11: The $5 + 9$ Split of the 14-Mode Multiplet, HCSM White Paper Series (2026).
- [6] S. Preschutti, HCSM-13: The Processor T and the Fermion/Boson-Class Correspondence, HCSM White Paper Series (2026).
- [7] S. Preschutti, HCSM-14: Winding Generator Spectrum and Class Partition, HCSM White Paper Series (2026).
- [8] S. Preschutti, HCSM-18: The Dynamical Bridge: DNLS on the Torus, HCSM White Paper Series (2026).
- [9] S. Preschutti, HCSM-21: The Time Theorem, HCSM White Paper Series (2026).
- [10] S. Preschutti, HCSM-23: The Midpoint Sorting Theorem, HCSM White Paper Series (2026).
- [11] S. Preschutti, HCSM-24: The Charge Theorem, HCSM White Paper Series (2026).
- [12] S. Preschutti, HCSM-26: The Observer Pair $\{W, t\}$, HCSM White Paper Series (2026).
- [13] S. Preschutti, HCSM-32: The Shell Assignment Rule, HCSM White Paper Series (2026).
- [14] S. Preschutti, HCSM-33: The Mass Map, HCSM White Paper Series (2026).
- [15] S. Preschutti, HCSM-54: Vacuum Stability and the Stability Coefficient, HCSM White Paper Series (2026).
- [16] R. Landauer, Irreversibility and heat generation in the computing process, IBM J. Res. Dev. **5**, 183 (1961).