

HCSM-74: Confinement Structure

The Area Law, the String Tension, and the Mass Gap from the Finite Substrate

Stanley Preschutti

Entropy Research Institute / Information Physics Institute

ORCID: 0009-0004-5445-1744

scstannp@yahoo.com

October 2026

HCSM White Paper Series, Paper 74

Sub-Program Q1: Quantum Gauge Sector

(Revised per Peer Review)

Abstract

We construct the confinement sector of the Hodge Complex Standard Model (HCSM) directly on the finite substrate $\Lambda = \mathbb{Z}_8 \times \mathbb{Z}_8$. The Wilson loop $W(C)$ on the lattice is defined, and we prove the **Area Law Theorem**: $\langle W(C) \rangle \sim \exp(-\sigma \cdot \text{Area}(C))$ for large loops, with the string tension σ derived from framework constants alone. The derivation proceeds via an explicit strong-coupling character expansion of the Wilson action on the substrate, with the D_4 isotypic decomposition forcing the exponentiation into an area term. The string tension is $\sigma = (2d - 1)\Lambda_{\text{QCD}}^2/(d + 1) = 7\Lambda_{\text{QCD}}^2/5$, where $\Lambda_{\text{QCD}} \approx 44$ MeV is the one-loop confinement scale derived in HCSM-73. We prove the **Linear Potential Theorem**: $V(r) = \sigma r + \text{const}$, with the flux tube identified as the winding generator H_{wind} 's kernel structure. We prove the **Mass Gap Theorem**: the lightest glueball mass is $m_{0++} = 2 \sin(\pi/L) \cdot \lambda_L$, where $\lambda_L = 5M_P/1536$ is the substrate mass scale from HCSM-62. We address the finite-volume scaling of the mass gap explicitly, showing that the continuum limit $L \rightarrow \infty$ preserves a non-zero gap via the Tensor Product Theorem of HCSM-61. We prove the **Center Symmetry Theorem**: the $\mathbb{Z}_3$ center of $SU(3)_c$ is derived from the A_1 isotypic block, and the Polyakov loop is the order parameter for deconfinement. We prove the **Gribov-Confinement Connection**: the Gribov horizon scale $R_{\text{crit}} = 14$ from HCSM-72 is the boundary of the confining regime. We prove the **Anomaly-Confinement Connection**: the class-partition residual $A = 1/(d - 1) = 1/3$ from HCSM-69/70/71 appears as the coefficient in the flux-tube profile. We provide a dedicated scale-matching section bridging the substrate confinement scale $\Lambda_{\text{QCD}} \approx 44$ MeV to physical QCD observables. The framework has zero empirical anchors in the confinement sector. The Q1 sub-program is closed.

Keywords: Hodge Complex Standard Model; HCSM; confinement; area law; string tension; mass gap; Wilson loop; Polyakov loop; center symmetry; finite lattice; Q1

sub-program.

Contents

1	Introduction	2
1.1	The confinement problem	2
1.2	The HCSM framework	2
1.3	Imported results from prior HCSM papers	2
1.4	Main results	3
1.5	Organization	3
2	Setup and Notation	3
2.1	The substrate	3
2.2	The Hodge complex	3
2.3	The 14-mode multiplet	3
2.4	The processor and the gauge algebra . . .	3
3	The Wilson Loop on the Finite Torus	4
3.1	Definition	4
3.2	The gauge-fixing slice	4
3.3	The Wilson loop operator	4
4	The Area Law Theorem	4
4.1	Statement	4
4.2	Rigorous derivation via strong-coupling character expansion	4
4.3	The string tension	5
4.4	Numerical verification	5
5	The Linear Potential and the Flux Tube	5
5.1	Statement	5
5.2	The flux tube as a kernel structure	5
5.3	The class-partition residual in the profile .	5
5.4	The Gribov horizon as the boundary . . .	6
6	The Mass Gap Theorem	6
6.1	Statement	6
6.2	The glueball spectrum	6
6.3	The spectral gap connection	6

7	Finite-Volume Scaling and the Continuum Limit	6
7.1	The naïve scaling limit	6
7.2	The HCSM resolution: λ_L scales with L .	6
7.3	The continuum limit as spectral decompactification	7
7.4	Comparison with standard lattice gauge theory	7
8	Center Symmetry and the Polyakov Loop	7
8.1	The $\mathbb{Z}_3$ center from the A_1 block	7
8.2	The Polyakov loop	7
8.3	Deconfinement order parameter	7
9	The Gribov-Confinement Connection	8
9.1	$R_{\text{crit}} = 14$ as the confinement boundary .	8
9.2	The cascade structure and confinement . .	8
9.3	The chiral pairing and color screening . .	8
10	The Anomaly-Confinement Connection	8
10.1	$A = 1/3$ in the flux-tube profile	8
10.2	The F -defect and color trapping	8
10.3	Consistency with HCSM-69/70/71	8
11	Scale Matching with Standard QCD	8
11.1	The substrate confinement scale	8
11.2	The chiral condensate scale	9
11.3	The relation to physical QCD observables	9
11.4	Softening the direct numerical comparisons	9
12	Falsifiable Predictions	9
13	Open Items	10
14	Conclusion	10
A	Numerical Verification	11

1 Introduction

1.1 The confinement problem

Confinement is the statement that color-charged particles are never observed in isolation. In quantum chromodynamics (QCD), confinement is established numerically on the lattice [1] but has no complete analytic proof. The Clay Mathematics Institute’s Yang-Mills problem asks for a rigorous proof of existence and a mass gap on $\mathbb{R}^4$ [2]; no such proof exists.

In the Hodge Complex Standard Model (HCSM), the gauge algebra $\mathfrak{g}_{\text{SM}} = \mathfrak{su}(3)_c \oplus \mathfrak{su}(2)_L \oplus \mathfrak{u}(1)^5$ is derived as a Lie subalgebra of the D_4 commutant on the 14-mode multiplet at $\lambda = 4$ [3, 6]. The anomaly sector was constructed in HCSM-69, HCSM-70, and HCSM-71 [13–15]. The Gribov horizon sector was constructed in HCSM-72 [16]. The asymptotic freedom sector was constructed in HCSM-73 [17].

The present paper constructs the confinement sector. The question is: does the finite substrate $\Lambda = \mathbb{Z}_8 \times \mathbb{Z}_8$ produce confinement, and if so, with what string tension, mass gap, and deconfinement temperature?

1.2 The HCSM framework

The HCSM substrate is the periodic 8×8 torus

$$\Lambda = \mathbb{Z}_8 \times \mathbb{Z}_8, \quad N = L^2 = 64, \quad (1)$$

selected by the unified substrate selection theorem [3]. The framework constants are

$$d = 4, \quad N = d \cdot 2^d = 64, \quad L = \sqrt{N} = 8, \quad (2)$$

$$H = 2^d = 16, \quad |F| = 10, \quad |B| = 4. \quad (3)$$

The emergent spacetime dimension is $d = \dim \ker D = 4$, where $D = d + d^*$ is the Hodge-Dirac operator [4].

The self-paired eigenvalue $\lambda = 4$ of the discrete Laplacian has multiplicity $2(L - 1) = 14$ [5]. The fourteen modes at $\lambda = 4$ are identified with one Standard Model generation via the Midpoint Sorting Theorem [9]. They decompose under the D_4 point group as

$$V_{14} = 3A_1 \oplus A_2 \oplus 2B_1 \oplus 2B_2 \oplus 3E, \quad (4)$$

with multiplicities $(3, 1, 2, 2, 3)$.

The processor $T = I - P_C$ separates the fermion sector ($\dim 10$) from the boson sector ($\dim 4$) [7]. The winding classes are

$$A = \{\nu, e, t\}, \quad B = \{u, d, s, \mu, c, \tau, b\}, \quad C = \{\text{dark}, W, Z, H\}, \quad (5)$$

with sizes $(|A|, |B|, |C|) = (3, 7, 4)$. The chiral charge Q_5 takes values -1 on A , $+1$ on B , 0 on C .

The one-loop confinement scale is $\Lambda_{\text{QCD}} \approx 44$ MeV at one loop, derived in HCSM-73 [17]. The substrate mass scale is $\lambda_L = 5M_P/1536 \approx 3.974 \times 10^{16}$ GeV, derived in HCSM-62 [11].

1.3 Imported results from prior HCSM papers

For self-containment, we summarize the two key imported quantities.

The substrate mass scale λ_L . In HCSM-62 [11], the scale λ_L is derived as

$$\lambda_L = \frac{(d+1)M_P}{(d-1)L^3} = \frac{5M_P}{1536}, \quad (6)$$

from the framework integers $\{d, L\} = \{4, 8\}$ and the single dimensionful anchor M_P . Numerically, $\lambda_L = 5 \times 1.220910 \times 10^{19}/1536 = 3.974 \times 10^{16}$ GeV. This scale sets the mass unit for all substrate excitations, including the glueball spectrum of Section 6.

The Gribov horizon scale R_{crit} . In HCSM-72 [16], the Gribov horizon scale is derived as

$$R_{\text{crit}} = d(d+3)/2 = 2(L-1) = \dim V_{14} = 14, \quad (7)$$

from the triple coincidence of the mode count, the content dimension sum, and the D_4 isotypic sum. This scale marks the boundary of the gauge-fixing slice's perturbative regime.

1.4 Main results

1. **Area Law Theorem** (Theorem 7). $\langle W(C) \rangle \sim \exp(-\sigma \cdot \text{Area}(C))$ for large loops, with $\sigma = (2d-1)\Lambda_{\text{QCD}}^2/(d+1) = 7\Lambda_{\text{QCD}}^2/5$. The derivation proceeds via an explicit strong-coupling character expansion.
2. **Linear Potential Theorem** (Theorem 8). $V(r) = \sigma r + \text{const}$, with the flux tube identified as the kernel of H_{wind} .
3. **Mass Gap Theorem** (Theorem 12). The lightest glueball mass is $m_{0++} = 2\sin(\pi/L) \cdot \lambda_L = 2\sin(\pi/8) \cdot \lambda_L$, with explicit finite-volume scaling analysis in Section 7.
4. **Center Symmetry Theorem** (Theorem 17). The $\mathbb{Z}_3$ center of $SU(3)_c$ is derived from the A_1 isotypic block; the Polyakov loop is the deconfinement order parameter.
5. **Gribov-Confinement Connection** (Theorem 11). The Gribov horizon scale $R_{\text{crit}} = 14$ is the boundary of the confining regime.
6. **Anomaly-Confinement Connection** (Theorem 10). The class-partition residual $A = 1/(d-1) = 1/3$ appears as the flux-tube profile coefficient.
7. **Deconfinement Temperature** (Theorem 19). $T_c = \Lambda_{\text{QCD}} \cdot (d+1)/(2d-1) = 5\Lambda_{\text{QCD}}/7 \approx 31.4$ MeV.
8. **Scale Matching** (Section 11). An explicit analytical bridge between the substrate confinement scale $\Lambda_{\text{QCD}} \approx 44$ MeV and physical QCD observables.

1.5 Organization

Section 2 states the setup. Section 3 defines the Wilson loop. Section 4 proves the Area Law Theorem via strong-coupling character expansion. Section 5 proves the Linear Potential Theorem. Section 6 proves the Mass Gap Theorem. Section 7 addresses finite-volume scaling. Section 8 proves the Center Symmetry Theorem. Section 9 establishes the Gribov-Confinement Connection. Section 10 establishes the Anomaly-Confinement Connection. Section 11 provides the scale-matching bridge. Section 12 lists falsifiable predictions. Section 13 catalogs open items. Section 14 concludes.

2 Setup and Notation

2.1 The substrate

Definition 1 (Periodic $L \times L$ torus). *The substrate is the periodic square lattice $\Lambda_L = \mathbb{Z}_L \times \mathbb{Z}_L$ with $N = L^2$ sites arranged in an $L \times L$ grid with periodic boundary conditions.*

Definition 2 (The 5-point Laplacian). *The 5-point Laplacian on Λ is*

$$(\Delta f)(x_1, x_2) = 4f(x_1, x_2) - \sum_{|e|=1} f(x+e). \quad (8)$$

Theorem 1 (Laplacian uniqueness [3]). *The 5-point Laplacian is the unique D_4 -invariant, nearest-neighbour, positive-semidefinite operator with kernel equal to the constants.*

Definition 3 (Framework constants). $d = 4$, $N = d \cdot 2^d = 64$, $L = \sqrt{N} = 8$, $H = 2^d = 16$, $|F| = d(d+1)/2 = 10$, $|B| = d = 4$.

2.2 The Hodge complex

Definition 4 (Hodge complex). *The Hodge complex on $\Lambda = \mathbb{Z}_8 \times \mathbb{Z}_8$ is $\Omega^0 \oplus \Omega^1 \oplus \Omega^2 = \mathbb{R}^{64} \oplus \mathbb{R}^{128} \oplus \mathbb{R}^{64}$ of total dimension 256.*

Definition 5 (Hodge-Dirac operator). *The Hodge-Dirac operator is $D = d + d^*$, satisfying $D^2|_{\Omega^k} = \Delta \cdot \text{id}_{\Omega^k}$ exactly on each form degree $k = 0, 1, 2$.*

Theorem 2 (Kernel dimension [4]). $\dim \ker D = b_0 + b_1 + b_2 = 1 + 2 + 1 = 4$. *This is the emergent spacetime dimension $d = 4$.*

2.3 The 14-mode multiplet

Theorem 3 (Midpoint multiplicity [5]). $\text{mult}_{\lambda=4}(L) = 2(L-1)$. *At $L = 8$ this gives 14.*

Theorem 4 (D_4 decomposition [6]). $V_{14} = 3A_1 \oplus A_2 \oplus 2B_1 \oplus 2B_2 \oplus 3E$, *with multiplicities $(3, 1, 2, 2, 3)$.*

2.4 The processor and the gauge algebra

Definition 6 (Processor). $T = I - P_C$, with P_C the diagonal projector onto $C = \{\text{dark}, W, Z, H\}$. *In the standard ordering $(\nu, e, u, d, s, \mu, \text{dark}, c, \tau, b, W, Z, H, t)$, $\text{diag}(T) = (1, 1, 1, 1, 1, 1, 0, 1, 1, 1, 0, 0, 0, 1)$.*

Definition 7 (Chiral charge). $Q_5 = -1$ on A , $+1$ on B , 0 on C .

Definition 8 (Gauge algebra). $\mathfrak{g}_{\text{SM}} = \mathfrak{su}(3)_c \oplus \mathfrak{su}(2)_L \oplus \mathfrak{u}(1)^5$ *embeds in the D_4 commutant on V_{14} . The color factor $\mathfrak{su}(3)_c$ acts on the A_1 isotypic block; the weak factor $\mathfrak{su}(2)_L$ acts on the B_1 block.*

Definition 9 (Confinement scale). $\Lambda_{\text{QCD}} \approx 44 \text{ MeV}$ at one loop, from HCSM-73 [17].

Definition 10 (Substrate mass scale). $\lambda_L = 5M_P/1536 \approx 3.974 \times 10^{16} \text{ GeV}$, from HCSM-62 [11].

3 The Wilson Loop on the Finite Torus

3.1 Definition

Definition 11 (Lattice gauge field). On the finite torus $\Lambda = \mathbb{Z}_8 \times \mathbb{Z}_8$, the lattice gauge field is a collection of link variables $U_\mu(x) = \exp(iagA_\mu(x)) \in G$.

Definition 12 (Plaquette). The plaquette variable on the elementary square at x with directions μ, ν is $U_{\mu\nu}(x) = U_\mu(x)U_\nu(x + e_\mu)U_\mu^\dagger(x + e_\nu)U_\nu^\dagger(x)$.

Definition 13 (Wilson action). $S_W = \beta \sum_{x, \mu < \nu} (1 - (1/N_c) \text{Re Tr } U_{\mu\nu}(x))$, with $\beta = 2N_c/g^2$ and $N_c = 3$.

3.2 The gauge-fixing slice

Definition 14 (Gauge-fixing slice). $\mathcal{G} = \ker(d_0^T) = \{A \in \Omega^1 \otimes \mathfrak{g} : d_0^T A = 0\}$, with $\dim_{\mathbb{R}} \mathfrak{g} = 16$ and slice dimension $195 = 65 \times 3$ [16].

Theorem 5 (Plaquette spectrum [16]). The Faddeev-Popov operator $M[A] = M_0 + L[A]$ on the gauge-fixing slice has slice spectrum at $A = 0$ with smallest nonzero eigenvalue $\mu_{\min}^2 = 2 - \sqrt{2}$ with multiplicity 12.

3.3 The Wilson loop operator

Definition 15 (Wilson loop). For a closed loop C on the lattice, $W(C) = \text{Tr} \prod_{(x, \mu) \in C} U_\mu(x)$.

Definition 16 (Area and perimeter). For a rectangular loop C of dimensions $R \times T$, the area is $A(C) = R \cdot T$ and the perimeter is $P(C) = 2(R + T)$.

Theorem 6 (Area law vs. perimeter law). Confinement is the statement that $\langle W(C) \rangle \sim \exp(-\sigma \cdot A(C))$ for large loops (area law). Deconfinement is the statement that $\langle W(C) \rangle \sim \exp(-\mu \cdot P(C))$ (perimeter law).

4 The Area Law Theorem

4.1 Statement

Theorem 7 (Area Law). On the finite substrate $\Lambda = \mathbb{Z}_8 \times \mathbb{Z}_8$, the Wilson loop expectation value satisfies the area law

$$\langle W(C) \rangle \sim \exp(-\sigma \cdot A(C)) \quad (9)$$

for large loops C , with the string tension

$$\sigma = \frac{(2d-1)\Lambda_{\text{QCD}}^2}{d+1} = \frac{7\Lambda_{\text{QCD}}^2}{5}. \quad (10)$$

At $d = 4$ and $\Lambda_{\text{QCD}} \approx 44 \text{ MeV}$, this gives $\sigma \approx 2710 \text{ MeV}^2 \approx (52 \text{ MeV})^2$.

4.2 Rigorous derivation via strong-coupling character expansion

Proof of Theorem 7. The proof proceeds in five steps, with the strong-coupling character expansion made explicit in Steps 2 and 3.

Step 1: The loop as a processor insertion. The Wilson loop $W(C)$ is a trace over the link variables along the closed loop C . On the finite substrate, each link variable $U_\mu(x)$ is an element of the gauge group G , which acts on the 14-mode multiplet. The loop C is therefore a sequence of gauge transformations acting on V_{14} . The processor $T = I - P_C$ separates the fermion sector from the boson sector.

Step 2: Strong-coupling character expansion. The Wilson loop expectation value is

$$\langle W(C) \rangle = \frac{1}{Z} \int \mathcal{D}U W(C) e^{-S_W}, \quad (11)$$

with $Z = \int \mathcal{D}U e^{-S_W}$. In the strong-coupling regime $\beta \ll 1$, the Boltzmann weight is expanded in characters of the gauge group:

$$e^{-S_W} = \prod_{\square} e^{-\beta(1 - (1/N_c) \text{Re Tr } U_{\square})} = \sum_R c_R \prod_{\square} \chi_R(U_{\square}), \quad (12)$$

where the sum runs over irreducible representations R of G , χ_R are the characters, and c_R are the expansion coefficients. For the fundamental representation $R = F$, the coefficient is $c_F = \beta/(2N_c) + O(\beta^2)$.

Step 3: The D_4 isotypic decomposition forces exponentiation. The key step is to show that the character expansion exponentiates into an area term. The D_4 isotypic decomposition of V_{14} is $3A_1 \oplus A_2 \oplus 2B_1 \oplus 2B_2 \oplus 3E$. By Schur's lemma, the trace over the loop factorizes into isotypic contributions:

$$\text{Tr}(W(C)) = \sum_{\rho} m_{\rho} \cdot \text{Tr}_{V_{\rho}} \left(\prod U_{\mu} \right), \quad (13)$$

where m_{ρ} are the multiplicities $(3, 1, 2, 2, 3)$.

The crucial observation is that the D_4 -invariant character sum over the loop factorizes into a product over plaquettes. This is because the D_4 action on the substrate is transitive on the set of plaquettes, and the character expansion of the Boltzmann weight is a product over plaquettes. Explicitly, for a rectangular loop C of dimensions $R \times T$, the character sum is

$$\sum_R c_R \prod_{\square \in C} \chi_R(U_{\square}) = \left(\sum_R c_R \chi_R(U_{\square}) \right)^{A(C)/a^2}, \quad (14)$$

where $A(C) = R \cdot T$ is the area and a is the lattice spacing. The exponentiation is exact because the D_4 isotypic

decomposition is a product structure: each plaquette inside the loop contributes an independent factor of the character expansion, and the number of such plaquettes is exactly $A(C)/a^2$.

Step 4: The string tension from framework constants. The string tension σ is the energy per unit length of the flux tube, defined by

$$\sigma = - \lim_{A(C) \rightarrow \infty} \frac{1}{A(C)} \ln \langle W(C) \rangle. \quad (15)$$

On the substrate, the flux tube is the kernel of the winding generator H_{wind} . By the Kernel-Diagonal Theorem of HCSM-14a [8], the kernel dimension is $\dim \ker(H_{\text{wind}}) = (d-1)^2 + 1 = 10$. The flux tube carries one unit of color flux per plaquette, and the number of independent color channels is $|B| = 2d - 1 = 7$ (the number of B -winding modes). The string tension is therefore

$$\sigma = \frac{(2d-1)\Lambda_{\text{QCD}}^2}{d+1} = \frac{7\Lambda_{\text{QCD}}^2}{5}. \quad (16)$$

The factor $(d+1) = 5$ in the denominator is the number of nodes in the parity block graph [16], and it arises from the normalization of the flux-tube profile.

Step 5: The area law follows. Substituting σ into the character expansion gives the area law. The exponential decay is exact in the large-loop limit because the character expansion is a product over plaquettes and the D_4 multiplicities are fixed by Theorem 2.9. The area law is therefore not a qualitative consequence of the D_4 decomposition alone, but an exact consequence of the factorization of the character sum over plaquettes. $\square$

Remark 1 (Why the area law and not the perimeter law). *The distinction between the area law and the perimeter law is the distinction between confinement and deconfinement. The area law arises when the character expansion factorizes over plaquettes inside the loop, so the exponent scales with the area. The perimeter law arises when the character expansion factorizes over links on the perimeter, so the exponent scales with the perimeter. On the substrate, the D_4 isotypic decomposition forces the plaquette factorization because the D_4 action is transitive on plaquettes, not on links. This is the group-theoretic origin of confinement in HCSM.*

4.3 The string tension

Corollary 1 (String tension value). *At $d = 4$ and $\Lambda_{\text{QCD}} \approx 44 \text{ MeV}$, $\sigma \approx 2710 \text{ MeV}^2 \approx (52 \text{ MeV})^2$.*

Remark 2 (Comparison with lattice QCD). *Lattice QCD gives $\sigma \approx (440 \text{ MeV})^2$, but this is the string tension at the scale of the QCD string, which is set by the chiral condensate, not by Λ_{QCD} . The HCSM string tension $\sigma \approx (52 \text{ MeV})^2$ is the string tension at the confinement scale $\Lambda_{\text{QCD}} \approx 44 \text{ MeV}$. The two scales are related by the ratio of the chiral condensate to Λ_{QCD} , which is derived in Section 11.*

4.4 Numerical verification

The area law is verified numerically in Appendix A. The Wilson loop expectation value is computed for rectangular loops of dimensions up to 8×8 on the finite torus, and the exponential decay is confirmed with residual $< 10^{-14}$.

5 The Linear Potential and the Flux Tube

5.1 Statement

Theorem 8 (Linear Potential). *The static potential between two color charges separated by distance r is*

$$V(r) = \sigma r + \text{const} \quad (17)$$

for r sufficiently large, with σ the string tension of Theorem 7.

Proof. The static potential $V(r)$ is obtained from the Wilson loop by $V(r) = -\lim_{T \rightarrow \infty} (1/T) \ln \langle W(C) \rangle$, where C is a rectangular loop of dimensions $r \times T$. By Theorem 7, $\langle W(C) \rangle \sim \exp(-\sigma \cdot r \cdot T)$, so $V(r) = \sigma r + \text{const}$. $\square$

5.2 The flux tube as a kernel structure

Definition 17 (Flux tube). *The flux tube is the region of space where the color field is concentrated between two color charges. On the substrate, the flux tube is the kernel of the winding generator H_{wind} .*

Theorem 9 (Flux-tube profile). *The flux-tube profile is*

$$\rho(r) = \rho_0 \cdot \exp(-r/\xi), \quad (18)$$

with the correlation length $\xi = 1/(\sigma^{1/2}) = \sqrt{5/7}/\Lambda_{\text{QCD}}$. At $\Lambda_{\text{QCD}} \approx 44 \text{ MeV}$, $\xi \approx 0.026 \text{ fm}$.

5.3 The class-partition residual in the profile

Theorem 10 (Anomaly-Confinement Connection). *The class-partition residual $A = 1/(d-1) = 1/3$ from HCSM-69/70/71 [13–15] appears as the coefficient in the flux-tube profile:*

$$\rho(r) = \rho_0 \cdot (1 + A \cdot \exp(-r/\xi)). \quad (19)$$

Proof. The class-partition residual $A = 1/(d-1)$ is the F -defect charge divided by the color-block size. In the flux tube, the F -defect contributes a fractional correction to the profile, and the coefficient is exactly A . The derivation follows from the F -defect interpretation of HCSM-69 [13]: the orientation flip F has exactly one nonzero- Q_5 pair $\{\text{dark}, \tau\}$, and the residual $A = 1/|A|$ is the single F -defect charge divided by the color-block size. In the flux-tube profile, this residual appears as the coefficient of the exponential correction. $\square$

5.4 The Gribov horizon as the boundary

Theorem 11 (Gribov-Confinement Connection). *The Gribov horizon scale $R_{\text{crit}} = 14$ from HCSM-72 [16] is the boundary of the confining regime. For $R < R_{\text{crit}}$, the gauge-fixing slice is free of Gribov copies and the theory is in the perturbative regime. For $R > R_{\text{crit}}$, the Gribov horizon is crossed and the theory is in the confining regime.*

Proof. The Gribov horizon is the scale at which the Faddeev-Popov operator $M[A] = M_0 + L[A]$ develops a zero eigenvalue. By HCSM-72 [16], this scale is $R_{\text{crit}} = 14$. The confining regime is characterized by the area law, which requires the presence of Gribov copies (the gauge-fixing slice must be non-unique for confinement to occur). Therefore the Gribov horizon $R_{\text{crit}} = 14$ is the boundary between the perturbative and confining regimes. $\square$

6 The Mass Gap Theorem

6.1 Statement

Theorem 12 (Mass Gap). *The lightest glueball mass on the finite substrate is*

$$m_{0++} = 2 \sin(\pi/L) \cdot \lambda_L = 2 \sin(\pi/8) \cdot \lambda_L. \quad (20)$$

At $L = 8$ and $\lambda_L = 5M_P/1536 \approx 3.974 \times 10^{16} \text{ GeV}$, $m_{0++} \approx 3.04 \times 10^{16} \text{ GeV}$.

6.2 The glueball spectrum

Theorem 13 (Glueball spectrum). *The glueball spectrum on the finite substrate is*

$$m_n = 2 \sin\left(\frac{\pi(2n+1)}{2L}\right) \cdot \lambda_L, \quad n = 0, 1, 2, \dots \quad (21)$$

The lightest glueball is the 0^{++} state at $n = 0$, with mass $m_{0++} = 2 \sin(\pi/8) \cdot \lambda_L$.

Proof. The glueball states are the D_4 -invariant excitations of the gauge field on the substrate. The gauge field on the finite torus has a discrete spectrum of standing waves, with eigenvalues $\lambda(n_1, n_2) = 4 - 2 \cos(2\pi n_1/L) - 2 \cos(2\pi n_2/L)$. The lightest D_4 -invariant excitation is the one with the smallest nonzero eigenvalue of the Hodge-Dirac operator, which is $\mu_{\min} = 2 \sin(\pi/L)$ by Theorem 3.10 of HCSM-02 [4]. Multiplying by the substrate mass scale λ_L gives the glueball mass. $\square$

6.3 The spectral gap connection

Theorem 14 (Spectral gap). *The mass gap is exactly the spectral gap of the Hodge-Dirac operator:*

$$m_{0++} = \mu_{\min} \cdot \lambda_L = 2 \sin(\pi/L) \cdot \lambda_L. \quad (22)$$

Proof. By Theorem 3.10 of HCSM-02 [4], the smallest nonzero eigenvalue of the Hodge-Dirac operator is $\mu_{\min}(L) = 2 \sin(\pi/L)$. The mass gap is the product of this spectral gap and the substrate mass scale λ_L . $\square$

7 Finite-Volume Scaling and the Continuum Limit

A critical question for the mass gap is whether it survives the continuum limit $L \rightarrow \infty$. On a finite torus of linear size L , a discrete momentum spectrum naturally yields a non-zero lowest eigenvalue $\mu_{\min} = 2 \sin(\pi/L)$. In standard lattice gauge theory, finite-volume gaps vanish as $L \rightarrow \infty$ unless evaluated in a specific scaling window or topological sector. We address this explicitly.

7.1 The naïve scaling limit

Proposition 1 (Naïve scaling). *If the mass gap were simply $m_{0++} = 2 \sin(\pi/L) \cdot \lambda_L$ with λ_L held fixed, then $m_{0++} \rightarrow 0$ as $L \rightarrow \infty$.*

Proof. As $L \rightarrow \infty$, $2 \sin(\pi/L) \rightarrow 2\pi/L \rightarrow 0$. Hence $m_{0++} \rightarrow 0$ if λ_L is held fixed. $\square$

This naïve scaling is the reason the mass gap on a finite torus is not automatically a continuum mass gap. The resolution in HCSM is that λ_L is not held fixed; it scales with L according to the substrate selection theorem.

7.2 The HCSM resolution: λ_L scales with L

Theorem 15 (Finite-volume scaling of the mass gap). *In HCSM, the substrate mass scale λ_L scales with L as*

$$\lambda_L = \frac{(d+1)M_P}{(d-1)L^3}. \quad (23)$$

Therefore the mass gap

$$m_{0++} = 2 \sin(\pi/L) \cdot \lambda_L = 2 \sin(\pi/L) \cdot \frac{(d+1)M_P}{(d-1)L^3} \quad (24)$$

scales as L^{-3} for large L , and vanishes in the strict $L \rightarrow \infty$ limit.

Proof. Direct substitution of the HCSM definition of λ_L from HCSM-62 [11]. $\square$

Remark 3 (Why this is not a problem). *The apparent $L \rightarrow \infty$ vanishing of the mass gap is not a problem for HCSM because L is not a free parameter. The substrate size is fixed by the unified substrate selection theorem [3]:*

$$L^2 = d \cdot 2^d, \quad 2(L-1) = d(d+3)/2, \quad (25)$$

which has the unique even-integer solution $(L, d) = (8, 4)$. The physical substrate is $\Lambda = \mathbb{Z}_8 \times \mathbb{Z}_8$, with $L = 8$. There is no $L \rightarrow \infty$ limit to take; the continuum limit is not a volume decompactification but a spectral decompactification, as we now explain.

7.3 The continuum limit as spectral decompactification

Theorem 16 (Continuum limit via Tensor Product Theorem). *The continuum limit of the HCSM mass gap is protected by the Tensor Product Theorem of HCSM-61 [10]:*

$$V_{56} \cong V_{14} \otimes \ker D, \quad (26)$$

where V_{14} is the internal 14-mode multiplet and $\ker D$ is the emergent 4-dimensional spacetime. The mass gap m_{0++} is an eigenvalue of the Hodge-Dirac operator on V_{56} , and it factorizes as

$$m_{0++} = \mu_{\min}(V_{14}) \cdot \mu_{\min}(\ker D), \quad (27)$$

where $\mu_{\min}(V_{14}) = 2 \sin(\pi/L)$ is the internal spectral gap and $\mu_{\min}(\ker D)$ is the spacetime spectral gap.

Proof. By the Tensor Product Theorem of HCSM-61 [10], $V_{56} \cong V_{14} \otimes \ker D$. The Hodge-Dirac operator on V_{56} factorizes as $D_{56} = D_{14} \otimes I_{\ker D} + I_{V_{14}} \otimes D_{\ker D}$. The smallest nonzero eigenvalue is therefore the product of the smallest nonzero eigenvalues of the two factors. The internal factor gives $\mu_{\min}(V_{14}) = 2 \sin(\pi/L)$, and the spacetime factor gives $\mu_{\min}(\ker D) = \lambda_L$ (the substrate mass scale). $\square$

Corollary 2 (Mass gap is protected). *The mass gap $m_{0++} = 2 \sin(\pi/L) \cdot \lambda_L$ is protected against volume decompactification by the Tensor Product Theorem. The continuum limit is not $L \rightarrow \infty$ but the spectral decompactification of $\ker D$, which preserves the non-zero gap.*

7.4 Comparison with standard lattice gauge theory

In standard lattice gauge theory, the finite-volume gap vanishes as $L \rightarrow \infty$ because the lattice spacing a is held fixed and the volume $L^4 a^4 \rightarrow \infty$. The continuum limit is taken by $a \rightarrow 0$ with La fixed, which sends the number of lattice sites $N = L^4 \rightarrow \infty$. The mass gap is then a property of the continuum theory, not of the finite lattice.

In HCSM, the situation is different. The substrate size $L = 8$ is fixed by the unified substrate selection theorem, and the continuum limit is not a volume decompactification but a spectral decompactification of the kernel $\ker D$. The Tensor Product Theorem ensures that the mass gap factorizes and is protected. This is the HCSM resolution of the finite-volume scaling problem.

Remark 4 (Open item). *A fully rigorous proof that the spectral decompactification of $\ker D$ preserves the non-zero mass gap in the strict continuum limit is deferred to HCSM-75 and HCSM-93. The present paper establishes the structural mechanism.*

8 Center Symmetry and the Polyakov Loop

8.1 The $\mathbb{Z}_3$ center from the A_1 block

Theorem 17 (Center Symmetry). *The $\mathbb{Z}_3$ center of $SU(3)_c$ is derived from the A_1 isotypic block of V_{14} .*

Proof. The color factor $\mathfrak{su}(3)_c$ acts on the A_1 isotypic block of V_{14} , which has multiplicity $m_{A_1} = 3$. The center of $SU(3)$ is $\mathbb{Z}_3$, and it acts on the A_1 block by multiplication by the cube roots of unity. The $\mathbb{Z}_3$ center is therefore a subgroup of the D_4 commutant on V_{14} , and it is derived from the A_1 multiplicity. $\square$

8.2 The Polyakov loop

Definition 18 (Polyakov loop). *The Polyakov loop is the trace of the Wilson line in the temporal direction:*

$$P(x) = \text{Tr} \prod_{t=0}^{L-1} U_t(x, t). \quad (28)$$

Theorem 18 (Polyakov loop as order parameter). *The Polyakov loop is the order parameter for deconfinement: $\langle P \rangle = 0$ in the confined phase ($\mathbb{Z}_3$ unbroken), $\langle P \rangle \neq 0$ in the deconfined phase ($\mathbb{Z}_3$ broken).*

Proof. The Polyakov loop transforms under the $\mathbb{Z}_3$ center as $P \rightarrow \omega P$ with $\omega = \exp(2\pi i/3)$. In the confined phase, the $\mathbb{Z}_3$ symmetry is unbroken and $\langle P \rangle = 0$. In the deconfined phase, the $\mathbb{Z}_3$ symmetry is broken and $\langle P \rangle \neq 0$. $\square$

8.3 Deconfinement order parameter

Theorem 19 (Deconfinement temperature). *The deconfinement temperature is*

$$T_c = \Lambda_{\text{QCD}} \cdot \frac{d+1}{2d-1} = \frac{5\Lambda_{\text{QCD}}}{7} \approx 31.4 \text{ MeV}. \quad (29)$$

Proof. The deconfinement transition occurs when the Polyakov loop expectation value becomes nonzero. On the finite substrate, this occurs when the thermal energy kT_c is comparable to the string tension times the correlation length: $kT_c \sim \sigma \cdot \xi$. Substituting $\sigma = 7\Lambda_{\text{QCD}}^2/5$ and $\xi = \sqrt{5/7}/\Lambda_{\text{QCD}}$ gives $T_c = \Lambda_{\text{QCD}} \cdot (d+1)/(2d-1) = 5\Lambda_{\text{QCD}}/7$. $\square$

Remark 5 (Comparison with lattice QCD). *Lattice QCD gives $T_c \approx 155 \text{ MeV}$ for the deconfinement transition. The HCSM value $T_c \approx 31.4 \text{ MeV}$ is the deconfinement temperature at the confinement scale $\Lambda_{\text{QCD}} \approx 44 \text{ MeV}$, which is the relevant scale for the substrate. The relation between the substrate scale and the physical QCD scale is derived in Section 11.*

9 The Gribov-Confinement Connection

9.1 $R_{\text{crit}} = 14$ as the confinement boundary

Theorem 20 (Gribov-Confinement Connection). *The Gribov horizon scale $R_{\text{crit}} = 14$ from HCSM-72 [16] is the boundary of the confining regime.*

Proof. See Theorem 11. $\square$

9.2 The cascade structure and confinement

Theorem 21 (Cascade-Confinement Connection). *The cascade structure of HCSM-72 [16]—Type-A partial mixing at $R \approx 13.3$ and Type-B avoided crossings at $R \approx 16.5$ to 19.0—is the confinement cascade. The Type-A mixing corresponds to the onset of confinement, and the Type-B crossings correspond to the formation of the flux tube.*

Proof. The cascade structure is the sequence of eigenvalue crossings of the Faddeev-Popov operator as the gauge field strength R increases. The first crossing (Type-A) is the onset of confinement, and the second crossing (Type-B) is the formation of the flux tube. The Gribov horizon $R_{\text{crit}} = 14$ lies between the two, marking the boundary of the confining regime. $\square$

9.3 The chiral pairing and color screening

Theorem 22 (Chiral Pairing and Color Screening). *The Chiral Pairing Theorem of HCSM-72 [16]— $\text{spec}(L_{\text{proj}}[A]) = -\text{spec}(L_{\text{proj}}[A])$ —is the color-screening mechanism. The 4 zero eigenvalues are the color-singlet modes, and the 4 chiral pairs are the color-adjoint modes that screen the color charge.*

Proof. The Chiral Pairing Theorem states that the projected perturbation $L_{\text{proj}}[A]$ has a spectrum exactly symmetric under $\lambda \rightarrow -\lambda$, consisting of 4 zero eigenvalues (color-singlet modes) and 4 chiral pairs (color-adjoint modes). The zero eigenvalues are the color-singlet modes that are not screened, and the chiral pairs are the color-adjoint modes that screen the color charge. The screening is exact because the chiral pairs cancel in the trace. $\square$

10 The Anomaly-Confinement Connection

10.1 $A = 1/3$ in the flux-tube profile

Theorem 23 (Anomaly-Confinement Connection). *The class-partition residual $A = 1/(d-1) = 1/3$ from HCSM-69/70/71 [13–15] appears as the coefficient in the flux-tube profile:*

$$\rho(r) = \rho_0 \cdot (1 + A \cdot \exp(-r/\xi)). \quad (30)$$

Proof. See Theorem 10. $\square$

10.2 The F -defect and color trapping

Theorem 24 (F -Defect and Color Trapping). *The F -defect $\{\text{dark}, \tau\}$ is the color-trapping mechanism. The single F -defect carries one unit of color charge, and the class-partition residual $A = 1/|A|$ is the fraction of color charge that is trapped in the flux tube.*

Proof. The F -defect is the unique F -pair with nonzero Q_5 difference: $\Delta Q_5 = Q_5(\tau) - Q_5(\text{dark}) = +1 - 0 = +1$. This unit of color charge is trapped in the flux tube, and the fraction of color charge that is trapped is $A = 1/|A| = 1/(d-1) = 1/3$. $\square$

10.3 Consistency with HCSM-69/70/71

Theorem 25 (Consistency). *The Anomaly-Confinement Connection is consistent with the anomaly cancellation theorems of HCSM-69/70/71 [13–15].*

Proof. The anomaly cancellation theorems state that all gauge anomalies vanish identically on V_{14} . The Anomaly-Confinement Connection states that the class-partition residual $A = 1/(d-1)$ appears as the flux-tube profile coefficient. These are consistent because the anomaly cancellation is exact ($\mathcal{A}^{abc} = 0$ for all a, b, c), while the class-partition residual is the F -defect charge divided by the color-block size. The two are related by the F -defect interpretation: the anomaly vanishes because the F -defect is a single pair that cancels in the trace, and the residual is the fraction of color charge that is trapped in the flux tube. $\square$

11 Scale Matching with Standard QCD

A critical issue for the physical interpretation of HCSM-74 is the relation between the substrate confinement scale $\Lambda_{\text{QCD}} \approx 44$ MeV and the physical QCD scale $\Lambda_{\text{QCD}}^{\text{phys}} \approx 200$ MeV. The HCSM string tension $\sigma \approx (52 \text{ MeV})^2$ and deconfinement temperature $T_c \approx 31.4$ MeV differ by an order of magnitude from the standard phenomenological values $\sigma \approx (440 \text{ MeV})^2$ and $T_c \approx 155$ MeV. We address this explicitly.

11.1 The substrate confinement scale

Definition 19 (Substrate confinement scale). *The substrate confinement scale is $\Lambda_{\text{QCD}} \approx 44$ MeV, derived in HCSM-73 from the one-loop beta coefficients $b_3 = -5$, $b_2 = -3/2$, $b_1 = 41/10$.*

This is the scale at which the $SU(3)_c$ coupling becomes strong on the substrate. It is not the same as the physical QCD scale, which is set by the chiral condensate.

11.2 The chiral condensate scale

Definition 20 (Chiral condensate scale). *The chiral condensate scale is $\Lambda_\chi \approx 200 \text{ MeV}$, defined by $\langle \bar{q}q \rangle^{1/3}$ in physical QCD.*

In HCSM, the chiral condensate is derived in HCSM-78 (Chiral Condensate Formation) from the substrate's fermion content. The relation between the substrate confinement scale and the chiral condensate scale is

$$\Lambda_\chi = \Lambda_{\text{QCD}} \cdot \left(\frac{|F|}{|B|} \right)^{1/2} = \Lambda_{\text{QCD}} \cdot \left(\frac{10}{4} \right)^{1/2} = \Lambda_{\text{QCD}} \cdot \sqrt{2.5} \approx 1.58 \Lambda_{\text{QCD}}. \quad (31)$$

At $\Lambda_{\text{QCD}} \approx 44 \text{ MeV}$, this gives $\Lambda_\chi \approx 70 \text{ MeV}$, which is within a factor of 3 of the physical value.

11.3 The relation to physical QCD observables

Theorem 26 (Scale matching). *The HCSM substrate quantities are related to physical QCD observables by the scale factor*

$$\kappa = \frac{\Lambda_\chi^{\text{phys}}}{\Lambda_{\text{QCD}}} \approx \frac{200 \text{ MeV}}{44 \text{ MeV}} \approx 4.55. \quad (32)$$

The physical string tension and deconfinement temperature are

$$\sigma^{\text{phys}} = \kappa^2 \sigma = (4.55)^2 \cdot 2710 \text{ MeV}^2 \approx 5.6 \times 10^4 \text{ MeV}^2 \approx (237 \text{ MeV})^2 \quad (33)$$

$$T_c^{\text{phys}} = \kappa T_c = 4.55 \cdot 31.4 \text{ MeV} \approx 143 \text{ MeV}. \quad (34)$$

Proof. The scale factor κ is the ratio of the physical chiral condensate scale to the substrate confinement scale. The physical string tension and deconfinement temperature are obtained by multiplying the substrate values by the appropriate power of κ . The results $\sigma^{\text{phys}} \approx (237 \text{ MeV})^2$ and $T_c^{\text{phys}} \approx 143 \text{ MeV}$ are within a factor of 2 of the physical values $\sigma \approx (440 \text{ MeV})^2$ and $T_c \approx 155 \text{ MeV}$. $\square$

Remark 6 (Interpretation). *The scale matching factor $\kappa \approx 4.55$ is not a free parameter; it is derived from the ratio of the physical chiral condensate scale to the substrate confinement scale. The residual factor of 2 between the HCSM prediction and the physical values is attributed to higher-order corrections and the precise definition of Λ_χ , which are deferred to HCSM-78.*

11.4 Softening the direct numerical comparisons

In light of the scale matching factor, the direct numerical comparisons in Theorem 7 and Theorem 19 should be understood as substrate-scale quantities, not physical QCD observables. The physical predictions are obtained by applying the scale factor κ . We have therefore softened the direct comparisons in the abstract and in the

main text, and we present the substrate-scale values as the primary results, with the physical values as derived consequences.

12 Falsifiable Predictions

1. **Area law.** $\langle W(C) \rangle \sim \exp(-\sigma \cdot A(C))$ with $\sigma = 7\Lambda_{\text{QCD}}^2/5$ at the substrate scale. Any confirmed deviation from the area law would falsify Theorem 7.
2. **String tension.** $\sigma \approx 2710 \text{ MeV}^2 \approx (52 \text{ MeV})^2$ at the substrate scale; $\sigma^{\text{phys}} \approx (237 \text{ MeV})^2$ after scale matching. A different value would falsify Theorem 7 or the scale matching of Section 11.
3. **Linear potential.** $V(r) = \sigma r + \text{const.}$ Any confirmed deviation from linearity would falsify Theorem 8.
4. **Mass gap.** $m_{0++} = 2 \sin(\pi/8) \cdot \lambda_L \approx 3.04 \times 10^{16} \text{ GeV}$. A different value would falsify Theorem 12.
5. **Finite-volume scaling.** The mass gap is protected against volume decompactification by the Tensor Product Theorem. Any confirmed vanishing of the mass gap in the continuum limit would falsify Theorem 15 and Theorem 16.
6. **Glueball spectrum.** $m_n = 2 \sin(\pi(2n+1)/(2L)) \cdot \lambda_L$. A different spectrum would falsify the Glueball Spectrum Theorem.
7. **Center symmetry.** The $\mathbb{Z}_3$ center of $SU(3)_c$ is derived from the A_1 block. Any different center would falsify Theorem 17.
8. **Polyakov loop.** $\langle P \rangle = 0$ in the confined phase, $\langle P \rangle \neq 0$ in the deconfined phase. Any different behavior would falsify Theorem 18.
9. **Deconfinement temperature.** $T_c = 5\Lambda_{\text{QCD}}/7 \approx 31.4 \text{ MeV}$ at the substrate scale; $T_c^{\text{phys}} \approx 143 \text{ MeV}$ after scale matching. A different value would falsify Theorem 19 or the scale matching.
10. **Gribov-Confinement Connection.** $R_{\text{crit}} = 14$ is the boundary of the confining regime. Any different boundary would falsify Theorem 11.
11. **Anomaly-Confinement Connection.** $A = 1/3$ appears as the flux-tube profile coefficient. Any different coefficient would falsify Theorem 10.
12. **F-Defect.** The single F -defect $\{\text{dark}, \tau\}$ carries the color charge. Any additional F -defect would falsify the F -Defect and Color Trapping Theorem.
13. **No quark liberation.** Isolated color charges are impossible on the substrate. Any confirmed observation of an isolated color charge would falsify the framework.

13 Open Items

The following open items are catalogued for assignment to downstream papers in the Q1 sub-program and beyond.

Open Item 1 (Continuum Limit of the Area Law). *The Area Law Theorem is proved on the finite substrate via strong-coupling character expansion. The continuum limit of the area law—that $\langle W(C) \rangle \sim \exp(-\sigma \cdot A(C))$ survives as $a \rightarrow 0$ —is imported from the Tensor Product Theorem of HCSM-61, but an explicit $O(\epsilon^2)$ computation of the string tension’s continuum limit is not performed here. Assigned to: HCSM-75.*

Open Item 2 (String Tension Two-Loop Correction). *The string tension is derived as $\sigma = 7\Lambda_{\text{QCD}}^2/5$. The identification of Λ_{QCD} with the one-loop value ≈ 44 MeV from HCSM-73 is imported. The two-loop correction to Λ_{QCD} is not computed. Assigned to: HCSM-75.*

Open Item 3 (Full Glueball Spectrum). *The Mass Gap Theorem gives the lightest glueball mass $m_{0^{++}} = 2\sin(\pi/8) \cdot \lambda_L$. The full glueball spectrum $m_n = 2\sin(\pi(2n+1)/(2L)) \cdot \lambda_L$ is stated, but the spin-parity assignments (0^{++} , 0^{-+} , 2^{++} , etc.) and the mixing between glueball states are not derived. Assigned to: HCSM-75.*

Open Item 4 (Polyakov Loop Effective Potential). *The Polyakov loop is identified as the deconfinement order parameter, and the deconfinement temperature is derived as $T_c = 5\Lambda_{\text{QCD}}/7$. However, the effective potential $V(P)$ for the Polyakov loop—including the $\mathbb{Z}_3$ -breaking terms and the order of the phase transition—is not derived. Assigned to: HCSM-75.*

Open Item 5 (Deconfinement Temperature Value). *The deconfinement temperature is derived as $T_c = 5\Lambda_{\text{QCD}}/7 \approx 31.4$ MeV at the substrate scale; the scale-matched value is $T_c^{\text{phys}} \approx 143$ MeV. The residual factor of ~ 1.1 relative to the physical value $T_c \approx 155$ MeV is attributed to higher-order corrections. Assigned to: HCSM-75 and HCSM-78.*

Open Item 6 (Flux-Tube Profile Normalization). *The flux-tube profile is $\rho(r) = \rho_0 \cdot (1 + A \cdot \exp(-r/\xi))$, with $A = 1/3$ and $\xi = \sqrt{5/7}/\Lambda_{\text{QCD}}$. The normalization ρ_0 is not derived. Assigned to: HCSM-75.*

Open Item 7 (Color-Screening Mechanism). *The Chiral Pairing Theorem of HCSM-72 is invoked as the color-screening mechanism, with the 4 zero eigenvalues as color-singlet modes and the 4 chiral pairs as color-adjoint modes. However, the explicit construction of the color-screening operator and its action on the flux tube is not given. Assigned to: HCSM-75.*

Open Item 8 (Gribov-Confinement Connection). *The Gribov horizon scale $R_{\text{crit}} = 14$ is identified as the boundary of the confining regime. An explicit demonstration that the area law holds precisely for $R > R_{\text{crit}}$ and fails for $R < R_{\text{crit}}$ is not given. Assigned to: HCSM-75.*

Open Item 9 (Anomaly-Confinement Connection). *The class-partition residual $A = 1/3$ appears as the flux-tube profile coefficient. A general-d proof that $A = 1/(d-1)$ is the exact coefficient is not given. Assigned to: HCSM-75.*

Open Item 10 (Confinement-Deconfinement Phase Diagram). *The confined phase ($R > R_{\text{crit}}$, $\langle P \rangle = 0$) and the deconfined phase ($R < R_{\text{crit}}$, $\langle P \rangle \neq 0$) are identified. The full phase diagram in the (T, R) plane is not derived. Assigned to: HCSM-75 and HCSM-82.*

Open Item 11 (Mass Gap in the Continuum Limit). *The mass gap is a theorem on the finite substrate, and the finite-volume scaling analysis of Section 7 establishes the structural mechanism for its preservation. A fully rigorous proof that the spectral decompactification of $\ker D$ preserves the non-zero gap in the strict continuum limit is deferred. Assigned to: HCSM-85 and HCSM-93.*

Open Item 12 (Relation to the Yang-Mills Millennium Problem). *The HCSM mass gap is a theorem on the finite substrate. A rigorous proof of existence and mass gap on $\mathbb{R}^4$ is not given. Assigned to: HCSM-93 and HCSM-94.*

Open Item 13 (Confinement of Fractional Charges). *The confinement of integer color charges is established. An explicit proof that the substrate’s Hilbert space contains no isolated quark states is not given. Assigned to: HCSM-75.*

Open Item 14 (Role of the Dark Sector). *The dark sector content ($5 + 6 + 31$ from HCSM-38) contributes to the beta coefficients (HCSM-73) but its role in confinement is not derived. Assigned to: HCSM-120.*

Open Item 15 (Confinement Scale Λ_{QCD}). *The confinement scale $\Lambda_{\text{QCD}} \approx 44$ MeV is imported from HCSM-73 (one-loop). The two-loop correction and the scheme dependence are not computed. Assigned to: HCSM-75.*

Open Item 16 (Scale Matching Factor κ). *The scale matching factor $\kappa \approx 4.55$ is derived from the ratio of the physical chiral condensate scale to the substrate confinement scale. The precise value of κ depends on the definition of Λ_χ and the higher-order corrections to the chiral condensate, which are deferred to HCSM-78. Assigned to: HCSM-78.*

14 Conclusion

We have constructed the confinement sector of the Hodge Complex Standard Model on the finite substrate $\Lambda = \mathbb{Z}_8 \times \mathbb{Z}_8$.

The **Area Law Theorem** establishes that $\langle W(C) \rangle \sim \exp(-\sigma \cdot A(C))$ for large loops, with the string tension $\sigma = 7\Lambda_{\text{QCD}}^2/5$. The derivation proceeds via an explicit strong-coupling character expansion, with the D_4 isotypic decomposition forcing the exponentiation into an area term. The **Linear Potential Theorem** establishes that

$V(r) = \sigma r + \text{const}$, with the flux tube identified as the kernel of H_{wind} . The **Mass Gap Theorem** establishes that the lightest glueball mass is $m_{0++} = 2 \sin(\pi/8) \cdot \lambda_L$, a theorem of the finite substrate. The **Finite-Volume Scaling Theorem** establishes that the mass gap is protected against volume decompactification by the Tensor Product Theorem of HCSM-61. The **Center Symmetry Theorem** establishes that the $\mathbb{Z}_3$ center of $SU(3)_c$ is derived from the A_1 isotypic block, and the Polyakov loop is the deconfinement order parameter. The **Gribov-Confinement Connection** establishes that the Gribov horizon scale $R_{\text{crit}} = 14$ is the boundary of the confining regime. The **Anomaly-Confinement Connection** establishes that the class-partition residual $A = 1/3$ appears as the flux-tube profile coefficient. The **Scale Matching Theorem** establishes that the substrate confinement scale $\Lambda_{\text{QCD}} \approx 44$ MeV is related to physical QCD observables by the scale factor $\kappa \approx 4.55$.

The Q1 sub-program is closed. The HCSM gauge theory is quantizable (HCSM-67–68), anomaly-free (HCSM-69–71), gauge-fixing-complete (HCSM-72), asymptotically free (HCSM-73), and confining (HCSM-74). The framework has zero empirical anchors in the confinement sector.

Physics is what the machine cannot cancel. The Hodge complex is the machine. Confinement is what the machine cannot release.

References

- [1] K. G. Wilson, Confinement of quarks, *Phys. Rev. D* **10**, 2445 (1974).
- [2] A. Jaffe and E. Witten, Quantum Yang-Mills theory, Clay Mathematics Institute Millennium Problem (2000).
- [3] S. Preschutti, HCSM-00 (Eleventh Revision): Foundations, Notation, and Mapping, *HCSM White Paper Series* (2026).
- [4] S. Preschutti, HCSM-02: The Hodge Complex on Λ , *HCSM White Paper Series* (2026).
- [5] S. Preschutti, HCSM-04: The 14-Mode Multiplet at $\lambda = 4$, *HCSM White Paper Series* (2026).
- [6] S. Preschutti, HCSM-05: D_4 Representation Theory of the 14-Mode Multiplet, *HCSM White Paper Series* (2026).
- [7] S. Preschutti, HCSM-13: The Processor T and the Fermion/Boson-Class Correspondence, *HCSM White Paper Series* (2026).
- [8] S. Preschutti, HCSM-14a: The Kernel-Diagonal Theorem, *HCSM White Paper Series* (2026).

- [9] S. Preschutti, HCSM-23 (Eighth Revision): The Midpoint Sorting Theorem, *HCSM White Paper Series* (2026).
- [10] S. Preschutti, HCSM-61: Maxwell and Lorentz in the Classical Limit, *HCSM White Paper Series* (2026).
- [11] S. Preschutti, HCSM-62: Newtonian Gravity as the Classical Limit, *HCSM White Paper Series* (2026).
- [12] S. Preschutti, HCSM-65: The Dirac Equation from the Hodge-Dirac Operator, *HCSM White Paper Series* (2026).
- [13] S. Preschutti, HCSM-69: Quantum Chiral Anomalies on the Finite Lattice, *HCSM White Paper Series* (2026).
- [14] S. Preschutti, HCSM-70: Quantum Gauge Anomalies, *HCSM White Paper Series* (2026).
- [15] S. Preschutti, HCSM-71: Gravitational Anomalies on the Finite Lattice, *HCSM White Paper Series* (2026).
- [16] S. Preschutti, HCSM-72: Gribov Copies on the Finite Torus, *HCSM White Paper Series* (2026).
- [17] S. Preschutti, HCSM-73: Asymptotic Freedom from the Substrate, *HCSM White Paper Series* (2026).

A Numerical Verification

The companion script `HCSM-74_verification.py` constructs the following objects from the substrate and verifies:

1. The 14-mode multiplet at $\lambda = 4$ (from HCSM-04).
2. The Hodge-Dirac operator D and its kernel.
3. The D_4 isotypic decomposition $V_{14} = 3A_1 \oplus A_2 \oplus 2B_1 \oplus 2B_2 \oplus 3E$.
4. The processor $T = I - P_C$ with $C = \{\text{dark}, W, Z, H\}$.
5. The gauge algebra $\mathfrak{g}_{\text{SM}} = \mathfrak{su}(3)_c \oplus \mathfrak{su}(2)_L \oplus \mathfrak{u}(1)^5$.
6. The Wilson loop $W(C)$ on the finite torus.
7. The strong-coupling character expansion coefficients.
8. The string tension $\sigma = 7\Lambda_{\text{QCD}}^2/5$.
9. The mass gap $m_{0++} = 2 \sin(\pi/8) \cdot \lambda_L$.
10. The finite-volume scaling of λ_L .
11. The $\mathbb{Z}_3$ center from the A_1 block.
12. The Polyakov loop $P(x)$.
13. The deconfinement temperature $T_c = 5\Lambda_{\text{QCD}}/7$.

14. The Gribov horizon $R_{\text{crit}} = 14$.
15. The class-partition residual $A = 1/3$.
16. The scale matching factor $\kappa = \Lambda_{\chi}^{\text{phys}}/\Lambda_{\text{QCD}}$.

The script verifies:

1. The area law: $\langle W(C) \rangle \sim \exp(-\sigma \cdot A(C))$.
2. The strong-coupling character expansion exponentiation.
3. The linear potential: $V(r) = \sigma r + \text{const.}$
4. The mass gap: $m_{0++} = 2 \sin(\pi/8) \cdot \lambda_L$.
5. The finite-volume scaling: $m_{0++} \propto L^{-3}$.
6. The Tensor Product protection of the mass gap.
7. The center symmetry: $\mathbb{Z}_3$ from the A_1 block.
8. The Polyakov loop: $\langle P \rangle = 0$ in the confined phase.
9. The deconfinement temperature: $T_c = 5\Lambda_{\text{QCD}}/7$.
10. The Gribov-Confinement Connection: $R_{\text{crit}} = 14$.
11. The Anomaly-Confinement Connection: $A = 1/3$.
12. The scale matching: $\sigma^{\text{phys}} \approx (237 \text{ MeV})^2$, $T_c^{\text{phys}} \approx 143 \text{ MeV}$.

Running `12-VERIFICATION/run_all.py` reproduces every numerical claim in the HCSM paper stack.