

HCSM-00 (Eighth Revision, Corrected)

Foundations, Notation, and Mapping

The Hodge Complex Standard Model:
Substrate, Hodge Complex, Spectral Midpoint, Processor,
Output Structures, Emergent Physics, and Cosmology

Stanley Preschutti

Entropy Research Institute / Information Physics Institute

ORCID: 0009-0004-5445-1744

scstamp@yahoo.com

September 2026

Abstract

The Hodge Complex Standard Model (HCSM) derives the gauge structure, matter content, particle spectrum, mixing matrices, spacetime dimension, graviton, Newton constant, and vacuum stability coefficient of the Standard Model from a single discrete substrate: the periodic 8×8 torus $\Lambda = \mathbb{Z}_8 \times \mathbb{Z}_8$. The substrate is selected by the unified substrate selection theorem (Section 2.6), and the emergent spacetime dimension is fixed by the topology of the Hodge-Dirac kernel. The Hodge complex of differential forms carries the operator $D = d + d^*$ with $D^2 = \Delta$ exactly on each form degree. The 14 modes at the self-paired eigenvalue $\lambda = 4$ are identified with one Standard Model generation.

This paper establishes the framework's foundational objects and states the notation. All seven rank blockers of the framework's derivation program are closed. The framework has a single dimensionful anchor (the Planck mass M_P) and derives every other quantity from the substrate at $d = 4$ and $L = 8$.

The paper establishes the following theorems. The 5-point Laplacian is the unique D_4 -invariant nearest-neighbour positive-semidefinite operator with kernel equal to the constants. The Hodge-Dirac operator satisfies $D^2|_{\Omega^k} = \Delta \cdot \text{id}$ on each form degree, with smallest nonzero eigenvalue $2 \sin(\pi/L)$ on Λ_L . The Lorentzian metric on $\ker D$ is the unique D_4 -invariant symmetric form that vanishes on pure-degree forms and has signature $(3, 1)$. The Kernel-Diagonal Theorem yields the EFT coefficient $c = 5/9$, the mass-map coefficient $(d-1)/2 = 3/2$, and the kernel dimension $10 = (d-1)^2 + 1$ as three traces of one projector. The Λ -promotion mechanism is a pure projection onto the timelike direction of $\ker D$, and the refined dark energy density matches observation to 0.010%.

The charge sector is a rigidity-class result: five structural constraints reduce the charge assignment from $5^{14} \approx 6 \times 10^9$ configurations to exactly two, both with total charge $-1e$. The residual binary choice is resolved by the Midpoint Sorting Theorem of HCSM-23 Rev. 7.

The present revision incorporates two structural sharpenings. First, the identification of the A_1 isotypic block with physical color is now a theorem (HCSM-06, Theorem 5.4), derived from a dimension-and-invariance argument; the identification of the B_1 block with weak isospin is now a labeling convention (HCSM-06, Theorem 5.10), forced by the outer automorphism of D_4 . Second, the Standard Model content structure and the hypercharge values are now derived (HCSM-07, Theorems 8.1 and 8.3) from the 14-mode charges, the D_4 isotypic multiplicities, the Hodge-lift chirality structure, and the anomaly cancellation conditions.

Contents

1	Introduction	3
1.1	What HCSM is	3
1.2	The single dimensionful anchor	3
1.3	Architectural levels	4
1.4	Continuum limit and emergent Lorentz invariance	4
2	Level 0: The Substrate	4
2.1	The periodic $L \times L$ torus	4
2.2	The 5-point Laplacian is forced	4
2.3	The framework relation $N = d \cdot 2^d$	5
2.4	Two derivations of $L = 8$	5
2.5	The 14-mode count	6
2.6	Unified substrate selection	6
2.6.1	The two framework identities	6
2.6.2	The selection theorem	6
2.6.3	The four prior derivations as corollaries	7
2.6.4	What this changes	7
3	Level 1: The Hodge Complex	7
3.1	The three form degrees	7
3.2	The exterior derivative and the codifferential	7
3.3	The Hodge-Dirac operator	8
3.4	The $\mathbb{Z}_2$ grading and the supercharge structure	8
3.5	The kernel and the emergent dimension	9
3.6	The spectral gap and absence of species doubling	9
4	Level 2: The Spectral Midpoint	9
4.1	The self-paired eigenvalue	9
4.2	The D_4 orbit decomposition	9
4.3	The 56-dimensional Hodge lift	9
4.4	The D_4 decomposition of the 56-dimensional sector	10
5	Level 3: The Processor and Its Dynamics	10
5.1	The winding generator	10
5.2	The winding classes and the processor	10
5.3	The Kernel-Diagonal Theorem	10
5.4	The five-layer loop	11
5.5	The zero state	11
6	Level 4: Output Structures	11
6.1	The charge functional	11
6.2	The rigidity class of the charge assignment	11
6.3	The mass map	12
6.4	The Midpoint Sorting Theorem (summary)	12
7	Level 5: Emergent Physics	13
7.1	Gauge algebra from the D_4 commutant	13
7.2	Hypercharge quantization	13
7.3	Flavor: the CKM matrix	13
7.4	Flavor: the PMNS matrix	13
7.5	Higgs sector	13

7.6	Gravity	13
7.7	Absence of a light dark photon	13
7.8	Spin chirality $\mathbb{Z}_2$ grading	14
7.9	Strong CP from orientation $\mathbb{Z}_2$	14
7.10	Proca sector and $\mathfrak{u}(1)_Y$ origin	14
8	Level 6: Cosmology	14
8.1	The kL crossover	14
8.2	Dark energy and the DME entropy-well normalization	14
8.3	Vacuum stability	14
8.4	The Λ -promotion mechanism	15
9	Falsifiable Predictions	15
10	Theorem Inventory	16
11	Reproducibility	17
12	Conclusion	17
A	Web Crawler Metadata	20
A.1	Dublin Core	20
A.2	BibTeX	21

1 Introduction

1.1 What HCSM is

The Hodge Complex Standard Model derives the Standard Model of particle physics from a single discrete substrate: the periodic 8×8 torus $\Lambda = \mathbb{Z}_8 \times \mathbb{Z}_8$. The framework’s central claim is that the Standard Model is not a fundamental description of nature but the residue left when a small, discrete, information-bearing lattice cancels against its own mirror.

Physics is what the machine cannot cancel. The Hodge complex is the machine.

HCSM does not ask what fields exist or what symmetry groups are broken. It asks a single question: what substrate could produce exactly the Standard Model, and nothing else? The answer is a discrete lattice whose mirror cancellation leaves exactly the observed particle content.

1.2 The single dimensionful anchor

The framework has exactly one dimensionful input: the Planck mass M_P . Every other dimensionful quantity is a derived multiple of an appropriate power of M_P . The dimensionless integers $d = 4$ and $N = 64$ are derived from the substrate (Section 2); the substrate size $L = 8$ is derived from the unified selection theorem (Section 2.6); and the emergent Lorentzian signature $(3, 1)$ is derived from the Hodge star on $\ker D$ (Section 3, Theorem 3.10).

At the particle level, the framework has zero empirical anchors. The identification of the 14 modes at $\lambda = 4$ with one Standard Model generation was until recently a disclosed input; it is now a theorem (HCSM-23 Rev. 7). The charge table was until recently a disclosed input up to a binary choice; it is now a theorem whose binary residue is resolved by the same sorting theorem.

As of the present revision, the gauge sector also has zero empirical anchors in the physical-identification step. The identification of the A_1 isotypic block with physical color $\mathfrak{su}(3)_c$ is a theorem (HCSM-06, Theorem 5.4): the A_1 block is the unique three-dimensional D_4 -invariant

subspace of V_{14} , so it is the unique candidate for the physical color triplet. The identification of the B_1 block with physical weak isospin $\mathfrak{su}(2)_L$ is a labeling convention (HCSM-06, Theorem 5.10): the two blocks B_1 and B_2 are exchanged by the outer automorphism of D_4 , and no D_4 -equivariant operator on V_{14} distinguishes them.

The hypercharge sector also has zero empirical anchors in the hypercharge-value step. The hypercharge values $-2/3, +1/6, +1/3, -1/2, +1, 0$ are now derived from the Standard Model content structure and the anomaly cancellation conditions (HCSM-07, Theorem 8.1). The content structure itself is derived from the 14-mode charges, the D_4 isotypic multiplicities, the Hodge-lift chirality structure, and the requirement of anomaly cancellation (HCSM-07, Theorem 8.3).

1.3 Architectural levels

The framework’s architecture is a chain from the substrate to observable physics, organized in six levels.

Level	Name	Content
0	Substrate	8×8 torus, 5-point Laplacian
1	Hodge complex	$\Omega^0 \oplus \Omega^1 \oplus \Omega^2$, $D = d + d^*$
2	Spectral midpoint	$\lambda = 4$, 14 modes, D_4 orbits (2, 8, 4)
3	Processor	$T = I - P_C$, zero state s , five-layer loop
4	Output structures	Charge q_3 , mass map, $N/4$ labels, (S, D) pairs
5	Emergent physics	Gauge, flavor, Higgs, gravity, dark photon
6	Cosmology	kL crossover, dark energy, Λ -promotion

1.4 Continuum limit and emergent Lorentz invariance

The substrate is not a discretization of an underlying continuum. The framework’s claim is the opposite: the continuum emerges from the substrate. The physical scale is set by the mass map and the kL crossover.

The 5-point Laplacian has the isotropic expansion

$$\lambda(n_1, n_2) = \frac{\pi^2}{16}(n_1^2 + n_2^2) + O(n^4). \quad (1)$$

2 Level 0: The Substrate

2.1 The periodic $L \times L$ torus

The substrate is the periodic square lattice

$$\Lambda_L = \mathbb{Z}_L \times \mathbb{Z}_L, \quad N = L^2, \quad (2)$$

consisting of N sites arranged in an $L \times L$ grid with periodic boundary conditions. Site (x_1, x_2) is identified with $(x_1 + L, x_2)$ and with $(x_1, x_2 + L)$; equivalently, the coordinates x_1 and x_2 are each defined modulo L . Topologically, Λ_L is the torus $T^2 = S^1 \times S^1$.

The D_4 point group of the square acts on Λ_L by the eight rotations and reflections of the lattice. Its presentation is $D_4 = \langle r, s \mid r^4 = s^2 = (rs)^2 = e \rangle$.

2.2 The 5-point Laplacian is forced

Theorem 2.1 (Laplacian uniqueness). *Let $\Delta : \mathbb{C}^N \rightarrow \mathbb{C}^N$ be a linear operator on the substrate $\Lambda_L = \mathbb{Z}_L \times \mathbb{Z}_L$ satisfying:*

- (1) D_4 -invariance: $[R, \Delta] = 0$ for every $R \in D_4$.
- (2) Nearest-neighbour locality: $(\Delta f)(x)$ depends only on $f(x)$ and f at the four nearest neighbours of x .
- (3) Positivity and kernel: Δ is positive-semidefinite and $\ker \Delta$ is exactly the set of constant functions.

Then Δ is uniquely determined up to an overall positive scale, and equals the 5-point Laplacian

$$(\Delta f)(x_1, x_2) = 4f(x_1, x_2) - \sum_{|e|=1} f(x + e). \quad (3)$$

Proof. Condition (2) writes $\Delta f(x) = a_0 f(x) + \sum_{\mu} a_{\mu} f(x + \mu)$. The four nearest-neighbour directions $\pm e_1, \pm e_2$ lie in a single D_4 orbit. Hence condition (1) forces $a_{\mu} = a$ for all four directions. Condition (3) forces $\Delta \mathbf{1} = 0$, so $a_0 + 4a = 0$, giving $a_0 = -4a$. Normalizing $a = 1$ gives the 5-point Laplacian. $\square$

Remark 2.2. The 5-point Laplacian annihilates the constants by construction. Its eigenvalue on the plane wave $\phi_{n_1, n_2}(x) = \exp(2\pi i(n_1 x_1 + n_2 x_2)/L)$ is

$$\lambda(n_1, n_2) = 4 - 2\cos(2\pi n_1/L) - 2\cos(2\pi n_2/L), \quad (4)$$

with $n_1, n_2 \in \{0, 1, \dots, L-1\}$. The spectrum is mirror-symmetric under $\lambda \leftrightarrow 8 - \lambda$.

2.3 The framework relation $N = d \cdot 2^d$

The framework uses the relation

$$N = d \cdot 2^d, \quad (5)$$

with N the number of substrate sites and d the emergent spacetime dimension. At $d = 4$ this gives $N = 4 \cdot 2^4 = 64$. Since $\Lambda_L = \mathbb{Z}_L \times \mathbb{Z}_L$ requires $N = L^2$, this gives $L = \sqrt{64} = 8$.

At $d = 4$, $N = 64$ is a perfect square. The integer solutions of $d \cdot 2^d = L^2$ form the infinite family $d \in \{4k^2 : k \geq 1\}$, with $d = 16, 36, 64, \dots$ also giving perfect squares. The physical selection of $d = 4$ comes from the unified selection theorem of Section 2.6.

2.4 Two derivations of $L = 8$

The substrate size $L = 8$ is selected by two independent derivations.

First derivation: equal entropy spacing. On the $L \times L$ torus, the standing-wave modes fall into discrete Shannon entropy classes.

Theorem 2.3 (Equal entropy spacing selects $L = 8$). *Among all integers $4 \leq L \leq 10^4$, the only L for which the $L \times L$ torus has exactly three equally spaced Shannon entropy classes is $L = 8$. At $L = 8$ the three classes are 6.0 bits, 5.5 bits, and 5.0 bits, with uniform spacing 0.5 bits.*

Second derivation: maximal winding cancellation. The winding generator generalizes to any even L as $H_{\text{wind}}(L) = i \sin(\pi M/L)$ on the self-paired mode set of Λ_L .

Theorem 2.4 (Maximal cancellation selects $L = 8$). *Let $\mathcal{G}_L$ be the parity block graph of $H_{\text{wind}}(L)$ on the five nodes $\{A_{\text{odd}}, A_{\text{even}}, B_{\text{odd}}, B_{\text{even}}, AB\}$. The number of block couplings that vanish identically is*

$$\#\text{zeros}(\mathcal{G}_L) = \begin{cases} 4, & L \equiv 2 \pmod{4}, \\ 7, & L \equiv 0 \pmod{4}, L \geq 12, \\ 10, & L \in \{4, 8\}. \end{cases} \quad (6)$$

Among the two maximally-cancelling sizes, only $L = 8$ carries the 14-mode multiplet.

The two derivations are independent: the first uses only the torus Laplacian's entropy classes; the second uses only the winding generator's parity block structure.

2.5 The 14-mode count

Theorem 2.5 (Midpoint multiplicity). *On the $L \times L$ torus with L even, the multiplicity of the self-paired eigenvalue $\lambda = 4$ is*

$$\text{mult}_{\lambda=4}(L) = 2(L - 1). \quad (7)$$

At $L = 8$ this gives 14.

Proof. The self-paired eigenvalue is the fixed point of the mirror symmetry $\lambda \leftrightarrow 8 - \lambda$. Its eigenspace consists of the modes (n_1, n_2) satisfying $\cos(2\pi n_1/L) + \cos(2\pi n_2/L) = 0$. Using the sum-to-product identity $\cos a + \cos b = 2 \cos((a+b)/2) \cos((a-b)/2)$, this becomes $2 \cos(\pi(n_1 + n_2)/L) \cos(\pi(n_1 - n_2)/L) = 0$. The solutions are $n_1 + n_2 \equiv L/2 \pmod{L}$ or $n_1 - n_2 \equiv L/2 \pmod{L}$. Each congruence has L solutions; the two congruences intersect in exactly two points. By inclusion-exclusion, the total is $2L - 2 = 2(L - 1)$. $\square$

2.6 Unified substrate selection

The substrate size $L = 8$ and the emergent dimension $d = 4$ are selected by four independent derivations in the current framework: equal entropy spacing on the torus Laplacian; maximal winding cancellation of the winding generator; Lorentzian signature on $\ker D$; and the Betti numbers of T^2 . This subsection shows that they are corollaries of a single selection theorem.

2.6.1 The two framework identities

Identity 1 (site-count relation). The number of substrate sites N and the emergent dimension d satisfy

$$N = d \cdot 2^d. \quad (8)$$

Identity 2 (mode-content relation). The self-paired mode count of the torus Laplacian at the midpoint eigenvalue equals the total particle content of one Standard Model generation:

$$\text{mult}_{\lambda=4}(L) = |F| + |B|. \quad (9)$$

The right-hand side is derived content: the fermion-sector dimension $|F| = d(d+1)/2$ (HCSM-13, HCSM-19) and the boson-sector dimension $|B| = d$ (HCSM-13, HCSM-19). At $d = 4$ they give $|F| = 10$, $|B| = 4$, and total content 14.

The left-hand side is HCSM-04's multiplicity theorem:

$$\text{mult}_{\lambda=4}(L) = 2(L - 1). \quad (10)$$

2.6.2 The selection theorem

Theorem 2.6 (Unified substrate selection). *The system*

$$L^2 = d \cdot 2^d, \quad 2(L - 1) = d(d + 3)/2, \quad (11)$$

has the unique even-integer solution $L = 8$, $d = 4$, $N = 64$.

Proof. By Identity 1, $L = \sqrt{d \cdot 2^d}$. For L to be an even integer, $d \cdot 2^d$ must be a perfect square. Since 2^d is a perfect square iff d is even, write $d = 2m$. Then $d \cdot 2^d = 2m \cdot 2^{2m}$, which is a perfect square iff $2m$ is a perfect square, i.e. $m = 2k^2$ for integer $k \geq 1$. Hence

$$d = 4k^2, \quad L = 2^{2k^2+1} \cdot k. \quad (12)$$

Substituting into Identity 2 gives

$$2^{2k^2+1} \cdot k - 1 = k^2(4k^2 + 3). \quad (13)$$

Case $k = 1$: Left side $2^3 \cdot 1 - 1 = 7$. Right side $1 \cdot 7 = 7$. Satisfied.

Case $k = 2$: Left side $2^9 \cdot 2 - 1 = 1023$. Right side $4 \cdot 19 = 76$. Left exceeds right.

Case $k \geq 3$: The left side grows as $k \cdot 2^{2k^2+1}$, super-exponentially in k^2 ; the right side grows as $4k^4 + 3k^2$, polynomially. The left strictly dominates for all $k \geq 2$.

Hence $k = 1$ is the unique solution, giving $d = 4$, $L = 8$, $N = 64$. $\square$

Corollary 2.7 (Uniqueness). *The pair $(L, d) = (8, 4)$ is the unique even-integer pair satisfying the two framework identities simultaneously.*

2.6.3 The four prior derivations as corollaries

Corollary 2.8 (Entropy spacing). *At $L = 8$, the substrate $\Lambda = \mathbb{Z}_8 \times \mathbb{Z}_8$ has exactly three equally spaced Shannon entropy classes: 6.0, 5.5, and 5.0 bits.*

Corollary 2.9 (Maximal winding cancellation). *At $L = 8$, the winding generator $H_{\text{wind}} = i \sin(\pi M/8)$ has cancellation depth 10 on the five-node parity block graph, with kernel dimension $d(d+1)/2 = 10$.*

Corollary 2.10 (Lorentzian signature). *At $L = 8$, the Hodge star on $\ker D = H^0 + H^1 + H^2$ has eigenvalues $\{+1, -1, +i, -i\}$, giving a metric of signature $(3, 1)$ on $\ker D$.*

Corollary 2.11 (Betti numbers). *At $L = 8$, the kernel dimension is $d = \dim \ker D = b_0 + b_1 + b_2 = 1 + 2 + 1 = 4$.*

2.6.4 What this changes

Item	Before	After
Derivations of $L = 8$	2 independent	1 theorem, 1 corollary
Derivations of $d = 4$	2 independent	1 theorem, 2 corollaries
Status of $L = 8$	4 consistent derivations	1 theorem, 3 consistency checks
Status of $d = 4$	4 consistent derivations	1 theorem, 3 consistency checks

After the unified selection, the framework has exactly one open item outside the particle sector: the continuum limit.

3 Level 1: The Hodge Complex

3.1 The three form degrees

On the 2-torus Λ_L , the Hodge complex of differential forms has three graded pieces:

$$\Omega^0 = \mathbb{R}^N, \quad \Omega^1 = \mathbb{R}^{2N}, \quad \Omega^2 = \mathbb{R}^N, \quad (14)$$

with total dimension $N + 2N + N = 4N$. At $N = 64$ this gives 256.

3.2 The exterior derivative and the codifferential

Let $e_1 = (1, 0)$ and $e_2 = (0, 1)$ be the two oriented lattice directions. The exterior derivative acts on 0-forms by

$$(d_0 f)(x, \mu) = f(x + e_\mu) - f(x), \quad \mu \in \{1, 2\}, \quad (15)$$

and on 1-forms by

$$(d_1 \omega)(x) = \omega_1(x) + \omega_2(x + e_1) - \omega_1(x + e_2) - \omega_2(x). \quad (16)$$

These satisfy $d_1 d_0 = 0$ exactly.

The codifferential $d^* = d^\dagger$ is defined as the adjoint with respect to the standard Euclidean inner product. Taking $d_0^* = d_0^T$ and $d_1^* = d_1^T$ as matrices gives the explicit formulas used throughout the stack.

3.3 The Hodge-Dirac operator

Definition 3.1 (Hodge-Dirac operator). *The Hodge-Dirac operator is $D = d + d^*$, acting on the full Hodge complex $\Omega^0 \oplus \Omega^1 \oplus \Omega^2$. In block form,*

$$D = \begin{pmatrix} 0 & d_0^* & 0 \\ d_0 & 0 & d_1^* \\ 0 & d_1 & 0 \end{pmatrix}, \quad (17)$$

a symmetric 256×256 real operator.

Theorem 3.2 (Hodge-Dirac square). *On the discrete torus Λ , the identity $D^2|_{\Omega^k} = \Delta \cdot \text{id}_{\Omega^k}$ holds exactly on each form degree $k = 0, 1, 2$.*

Proof. On Ω^0 , $D^2 = d_0^* d_0 = \Delta$. On Ω^1 , $D^2 = d_0 d_0^* + d_1^* d_1 = \Delta$ by direct component expansion. On Ω^2 , $D^2 = d_1 d_1^* = \Delta$. Cross terms are removed by nilpotency $d_1 d_0 = 0$. All three identities are verified at machine precision. $\square$

Remark 3.3 (No species doubling). *The Hodge-Dirac operator on the discrete torus has no species doubling: its square equals the scalar Laplacian on each form degree. The low-energy spectrum is bounded away from zero by $2 \sin(\pi/L)$ except for the four zero modes.*

3.4 The $\mathbb{Z}_2$ grading and the supercharge structure

Definition 3.4 (Parity operator). *The parity operator $(-1)^F$ on the Hodge complex is $+1$ on $\Omega^0 \oplus \Omega^2$ and -1 on Ω^1 . It $\mathbb{Z}_2$ -grades the complex into a bosonic sector $\Omega^0 \oplus \Omega^2$ and a fermionic sector Ω^1 .*

Proposition 3.5. $D(-1)^F = -(-1)^F D$.

Theorem 3.6 (Supercharge structure). *In the $\mathbb{Z}_2$ grading, the Hodge-Dirac operator decomposes as*

$$D = Q + Q^\dagger, \quad (18)$$

where Q is the off-diagonal block of D mapping the bosonic sector to the fermionic sector, and $Q^\dagger$ is its adjoint. The supercharges satisfy

$$Q^2 = (Q^\dagger)^2 = 0, \quad \{Q, Q^\dagger\} = D^2 = \Delta, \quad (19)$$

and the Hamiltonian $H = D^2/2 = \Delta/2$ satisfies

$$\{Q, Q^\dagger\} = 2H. \quad (20)$$

This is the standard $N = 1$ supersymmetric quantum mechanics algebra.

Proof. In the $\mathbb{Z}_2$ grading, D is off-diagonal:

$$D = \begin{pmatrix} 0 & Q^\dagger \\ Q & 0 \end{pmatrix}, \quad (21)$$

with $Q : \Omega^0 \oplus \Omega^2 \rightarrow \Omega^1$ and $Q^\dagger : \Omega^1 \rightarrow \Omega^0 \oplus \Omega^2$. Then

$$D^2 = \begin{pmatrix} Q^\dagger Q & 0 \\ 0 & Q Q^\dagger \end{pmatrix}. \quad (22)$$

By Theorem 3.2, $D^2 = \Delta$ on each form degree, so $Q^\dagger Q = \Delta|_{\text{bosonic}}$ and $QQ^\dagger = \Delta|_{\text{fermionic}}$. The nilpotency $Q^2 = 0$ follows from the fact that Q raises form degree by 1: Q maps $\Omega^0 \rightarrow \Omega^1$ and $\Omega^2 \rightarrow 0$. Similarly $(Q^\dagger)^2 = 0$. $\square$

Corollary 3.7 (Boson-fermion pairing). *Every bosonic mode at energy λ_n has a fermionic partner at the same energy, unless the mode is annihilated by Q or $Q^\dagger$. In particular, at $\lambda = 4$: 28 bosons + 28 fermions.*

3.5 The kernel and the emergent dimension

Theorem 3.8 (Kernel dimension). *The kernel of the Hodge-Dirac operator is isomorphic to the de Rham cohomology of the torus: $\ker D \cong \bigoplus_{k=0}^2 H^k(T^2; \mathbb{R})$, with*

$$\dim \ker D = b_0 + b_1 + b_2 = 1 + 2 + 1 = 4. \quad (23)$$

Corollary 3.9 (Emergent dimension). $d = \dim \ker D = 4$.

Theorem 3.10 (Lorentzian signature). *The Hodge star $\star$ restricted to $\ker D$ has eigenvalues $\{+1, -1, +i, -i\}$: the ± 1 pair from the $(H^0 + H^2)$ sector, the $\pm i$ pair from the H^1 sector. The intersection form $Q(\alpha, \beta) = \int_\Lambda \alpha \wedge \beta$ on the even-degree sector and the Hodge inner product $\langle \alpha, \beta \rangle = \int_\Lambda \alpha \wedge \star \beta$ on the odd-degree sector combine into a metric*

$$g = Q|_{H^0 + H^2} + \langle \cdot, \cdot \rangle|_{H^1} \quad (24)$$

of signature $(3, 1)$ on $\ker D$.

3.6 The spectral gap and absence of species doubling

Theorem 3.11 (Smallest nonzero eigenvalue of D). *On $\Lambda_L = \mathbb{Z}_L \times \mathbb{Z}_L$ with L even, the smallest nonzero eigenvalue of the Hodge-Dirac operator is exactly $\mu_{\min}(L) = 2 \sin(\pi/L)$.*

Corollary 3.12. *For every finite even L , the Hodge-Dirac operator has exactly four zero eigenvalues and the next eigenvalue is bounded away from zero by $2 \sin(\pi/L)$.*

4 Level 2: The Spectral Midpoint

4.1 The self-paired eigenvalue

The mirror symmetry $\lambda \leftrightarrow 8 - \lambda$ of the scalar Laplacian spectrum has a unique fixed point at $\lambda = 4$. The eigenspace at $\lambda = 4$ has multiplicity $2(L - 1) = 14$ at $L = 8$. The 14 modes are

$$\mathcal{O}_0 = \{(0, 4), (4, 0)\}, \quad (25)$$

$$\mathcal{O}_1 = \{(1, 3), (1, 5), (3, 1), (3, 7), (5, 1), (5, 7), (7, 3), (7, 5)\}, \quad (26)$$

$$\mathcal{O}_2 = \{(2, 2), (2, 6), (6, 2), (6, 6)\}. \quad (27)$$

4.2 The D_4 orbit decomposition

Proposition 4.1 (Orbit stabilizers). *Under the D_4 point group, the 14 modes decompose into the three orbits above of sizes $(2, 8, 4)$. The stabilizer orders of the three orbits are 4, 1, 2 respectively, satisfying the orbit-stabilizer identity $|\mathcal{O}_k| = |D_4|/|\text{Stab}(\mathcal{O}_k)| = 8/|\text{Stab}(\mathcal{O}_k)|$.*

4.3 The 56-dimensional Hodge lift

Theorem 4.2 (Fourteen-fold lift). *The 14-fold degeneracy of the scalar Laplacian at $\lambda = 4$ lifts to the Hodge complex as $14\Omega^0 + 28\Omega^1 + 14\Omega^2 = 56$.*

4.4 The D_4 decomposition of the 56-dimensional sector

Theorem 4.3 (D_4 decomposition). *As D_4 representations,*

$$\Omega^0|_{\lambda=4} = 3A_1 \oplus A_2 \oplus 2B_1 \oplus 2B_2 \oplus 3E, \quad (28)$$

$$\Omega^1|_{\lambda=4} = \Omega^0|_{\lambda=4} \otimes E = 3A_1 \oplus 3A_2 \oplus 3B_1 \oplus 3B_2 \oplus 8E, \quad (29)$$

$$\Omega^2|_{\lambda=4} = 3A_1 \oplus A_2 \oplus 2B_1 \oplus 2B_2 \oplus 3E. \quad (30)$$

The full 56-dimensional sector decomposes as $56 = 9A_1 \oplus 5A_2 \oplus 7B_1 \oplus 7B_2 \oplus 14E$.

5 Level 3: The Processor and Its Dynamics

5.1 The winding generator

Definition 5.1 (Winding matrix and winding generator). *Let $n_1, n_2 \in \mathbb{R}^{14}$ be the coordinate vectors of the 14 modes. The winding matrix is the rank-2 antisymmetric bilinear $M_{ij} = n_1^{(i)} n_2^{(j)} - n_1^{(j)} n_2^{(i)}$. The winding generator is $H_{\text{wind}} = i \sin(\pi M/8)$, with $\sin$ applied elementwise.*

Theorem 5.2 (Characteristic polynomial). *The characteristic polynomial of H_{wind} is*

$$p(\lambda) = \det(\lambda \text{id}_{14} - H_{\text{wind}}) = \lambda^{10}(\lambda^2 - 8)(\lambda^2 - 32). \quad (31)$$

Its spectrum is $\{-4\sqrt{2}, -2\sqrt{2}, 0^{(10)}, +2\sqrt{2}, +4\sqrt{2}\}$.

5.2 The winding classes and the processor

Definition 5.3 (Winding class partition). *The 14 modes partition into three winding classes:*

$$A = \{\nu, e, t\}, |A| = 3; \quad B = \{u, d, s, \mu, c, \tau, b\}, |B| = 7; \quad C = \{\text{dark}, W, Z, H\}, |C| = 4. \quad (32)$$

The partition is the joint eigenspace decomposition of the n_1 -parity D and the processor T .

Definition 5.4 (Processor). *The processor T is the rank-10 orthogonal projection $T = I - P_C$ with P_C the diagonal projector onto winding class C .*

Proposition 5.5 (Fermion/boson split). *$T = I - P_C = P_A + P_B$, with $\text{im}(T) = F = A \cup B$ and $\ker(T) = C$. The fermion and boson sector dimensions are $|F| = d(d+1)/2 = 10$ and $|B| = d = 4$ (HCSM-13, HCSM-19). The internal ratios are $(|F| - |B|)/|B| = (d-1)/2 = 3/2$ and $|F|/|B| = (d+1)/2 = 5/2$.*

5.3 The Kernel-Diagonal Theorem

Theorem 5.6 (Kernel-Diagonal). *Let $P_{\ker}$ be the orthogonal projector onto $\ker(H_{\text{wind}})$. The diagonal of $P_{\ker}$ in the standard mode basis takes exactly two values:*

$$\text{diag}(P_{\ker})_i = \begin{cases} 2/d = 1/2, & i \in \mathcal{O}_0, \\ (d-1)/d = 3/4, & i \in \mathcal{O}_1 \cup \mathcal{O}_2. \end{cases} \quad (33)$$

Corollary 5.7 (Three coefficients from one diagonal). *With $P_{\ker}$ as above,*

$$c = (d+1)/\text{Tr}(P_{\ker}|_{\mathcal{O}_1 \cup \mathcal{O}_2}) = 5/9, \quad (34)$$

$$\text{diag}(P_{\ker})|_{\mathcal{O}_1 \cup \mathcal{O}_2} / \text{diag}(P_{\ker})|_{\mathcal{O}_0} = (d-1)/2 = 3/2, \quad (35)$$

$$\dim \ker(H_{\text{wind}}) = 1 + (d-1)^2 = 10. \quad (36)$$

5.4 The five-layer loop

The five-layer loop acts diagonally on the 14-mode multiplet: $M_{\text{loop}} = T \cdot L_{\text{EFT}} \cdot L_{\text{Core}} \cdot L_{\text{DM}} \cdot L_{\text{FDLM}}$.

Theorem 5.8 (Loop-pass duration). *The loop-pass duration is selected by the deterministic Landauer bound: $T_{\text{pass}} = T_{\text{DM}} = 1/(2\gamma) = 0.7114000035169658\dots$*

Theorem 5.9 (Exact ratio). $|M_A|/|M_B| = 3 = d - 1$.

5.5 The zero state

The zero state s is the unique nullspace vector of a 14×19 measurement matrix M of vacuum-native observables. The matrix has rank 13, nullity 1.

Theorem 5.10 (Zero-state structure). *The zero state satisfies $s(rs(m)) = -s(m)$ for all modes m , where rs is the diagonal reflection $(n_1, n_2) \mapsto (n_2, n_1)$. The two rs -fixed modes $W = (2, 2)$ and $t = (6, 6)$ have $s(W) = s(t) = 0$ exactly. The six independent amplitudes satisfy $a_3 = 2a_4$ and $a_5 = (2 - \sqrt{2})a_4$. Weight fractions: 3.3% on $\mathcal{O}_0$, 86.5% on $\mathcal{O}_1$, 10.2% on $\mathcal{O}_2$.*

Remark 5.11 (Zero state as sort key). *The zero state is not only the unique nullvector of the measurement matrix; it is also the sort key of the charge assignment and one of the three native operators used in the Midpoint Sorting Theorem (HCSM-23 Rev. 7).*

6 Level 4: Output Structures

6.1 The charge functional

The electric charge in units of $e/3$ is given by the functional

$$q_3 = F(n_1, \sigma), \quad \sigma = \text{sign}(s), \quad (37)$$

where n_1 is the mode's first momentum index and s is the zero-state amplitude.

Theorem 6.1 (Charge bijection). *The 14 modes have 14 pairwise distinct (n_1, σ) tuples. The map F is well-defined on the realized pairs and is a bijection onto the five Standard Model charge eigenvalues $\{-3, -1, 0, +2, +3\}$.*

Theorem 6.2 (σ -line sums). *The charge sums over the three σ -lines are*

$$\sum_{\sigma=-1} q_3 = 0, \quad \sum_{\sigma=0} q_3 = +5, \quad \sum_{\sigma=+1} q_3 = -8. \quad (38)$$

The total vacuum charge is $\sum_i q_{3i} = -3 = -1e$.

6.2 The rigidity class of the charge assignment

Theorem 6.3 (Charge rigidity class). *Among the 5^{14} assignments of charge values $\{-3, -1, 0, +2, +3\}$ to the 14 modes at $\lambda = 4$, exactly two assignments simultaneously satisfy the following five structural constraints:*

(C1) *SM multiset. The multiset of charges is $\{-3, -1, 3, 0, 4, +2, 3, +3, 1\}$.*

(C2) *σ -line sums. The charge sums over the three zero-state sign lines are 0 , $d + 1 = 5$, and $-2d = -8$.*

(C3) *F -pair sum rules. Within each $|\Delta n|$ shell, the pair charge sum is rank- d (for $|\Delta n| = 4$) or $(d + 1) - d \cdot \text{rank}$ (for $|\Delta n| = 2$).*

(C4) $n_1 + n_2 = 4$ line. The five modes on the line $n_1 + n_2 = 4$ carry the five charge values exactly once each.

(C5) Processor bookkeeping. The image-of- T charge sum is -2 and the kernel-of- T charge sum is $+1$.

Proof. Exhaustive enumeration. The F -pair sum rules (C3) reduce the assignment space to 192 configurations. Constraints (C1), (C2), and (C4) reduce this to 3; constraint (C5) reduces this to exactly 2. $\square$

Theorem 6.4 (Charge assignment solutions). *The two rigid solutions are*

$$Q_3^{(1)} = (-3, 0, -1, -1, 2, -3, 0, 2, -3, -1, 3, 0, 0, 2), \quad (39)$$

$$Q_3^{(2)} = (0, -3, 2, -1, -1, -3, 0, 2, -3, -1, 3, 0, 0, 2). \quad (40)$$

The two solutions differ only by the linked swap on the pair $\{\nu, e\}$ and the pair $\{u, s\}$.

Theorem 6.5 (Total charge rigidity). *Both solutions have total charge $\sum_i q_{3i} = -3 = -1e$. Consequently, every assignment satisfying the five structural constraints has total charge $-1e$:*

$$\sum_i q_{3i} = -(d-1) = -1e. \quad (41)$$

The total vacuum charge is a rigidity-class invariant.

Corollary 6.6 (Empirical residue of the charge sector). *The charge assignment is determined up to exactly one binary choice: which member of the pair $\{\nu, e\}$ carries charge 0 and which carries -3 . The Standard Model content resolves this choice: the electron e carries $-1e$ and the neutrino ν carries 0. This binary choice is now derived, not disclosed: HCSM-23 Rev. 7 Corollary 4.4 shows that the within-pair ordering rule of the Midpoint Sorting Theorem forces $\nu = (0, 4)$ and $e = (4, 0)$.*

Remark 6.7 (Physical reading: what the machine must not cancel). *The rigidity theorem states that the machine cannot cancel its residual charge. Any attempt to modify the charge assignment away from the two-element family violates one of the five structural constraints. The total charge $-1e$ is not a choice made by the machine; it is the unique value permitted by the machine's own structure.*

6.3 The mass map

The mass map is $m_i = A_{\text{mass}} \exp(-B_{\text{mass}} N/4_i)$, with

$$A_{\text{mass}} = (2d-1)N/2 = 224\text{GeV}, \quad B_{\text{mass}} = (d-1)kL^2/(2N^2) = 0.5411967\dots \quad (42)$$

The map reproduces the twelve charged fermion masses and the W , Z , H , and t masses at RMS 0.0449 dex across thirteen orders of magnitude in mass. The coefficient $(d-1)/2$ in B_{mass} is derived in HCSM-19 by two independent routes (the processor F/B split and the DNLS Hartree orbit trace). The mass map itself is derived in HCSM-33.

6.4 The Midpoint Sorting Theorem (summary)

The identification of the 14 modes with one Standard Model generation is now a theorem (HCSM-23 Rev. 7). The sorting uses the T_{half} involution $(n_1, n_2) \mapsto (n_1 + 4, n_2 + 4)$, the pair index $k = \min(n_1)$, the parity-block node from the sign of the signed coordinate v of the smaller- n_1 member, the (S, D) table of HCSM-23 Rev. 6, and the within-pair ordering rule.

7 Level 5: Emergent Physics

7.1 Gauge algebra from the D_4 commutant

The gauge algebra $\mathfrak{su}(3) \oplus \mathfrak{su}(2) \oplus \mathfrak{u}(1)^5$ is derived as a Lie subalgebra of the D_4 commutant on the 14-mode multiplet (HCSM-06). The A_1 isotypic block is uniquely identified with physical color by a dimension-and-invariance theorem (HCSM-06, Theorem 5.4). The B_1 block is identified with physical weak isospin up to the outer-automorphism labeling convention (HCSM-06, Theorem 5.10).

7.2 Hypercharge quantization

The hypercharge functional is a specific element of the 15-dimensional sector-coarse space (HCSM-06, Theorem 6.3; HCSM-07). The A_1 coefficients are $(0, -1/6, -1/2)$ and the B_1 coefficients are $(0, +1/2, +1/2)$, uniquely determined by the Standard Model content and the sector assignment (HCSM-07, Theorems 5.1 and 6.1). The hypercharge values are derived from the content structure and the anomaly cancellation conditions (HCSM-07, Theorem 8.1). The content structure is derived from the 14-mode charges, the D_4 isotypic multiplicities, and the Hodge-lift chirality structure (HCSM-07, Theorem 8.3). All values lie in $(1/6)\mathbb{Z}$.

7.3 Flavor: the CKM matrix

The Cabibbo angle satisfies $\sin \theta_{12} = \pi/14 = 0.2243995\dots$, matching observation to 0.044% (HCSM-34, Theorem 1). The winding amplitude $a = 3/5$ is topological: $a = (|F| - |B|)/|F| = 6/10$. The Wolfenstein parameters are $\lambda = \pi/14$, $A = (d/(d+1))(1 + \delta_{\text{edge}})$, and $\sqrt{\bar{\rho}^2 + \bar{\eta}^2} = (d+1)/(d(d-1)) = 5/12$ (HCSM-34, Theorem 3). The full CKM hierarchy is derived in HCSM-34, Theorems 5 and 9.

7.4 Flavor: the PMNS matrix

The neutrino $N/4$ spectrum is $(60, 60, 58)$, derived from the four spin structures of the torus with Dirac indices $\pm 1/4$ (HCSM-36, Theorem 4.1). The PMNS magnitudes match at RMS 0.035 (HCSM-36, Theorem 6.1). The mixing angles are $\theta_{12} = 34.2^\circ$, $\theta_{23} = 47.9^\circ$, $\theta_{13} = 8.5^\circ$; the Dirac CP phase is $\delta_{CP} \approx 197^\circ$. The generation phase $\phi_{\text{gen}} = (3/5)\pi$ is exact (HCSM-36, Section 5).

7.5 Higgs sector

The top-Yukawa coefficient is $y_t^2 = 1 - (4/3)\delta_{\text{edge}}$ with $4/3 = |\mathcal{O}_2|/(d-1)$ (HCSM-40). The top mass is $m_t = 172.688398$ GeV (PDG 172.690 ± 0.30 GeV). The tree-level quartic coupling is $\lambda_{\text{tree}} = \ln(kL)/[(d-1)\pi^2]$, and the electroweak vacuum expectation value is $v_{\text{EW}} = M_P \exp(-kL_{\text{phys}})$.

7.6 Gravity

The graviton is the self-paired $B_1 \oplus B_2$ curvature mode of $\text{Sym}^2(\Omega^1|_{\lambda=4})$, consisting of $54B_1 \oplus 54B_2 = 108$ modes (HCSM-45, HCSM-46). It is exactly massless at all orders by D_4 Schur orthogonality (HCSM-46). The Newton constant is $G_N = (d+1)^2/(8\pi M_P^2) = 25/(8\pi M_P^2)$ (HCSM-48).

7.7 Absence of a light dark photon

Theorem 7.1 (Absence of a light dark photon). *The kinetic mixing parameter satisfies $\epsilon \equiv 0$ exactly over all mass scales.*

Proof. The HCSM gauge algebra $\mathfrak{su}(3)_c \oplus \mathfrak{su}(2)_L \oplus \mathfrak{u}(1)^5$ contains no hidden Abelian factor coupling to electrically charged matter. The hidden sector $5 + 6 + 31$ is $U(1)_Y$ -neutral (HCSM-41). $\square$

7.8 Spin chirality $\mathbb{Z}_2$ grading

The 56-dimensional Hodge lift at $\lambda = 4$ carries three commuting $\mathbb{Z}_2$ operators: $\gamma_5 = P_{\Omega^0} - P_{\Omega^1} + P_{\Omega^2}$ (Hodge form-parity), χ_{spin} (spin-structure parity), and $F : (n_1, n_2) \mapsto (n_2, n_1)$ (orientation flip). All three square to I and pairwise commute, generating $\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2$ (HCSM-42).

7.9 Strong CP from orientation $\mathbb{Z}_2$

Theorem 7.2 (Strong CP). $\theta_{\text{QCD}} = \theta_{\text{eff}} = 0$ *exactly*.

Proof. On the 14-mode multiplet at $\lambda = 4$, $[F, \Delta] = 0$ and $\{F, H_{\text{wind}}\} = 0$ exactly. The winding matrix M is antisymmetric under F ; the odd function $\sin(\pi M/8)$ preserves the antisymmetry. The physical identification $H_{\text{wind}} \leftrightarrow F_{\mu\nu} \tilde{F}^{\mu\nu}$ gives $\theta_{\text{eff}} = 0$ exactly (HCSM-43). $\square$

7.10 Proca sector and $\mathfrak{u}(1)_Y$ origin

The $\mathfrak{u}(1)_Y$ gauge symmetry is generated by the hypercharge functional $\hat{Y}$ on $\Omega^0|_{\lambda=4}$, with $[\hat{Y}, T] = 0$, $[\hat{Y}, F] = 0$, and eigenvalues in $(1/6)\mathbb{Z}$. The Proca Lagrangian is

$$\mathcal{L}_{\text{Proca}} = -\frac{1}{4}W_{\mu\nu}^a W^{a\mu\nu} - \frac{1}{4}B_{\mu\nu}B^{\mu\nu} + \frac{1}{2}m_W^2 W_\mu^+ W^{-\mu} + \frac{1}{2}m_Z^2 Z_\mu Z^\mu, \quad (43)$$

with $m_W = gv_{\text{EW}}/2$ and $m_Z = \sqrt{g^2 + g'^2}v_{\text{EW}}/2$. The mass ratio $m_W/m_Z = \cos\theta_W = 0.8769$ matches PDG 0.8814 at 0.5%. Tree-level $\rho = 1$ exactly (HCSM-44).

8 Level 6: Cosmology

8.1 The kL crossover

The bare warp factor is exact: $kL_{\text{bare}} = (d-1)N/(d+1) = 3 \cdot 64/5 = 192/5 = 38.4$. The physical warp factor is $kL_{\text{phys}} = 38.442527$, matching $\ln(M_P/v_{\text{EW}})$ to 0.004%. The correction is $kL_{\text{phys}} - kL_{\text{bare}} = ((|D_4| - 1)/2)\delta_{\text{edge}} = (7/2)\delta_{\text{edge}}$ (HCSM-51).

8.2 Dark energy and the DME entropy-well normalization

The bare dark energy density is $\rho_{DE}^{\text{bare}} = (|F|/|B|)m_s^4 = ((d+1)/2)m_s^4 = 2.4809 \times 10^{-47}\text{GeV}^4$ matching observation to 0.76%. The refined formula is $\rho_{DE} = ((d+1)/2)(1+c\delta)m_s^4$ with $c = (d+1)/(d-1)^2 = 5/9$ and $\delta = (1-\gamma)^2/(2\pi)$, giving $\rho_{DE} = 2.5003 \times 10^{-47}\text{GeV}^4$, matching observation to 0.010% (HCSM-52, HCSM-53).

The DME entropy-well normalization is $S_0 = v_{\text{EW}}^2 \cdot (7/40) \cdot (1 + \delta_{\text{edge}}/6) = 10630.722\text{GeV}^2$, matching the implied value at relative residual 2.44×10^{-6} (HCSM-53).

8.3 Vacuum stability

The stability coefficient is $\gamma = c_2/I(2) = 0.702839467990\dots$, with $I(2) = 1 - \frac{1}{2}\text{AGM}(1, \sqrt{2})$, $\text{AGM}(1, \sqrt{2}) = 1/(\sqrt{2}\eta(i)^2)$, and $\eta(i) = \Gamma(1/4)/(2\pi^{3/4})$. The stability theorem states $S''(0)/S_0 = -2\gamma < 0$: the equilibrium $D = 4$ is a strict local maximum of the vacuum entropy (HCSM-54).

8.4 The Λ -promotion mechanism

The residual vacuum energy after mirror cancellation is promoted to the observed cosmological constant by pure projection, not by gravitational coupling.

The five-layer loop $M_{\text{loop}} = T \cdot L_{\text{EFT}} \cdot L_{\text{Core}} \cdot L_{\text{DME}} \cdot L_{\text{FDLM}}$ acts on the 14-mode multiplet. The EFT layer L_{EFT} acts as a soft projector with weight $1 + c\delta$, where $c = 5/9$ is set by the Kernel-Diagonal Theorem. The residual energy projects onto the timelike direction $(v_0 - v_2)/\sqrt{2}$ of $\ker D$, with unit projection weight. The hidden $\mathfrak{u}(1)$ factors contribute zero.

The mechanism is therefore a projection, not a coupling. There is no gravitational suppression factor $1/M_P^2$; the residual vacuum energy is already at the right scale because it is topological. Combined with the absence of a light dark photon, this confirms that no gauge mediator exists between the hidden and visible sectors (HCSM-56.a).

9 Falsifiable Predictions

- (1) **No light dark photon.** The kinetic mixing parameter satisfies $\epsilon \equiv 0$ exactly. Falsifiable by SHiP, FASER2, MATHUSLA, CODEX-b, LDMX, NA64, Belle II, and HL-LHC at $\epsilon > 10^{-12}$.
- (2) **Strong CP.** $\theta_{\text{QCD}} = 0$ exactly. Any non-zero measurement of the neutron electric dipole moment at the level predicted by a generic θ_{QCD} would falsify the framework.
- (3) **ρ parameter.** Tree-level $\rho = m_W^2/(m_Z^2 \cos^2 \theta_W) = 1$ exactly. Any measured deviation of ρ from 1 at tree level would falsify the framework.
- (4) **Fermion content.** One Standard Model generation contains exactly 14 particles. A fourth generation would falsify the framework.
- (5) **Kernel structure.** The winding kernel has dimension 10 on Λ_8 and the kernel-diagonal values are $\{1/2, 3/4\}$. Any modification of the spectrum of H_{wind} that changes these values would falsify the framework.
- (6) **Dark energy.** The dark energy density is $\rho_{DE} = ((d+1)/2)(1+c\delta)m_\nu^4$, matching observation to 0.010%. Any future measurement of ρ_{DE} that deviates from this formula at more than 0.1% would falsify the framework.
- (7) **Charge rigidity.** The charge assignment on the 14-mode multiplet satisfies exactly two structurally consistent configurations, both with total charge $-1e$. A measurement of the vacuum's total electromagnetic charge that differs from $-1e$ (per one Standard Model generation) would falsify the rigidity theorem.
- (8) **Identification.** The Midpoint Sorting Theorem (HCSM-23 Rev. 7) assigns the 14 modes uniquely. A Standard Model particle with an $N/4$ label outside the set $\{60, 24, 21, 20, 14, 14, 12, 9.5, 9, 7.5, 2\}$ would falsify the framework.
- (9) **A_1 uniqueness.** The physical color algebra is the $\mathfrak{su}(3)$ subalgebra on the A_1 isotypic block, forced by the dimension-and-invariance theorem. A different embedding of $\mathfrak{su}(3)_c$ in the D_4 commutant would falsify Theorem 5.4 of HCSM-06.
- (10) **Hypercharge values.** The hypercharge values are uniquely determined by the content structure and anomaly cancellation. A different set of hypercharge values satisfying the same content structure and anomaly conditions would falsify Theorem 8.1 of HCSM-07.

10 Theorem Inventory

The framework's theorems are catalogued by their home paper. New theorems from the present revision cycle are marked with $\star$.

Theorem	Location
Laplacian uniqueness	HCSM-01 Thm 2.1
$d \cdot 2^d = 64$, $L = 8$	HCSM-01 Cor 4.1
Two derivations of $L = 8$	HCSM-01 §4, HCSM-14.b
$2(L - 1) = 14$	HCSM-01 Thm 2.5
Unified substrate selection	HCSM-00 §2.6 (Rev. 8)
Hodge-Dirac square	HCSM-02 Thm 3.2
Supercharge algebra	HCSM-02 Thm 4.4 (corrected)
Kernel dimension = 4	HCSM-02 Thm 3.8
Spectral gap $2 \sin(\pi/L)$	HCSM-02 Thm 3.12
Lorentzian metric uniqueness	HCSM-03 Thm 3.10
Self-paired eigenvalue $\lambda = 4$	HCSM-04 Thm 3.2
D_4 orbit decomposition $(2, 8, 4)$	HCSM-04 Thm 5.1
Support classes $\{64, 48, 32\}$	HCSM-04 Prop 6.2
IPRs $\{2/128, 3/128, 4/128\}$	HCSM-04 Prop 7.2
Entropy classes $\{6.0, 5.5, 5.0\}$	HCSM-04 Cor 8.3
D_4 character $\chi_{14} = (14, 2, 0, 2, 2)$	HCSM-05 Thm 4.1
Irreducible decomposition	HCSM-05 Thm 6.1
n_1 -parity orbit indicator	HCSM-05 Thm 7.3
Joint (D, T) class partition	HCSM-05 Thm 7.7
Sector-coarse gauge algebra	HCSM-06 Thm 5.10
$\star A_1$ uniqueness for physical color	HCSM-06 Thm 5.4
$\star B_1/B_2$ outer automorphism	HCSM-06 Thms 5.8, 5.10
Hypercharge coefficients	HCSM-07 Thms 5.1, 6.1
$\star$ Hypercharge values from anomaly cancellation	HCSM-07 Thm 8.1
$\star$ Content structure from HCSM data	HCSM-07 Thm 8.3
Orientation-flip $\mathbb{Z}_2$	HCSM-09
Winding class partition $(3, 7, 4)$	HCSM-14 Thm 15
Kernel-Diagonal Theorem	HCSM-14.a Thm 1
Three coefficients from one diagonal	HCSM-14.a Cor 2
Maximal cancellation selects $L = 8$	HCSM-14.b Thm 4.4.1
Hartree orbit trace identity	HCSM-19 Prop 9.2
Loop-pass duration $T_{\text{DM}} = 1/(2\gamma)$	HCSM-21 Thm 12.1
Exact ratio $ M_A / M_B = 3$	HCSM-20 Thm 9.4
Zero-state structure	HCSM-22 Thm 12
Midpoint Sorting Theorem	HCSM-23 Rev. 7 Thm 5.3
Charge bijection	HCSM-24 Thm 4.1
σ -line sums	HCSM-24 Thm 5.1
Non-existence of coordinate closed form	HCSM-24 Thm 9.1
Sorting-machine interpretation	HCSM-24 Thm 10.1
Framework-constant σ -line sums	HCSM-24 Thm 10.6
F -pair sum rules by shell	HCSM-24 Thm 10.7
Charge rigidity class (5 constraints)	HCSM-24 Thm 11.1
Charge assignment solutions (2)	HCSM-24 Thm 11.2
Total charge rigidity	HCSM-24 Thm 11.3

Theorem	Location
Empirical residue of the charge sector	HCSM-24 Cor 11.4
Top-Yukawa coefficient $4/3$	HCSM-40
Absence of light dark photon	HCSM-41 Thm 5
Spin chirality $\mathbb{Z}_2$ grading	HCSM-42
Strong CP $\theta_{\text{QCD}} = 0$	HCSM-43 Thm 6
Proca sector	HCSM-44
Graviton masslessness by Schur	HCSM-45, 46
$G_N = (d+1)^2/(8\pi M_P^2)$	HCSM-48
kL correction $(7/2)\delta_{\text{edge}}$	HCSM-51
EFT coefficient $c = 5/9$	HCSM-52
S_0 normalization	HCSM-53
Vacuum stability $\gamma = c_2/I(2)$	HCSM-54
Λ -promotion is a projection	HCSM-56.a

All seven rank blockers of the framework’s derivation program are closed. As of the present revision, the physical identification of the A_1 and B_1 blocks is derived (color) or a labeling convention (weak isospin), and the hypercharge values and content structure are derived. The framework has zero empirical anchors in the particle, gauge, and hypercharge sectors.

11 Reproducibility

All numerical claims in the HCSM paper stack are verified at machine precision in IEEE-754 double precision. The verification scripts use only standard libraries (NumPy, SciPy, SymPy, mpmath) and construct the objects they need from HCSM-native definitions. Running `12-VERIFICATION/run_all.py` reproduces every numerical claim in the stack.

The complete HCSM paper stack, verification scripts, and source are available at

<https://github.com/007STAN/HODGE-COMPLEX-STANDARD-MODEL-HCSM>

12 Conclusion

This paper has established the foundation of the Hodge Complex Standard Model. The substrate is the periodic 8×8 torus $\Lambda = \mathbb{Z}_8 \times \mathbb{Z}_8$, selected by the unified substrate selection theorem (Section 2.6). The substrate carries the Hodge complex $\Omega^0 \oplus \Omega^1 \oplus \Omega^2$ with $D^2 = \Delta$ exactly on each form degree. The 14 modes at the self-paired eigenvalue $\lambda = 4$ are identified with one Standard Model generation. The processor T and its five-layer loop are the machine’s operation. The output structures (charge, mass, flavor, gravity, cosmology) are the residue after mirror cancellation.

The framework’s most consequential structural results are five. First, the Kernel-Diagonal Theorem unifies the framework’s three leading coefficients under a single projector: the EFT coefficient $c = 5/9$, the mass-map coefficient $(d-1)/2 = 3/2$, and the kernel dimension $10 = (d-1)^2 + 1$ are three traces of the diagonal of the projector onto $\ker(H_{\text{wind}})$. Second, the Λ -promotion mechanism is closed: the residual vacuum energy is promoted by pure projection onto the timelike direction of $\ker D$, and the hidden $\mathfrak{u}(1)$ VEVs vanish. Third, the framework’s coverage of the Standard Model gauge content (absence of a light dark photon, spin-chirality $\mathbb{Z}_2$ grading, strong-CP solution, Proca sector, topological protection of ϕ_{gen}) is complete. Fourth, the charge sector is a rigidity-class result: five structural constraints reduce the charge assignment from 5^{14} configurations to exactly two, both with total charge $-1e$, and the residual binary choice is resolved by the Midpoint Sorting Theorem. Fifth, the identification thesis is closed: the 14

modes are uniquely sorted into the 14 Standard Model particles by native HCSM data, with zero empirical anchors remaining in the particle sector.

The present revision closes the physical identification of the gauge and hypercharge sectors. The A_1 isotypic block is uniquely identified with physical color by a dimension-and-invariance theorem (HCSM-06, Theorem 5.4). The B_1 block is identified with physical weak isospin up to the outer-automorphism labeling convention (HCSM-06, Theorem 5.10). The Standard Model content structure and hypercharge values are derived from HCSM-native data and anomaly cancellation (HCSM-07, Theorems 8.1 and 8.3). The framework now has zero empirical anchors in the particle, gauge, and hypercharge sectors.

The framework has a single dimensionful anchor (the Planck mass M_P) and derives every other quantity from the substrate at $d = 4$ and $L = 8$.

The framework's claim is that the Standard Model is what the substrate looks like when you observe it.

Physics is what the machine cannot cancel. The Hodge complex is the machine. The $-1e$ is what the machine must not cancel.

References

- [1] S. Preschutti, *HCSM-01: The 8×8 Torus Substrate from $d = 4$* , HCSM White Paper Series (2026).
- [2] S. Preschutti, *HCSM-02: The Hodge Complex on Λ* , HCSM White Paper Series (2026).
- [3] S. Preschutti, *HCSM-03: Lorentzian Signature from the Hodge Star*, HCSM White Paper Series (2026).
- [4] S. Preschutti, *HCSM-04: The 14-Mode Multiplet at $\lambda = 4$* , HCSM White Paper Series (2026).
- [5] S. Preschutti, *HCSM-05: D_4 Representation Theory of the 14-Mode Multiplet*, HCSM White Paper Series (2026).
- [6] S. Preschutti, *HCSM-06 (Revised): Sector-Coarse Correspondence*, HCSM White Paper Series (2026).
- [7] S. Preschutti, *HCSM-07 (Revised): Hypercharge Quantization*, HCSM White Paper Series (2026).
- [8] S. Preschutti, *HCSM-08: Anomaly Cancellation as a Theorem*, HCSM White Paper Series (2026).
- [9] S. Preschutti, *HCSM-09: Gauge Topology and Orientation-Flip $\mathbb{Z}_2$* , HCSM White Paper Series (2026).
- [10] S. Preschutti, *HCSM-13: The Processor T and the Fermion/Boson-Class Correspondence*, HCSM White Paper Series (2026).
- [11] S. Preschutti, *HCSM-14: Winding Generator Spectrum and Class Partition*, HCSM White Paper Series (2026).
- [12] S. Preschutti, *HCSM-14.a: The Kernel-Diagonal Theorem*, HCSM White Paper Series (2026).
- [13] S. Preschutti, *HCSM-14.b: Maximal Cancellation Selects $L = 8$* , HCSM White Paper Series (2026).

- [14] S. Preschutti, *HCSM-19: DNLS Hartree Mass-Map Coefficient*, HCSM White Paper Series (2026).
- [15] S. Preschutti, *HCSM-20: The Motion Theorem*, HCSM White Paper Series (2026).
- [16] S. Preschutti, *HCSM-21: The Time Theorem*, HCSM White Paper Series (2026).
- [17] S. Preschutti, *HCSM-22: The Zero State*, HCSM White Paper Series (2026).
- [18] S. Preschutti, *HCSM-23 (Seventh Revision): The Midpoint Sorting Theorem*, HCSM White Paper Series (2026).
- [19] S. Preschutti, *HCSM-24 (Revised): The Charge Theorem*, HCSM White Paper Series (2026).
- [20] S. Preschutti, *HCSM-33: Mass Map*, HCSM White Paper Series (2026).
- [21] S. Preschutti, *HCSM-34: The CKM Matrix*, HCSM White Paper Series (2026).
- [22] S. Preschutti, *HCSM-36: The PMNS Matrix*, HCSM White Paper Series (2026).
- [23] S. Preschutti, *HCSM-40: The Higgs Mechanism*, HCSM White Paper Series (2026).
- [24] S. Preschutti, *HCSM-41: Absence of a Light Dark Photon*, HCSM White Paper Series (2026).
- [25] S. Preschutti, *HCSM-42: Spin Chirality $\mathbb{Z}_2$ Grading*, HCSM White Paper Series (2026).
- [26] S. Preschutti, *HCSM-43: Strong CP from the Orientation $\mathbb{Z}_2$* , HCSM White Paper Series (2026).
- [27] S. Preschutti, *HCSM-44: The Proca Sector*, HCSM White Paper Series (2026).
- [28] S. Preschutti, *HCSM-45: Graviton as Self-Paired Curvature Mode*, HCSM White Paper Series (2026).
- [29] S. Preschutti, *HCSM-46: Graviton Exact Masslessness*, HCSM White Paper Series (2026).
- [30] S. Preschutti, *HCSM-48: Faddeev-Popov G_N Normalization*, HCSM White Paper Series (2026).
- [31] S. Preschutti, *HCSM-51: The kL Crossover*, HCSM White Paper Series (2026).
- [32] S. Preschutti, *HCSM-52: The EFT Coefficient in the Dark Energy Density*, HCSM White Paper Series (2026).
- [33] S. Preschutti, *HCSM-53: Dark Energy Density and the DME Entropy-Well Normalization*, HCSM White Paper Series (2026).
- [34] S. Preschutti, *HCSM-54: Vacuum Stability and the Stability Coefficient*, HCSM White Paper Series (2026).
- [35] S. Preschutti, *HCSM-56.a: Λ -Promotion as Pure Projection*, HCSM White Paper Series (2026).
- [36] J. Dodziuk, *Finite-difference approach to the Hodge theory of harmonic forms*, Amer. J. Math. **98**, 79 (1976).
- [37] M. Nakahara, *Geometry, Topology and Physics*, 2nd ed., CRC Press (2003).

- [38] C. Voisin, *Hodge Theory and Complex Algebraic Geometry I*, Cambridge University Press (2002).
- [39] Particle Data Group, Phys. Rev. D **110**, 030001 (2024).

A Web Crawler Metadata

A.1 Dublin Core

- **DC.title:** HCSM-00 (Eighth Revision, Corrected): Foundations, Notation, and Mapping
- **DC.creator:** Preschutti, Stanley
- **DC.creator.orcid:** 0009-0004-5445-1744
- **DC.date.issued:** 2026-09
- **DC.date.modified:** 2026-09
- **DC.type:** Text / Preprint / White Paper
- **DC.format:** application/pdf; text/x-latex
- **DC.language:** en
- **DC.publisher:** Entropia Research Institute
- **DC.rights:** All rights reserved
- **DC.subject:** Hodge Complex Standard Model; HCSM; foundations; notation; mapping; 8×8 torus; Hodge complex; spectral midpoint; processor; output structures; emergent physics; cosmology; Laplacian uniqueness; Hodge-Dirac square; Lorentzian signature; Kernel-Diagonal Theorem; charge rigidity class; total charge invariance; Midpoint Sorting Theorem; identification thesis; zero empirical anchors; A_1 uniqueness; B_1/B_2 outer automorphism; hypercharge values; content structure; Λ -promotion; falsifiable predictions
- **DC.description:** This paper establishes the foundational objects of the Hodge Complex Standard Model and states the notation. The substrate is the periodic 8×8 torus $\Lambda = \mathbb{Z}_8 \times \mathbb{Z}_8$, selected by the unified substrate selection theorem. The Hodge-Dirac operator $D = d + d^*$ satisfies $D^2 = \Delta$ on each form degree. The 14 modes at $\lambda = 4$ are identified with one Standard Model generation. The paper derives the Kernel-Diagonal Theorem, the Lorentzian metric uniqueness, the spectral gap $2 \sin(\pi/L)$, the three coefficients $c = 5/9$, $(d - 1)/2 = 3/2$, and $\dim \ker(H_{\text{wind}}) = 10$, the Λ -promotion mechanism, the absence of a light dark photon, and the rigidity class of the charge assignment. The eighth revision incorporates two structural sharpenings: the A_1 color identification is now a theorem (HCSM-06 Thm 5.4), the B_1/B_2 choice is a labeling convention (HCSM-06 Thm 5.10), and the hypercharge values and content structure are derived (HCSM-07 Thms 8.1, 8.3). The framework now has zero empirical anchors in the particle, gauge, and hypercharge sectors. This corrected version fixes the supercharge algebra statement (Theorem 3.6) and updates derivation-status citations.
- **DC.identifier:** HCSM-00-2026-09-Rev8-Corr

A.2 BibTeX

```
@article{PreschuttiHCSM00Rev8Corr,
  author = {Preschutti, Stanley},
  title = {{HCSM-00} (Eighth Revision, Corrected): Foundations, Notation, and Mapping},
  journal = {HCSM White Paper Series},
  year = {2026},
  month = {September},
  note = {ORCID: 0009-0004-5445-1744},
  keywords = {Hodge Complex Standard Model, HCSM, foundations, notation,
    mapping, 8x8 torus, Hodge complex, spectral midpoint, processor,
    output structures, emergent physics, cosmology, Laplacian
    uniqueness, Hodge-Dirac square, Lorentzian signature,
    Kernel-Diagonal Theorem, charge rigidity class, total charge
    invariance, Midpoint Sorting Theorem, identification thesis,
    zero empirical anchors, A1 uniqueness, B1/B2 outer automorphism,
    hypercharge values, content structure, Lambda-promotion,
    falsifiable predictions},
  abstract = {This paper establishes the foundational objects of the Hodge
    Complex Standard Model and states the notation. The substrate is
    the periodic 8x8 torus  $\Lambda = \mathbb{Z}^8 \times \mathbb{Z}^8$ , selected by the unified
    substrate selection theorem. The Hodge-Dirac operator  $D = d + d^*$ 
    satisfies  $D^2 = \Delta$  on each form degree. The 14 modes at
     $\lambda = 4$  are identified with one Standard Model generation.
    The paper derives the Kernel-Diagonal Theorem, the Lorentzian
    metric uniqueness, the spectral gap  $2 \sin(\pi/L)$ , the three
    coefficients  $c = 5/9$ ,  $(d-1)/2 = 3/2$ , and  $\dim \ker(H_{\text{wind}}) = 10$ ,
    the Lambda-promotion mechanism, the absence of a light dark
    photon, and the rigidity class of the charge assignment. This
    corrected version fixes the supercharge algebra statement and
    updates derivation-status citations.}
}
```