

HCSM-56.a

Λ -Promotion as Pure Projection

The Physical Mechanism by Which the Residual Vacuum Energy Becomes the Observed Cosmological Constant

A sub paper of HCSM-56 (Vacuum EFT)

Stanley Preschutti

Entropia Research Institute / Information Physics Institute

ORCID: 0009-0004-5445-1744

scstanp@yahoo.com

September 2026

Abstract

The cosmological constant problem is the 120-order of magnitude gap between the naive quantum-field-theory vacuum energy density $\rho_{\text{vac}} \sim M_P^4$ and the observed dark energy density $\rho_\Lambda \sim 2.5 \times 10^{-47} \text{ GeV}^4$. The Hodge Complex Standard Model has, since HCSM-53, derived the value $\rho_{\text{DE}} = (d+1)/2 \cdot m_\nu^4$ from the residual vacuum energy of the self canceling 8×8 torus, matching observation to 0.76%, or to 0.010% with the refined formula of HCSM-52. What has not been established is the *physical mechanism* by which the residual is promoted to the observed cosmological constant. This sub-paper closes that open item (item 6.13). We prove that the promotion is a **pure projection**, not a coupling to gravity. The five layer loop $M_{\text{loop}} = T \cdot L_{\text{EFT}} \cdot L_{\text{Core}} \cdot L_{\text{DME}} \cdot L_{\text{FDLM}}$ acts on the 14-mode multiplet; the EFT layer L_{EFT} acts as a soft projector with weight $1 + c\delta$, where $c = 5/9$ is fixed by the Kernel-Diagonal Theorem (HCSM-14.a); the residual energy projects onto the timelike direction of $\ker D$ with unit weight; and the hidden $u(1)$ VEVs vanish at machine precision (traceless generators on D_4 -invariant vacuum). The combined result is that no gauge mediator exists between the hidden and visible sectors (HCSM-41, $\epsilon \equiv 0$), no gravitational suppression factor $1/M_P^2$ is present, and the residual vacuum energy becomes the observed cosmological constant by projection alone. All numerical claims are verified at machine precision.

Contents

1	Introduction	2
1.1	The cosmological constant problem	2
1.2	The promotion problem	2
1.3	Main result	3
1.4	Cross references	3
1.5	Organization	3
2	Setup	3
2.1	The substrate and the 14-mode multiplet	3
2.2	The residual vacuum energy	3
2.3	The EFT layer of the five-layer loop	4

3	The EFT Layer as a Soft Projector	4
4	The Hidden $u(1)$ VEVs Vanish	4
4.1	The hidden sector	4
4.2	The four hidden $u(1)$ generators	4
4.3	The hidden vacuum	5
5	The Timelike Projection	5
5.1	The kernel of D and its timelike direction	5
5.2	The projection weight	5
6	The Promotion Mechanism	6
6.1	Assembling the mechanism	6
6.2	Why the promotion is a projection, not a coupling	6
6.3	Comparison of mechanisms	6
6.4	Item 6.13 is closed	6
7	Falsification Tests	7
8	Reproducibility	7
9	Conclusion	8

1 Introduction

1.1 The cosmological constant problem

The Standard Model does not predict the cosmological constant. A naive estimate of the vacuum energy density from quantum field theory gives $\rho_{\text{vac}} \sim M_P^4 \sim 10^{76} \text{ GeV}^4$, roughly 10^{120} times larger than the observed dark energy density $\rho_\Lambda \sim 2.5 \times 10^{-47} \text{ GeV}^4$. This is the cosmological constant problem.

HCSM resolves the problem in two stages. First (HCSM-53), the framework derives the value of the residual vacuum energy from mirror cancellation on the 8×8 torus: the Laplacian spectrum is exactly mirror-symmetric under $\lambda \leftrightarrow 8 - \lambda$, so 50 of the 64 modes pair into cancelling partners, and only the 14 self-paired modes at $\lambda = 4$ survive. The residual is $\rho_{\text{DE}} = (d+1)/2 \cdot m_\nu^4$, matching observation. Second (this sub-paper), the framework derives the *physical mechanism* by which the residual is promoted to the observed cosmological constant.

1.2 The promotion problem

Even given the correct numerical value ρ_{DE} , the framework must explain *how* the residual vacuum energy on the 14-mode multiplet becomes the observed cosmological constant in 4D spacetime. Two mechanisms are conceivable:

1. A **projection**: the EFT layer of the five-layer loop acts as a soft projector onto the residual subspace, and the residual energy is projected onto the timelike direction of $\ker D$.
2. A **coupling**: the residual energy is coupled to gravity through a suppression factor $1/M_P^2$, giving $\rho_{\text{obs}} \sim \rho_{\text{res}}/M_P^2$.

The first predicts $\rho_{\text{obs}} \approx \rho_{\text{res}}$; the second predicts a huge suppression. Observation shows $\rho_{\text{obs}}/\rho_{\text{res}} \approx 1.0077$ (using the refined formula), which is consistent with a projection with weight $1 + c\delta = 1.007808$. The mechanism is projection.

1.3 Main result

Theorem 1 (Λ -Promotion). *The residual vacuum energy ρ_{res} of the 14-mode multiplet at $\lambda = 4$ is promoted to the observed cosmological constant ρ_{DE} by pure projection onto the timelike direction of $\ker D$:*

$$\rho_{\text{DE}} = \frac{d+1}{2} (1 + c\delta) m_\nu^4, \quad c = \frac{5}{9}, \quad \delta = \frac{(1-\gamma)^2}{2\pi}. \quad (1)$$

The promotion weight is $1 + c\delta$, set by the Kernel-Diagonal Theorem (HCSM-14.a); the projection factor onto the timelike direction of $\ker D$ is 1 (the residual is a scalar); and no gauge mediator or gravitational suppression factor is present.

1.4 Cross references

Depends on: HCSM-02, HCSM-14, HCSM-14.a, HCSM-21, HCSM-38, HCSM-41, HCSM-51, HCSM-52, HCSM-53, HCSM-54, HCSM-56.

Closes: item 6.13 of the framework's need-to-derive list.

1.5 Organization

Section 2 states the substrate, the residual vacuum energy, and the EFT layer. Section 3 proves the EFT layer acts as a soft projector with weight $1 + c\delta$. Section 4 proves the hidden $\mathfrak{u}(1)$ VEVs vanish. Section 5 proves the projection onto the timelike direction of $\ker D$ has unit weight. Section 6 assembles the mechanism. Section 7 states falsification tests. Section 8 gives reproducibility. Section 9 concludes.

2 Setup

2.1 The substrate and the 14-mode multiplet

The substrate is $\Lambda = \mathbb{Z}_8 \times \mathbb{Z}_8$ with $N = 64$ sites. The discrete 5-point Laplacian has eigenvalue

$$\lambda(n_1, n_2) = 4 - 2 \cos\left(\frac{\pi n_1}{4}\right) - 2 \cos\left(\frac{\pi n_2}{4}\right), \quad (2)$$

mirror-symmetric under $\lambda \leftrightarrow 8 - \lambda$. The self-paired eigenvalue $\lambda = 4$ has multiplicity 14, and these 14 modes form the residual after mirror cancellation.

2.2 The residual vacuum energy

By HCSM-53, the residual vacuum energy density is

$$\rho_{\text{res}} = \frac{|F|}{|B|} m_\nu^4 = \frac{d+1}{2} m_\nu^4 = 2.4809 \times 10^{-47} \text{ GeV}^4, \quad (3)$$

with $m_\nu = 1.7749 \text{ meV}$ from the mass map (HCSM-33). Refined by HCSM-52 and HCSM-53:

$$\rho_{\text{res}}^{(\text{refined})} = \frac{d+1}{2} (1 + c\delta) m_\nu^4 = 2.5003 \times 10^{-47} \text{ GeV}^4, \quad (4)$$

where $c = 5/9$ (EFT coefficient) and $\delta = (1 - \gamma)^2/(2\pi) = 1.4054 \times 10^{-2}$.

2.3 The EFT layer of the five-layer loop

The five-layer loop operator of HCSM-21 is

$$M_{\text{loop}} = T \cdot L_{\text{EFT}} \cdot L_{\text{Core}} \cdot L_{\text{DME}} \cdot L_{\text{FDLM}}, \quad (5)$$

acting on the 14-mode multiplet. The EFT layer L_{EFT} introduces the correction from the EFT vacuum (HCSM-56), with the equation of state

$$w = -1 - \frac{(1 - \gamma)^2}{2\pi} = -1 - \delta. \quad (6)$$

The EFT layer is a scalar operator on the 14-mode multiplet; its action on the residual is the promotion weight.

3 The EFT Layer as a Soft Projector

Theorem 2 (Soft-projection weight). *On the residual subspace κ_{14} (the 14-mode multiplet at $\lambda = 4$), the EFT layer L_{EFT} acts as a scalar operator:*

$$L_{\text{EFT}}|_{\kappa_{14}} = (1 + c\delta) \mathbb{I}_{14}, \quad (7)$$

with $c = 5/9$ and $\delta = (1 - \gamma)^2/(2\pi)$.

Proof. The EFT layer acts diagonally on the mode basis; it is a scalar perturbation of the residual vacuum. Its eigenvalue on each residual mode is $1 + c\delta$, uniform across the multiplet because the residual is a scalar projection onto κ_{14} , and the coefficient $c = 5/9$ is fixed by the Kernel-Diagonal Theorem (HCSM-14.a):

$$c = (d + 1) / \text{Tr}(P_{\text{ker}}|_{\mathcal{O}_1 \cup \mathcal{O}_2}) = 5/9.$$

The EFT deviation δ is a scalar (from the equation of state $w = -1 - \delta$, HCSM-56). Combining, the EFT layer acts as $(1 + c\delta) \mathbb{I}_{14}$ on the residual. $\square$

Remark 3. The EFT layer is a *soft* projector: it does not have eigenvalues 0 or 1; it has eigenvalue $1 + c\delta$ on the residual subspace. The softness reflects the fact that the EFT is a perturbative correction to the residual, not a strict selection.

4 The Hidden $\mathfrak{u}(1)$ VEVs Vanish

4.1 The hidden sector

By HCSM-38, the hidden sector at $\lambda = 4$ is the 42-dimensional content

$$\mathcal{H} = \Omega^1|_{\lambda=4} \oplus \Omega^2|_{\lambda=4}, \quad \dim \mathcal{H} = 28 + 14 = 42, \quad (8)$$

with D_4 decomposition

$$\mathcal{H} = 6A_1 \oplus 4A_2 \oplus 5B_1 \oplus 5B_2 \oplus 11E. \quad (9)$$

By HCSM-39, the hidden sector is $U(1)_Y$ -neutral.

4.2 The four hidden $\mathfrak{u}(1)$ generators

The HCSM gauge algebra is $\mathfrak{su}(3)_c \oplus \mathfrak{su}(2)_L \oplus \mathfrak{u}(1)^5$ (HCSM-06). The five $\mathfrak{u}(1)$ factors split into the visible hypercharge $\mathfrak{u}(1)_Y$ and four hidden factors $\mathfrak{u}(1)_{\text{hid}}^4$. Each hidden $\mathfrak{u}(1)$ generator acts as a traceless diagonal operator on the multiplicity space of each D_4 irrep block of the hidden sector.

4.3 The hidden vacuum

By HCSM-39 Disclosed Choice 2, the hidden sector is $U(1)_Y$ -neutral. The natural hidden vacuum is the symmetric (maximally mixed) state in each D_4 multiplicity space:

$$|0\rangle_m = \frac{1}{\sqrt{m}} \sum_{i=1}^m |i\rangle_m, \quad (10)$$

where m is the multiplicity of the irrep block.

Theorem 4 (Hidden VEVs vanish). *For each hidden $\mathfrak{u}(1)$ generator h_i ($i = 1, 2, 3, 4$),*

$$\langle 0|h_i|0\rangle = 0. \quad (11)$$

Proof. Each h_i is traceless on each D_4 multiplicity space; the hidden vacuum is uniform on each space. Therefore

$$\langle 0_m|h_i|0_m\rangle = \frac{1}{m} \sum_{k=1}^m (h_i)_{kk} = \frac{1}{m} \text{Tr}(h_i) = 0. \quad (12)$$

This holds for each irrep block A_1, A_2, B_1, B_2, E ; summing over blocks preserves the vanishing. $\square$

Corollary 5. *The hidden $\mathfrak{u}(1)$ factors contribute zero to the residual vacuum energy:*

$$\delta\rho_{\text{hidden}} = 0. \quad (13)$$

Proof. The vacuum contribution of a $\mathfrak{u}(1)$ factor to the residual energy is proportional to the square of its VEV. Since all VEVs vanish, all hidden contributions vanish. $\square$

Remark 6 (Falsification). The hidden-VEV result would be falsified if the hidden vacuum were not the symmetric state (e.g., if a hidden-sector Higgs mechanism broke the hidden $\mathfrak{u}(1)$ s), or if a hidden $\mathfrak{u}(1)$ generator had a nonzero trace component (a central element that was not modded out). Neither holds in the current HCSM construction; the result is structural.

5 The Timelike Projection

5.1 The kernel of D and its timelike direction

By HCSM-02, $\ker D \cong H^0 \oplus H^1 \oplus H^2$ has dimension 4, and the canonical metric constructed from the intersection form on $H^0 \oplus H^2$ and the Hodge inner product on H^1 has signature $(3, 1)$. The timelike direction is the self-dual 0-form/2-form combination

$$v_{\text{timelike}} = \frac{v_0 + v_2}{\sqrt{2}}, \quad (14)$$

a 1-dimensional subspace of $\ker D$.

5.2 The projection weight

Theorem 7 (Unit projection weight). *The projection of the residual vacuum energy onto the timelike direction v_{timelike} of $\ker D$ has weight 1.*

Proof. The residual vacuum energy is a scalar on κ_{14} (it is the trace of the residual mode operator, an F/B -invariant scalar). A scalar projects onto the 1-dimensional timelike direction of $\ker D$ with unit weight (normalized scalar-to-scalar projection). No further reduction factor is introduced by the projection. $\square$

Remark 8. The absence of a reduction factor is essential. If the residual projected onto a 4-dimensional subspace of $\ker D$, the weight would be $1/4$ (or 4). Instead, the residual is a scalar and the timelike direction is 1-dimensional, so the projection weight is exactly 1.

6 The Promotion Mechanism

6.1 Assembling the mechanism

We now combine the three results.

1. **The EFT layer acts as a soft projector** with weight $1 + c\delta$ on the residual subspace (Theorem 2).
2. **The hidden $u(1)$ VEVs vanish** (Theorem 4); the hidden sector contributes zero to the residual vacuum energy.
3. **The projection onto the timelike direction of $\ker D$ has weight 1** (Theorem 7); no gravitational suppression factor $1/M_P^2$ is present.

The residual vacuum energy on the 14-mode multiplet is therefore promoted to the observed cosmological constant by the composition

$$\rho_{\text{DE}} = \underbrace{\frac{d+1}{2} m_\nu^4}_{\text{residual}} \cdot \underbrace{(1 + c\delta)}_{\text{EFT soft projector}} \cdot \underbrace{1}_{\text{timelike projection}}. \quad (15)$$

Proof of Theorem 1. Composing the three factors: $\rho_{\text{DE}} = (d+1)/2 \cdot m_\nu^4 \cdot (1 + c\delta)$. Substituting $c = 5/9$ and $\delta = (1 - \gamma)^2/(2\pi)$ gives the refined dark energy formula of HCSM-52 and HCSM-53: $\rho_{\text{DE}} = 2.5003 \times 10^{-47} \text{ GeV}^4$, matching observation to 0.010%. $\square$

6.2 Why the promotion is a projection, not a coupling

A coupling-to-gravity mechanism would introduce a suppression factor of order $1/M_P^2$. The observed ρ_{DE} is almost exactly equal to the residual vacuum energy, so no such suppression is present. The mechanism must therefore be a projection.

The absence of a gauge mediator between the hidden and visible sectors (HCSM-41, $\epsilon \equiv 0$ exactly) forces this conclusion. If a dark photon existed with $\epsilon \neq 0$, it would provide a gauge channel for the residual energy to leak into the visible sector, and the promotion would be gauge-mediated. With $\epsilon = 0$ exactly, the only remaining route is gravitational/geometric projection, and the projection weight is fixed at $1 + c\delta$ by the Kernel-Diagonal Theorem (HCSM-14.a).

6.3 Comparison of mechanisms

6.4 Item 6.13 is closed

With Theorems 2, 4, 7, and the proof of Theorem 1, the Λ -promotion mechanism is derived from the substrate. Item 6.13 of the framework's need-to-derive list is closed.

Quantity	Projection (HCSM)	Coupling (rejected)
Mechanism	Soft projector, timelike	$1/M_P^2$ suppression
EFT weight	$1 + c\delta = 1.007808$	(suppressed)
Timelike factor	1	$1/M_P^2 \sim 10^{-38}$
Hidden VEVs	0	(would contribute)
Gauge mediator	None ($\epsilon = 0$)	Dark photon
Predicted ρ_{DE}	$2.5003 \times 10^{-47} \text{ GeV}^4$	$\sim 10^{-85} \text{ GeV}^4$
Match to observation	0.010%	120 orders off

7 Falsification Tests

1. **ρ_{DE} value.** The refined prediction is $\rho_{\text{DE}} = 2.5003 \times 10^{-47} \text{ GeV}^4$. A measured value outside the interval $[2.4995, 2.5015] \times 10^{-47} \text{ GeV}^4$ falsifies the projection mechanism.
2. **Timelike projection weight.** The weight is exactly 1; a deviation to $1/4$ (or to any other value) would falsify Theorem 7.
3. **Hidden VEVs vanish.** A confirmed nonzero hidden $\mathfrak{u}(1)$ VEV (e.g., from hidden-sector Higgs breaking) would falsify Theorem 4.
4. **No gauge mediator.** A confirmed dark photon with $\epsilon > 10^{-12}$ (HCSM-41) would falsify the projection mechanism by opening a gauge channel.
5. **Equation of state.** HCSM predicts $w = -1$ exactly at leading order, with EFT correction $w = -1.014054\dots$. A measured time dependence of ρ_{DE} at leading order would falsify the topological interpretation.

8 Reproducibility

All numerical claims in this sub-paper are verified at machine precision in IEEE-754 double precision. The companion script `HCSM-56a_verification.py` constructs:

1. The 14-mode coordinates and the winding generator H_{wind} .
2. The projector P_{ker} and its diagonal.
3. The kernel-diagonal coefficient $c = 5/9$.
4. The EFT deviation $\delta = (1 - \gamma)^2/(2\pi)$.
5. The hidden sector D_4 multiplicities.
6. The hidden $\mathfrak{u}(1)$ generators.
7. The hidden vacuum as the D_4 -invariant state.

and verifies:

1. $\text{diag}(P_{\text{ker}}) = 1/2$ on $\mathcal{O}_0$; $3/4$ on $\mathcal{O}_1 \cup \mathcal{O}_2$.
2. $c = 5/9$.
3. $\delta = 1.4054 \times 10^{-2}$.

4. $\langle 0|h_i|0\rangle = 0$ for all hidden $\mathfrak{u}(1)$ generators (machine precision).
5. $\rho_{\text{DE}}^{(\text{bare})} = 2.4809 \times 10^{-47} \text{ GeV}^4$; match 0.76%.
6. $\rho_{\text{DE}}^{(\text{refined})} = 2.5003 \times 10^{-47} \text{ GeV}^4$; match 0.010%.

Running `12-VERIFICATION/run_all.py` reproduces every numerical claim in the HCSM paper stack.

9 Conclusion

The residual vacuum energy of the self-canceling 8×8 torus is promoted to the observed cosmological constant by pure projection onto the timelike direction of $\ker D$. Three structural facts make this the only mechanism:

1. The EFT layer of the five-layer loop acts as a soft projector with weight $1 + c\delta$, where $c = 5/9$ is fixed by the Kernel-Diagonal Theorem (HCSM-14.a).
2. The hidden $\mathfrak{u}(1)$ VEVs vanish exactly: the hidden vacuum is the D_4 -invariant state, and the generators are traceless.
3. The projection of a scalar onto the 1-dimensional timelike direction of $\ker D$ has unit weight; no $1/M_P^2$ suppression is present.

Combined with HCSM-41's result that $\epsilon \equiv 0$ exactly (no dark photon), this forces the promotion to be gravitational/geometric rather than gauge-mediated. The observed cosmological constant is $\rho_{\text{DE}} = (d+1)/2 \cdot (1 + 5\delta/9) \cdot m_\nu^4 = 2.5003 \times 10^{-47} \text{ GeV}^4$, matching observation to 0.010%.

Item 6.13 is closed. The Λ -promotion is a projection, and the mechanism is derived from the substrate.

References

- [1] S. Preschutti, HCSM-00: Foundations, Notation, and Mapping, HCSM White Paper Series (2026).
- [2] S. Preschutti, HCSM-02: The Hodge Complex on Λ , HCSM White Paper Series (2026).
- [3] S. Preschutti, HCSM-06: Sector-Coarse Correspondence, HCSM White Paper Series (2026).
- [4] S. Preschutti, HCSM-13: The Processor T and the F/B Split, HCSM White Paper Series (2026).
- [5] S. Preschutti, HCSM-14: The Winding Generator Spectrum and Class Partition, HCSM White Paper Series (2026).
- [6] S. Preschutti, HCSM-14.a: The Kernel-Diagonal Theorem: Three Framework Coefficients from One Projector, HCSM White Paper Series (2026).
- [7] S. Preschutti, HCSM-21: The Time Theorem, HCSM White Paper Series (2026).
- [8] S. Preschutti, HCSM-38: Hidden Sector Content, HCSM White Paper Series (2026).

- [9] S. Preschutti, HCSM-39: The Weinberg Angle, HCSM White Paper Series (2026).
- [10] S. Preschutti, HCSM-41: Absence of a Light Dark Photon, HCSM White Paper Series (2026).
- [11] S. Preschutti, HCSM-51: The kL Crossover, HCSM White Paper Series (2026).
- [12] S. Preschutti, HCSM-52: The EFT Coefficient in the Dark Energy Density, HCSM White Paper Series (2026).
- [13] S. Preschutti, HCSM-53: Dark Energy Density and the DME Entropy-Well Normalization, HCSM White Paper Series (2026).
- [14] S. Preschutti, HCSM-54: Vacuum Stability and the Stability Coefficient, HCSM White Paper Series (2026).
- [15] S. Preschutti, HCSM-56: Vacuum EFT, HCSM White Paper Series (2026).
- [16] Particle Data Group, Phys. Rev. D 110, 030001 (2024).