

HCSM-26

The Observer Pair $\{W, t\}$

Fixed Locus, Chirality Selection, Uniqueness, and the Klein Orbit Structure
The Hodge Complex Standard Model

Stanley Preschutti¹

¹Entropia Research Institute / Information Physics Institute, ORCID:
0009-0004-5445-1744, scstannp@yahoo.com

September 2026

Abstract

The Hodge Complex Standard Model assigns the fourteen modes at the self paired eigenvalue of the discrete Laplacian on the periodic 8×8 torus to one Standard Model generation. Among these fourteen modes, the W boson and the top quark form a distinguished pair: the *observer pair* $\{W, t\}$. This paper proves six theorems that together characterize the observer pair completely within the HCSM framework. First, the observer pair is the fixed locus of the unique chiral reflection of the D_4 point group of the square. Second, it is the unique reflection fixed pair straddling the fermion/boson boundary. Third, it is the unique pair satisfying five independent structural constraints simultaneously. Fourth, its $N/4$ labels are the unique solution to the linear system determined by the dark energy coefficient and the dimensional rigidity constant. Fifth, the framework constant $(d + 1)/2$ manifests in each of the three D_4 orbits through a distinct binary operation (ratio, difference, sum), and the observer pair realizes the sum signature. Sixth, the orbit-class swap pair $\{\text{dark}, t\}$ is the unique two-element subset of the fourteen-mode multiplet whose four symmetric-function values (sum, difference, product, ratio) are simultaneously all framework constants. The paper also derives a dimensional-rigidity chain: the observer pair's separation, the kL crossover identity, and the dark energy coefficient all reduce to the same polynomial equation that has the unique positive integer solution $d = 4$. All theorems are HCSM native; all numerical claims are verified at machine precision.

Contents

1	Introduction	3
1.1	The observer pair in HCSM	3
1.2	The one line summary	3
1.3	The six theorems of this paper	3
1.4	Cross references to the HCSM paper stack	4
1.5	Organization	4
2	Setup	4
2.1	The substrate and the fourteen-mode multiplet	4
2.2	The D_4 action	5
2.3	The winding generator and the class partition	5
2.4	The processor T	5

2.5	The zero state	6
2.6	Charge, $N/4$ labels, and the σ -lines	6
2.7	Notation	6
3	The Fixed Locus Theorem	7
3.1	The fixed set of rs	7
3.2	The zero state vector is odd under rs	7
3.3	The observer pair and the σ -lines	7
4	The Chirality Selection Theorem	7
4.1	Commutation and anticommutation with H_{wind}	8
4.2	The Chirality Selection Theorem	8
5	The D_4-Fixed Pair Taxonomy	8
5.1	The three fixed pairs	8
5.2	Structural fingerprints	9
6	The $(d+1)/2$ Orbit Signature	9
6.1	The $N/4$ values by orbit	9
6.2	The three operations	10
6.3	Specificity of the signature	10
7	The Observer-Pair Uniqueness Theorem	10
7.1	The five constraints	10
7.2	Uniqueness	11
8	The Swap-Pair Fingerprint Theorem	11
8.1	The orbit-class swap	11
8.2	The symmetric-function fingerprint	12
9	The Dimensional-Rigidity Chain	13
9.1	The three fingerprints of $d = 4$	13
9.2	The dark energy coefficient identity	14
10	Discussion	14
10.1	What the observer pair is	14
10.2	The three structural regions	14
10.3	The role of the swap	14
11	Reproducibility	15
12	Conclusion	15
A	Web Crawler Metadata	16
A.1	Dublin Core	16
A.2	Schema.org JSON-LD	17
A.3	BibTeX for Automated Indexing	18

1 Introduction

1.1 The observer pair in HCSM

The Hodge Complex Standard Model (HCSM) derives the gauge structure, matter content, particle spectrum, and mixing matrices of the Standard Model from a single discrete substrate: the periodic 8×8 torus $\Lambda = \mathbb{Z}_8 \times \mathbb{Z}_8$ [1]. The fourteen modes at the self paired eigenvalue $\lambda = 4$ of the discrete Laplacian are identified with one Standard Model generation. Their D_4 representation theory determines the gauge algebra; their $N/4$ labels determine the mass spectrum; the zero-state vector s from the 14×19 vacuum-native measurement matrix determines the flavor structure.

Among the fourteen modes, the pair $\{W, t\}$ the W boson and the top quark carries a distinctive structural status. This paper proves that the pair is the fixed locus of a unique chiral reflection of the D_4 point group of the square, and that the entire residual structure of the fourteen-mode multiplet organizes around it.

1.2 The one line summary

In HCSM, physics is what the machine cannot cancel. The observer pair is the pair of modes that the machine cannot cancel without breaking its own chirality. The twelve other modes can be paired up and cancelled in mirror pairs; the pair $\{W, t\}$ cannot, because both members are fixed by the reflection that the machine is built to be symmetric under. The pair is therefore the framework's *observer pair*: the boundary through which the substrate reads itself.

1.3 The six theorems of this paper

The paper is organized around six theorems.

1. **Fixed-Locus Theorem** (Section 3). The observer pair $\{W, t\}$ is the fixed locus of the diagonal reflection rs of D_4 , and coincides with the zero locus of the zero-state vector s and with the $\sigma = 0$ line of the charge theorem.
2. **Chirality Selection Theorem** (Section 4). The diagonal reflection rs is the unique element of D_4 that both fixes exactly the pair $\{(2, 2), (6, 6)\}$ and anticommutes with the winding generator H_{wind} .
3. **D_4 -Fixed Pair Taxonomy** (Section 5). The fourteen mode multiplet contains exactly three D_4 -involution-fixed two-element subsets, and they are distinguished in every structural feature: winding class, fermion/boson content, σ -line membership, and $N/4$ ratio.
4. **$(d+1)/2$ Orbit Signature** (Section 6). The framework constant $(d+1)/2 = 5/2$ manifests in each of the three D_4 orbits via a distinct binary operation: as a ratio in $\mathcal{O}_0$, a difference in $\mathcal{O}_1$, and a sum in $\mathcal{O}_2$.
5. **Observer-Pair Uniqueness Theorem** (Section 7). The observer pair $\{W, t\}$ is the unique pair of $N/4$ values satisfying the two numeric constraints $N/4(W) + N/4(t) = (d+1)/2$ and $N/4(W) - N/4(t) = (d-1)/2$ simultaneously with the reflection-fixed and $\sigma = 0$ conditions.
6. **Swap-Pair Fingerprint Theorem** (Section 8). The orbit class swap pair $\{\text{dark}, t\}$ is the unique two element subset of the fourteen-mode multiplet whose four symmetric function values (*sum*, *diff*, *product*, *ratio*) are simultaneously all framework constants.

Section 9 derives the dimensional-rigidity chain that links the observer pair to the kL crossover and the dark energy coefficient.

1.4 Cross references to the HCSM paper stack

The paper depends on three companion papers in the HCSM stack, whose numbering follows the master index in HCSM-00:

- **HCSM-13** [2]: the processor T and the F/B -class correspondence. Used for the class partition, the fermion/boson split, and the exact correspondence $T = I - P_C$.
- **HCSM-24** [7]: the charge theorem $q_3 = F(n_1, \sigma)$. Used for the definition of the σ -lines and the charge eigenvalues.
- **HCSM-53** [9]: the EFT coefficient in the dark energy density. Used for the dark energy coefficient $(d + 1)/2$.

Additional references to HCSM-14 (winding generator spectrum), HCSM-23 (the δ function and $N/4$ labels), and HCSM-22 (the zero state) appear where needed.

1.5 Organization

Section 2 states the setup: the fourteen-mode multiplet, the D_4 action, the winding generator, the reflection rs , and the zero state. Sections 3–8 prove the six theorems. Section 9 derives the dimensional rigidity chain. Section 11 gives reproducibility notes. Section 12 concludes.

2 Setup

This section states the HCSM primitives on which the paper depends. All statements are HCSM-native; the reader is referred to HCSM-00 [1] for the full foundations.

2.1 The substrate and the fourteen-mode multiplet

The substrate is the periodic eight-by-eight torus $\Lambda = \mathbb{Z}_8 \times \mathbb{Z}_8$ with $N = 64$ sites, selected by the photon helicity anchor and the equal-entropy-spacing anchor [1, 10, 11]. The discrete five point Laplacian

$$(\Delta f)(x, y) = 4f(x, y) - f(x + 1, y) - f(x - 1, y) - f(x, y + 1) - f(x, y - 1) \quad (1)$$

has spectrum $\lambda(n_1, n_2) = 4 - 2\cos(\pi n_1/4) - 2\cos(\pi n_2/4)$ for $(n_1, n_2) \in \{0, \dots, 7\}^2$. The self-paired eigenvalue $\lambda = 4$ has multiplicity $2(L - 1) = 14$ with $L = 8$.

The fourteen modes at $\lambda = 4$ decompose under the D_4 point group of the square into three orbits:

$$\mathcal{O}_0 = \{(0, 4), (4, 0)\}, \quad \text{size } 2, \quad (2)$$

$$\mathcal{O}_1 = \{(1, 3), (1, 5), (3, 1), (3, 7), (5, 1), (5, 7), (7, 3), (7, 5)\}, \quad \text{size } 8, \quad (3)$$

$$\mathcal{O}_2 = \{(2, 2), (2, 6), (6, 2), (6, 6)\}, \quad \text{size } 4. \quad (4)$$

Following HCSM-04, we identify these modes with one Standard Model generation. In the standard mode order used throughout HCSM (matching the zero-state vector of Section 2.5), the fourteen modes are

$$(\nu, e, u, d, s, \mu, \text{dark}, c, \tau, b, W, Z, H, t). \quad (5)$$

2.2 The D_4 action

The D_4 point group is generated by the rotation

$$r : (n_1, n_2) \mapsto (-n_2, n_1) \pmod{8} \quad (6)$$

and the reflection

$$s : (n_1, n_2) \mapsto (n_1, -n_2) \pmod{8}, \quad (7)$$

subject to the presentation $r^4 = s^2 = (rs)^2 = e$. There are five order-two elements (involutions): r^2, s, rs, r^2s, r^3s . Explicitly,

$$r^2 : (n_1, n_2) \mapsto (-n_1, -n_2), \quad (8)$$

$$s : (n_1, n_2) \mapsto (n_1, -n_2), \quad (9)$$

$$rs : (n_1, n_2) \mapsto (n_2, n_1), \quad (10)$$

$$r^2s : (n_1, n_2) \mapsto (-n_1, n_2), \quad (11)$$

$$r^3s : (n_1, n_2) \mapsto (-n_2, -n_1). \quad (12)$$

The reflection rs is called the *diagonal reflection* because it swaps the two mode coordinates.

2.3 The winding generator and the class partition

The winding generator on the fourteen-mode multiplet is

$$H_{\text{wind}} = i \sin\left(\frac{\pi}{8}M\right), \quad M_{ij} = n_1^{(i)}n_2^{(j)} - n_1^{(j)}n_2^{(i)}, \quad (13)$$

where $(n_1^{(i)}, n_2^{(i)})$ are the momentum labels of mode i . Its spectrum is

$$\text{spec}(H_{\text{wind}}) = \{-4\sqrt{2}, -2\sqrt{2}, 0^{(10)}, +2\sqrt{2}, +4\sqrt{2}\}. \quad (14)$$

The fourteen modes partition into three *winding classes* by frequency:

$$A = \{(0, 4), (4, 0), (6, 6)\}, \quad \omega_A = 2\sqrt{2}, \quad (15)$$

$$B = \{(1, 3), (1, 5), (3, 1), (3, 7), (7, 3), (7, 5), (5, 1)\}, \quad \omega_B = 6\sqrt{2}, \quad (16)$$

$$C = \{(5, 7), (2, 2), (2, 6), (6, 2)\}, \quad \omega_C = 0. \quad (17)$$

See HCSM-14 for the full derivation of the winding spectrum.

2.4 The processor T

The processor T is the rank-ten orthogonal projection onto the fermion sector F :

$$T = P_{\text{mesh}} + P_\tau, \quad F = \{\delta \neq 0\} \cup \{s = +1\}, \quad B = \{\delta = 0\} \setminus \{s = +1\}, \quad (18)$$

with $|F| = 10$ and $|B| = 4$. The class partition and the F/B split are related by the exact correspondence (proved in HCSM-13 [2])

$$A \cup B = \text{image}(T) = F, \quad C = \ker(T) = B, \quad (19)$$

equivalently $T = I - P_C$.

2.5 The zero state

A 14×19 measurement matrix of vacuum-native observables has rank 13 and nullity 1 [12]. Its unique nullspace vector s , normalized so that $\max_i |s_i| = 1$, is the zero state. In the standard mode order,

$$s = \begin{pmatrix} -0.105198199128, & +0.105198199128, & +0.937726561300, \\ -0.192806690737, & -0.937726561300, & +0.658283218429, \\ -1.000000000000, & -0.658283218429, & +1.000000000000, \\ +0.192806690737, & 0.000000000000, & -0.329141609215, \\ +0.329141609215, & 0.000000000000 \end{pmatrix}. \quad (20)$$

The two modes with $s = 0$ exactly are the W boson at $(2, 2)$ and the top quark at $(6, 6)$.

2.6 Charge, $N/4$ labels, and the σ -lines

The charge theorem (HCSM-24 [7]) states that the electric charge in units of $e/3$ is a bijection

$$q_3 = F(n_1, \sigma), \quad n_1 \in \mathbb{Z}_8, \quad \sigma = \text{sign}(s) \in \{-1, 0, +1\}, \quad (21)$$

from the fourteen realized pairs to the five Standard Model charge eigenvalues $\{-3, -1, 0, +2, +3\}$. The three σ -lines are

$$\sigma = -1 : \{\nu, d, s, \text{dark}, c, Z\}, \quad \text{charge sum } 0, \quad (22)$$

$$\sigma = 0 : \{W, t\}, \quad \text{charge sum } +5, \quad (23)$$

$$\sigma = +1 : \{e, \mu, \tau, b, H, u\}, \quad \text{charge sum } -8. \quad (24)$$

The grand total is $-3 = -1e$.

The $N/4$ labels are derived from the mode coordinates by the δ function (HCSM-23 [13]). In the standard mode order,

$$(N/4)_i = (60, 24, 21, 20, 14, 14, 12, 9.5, 9, 7.5, 2, 2, 1, 0.5). \quad (25)$$

2.7 Notation

Throughout the paper, we use the following notation.

- $\mathcal{M}$ denotes the fourteen-mode multiplet at $\lambda = 4$.
- $\mathcal{O}_0, \mathcal{O}_1, \mathcal{O}_2$ denote the three D_4 orbits.
- A, B, C denote the three winding classes.
- F, B denote the fermion and boson sectors (image and kernel of T).
- $\sigma = \text{sign}(s)$ denotes the sign of the zero-state amplitude.
- $N/4$ denotes the mass-ordering label of a mode.
- $d = 4$ denotes the spacetime dimension.
- $H = 2^d = 16$ denotes the hidden-sector dimension parameter.
- $\mathcal{B}(\alpha, \beta)$ denotes the number of B_1 isotypic states.

3 The Fixed Locus Theorem

This section proves that the observer pair is the fixed locus of the diagonal reflection rs .

3.1 The fixed set of rs

Lemma 3.1 (Fixed set of rs). *The diagonal reflection $rs : (n_1, n_2) \mapsto (n_2, n_1)$ acts on the fourteen mode multiplet with fixed set exactly*

$$\text{Fix}(rs) = \{(2, 2), (6, 6)\} = \{W, t\}. \quad (26)$$

Proof. A mode (n_1, n_2) is fixed by rs if and only if $n_1 = n_2 \pmod{8}$. Among the fourteen modes at $\lambda = 4$, the diagonal pairs (n, n) with $n \in \{0, \dots, 7\}$ are $(0, 0), (1, 1), \dots, (7, 7)$. Of these, only $(2, 2)$ and $(6, 6)$ satisfy the eigenvalue equation $\lambda = 4$, i.e., $\cos(\pi n/4) = 0$, which requires $n \in \{2, 6\}$. Hence the fixed set is exactly $\{(2, 2), (6, 6)\}$. $\square$

3.2 The zero state vector is odd under rs

Lemma 3.2 (Oddness of s under rs). *The zero-state vector s satisfies $s(rs(m)) = -s(m)$ for every mode m in the fourteen-mode multiplet.*

Proof. Direct computation on the explicit vector of Section 2.5. For example, $s(\nu) = -0.105198199128$, $s(e) = +0.105198199128$, and $rs(\nu) = e$; $s(u) = +0.937726561300$, $s(s) = -0.937726561300$, and $rs(u) = s$. In every case, $s(rs(m)) = -s(m)$. $\square$

Corollary 3.3 (Vanishing on the fixed set). *The zero state vector satisfies $s(W) = s(t) = 0$ exactly.*

Proof. By Lemma 3.1, $rs(W) = W$ and $rs(t) = t$. By Lemma 3.2, $s(rs(W)) = -s(W)$, hence $s(W) = -s(W)$, hence $s(W) = 0$. The same argument gives $s(t) = 0$. $\square$

3.3 The observer pair and the σ -lines

Theorem 3.4 (Fixed Locus Theorem). *The following three characterizations of the observer pair coincide:*

$$\{W, t\} = \text{Fix}(rs) = \{m \in \mathcal{M} : s(m) = 0\} = \{\sigma = 0 \text{ line}\}. \quad (27)$$

Moreover, the reflection rs swaps the $\sigma = +1$ and $\sigma = -1$ lines, and fixes the $\sigma = 0$ line pointwise.

Proof. The first equality is Lemma 3.1. The second is Corollary 3.3. For the third, note that by definition $\sigma(m) = \text{sign}(s(m))$, so $\sigma(m) = 0$ iff $s(m) = 0$; this gives $\{m : s(m) = 0\} = \{\sigma = 0 \text{ line}\}$.

For the swap property: by Lemma 3.2, $s(rs(m)) = -s(m)$ for all m . Therefore $\sigma(rs(m)) = \text{sign}(s(rs(m))) = \text{sign}(-s(m)) = -\sigma(m)$ whenever $\sigma(m) \neq 0$. On the fixed set, $\sigma(W) = \sigma(t) = 0$, so the $\sigma = 0$ line is fixed pointwise. $\square$

Remark 3.5. The theorem gives three independent characterizations of the observer pair: as the fixed locus of the chiral reflection rs , as the zero locus of the zero state vector, and as the $\sigma = 0$ line of the charge theorem. That these three characterizations coincide is not a coincidence; it is a consequence of the fact (proved in Section 4) that rs is the unique chiral element of D_4 .

4 The Chirality Selection Theorem

This section proves that the diagonal reflection rs is the unique chiral element of D_4 , in the sense of anticommuting with the winding generator.

4.1 Commutation and anticommutation with H_{wind}

Definition 4.1 (Chiral element). An element $g \in D_4$ is called *chiral* if $gH_{\text{wind}}g^{-1} = -H_{\text{wind}}$, i.e., if g anticommutes with the winding generator.

Lemma 4.2 (Chirality of D_4 elements). *Among the five order two elements of D_4 , only rs anticommutes with H_{wind} .*

Proof. Direct computation of the commutator $[g, H_{\text{wind}}]$ and the anticommutator $\{g, H_{\text{wind}}\}$ for each of the five involutions r^2, s, rs, r^2s, r^3s . The Frobenius norms are:

Element g	$\ [g, H_{\text{wind}}]\ _F$	$\ \{g, H_{\text{wind}}\}\ _F$
r^2	1.131×10^1	1.386×10^1
s	1.131×10^1	1.386×10^1
rs	1.789×10^1	0.000×10^0
r^2s	1.131×10^1	1.386×10^1
r^3s	1.386×10^1	1.131×10^1

Only rs has $\|\{g, H_{\text{wind}}\}\|_F = 0$ exactly (at machine precision), and its commutator norm is nonzero. Hence rs is the unique element that anticommutes with H_{wind} . $\square$

4.2 The Chirality Selection Theorem

Theorem 4.3 (Chirality Selection Theorem). *The diagonal reflection $rs \in D_4$ is the unique element that (i) fixes exactly the pair $\{(2, 2), (6, 6)\}$ on the fourteen-mode multiplet, and (ii) anticommutes with the winding generator H_{wind} .*

Proof. Uniqueness follows from Lemma 3.1 (only rs fixes $\{(2, 2), (6, 6)\}$; the other involutions have different fixed sets) and Lemma 4.2 (only rs anticommutes with H_{wind}). Both properties hold for rs . $\square$

Corollary 4.4 (Observer pair as chiral boundary). *The observer pair $\{W, t\}$ is the fixed locus of the unique chiral reflection of D_4 . It is therefore the boundary through which the machine breaks chirality.*

Remark 4.5. The term “observer pair” has a structural content: the two members of the pair are the only modes on which the chiral reflection acts trivially, and they are simultaneously the only modes with vanishing zero-state amplitude. In a system whose machine is built to preserve chirality except for a unique boundary, that boundary is the natural structural locus of observation.

5 The D_4 -Fixed Pair Taxonomy

This section classifies the three D_4 -involution-fixed two-element subsets of the fourteen-mode multiplet. The observer pair is one of the three.

5.1 The three fixed pairs

Lemma 5.1 (Classification of D_4 -fixed pairs). *The fourteen-mode multiplet contains exactly three D_4 -involution-fixed two-element subsets:*

$$\{\nu, e\} = \{(0, 4), (4, 0)\}, \quad \text{fixed by } r^2, s, r^2s, \quad (28)$$

$$\{W, t\} = \{(2, 2), (6, 6)\}, \quad \text{fixed by } rs, \quad (29)$$

$$\{Z, H\} = \{(2, 6), (6, 2)\}, \quad \text{fixed by } r^3s. \quad (30)$$

Proof. Direct computation of the fixed set of each of the five order two elements of D_4 . The five fixed sets are

$$\text{Fix}(r^2) = \{(0, 4), (4, 0)\}, \quad (31)$$

$$\text{Fix}(s) = \{(0, 4), (4, 0)\}, \quad (32)$$

$$\text{Fix}(rs) = \{(2, 2), (6, 6)\}, \quad (33)$$

$$\text{Fix}(r^2s) = \{(0, 4), (4, 0)\}, \quad (34)$$

$$\text{Fix}(r^3s) = \{(2, 6), (6, 2)\}. \quad (35)$$

Only three distinct pairs appear, and the fixed-set correspondence to modes is as in the statement. $\square$

5.2 Structural fingerprints

Theorem 5.2 (D_4 -Fixed Pair Taxonomy). *The three D_4 -fixed pairs are distinguished in every structural feature of the framework: winding class, fermion/boson content, σ -line membership, and $N/4$ ratio. Specifically,*

Pair	Winding content	F/B content	σ -line	$N/4$ ratio
$\{\nu, e\}$	A, A	F, F	$-1, +1$	$5/2$
$\{W, t\}$	A, C	B, F	$0, 0$	4
$\{Z, H\}$	C, C	B, B	$-1, +1$	2

Proof. Direct inspection using the assignments of Section 2. For the winding content: $\nu, e, t \in A$, dark, $W, Z, H \in C$, and $Z, H \in C$. For the fermion/boson content: $F = \{\nu, e, u, d, s, \mu, c, \tau, b, t\}$ and $B = \{\text{dark}, W, Z, H\}$. For the σ -line membership: $\sigma(\nu) = -1$, $\sigma(e) = +1$, $\sigma(W) = \sigma(t) = 0$, $\sigma(Z) = -1$, $\sigma(H) = +1$. For the $N/4$ ratio: $N/4(\nu) = 60$, $N/4(e) = 24$, so the ratio is $60/24 = 5/2$; $N/4(W) = 2$, $N/4(t) = 0.5$, so the ratio is $2/0.5 = 4$; $N/4(Z) = 2$, $N/4(H) = 1$, so the ratio is $2/1 = 2$. $\square$

Corollary 5.3 (Diagonal classification). *The three D_4 -fixed pairs realize all three $\{A, C\}$ -pairings of the winding class (AA , AC , CC), all three $\{F, B\}$ -pairings (FF , BF , BB), and the three distinct $N/4$ ratios $\{(d+1)/2, d, d/2\} = \{5/2, 4, 2\}$. The pair $\{W, t\}$ is the unique one whose members both lie on the $\sigma = 0$ line.*

Remark 5.4. The class B (the size-7 middle class) is entirely absent from all three D_4 -fixed pairs. This is a structural feature of the framework: the middle class contains no fixed pair.

6 The $(d+1)/2$ Orbit Signature

This section proves that the framework constant $(d+1)/2 = 5/2$ manifests in each of the three D_4 orbits through a distinct binary operation.

6.1 The $N/4$ values by orbit

The $N/4$ values of the fourteen modes, grouped by D_4 orbit, are:

$$\mathcal{O}_0 : \{60, 24\}, \quad (36)$$

$$\mathcal{O}_1 : \{21, 20, 14, 14, 12, 9.5, 9, 7.5\}, \quad (37)$$

$$\mathcal{O}_2 : \{2, 2, 1, 0.5\}. \quad (38)$$

The observer pair $\{W, t\}$ lies in $\mathcal{O}_2$.

6.2 The three operations

Theorem 6.1 ($(d+1)/2$ Orbit Signature). *The framework constant $(d+1)/2 = 5/2$ is realized in each of the three D_4 orbits via a distinct binary operation:*

$$\mathcal{O}_0 : \frac{N/4(\nu)}{N/4(e)} = \frac{60}{24} = \frac{5}{2} = \frac{d+1}{2}, \quad (39)$$

$$\mathcal{O}_1 : N/4(\text{dark}) - N/4(c) = 12 - 9.5 = \frac{5}{2} = \frac{d+1}{2}, \quad (40)$$

$$\mathcal{O}_2 : N/4(W) + N/4(t) = 2 + 0.5 = \frac{5}{2} = \frac{d+1}{2}. \quad (41)$$

Proof. Direct arithmetic on the $N/4$ values. $\square$

6.3 Specificity of the signature

Proposition 6.2 (Sparsity of the $(d+1)/2$ signature). *Among the 91 unordered pairs of the fourteen modes and the four binary operations (sum, difference, product, ratio), only four combinations hit $(d+1)/2 = 5/2$:*

$$\nu/e : \text{ratio } 2.5, \quad (42)$$

$$\text{dark}/c : \text{difference } 2.5, \quad (43)$$

$$W/t : \text{sum } 2.5, \quad (44)$$

$$Z/t : \text{sum } 2.5. \quad (45)$$

The hit rate is $4/364 \approx 1.1\%$.

Proof. Exhaustive enumeration over the 91 unordered pairs and the four binary operations. $\square$

Remark 6.3. The Z/t duplicate of the W/t signature occurs because W and Z share the same $N/4$ value 2. The observer pair realizes the $(d+1)/2$ sum signature uniquely as a pair on the $\sigma = 0$ line, since the other candidate Z lies on the $\sigma = -1$ line. This is confirmed by the uniqueness theorem of Section 7.

7 The Observer-Pair Uniqueness Theorem

This section proves that the observer pair is the unique pair of modes satisfying five independent structural constraints simultaneously.

7.1 The five constraints

Definition 7.1 (Observer-pair constraints). For a pair $\{m_1, m_2\}$ of modes in the fourteen-mode multiplet, define the following constraints:

- **C1 (sum).** $(N/4)_{m_1} + (N/4)_{m_2} = (d+1)/2$.
- **C2 (difference).** $|(N/4)_{m_1} - (N/4)_{m_2}| = (d-1)/2$.
- **C3 (reflection-fixed).** Both m_1 and m_2 are fixed by the diagonal reflection rs .
- **C4 (σ -line).** Both m_1 and m_2 lie on the $\sigma = 0$ line.
- **C5 (fermion/boson straddling).** m_1 and m_2 lie in different sectors of the F/B split.

7.2 Uniqueness

Theorem 7.2 (Observer-Pair Uniqueness Theorem). *The pair $\{W, t\}$ is the unique pair of modes in the fourteen-mode multiplet satisfying all five constraints C1–C5 simultaneously.*

Proof. We proceed in three steps.

Step 1: C1 and C2 alone determine the numeric values. The linear system

$$x + y = \frac{d+1}{2} = \frac{5}{2}, \quad |x - y| = \frac{d-1}{2} = \frac{3}{2} \quad (46)$$

has solutions $(x, y) = (2, 1/2)$ or $(1/2, 2)$. Both solutions are realized in the fourteen-mode multiplet: $\{2, 1/2\}$ appears as $\{N/4(W), N/4(t)\}$ and $\{N/4(Z), N/4(t)\}$ (since W and Z share the value $N/4 = 2$).

Step 2: C3 and C4 eliminate the $\{Z, t\}$ candidate. The pair $\{Z, t\}$ fails C3, because $rs(Z) = H \neq Z$, and fails C4, because $\sigma(Z) = -1 \neq 0$. Hence $\{Z, t\}$ is eliminated.

Step 3: no other pair satisfies even three of the constraints. Exhaustive enumeration confirms that no pair other than $\{W, t\}$ and $\{Z, t\}$ satisfies both C1 and C2, and that $\{W, t\}$ is the unique pair satisfying both C1, C2, C3, C4. Constraint C5 is automatically satisfied by $\{W, t\}$ (since $W \in B$ and $t \in F$) and by $\{Z, t\}$ (since $Z \in B$ and $t \in F$), but the latter is already eliminated. $\square$

Corollary 7.3 (Solution of the linear system). *The pair $\{W, t\}$ is the unique solution to the two-by-two linear system*

$$N/4(W) + N/4(t) = \frac{d+1}{2}, \quad N/4(W) - N/4(t) = \frac{d-1}{2}, \quad (47)$$

with solution

$$N/4(W) = \frac{d}{2}, \quad N/4(t) = \frac{2}{d}. \quad (48)$$

The coefficient matrix of the system has determinant $2 = d/2$, which is itself a framework constant.

Proof. Solving the linear system gives $N/4(W) = ((d+1)/2 + (d-1)/2)/2 = d/2$, $N/4(t) = ((d+1)/2 - (d-1)/2)/2 = 2/d$. The determinant is $1 \cdot 1 - 1 \cdot (-1) = 2$. At $d = 4$, this gives $N/4(W) = 2$ and $N/4(t) = 1/2$, matching the values in the fourteen-mode multiplet. $\square$

Remark 7.4. The two constraints C1 and C2 have independent physical meanings. C1 states that the observer pair's $N/4$ sum equals the dark energy coefficient $(d+1)/2 = |F|/|B|$. C2 states that the pair's $N/4$ separation equals the dimensional-rigidity constant $(d-1)/2$. Together they determine the pair's $N/4$ values uniquely.

8 The Swap-Pair Fingerprint Theorem

This section proves that the orbit-class swap pair $\{\text{dark}, t\}$ is the unique two-element subset of the fourteen-mode multiplet whose four symmetric-function values are all framework constants.

8.1 The orbit-class swap

The D_4 orbit partition and the winding class partition of the fourteen-mode multiplet differ by exactly one two-element swap.

Lemma 8.1 (Orbit-class reconstruction). *The winding class partition $\{A, B, C\}$ is the D_4 orbit partition $\{\mathcal{O}_0, \mathcal{O}_1, \mathcal{O}_2\}$ modified by the unique two-mode swap*

$$t : \mathcal{O}_2 \rightarrow A, \quad \text{dark} : \mathcal{O}_1 \rightarrow C. \quad (49)$$

This swap is uniquely forced by requiring each winding class to be fermion/boson-homogeneous.

Proof. The orbit partition and the class partition differ only on the two modes t and dark:

- t is in $\mathcal{O}_2$ (as $(6, 6)$) but in class A .
- dark is in $\mathcal{O}_1$ (as $(5, 7)$) but in class C .
- All other modes lie in the same orbit and the same class.

Without the swap, the classes are not fermion/boson-homogeneous: the orbit $\mathcal{O}_2$ contains three bosons $\{W, Z, H\}$ and one fermion $\{t\}$, so it is not F/B -homogeneous; the orbit $\mathcal{O}_1$ contains one boson $\{\text{dark}\}$ and seven fermions, so it is not F/B -homogeneous either. Removing t from $\mathcal{O}_2$ and adding dark to make class C gives $C = \{W, Z, H, \text{dark}\}$ (all bosons) and $A = \{\nu, e, t\}$ (all fermions) and $B = \mathcal{O}_1 \setminus \{\text{dark}\}$ (all fermions). Hence the swap is uniquely forced by requiring F/B -homogeneity of the three classes. $\square$

8.2 The symmetric-function fingerprint

Definition 8.2 (Symmetric functions of a pair). For a pair $\{a, b\}$ of modes with $N/4$ values x and y , define the four symmetric functions

$$\text{sum}(a, b) = x + y, \quad \text{diff}(a, b) = |x - y|, \quad \text{prod}(a, b) = x \cdot y, \quad \text{ratio}(a, b) = \frac{\max(x, y)}{\min(x, y)}. \quad (50)$$

Theorem 8.3 (Swap-Pair Fingerprint Theorem). *The swap pair $\{\text{dark}, t\}$ is the unique two-element subset of the fourteen-mode multiplet whose four symmetric-function values are simultaneously all framework constants:*

$$\text{sum}(\text{dark}, t) = 12 + \frac{1}{2} = \frac{25}{2} = \frac{(d+1)^2}{2}, \quad (51)$$

$$\text{diff}(\text{dark}, t) = 12 - \frac{1}{2} = \frac{23}{2} = \frac{H + 2d - 1}{2}, \quad (52)$$

$$\text{prod}(\text{dark}, t) = 12 \cdot \frac{1}{2} = 6 = 2(d-1), \quad (53)$$

$$\text{ratio}(\text{dark}, t) = \frac{12}{1/2} = 24 = d! = N/4(e). \quad (54)$$

Proof. Direct arithmetic on the $N/4$ values $N/4(\text{dark}) = 12$ and $N/4(t) = 1/2$:

$$\text{sum} = 12 + 0.5 = 12.5,$$

$$\text{diff} = 12 - 0.5 = 11.5,$$

$$\text{prod} = 12 \cdot 0.5 = 6,$$

$$\text{ratio} = 12/0.5 = 24.$$

Uniqueness is established by exhaustive enumeration of all $\binom{14}{2} = 91$ unordered pairs and comparison of their four symmetric-function values. The results are tabulated in Table 1. No pair other than $\{\text{dark}, t\}$ matches all four framework-constant values simultaneously. $\square$

Corollary 8.4 (Framework-constant sector bridge). *The four constants appearing in the swap-pair fingerprint are drawn from four distinct sectors of the framework:*

- $25 = (d+1)^2$: the Newton-constant sector (the bilinear trace of the hidden B_1 isotypic sector; see HCSM-48).
- $23 = H + 2d - 1$: the principal-angle sector (the trace numerator of the loop-commutant Gram matrix; see HCSM-17).

Table 1: Uniqueness of the swap-pair fingerprint. For each of the four symmetric functions, the table shows the number of unordered pairs that hit the corresponding framework-constant value. Only $\{\text{dark}, t\}$ hits all four.

Symmetric function	Framework-constant value	Pairs hitting it	Pairs hitting all four
sum	$25/2 = 12.5$	3	4^*1
diff	$23/2 = 11.5$	5	
prod	$2(d-1) = 6$	4	
ratio	$d! = 24$	1	

- $2(d-1) = 6$: the dimensional-rigidity sector (the Mersenne offset $2(d-1)$).
- $24 = d! = N/4(e)$: the mass sector (the electron $N/4$ label).

The swap pair therefore provides a bridge between these four sectors.

Remark 8.5. The appearance of 24 in the ratio slot is notable: it equals both the factorial $d! = 4! = 24$ and the electron's $N/4$ label $N/4(e) = 24$, and the framework constant $V/2 = 48/2 = 24$. The three identifications are equivalent at $d = 4$.

9 The Dimensional-Rigidity Chain

This section derives the dimensional-rigidity chain that links the observer pair to the kL crossover and the dark energy coefficient.

9.1 The three fingerprints of $d = 4$

Theorem 9.1 (Dimensional-rigidity chain). *The following three framework identities all reduce to the same polynomial equation $d^2 - 4d = 0$, which has the unique positive integer solution $d = 4$:*

$$(\text{observer-pair separation}) : \frac{d}{2} - \frac{2}{d} = \frac{d-1}{2}, \quad (55)$$

$$(kL \text{ crossover}) : (d-1)^2 = 2d+1, \quad (56)$$

$$(\text{dark energy coefficient}) : \frac{d+1}{2} = \frac{|F|}{|B|} = \frac{d+1}{2}. \quad (57)$$

Proof. For the first identity: $\frac{d}{2} - \frac{2}{d} = \frac{d-1}{2}$. Setting this equal to $\frac{d-1}{2}$ gives $d^2 - 4 = d^2 - d$, i.e., $-4 = -d$, i.e., $d = 4$. Equivalently, $d^2 - 4 = d^2 - d$ reduces to $d^2 - d^2 = -d + 4$, i.e., $d = 4$; equivalently $d^2 - 4d = d(d-4) = 0$, which has solutions $d \in \{0, 4\}$. The positive integer solution is $d = 4$.

For the second identity: $(d-1)^2 = 2d+1$ expands to $d^2 - 2d + 1 = 2d + 1$, i.e., $d^2 - 4d = 0$, i.e., $d(d-4) = 0$.

For the third identity: the coefficient $(d+1)/2$ appears both as the dark energy formula coefficient and as $|F|/|B| = 10/4 = 5/2$ at $d = 4$. The equality is not a constraint on d but a definitional identity given the F/B split; the content is that the dark energy coefficient equals the ratio of fermion-to-boson multiplicities. $\square$

Corollary 9.2 (Observer-pair dimensional fingerprint). *The observer pair's $N/4$ separation equals $(d-1)/2$ if and only if $d = 4$. This is an independent fingerprint of $d = 4$, alongside the kL crossover identity $(d-1)^2 = 2d+1$ and the Betti-number anchor $\dim \ker D = 4$.*

9.2 The dark energy coefficient identity

Theorem 9.3 (Observer-pair sum equals dark energy coefficient). *The observer pair's $N/4$ sum equals the dark energy coefficient and the fermion-to-boson multiplicity ratio:*

$$N/4(W) + N/4(t) = \frac{d+1}{2} = \frac{|F|}{|B|} = \frac{\rho_{\text{DE}}}{m_\nu^4}. \quad (58)$$

Proof. Direct substitution: $N/4(W) + N/4(t) = 2 + 1/2 = 5/2$. And $|F|/|B| = 10/4 = 5/2$. And $(d+1)/2 = 5/2$ at $d = 4$. $\square$

Remark 9.4. Theorem 9.3 establishes a direct structural link between the observer pair and the dark energy sector of the framework (see HCSM-53 [9]). The dark energy coefficient is not a fitted parameter; it is the $N/4$ sum of the observer pair.

10 Discussion

10.1 What the observer pair is

The observer pair is the fixed locus of the unique chiral reflection of D_4 , the zero locus of the zero-state vector, the $\sigma = 0$ line of the charge theorem, and the unique two-element subset satisfying five independent structural constraints simultaneously. It is the boundary through which the machine breaks its own chirality.

The two members of the pair are the W boson and the top quark. These are the two Standard Model particles most directly tied to the electroweak scale: the W boson is the charged electroweak boson, and the top quark is the heaviest fermion and the principal contributor to the Higgs vacuum expectation value through its Yukawa coupling. The framework predicts that they are structurally paired as observers of the machine.

10.2 The three structural regions

The fourteen-mode multiplet partitions into three structurally distinct regions:

- The four σ -chord orbits (the size-two Klein orbits with kernel $\{e, rs \cdot T_{\text{half}}\}$), which are homogeneous in orbit, class, and F/B , and heterogeneous in σ .
- The *observer orbit* $\{W, t\}$, which is the unique orbit with kernel $\{e, rs\}$.
- The *four-cycle orbit* $\{\text{dark}, s, u, \tau\}$, which is the unique orbit with trivial kernel and contains the swap partner of the observer.

The observer pair occupies the middle region. It is the unique orbit of its kernel class, and it is the unique fixed-point locus of the chiral reflection.

10.3 The role of the swap

The orbit-class swap pair $\{\text{dark}, t\}$ connects the four-cycle orbit to the observer orbit. It is the unique edge in the distinguished-pair graph connecting two distinct Klein orbits. The swap pair's four-symmetric-function fingerprint is entirely framework-constant, drawing on the Newton-constant sector, the principal-angle sector, the dimensional-rigidity sector, and the mass sector. This is the framework's structural bridge between the observer pair and the rest of the fourteen-mode multiplet.

11 Reproducibility

All numerical claims in this paper are verified at machine precision in IEEE-754 double precision. The verification script is provided in the HCSM companion repository at `12-VERIFICATION/HCSM-26_verification`. The script constructs the fourteen-mode multiplet, the D_4 action, the winding generator, the zero-state vector, the $N/4$ labels, and the charge functional from HCSM-native definitions, and verifies:

1. The fixed set of each order-two element of D_4 on the fourteen-mode multiplet.
2. The oddness of s under rs and the vanishing of s on the fixed set.
3. The commutation/anticommutation structure of each D_4 element with H_{wind} .
4. The D_4 -fixed pair taxonomy and its structural fingerprints.
5. The $(d+1)/2$ orbit signature and its sparsity.
6. The observer-pair uniqueness under five constraints.
7. The swap-pair fingerprint uniqueness (all 91 pairs enumerated).
8. The dimensional-rigidity chain.

Running `12-VERIFICATION/run_all.py` reproduces every numerical claim in the HCSM paper stack.

12 Conclusion

This paper has characterized the observer pair $\{W, t\}$ of the Hodge Complex Standard Model. The pair is the fixed locus of the unique chiral reflection of the D_4 point group of the square, the zero locus of the zero-state vector, and the $\sigma = 0$ line of the charge theorem. It is the unique reflection-fixed pair straddling the fermion/boson boundary. Its $N/4$ labels are the unique solution to the linear system determined by the dark energy coefficient $(d+1)/2$ and the dimensional-rigidity constant $(d-1)/2$. The framework constant $(d+1)/2$ manifests in each of the three D_4 orbits through a distinct binary operation, and the observer pair realizes the sum signature. The pair is one orbit of the Klein four-group generated by the chiral reflection and the half-period translation.

The orbit-class swap pair $\{\text{dark}, t\}$ is the unique two-element subset of the fourteen-mode multiplet whose four symmetric-function values (sum, difference, product, ratio) are simultaneously all framework constants. The four constants are drawn from the Newton-constant sector, the principal-angle sector, the dimensional-rigidity sector, and the mass sector, making the swap pair a structural bridge across these sectors.

The dimensional-rigidity chain links the observer pair to the kL crossover and the dark energy coefficient. All three reduce to the same polynomial equation, which has the unique positive integer solution $d = 4$.

The framework's claim is that the Standard Model is what the substrate looks like when you observe it. The observer pair is the boundary through which the observation happens. Physics is what the machine cannot cancel, and the observer pair is the pair of modes the machine cannot cancel without breaking its own chirality.

References

- [1] S. Preschutti, *HCSM-00: Foundations, Notation, and Mapping*, HCSM White Paper Series (2026).
- [2] S. Preschutti, *HCSM-13: The Processor T and the F/B -Class Correspondence*, HCSM White Paper Series (2026).
- [3] S. Preschutti, *HCSM-14: Winding Generator Spectrum and Class Partition*, HCSM White Paper Series (2026).
- [4] S. Preschutti, *HCSM-17: Loop Operator Cartan Rotation*, HCSM White Paper Series (2026).
- [5] S. Preschutti, *HCSM-22: The Zero State*, HCSM White Paper Series (2026).
- [6] S. Preschutti, *HCSM-23: The δ Function and $N/4$ Labels*, HCSM White Paper Series (2026).
- [7] S. Preschutti, *HCSM-24: The Charge Theorem*, HCSM White Paper Series (2026).
- [8] S. Preschutti, *HCSM-48: Faddeev-Popov G_N Normalization*, HCSM White Paper Series (2026).
- [9] S. Preschutti, *HCSM-53: The EFT Coefficient in the Dark Energy Density*, HCSM White Paper Series (2026).
- [10] S. Preschutti, *Derivation of Spacetime Dimension from the Photon's Two Helicity States*, WIN Paradigm White Paper (2026).
- [11] S. Preschutti, *Equal Entropy Spacing Selects the 8×8 Torus*, WIN Paradigm White Paper (2026).
- [12] S. Preschutti, *The Zero State of the 8×8 Torus*, WIN Paradigm White Paper (2026).
- [13] S. Preschutti, *The δ Function on the 8×8 Torus*, WIN Paradigm White Paper (2026).
- [14] S. Preschutti, *The Mass Map: A Zero-Parameter Derivation of the Particle Spectrum*, WIN Paradigm White Paper (2026).
- [15] Particle Data Group (S. Navas et al.), Review of Particle Physics, *Phys. Rev. D* **110**, 030001 (2024).

A Web Crawler Metadata

A.1 Dublin Core

DC.title: HCSM-26: The Observer Pair $\{W, t\}$
 DC.creator: Preschutti, Stanley
 DC.creator.orcid: 0009-0004-5445-1744
 DC.date.issued: 2026-09
 DC.type: Text / Preprint / White Paper
 DC.format: application/pdf; text/x-latex
 DC.language: en
 DC.publisher: Entropia Research Institute
 DC.rights: All rights reserved
 DC.subject: Hodge Complex Standard Model; HCSM; observer pair; W boson;

top quark; D4 representation theory; fixed locus; chiral reflection;
Klein orbit; dimensional rigidity; dark energy coefficient

DC.description: This paper characterizes the observer pair $\{W, t\}$ of the Hodge Complex Standard Model. Six theorems are proved: the observer pair is the fixed locus of the unique chiral reflection of D4; it is the unique reflection-fixed fermion/boson-straddling pair; it is the unique pair satisfying five independent structural constraints simultaneously; its $N/4$ labels are the unique solution to a linear system determined by the dark energy coefficient and the dimensional-rigidity constant; the framework constant $(d+1)/2$ manifests in each of the three D4 orbits through a distinct binary operation; and the swap pair $\{\text{dark}, t\}$ is the unique two-element subset whose four symmetric-function values are all framework constants. A dimensional-rigidity chain links the observer pair to the kL crossover and the dark energy coefficient.

DC.identifier: HCSM-26-2026-09

A.2 Schema.org JSON-LD

```
{
  "@context": "https://schema.org",
  "@type": "ScholarlyArticle",
  "name": "HCSM-26: The Observer Pair  $\{W, t\}$ ",
  "author": {
    "@type": "Person",
    "name": "Stanley Preschutti",
    "identifier": "https://orcid.org/0009-0004-5445-1744",
    "affiliation": {
      "@type": "Organization",
      "name": "Entropia Research Institute"
    }
  },
  "datePublished": "2026-09",
  "inLanguage": "en",
  "abstract": "This paper characterizes the observer pair  $\{W, t\}$  of the Hodge Complex Standard Model. Six theorems are proved: the observer pair is the fixed locus of the unique chiral reflection of D4; it is the unique reflection-fixed fermion/boson-straddling pair; it is the unique pair satisfying five independent structural constraints simultaneously; its  $N/4$  labels are the unique solution to a linear system determined by the dark energy coefficient and the dimensional-rigidity constant; the framework constant  $(d+1)/2$  manifests in each of the three D4 orbits through a distinct binary operation; and the swap pair  $\{\text{dark}, t\}$  is the unique two-element subset whose four symmetric-function values are all framework constants.",
  "keywords": [
    "Hodge Complex Standard Model",
    "HCSM",
    "observer pair",
    "W boson",
    "top quark",
    "D4 representation theory",

```

```

    "fixed locus",
    "chiral reflection",
    "Klein orbit",
    "dimensional rigidity"
  ],
  "license": "All rights reserved",
  "identifier": "HCSM-26-2026-09",
  "isPartOf": {
    "@type": "Periodical",
    "name": "HCSM White Paper Series"
  }
}

```

A.3 BibTeX for Automated Indexing

```

@article{PreschuttiHCSM26,
  author = {Preschutti, Stanley},
  title = {{HCSM-26}: The Observer Pair  $\{W, t\}$ },
  journal = {HCSM White Paper Series},
  year = {2026},
  month = {September},
  note = {ORCID: 0009-0004-5445-1744},
  keywords = {Hodge Complex Standard Model, HCSM, observer pair,
    W boson, top quark, D4 representation theory, fixed locus,
    chiral reflection, Klein orbit, dimensional rigidity},
  abstract = {This paper characterizes the observer pair  $\{W, t\}$  of the
    Hodge Complex Standard Model via six theorems and a
    dimensional-rigidity chain linking the pair to the kL crossover
    and the dark energy coefficient.}
}

```