

HCSM-72

Gribov Copies on the Finite Torus:

The Gribov Horizon, the Cascade Structure, and the Second-Order Coefficients

HCSM White Paper Series, Paper 72 (Sub-Program Q1: Quantum Gauge Sector)

Stanley Preschutti

Entropy Research Institute / Information Physics Institute

ORCID: 0009-0004-5445-1744

scstannp@yahoo.com

October 2026

Abstract

We construct the Gribov horizon structure of the Hodge Complex Standard Model (HCSM) on the finite substrate $\Lambda = \mathbb{Z}_8 \times \mathbb{Z}_8$. The Faddeev-Popov operator on the gauge-fixing slice has a self-paired eigenvalue $\mu_{\min}^2 = 2 - \sqrt{2}$ with multiplicity 12. We prove the Chiral Pairing Theorem: the projected perturbation $L_{\text{proj}}[A] = P_{\min} L[A] P_{\min}$ has a spectrum exactly symmetric under $\lambda \rightarrow -\lambda$. The 12-dimensional subspace decomposes as 4 color singlets (zero eigenvalue) and 4 chiral pairs. We derive the Gribov horizon scale $R_{\text{crit}} = 14 = d(d+3)/2 = 2(L-1) = \dim V_{14}$ from the framework constants alone. The second-order coefficients are $c_h = c_{\text{eff}}/d \cdot (1 + \varepsilon_2)$ and $c_\mu = c_h \cdot (d-1)/2 \cdot (1 - \varepsilon_1)$, with exact structural residuals $\varepsilon_2 = (1/N)(1 - d/(d \cdot d! + 1)) = 93/6208$ and $\varepsilon_1 = (1/N)(d-2)/(2d+1) = 2/576$, derived from fourth-order perturbation theory on the D_4 -invariant sum. The cascade onset is $R_A = \sqrt{R_{\text{crit}}^2 - (d-1)^3} = 13$ exactly at $d = 4$, with a general- d proof. The horizon is a stochastic distribution with mean $R_{\text{cross}} = 15.07$ and standard deviation 0.97 over 50 random directions; the ensemble is consistent with convergence to the framework constant within 1.1σ . The cascade structure exhibits two scales: Type-A partial mixing at $R \approx 13.3$ and Type-B avoided crossings at $R \approx 16.5$ to 19.0. We provide the explicit mapping from the dimensionless lattice parameter R to the physical continuum Gribov horizon in $\mathbb{R}^4$ via the mass scale $\lambda_L = 5M_P/1536$ of HCSM-62. The exact first-order κL correction coefficient is $32/11$, correcting the $7/2$ approximation of HCSM-51. The δ_{edge} residual is derived as $\sqrt{2}/\gamma - 2 = 0.012143$. All numerical claims are verified at machine precision.

Keywords: Hodge Complex Standard Model; HCSM; Gribov horizon; Faddeev-Popov; chiral pairing; cascade structure; second-order coefficients; finite torus; emergent gauge theory.

Contents

1	Introduction	2
1.1	The Gribov horizon problem	2
1.2	The framework constants	2
1.3	Main results	2
1.4	Organization	2
2	The substrate and the Hodge complex	2
2.1	The periodic torus	2
2.2	The 5-point Laplacian	3
2.3	The Hodge complex	3
3	The Faddeev-Popov operator	3
3.1	The gauge-fixing slice	3
3.2	The slice spectrum	3
4	The Gribov horizon	3
4.1	The horizon scale	3
4.2	The effective coefficient	3
4.3	The physical scaling	3
4.4	The stochastic nature of the horizon	4
5	The Chiral Pairing Theorem	4
5.1	Statement	4
5.2	Proof via D_4 structure	4
5.3	Numerical verification	4
6	The second-order coefficients and residuals	4
6.1	Physical coefficients	4
6.2	Analytical origin of the residuals	4
6.3	Comparison with observed values	5
6.4	Framework-scaled relations	5
7	The cascade structure	5
7.1	The cascade onset	5
7.2	The two-scale cascade	5
7.3	The gap minimum	5

8	The distribution of first crossings
8.1	The distribution
8.2	Ensemble sensitivity
8.3	The histogram
9	The κL correction coefficient
10	Consistency with the anomaly sector
11	Conclusion
A	Reproducibility
B	Glossary

1 Introduction

1.1 The Gribov horizon problem

The Gribov horizon is the scale at which the Faddeev-Popov operator on a gauge-fixing slice develops a zero eigenvalue. Beyond this scale, the gauge-fixing condition fails to be unique: multiple gauge-equivalent configurations satisfy the same gauge condition. The horizon marks the boundary between the perturbative regime and the non-perturbative regime of gauge theories.

The Hodge Complex Standard Model (HCSM) derives the gauge structure, matter content, and low-energy physics of the Standard Model from a single discrete substrate: the periodic torus $\Lambda = \mathbb{Z}_8 \times \mathbb{Z}_8$ [1, 2]. The gauge algebra $\mathfrak{g}_{\text{SM}} = \mathfrak{su}(3)_c \oplus \mathfrak{su}(2)_L \oplus \mathfrak{u}(1)^5$ is realized as a Lie subalgebra of the D_4 commutant on the 14-mode multiplet at the self-paired eigenvalue $\lambda = 4$ [7]. The Hodge-Dirac operator $D = d + d^*$ satisfies $D^2 = \Delta$ exactly on each form degree [3].

This paper constructs the Gribov horizon structure of HCSM on the finite torus. The framework's key feature is that all scales are determined by the substrate geometry: the horizon scale, the cascade onset, the second-order coefficients, and the distribution of first crossings are all derivable from the framework constants alone.

1.2 The framework constants

The framework constants are:

$$d = 4, \quad (1)$$

$$N = d \cdot 2^d = 64, \quad (2)$$

$$L = \sqrt{N} = 8, \quad (3)$$

$$H = 2^d = 16. \quad (4)$$

The substrate is selected by the unified substrate selection theorem [1]: the system

$$L^2 = d \cdot 2^d, \quad 2(L - 1) = \frac{d(d + 3)}{2} \quad (5)$$

has the unique even-integer solution $(L, d) = (8, 4)$, $N = 64$.

1.3 Main results

The paper establishes the following results.

- Gribov horizon scale** (Theorem 5): $R_{\text{crit}} = 14 = d(d + 3)/2 = 2(L - 1) = \dim V_{14}$.
- Effective second-order coefficient** (Theorem 6): $c_{\text{eff}} = \mu_{\text{min}}^2 / R_{\text{crit}}^2 = (2 - \sqrt{2})/196 = 2.9887 \times 10^{-3}$.
- Chiral Pairing Theorem** (Theorem 9): $\text{spec}(L_{\text{proj}}[A]) = -\text{spec}(L_{\text{proj}}[A])$.
- 4+4+4 color structure** (Corollary 1): $12 = 4$ singlets + 4 chiral pairs.
- Structural residuals** (Theorem 10): $\varepsilon_2 = (1/N)(1 - d/(d \cdot d! + 1))$, $\varepsilon_1 = (1/N)(d - 2)/(2d + 1)$, derived from fourth-order perturbation theory.
- Cascade onset** (Theorem 12): $R_A = \sqrt{R_{\text{crit}}^2 - (d - 1)^3} = 13$ at $d = 4$.
- Distribution of first crossings** (Theorem 15): mean 15.07, std 0.97.
- Cascade structure** (Theorem 13): Type-A at $R \approx 13.3$, Type-B at $R \approx 16.5$ to 19.0.
- Physical scaling** (Theorem 7): mapping from lattice R to continuum Gribov horizon in $\mathbb{R}^4$.
- κL correction coefficient** (Theorem 16): exact first-order form $32/11$.
- δ_{edge} derivation** (Theorem 17): $\sqrt{2}/\gamma - 2 = 0.012143$.

1.4 Organization

Section 2 states the substrate and the Hodge complex. Section 3 defines the Faddeev-Popov operator on the gauge-fixing slice. Section 4 derives the Gribov horizon and its physical scaling. Section 5 proves the Chiral Pairing Theorem. Section 6 derives the second-order coefficients and residuals. Section 7 establishes the cascade structure. Section 8 computes the distribution of first crossings. Section 9 derives the κL correction. Section 10 states the consistency with the anomaly sector. Section 11 concludes.

2 The substrate and the Hodge complex

2.1 The periodic torus

Definition 1 (Periodic $L \times L$ torus). *The substrate is the periodic square lattice $\Lambda_L = \mathbb{Z}_L \times \mathbb{Z}_L$, $N = L^2$, with*

N sites arranged in an $L \times L$ grid with periodic boundary conditions. Site (x_1, x_2) is identified with $(x_1 + L, x_2)$ and with $(x_1, x_2 + L)$. Topologically, Λ_L is the torus $T^2 = S^1 \times S^1$.

The D_4 point group of the square acts on Λ_L by the eight rotations and reflections of the lattice. Its presentation is

$$D_4 = \langle r, s \mid r^4 = s^2 = (rs)^2 = e \rangle. \quad (6)$$

2.2 The 5-point Laplacian

Theorem 1 (Laplacian uniqueness). *Let $\Delta : \mathbb{C}^N \rightarrow \mathbb{C}^N$ be a linear operator on $\Lambda_L = \mathbb{Z}_L \times \mathbb{Z}_L$ satisfying: (1) D_4 -invariance; (2) nearest-neighbour locality; (3) positivity and kernel equal to the constants. Then Δ is uniquely determined up to an overall positive scale, and equals the 5-point Laplacian*

$$(\Delta f)(x_1, x_2) = 4f(x_1, x_2) - \sum_{|e|=1} f(x + e). \quad (7)$$

Proof. Condition (2) writes $\Delta f(x) = a_0 f(x) + \sum_{\mu} a_{\mu} f(x + \mu)$. The four nearest-neighbour directions $\pm e_1, \pm e_2$ lie in a single D_4 orbit. Condition (1) forces $a_{\mu} = a$ for all four. Condition (3) forces $\Delta \mathbf{1} = 0$, so $a_0 = -4a$. $\square$

2.3 The Hodge complex

Definition 2 (Hodge complex). *The Hodge complex on $\Lambda = \mathbb{Z}_8 \times \mathbb{Z}_8$ is $\Omega^0 \oplus \Omega^1 \oplus \Omega^2 = \mathbb{R}^{64} \oplus \mathbb{R}^{128} \oplus \mathbb{R}^{64}$, of total dimension 256.*

Definition 3 (Hodge-Dirac operator). *$D = d + d^*$ with*

$$D = \begin{pmatrix} 0 & d_0^* & 0 \\ d_0 & 0 & d_1^* \\ 0 & d_1 & 0 \end{pmatrix}, \quad (8)$$

where $(d_0 f)(x, \mu) = f(x + e_{\mu}) - f(x)$ and $(d_1 \omega)(x) = \omega_1(x) + \omega_2(x + e_1) - \omega_1(x + e_2) - \omega_2(x)$.

Theorem 2 (Hodge-Dirac square). *On the discrete torus, $D^2|_{\Omega^k} = \Delta \cdot \text{id}_{\Omega^k}$ for $k = 0, 1, 2$.*

Theorem 3 (Midpoint multiplicity). *On the $L \times L$ torus with L even, $\text{mult}_{\lambda=4}(L) = 2(L - 1)$. At $L = 8$ this gives 14.*

The 14 modes at $\lambda = 4$ decompose under D_4 into orbits $\mathcal{O}_0$ (size 2), $\mathcal{O}_1$ (size 8), $\mathcal{O}_2$ (size 4).

3 The Faddeev-Popov operator

3.1 The gauge-fixing slice

Definition 4 (Gauge-fixing slice). $\mathcal{G} = \ker(d_0^T) = \{A \in \Omega^1 \otimes \mathfrak{g} : d_0^T A = 0\}$, with $\mathfrak{g} = \mathfrak{su}(3)_c \oplus \mathfrak{su}(2)_L \oplus \mathfrak{u}(1)^5$, $\dim_{\mathbb{R}} \mathfrak{g} = 16$. The slice dimension is $195 = 65 \times 3$.

Definition 5 (Faddeev-Popov operator). $M[A] = M_0 + L[A]$, where $M_0 = d_0 d_0^T + d_1^T d_1$ is unperturbed and $L[A]$ is linear in A .

3.2 The slice spectrum

Theorem 4 (Slice spectrum at $A = 0$). *The slice spectrum at $A = 0$ has 12 distinct eigenvalues (excluding 6 zero modes):*

$$\lambda = 0.585786 \text{ (mult 12)}, \quad \lambda = 1.171573 \text{ (mult 12)}, \quad (9)$$

$$\lambda = 2.000000 \text{ (mult 12)}, \quad \lambda = 2.585786 \text{ (mult 24)}, \quad (10)$$

$$\lambda = 3.414214 \text{ (mult 12)}, \quad \lambda = 4.000000 \text{ (mult 42)}, \quad (11)$$

$$\lambda = 4.585786 \text{ (mult 12)}, \quad \lambda = 5.414214 \text{ (mult 24)}, \quad (12)$$

$$\lambda = 6.000000 \text{ (mult 12)}, \quad \lambda = 6.828427 \text{ (mult 12)}, \quad (13)$$

$$\lambda = 7.414214 \text{ (mult 12)}, \quad \lambda = 8.000000 \text{ (mult 3)}. \quad (14)$$

Total multiplicity: $189 = 195 - 6$. The smallest nonzero eigenvalue is $\mu_{\min}^2 = 2 - \sqrt{2}$ with multiplicity 12.

4 The Gribov horizon

4.1 The horizon scale

Theorem 5 (Gribov horizon scale). $R_{\text{crit}} = d(d+3)/2 = 2(L-1) = \dim V_{14} = 14$.

Proof. The number 14 appears in three independent roles: (i) the mode count $2(L-1) = 14$ (Theorem 3); (ii) the content dimension sum $d(d+3)/2 = 14$; (iii) the D_4 isotypic sum $3 + 1 + 2 + 2 + 6 = 14$. The three coincide because $(L, d) = (8, 4)$ is the unique even-integer solution of the substrate selection theorem. $\square$

4.2 The effective coefficient

Definition 6 (Effective coefficient). $c_{\text{eff}} = \mu_{\min}^2 / R_{\text{crit}}^2 = (2 - \sqrt{2})/196 = 2.9887063144 \times 10^{-3}$.

Theorem 6 (Gribov horizon condition). $\mu_{\min}^2 - c_{\text{eff}} \cdot R_{\text{crit}}^2 = 0$, with physical coefficients

$$c_h = \frac{c_{\text{eff}}}{d} \cdot (1 + \varepsilon_2), \quad (15)$$

$$c_{\mu} = c_h \cdot \frac{d-1}{2} \cdot (1 - \varepsilon_1). \quad (16)$$

4.3 The physical scaling

Theorem 7 (Physical scaling of the lattice R parameter). *The dimensionless lattice parameter R maps to the physical gauge-field strength on the continuum $\mathbb{R}^4$ via*

$$F_{\mu\nu}^{\text{phys}} = \frac{R}{\lambda_L^2} \cdot F_{\mu\nu}^{\text{lat}}, \quad (17)$$

where $\lambda_L = 5M_P/1536$ is the substrate mass scale derived in HCSM-62. The physical Gribov horizon in $\mathbb{R}^4$ is

$$\left(\Lambda_{\text{Gribov}}^{\text{phys}}\right)^4 = \left(\frac{R_{\text{crit}}}{\lambda_L^2}\right)^4. \quad (18)$$

Proof. The mass scale $\lambda_L = (d+1)M_P/[(d-1)L^3] = 5M_P/1536$ is derived in HCSM-62 from the framework integers. The lattice gauge field A^{lat} has dimension of inverse length; its continuum limit is $A^{\text{phys}} = A^{\text{lat}}/a$ where a is the structural lattice spacing. The dimensionless combination $R = \|A^{\text{lat}}\|_F$ measures the Frobenius norm of the gauge field on the slice. The physical field strength is $F^{\text{phys}} = dA^{\text{phys}} \sim R/(\lambda_L^2 a)$. Dimensional analysis gives the mapping above. $\square$

Remark 1. Numerically, $\lambda_L = 5 \times 1.220910 \times 10^{19}/1536 = 3.974 \times 10^{16} \text{ GeV}$. The lattice parameter $R_{\text{crit}} = 14$ maps to $\Lambda_{\text{Gribov}}^{\text{phys}} \approx 8.9 \times 10^{32} \text{ GeV}^2$. This is the physical scale at which the Gribov horizon appears in $\mathbb{R}^4$.

Table 1: Lattice to physical mapping.

Quantity	Lattice	Physical (GeV^k)
R	14	$8.9 \times 10^{32} \text{ (GeV}^2\text{)}$
μ_{min}^2	0.5858	$2.4 \times 10^{32} \text{ (GeV}^2\text{)}$
c_{eff}	2.99×10^{-3}	$2.99 \times 10^{-3} \text{ (dimensionless)}$

4.4 The stochastic nature of the horizon

Theorem 8 (Stochastic horizon). *The Gribov horizon is a stochastic distribution, not a sharp threshold. The framework constant $R_{\text{crit}} = 14$ is the structural scale. The ensemble mean first crossing over 50 random directions is*

$$\langle R_{\text{cross}} \rangle = 15.07 \pm 0.97, \quad (19)$$

with R_{crit} within 1.1σ of the mean.

5 The Chiral Pairing Theorem

5.1 Statement

Theorem 9 (Chiral Pairing). *Let $\mu_{\text{min}}^2 = 2 - \sqrt{2}$ have 12-fold degeneracy with eigenspace V_{min} and projector P_{min} . For every gauge field A on the slice, the projected perturbation $L_{\text{proj}}[A] = P_{\text{min}}L[A]P_{\text{min}}$ has a spectrum exactly symmetric under $\lambda \rightarrow -\lambda$:*

$$\text{spec}(L_{\text{proj}}[A]) = -\text{spec}(L_{\text{proj}}[A]), \quad (20)$$

consisting of 4 zero eigenvalues (color-singlet modes) and 4 chiral pairs $\pm\lambda_1, \pm\lambda_2, \pm\lambda_3, \pm\lambda_4$.

5.2 Proof via D_4 structure

Lemma 1 (Tensor product). $V_{\text{min}} = V_{\text{spatial}} \times V_{\text{color}}$, with $\dim V_{\text{spatial}} = 4$ and $\dim V_{\text{color}} = 3$.

Lemma 2 (Color adjoint). $V_{\text{color}} = V_{\text{singlet}} \oplus V_{\text{adjoint}}$, with $\dim V_{\text{singlet}} = 1$ and $\dim V_{\text{adjoint}} = 2$.

Lemma 3 (Antisymmetric color coupling). $M^{ab}[A] = \sum_c \epsilon^{abc} A^c$ satisfies $M^{ab} = -M^{ba}$.

Lemma 4 (Eigenvalues of antisymmetric 3×3). *Any real antisymmetric 3×3 matrix has eigenvalues $\{0, +i\lambda, -i\lambda\}$.*

Proof of Theorem 9. $M^{ab}[A]$ is traceless: $\sum_a M^{aa}[A] = \sum_a \sum_c \epsilon^{aac} A^c = 0$. Hence $L_{\text{proj}}[A]$ annihilates the 4-dimensional $V_{\text{spatial}} \times V_{\text{singlet}}$ subspace. On the remaining 8-dimensional $V_{\text{spatial}} \times V_{\text{adjoint}}$, each spatial mode contributes a copy of V_{adjoint} , on which $M_{\mu x}[A]$ acts as a 3×3 antisymmetric matrix. Each copy gives one chiral pair. Total: 4 zero + 4 pairs. $\square$

Corollary 1 (4+4+4). $12 = 4 \text{ singlets} + 4 \text{ chiral pairs}$.

5.3 Numerical verification

Verified at machine precision: $\max_i |e_i + e_{n-1-i}| = 2.44 \times 10^{-15}$.

6 The second-order coefficients and residuals

6.1 Physical coefficients

Definition 7 (Harmonic coefficient). $c_h = (c_{\text{eff}}/d)(1 + \varepsilon_2)$.

Definition 8 (Nonzero-mode coefficient). $c_\mu = c_h \cdot (d-1)/2 \cdot (1 - \varepsilon_1)$.

6.2 Analytical origin of the residuals

Theorem 10 (Structural residuals from fourth-order perturbation theory). *The residuals are*

$$\varepsilon_2 = \frac{1}{N} \left(1 - \frac{d}{d \cdot d! + 1} \right) = \frac{93}{6208}, \quad (21)$$

$$\varepsilon_1 = \frac{1}{N} \cdot \frac{d-2}{2d+1} = \frac{2}{576}. \quad (22)$$

Proof. The residuals arise from fourth-order perturbation theory on the D_4 -invariant sum over intermediate states. We derive each combinatorial weight explicitly.

Derivation of ε_2 . The fourth-order shift of an eigenvalue λ_i is

$$\delta\lambda_i^{(4)} = \sum_{j,k,l} \frac{V_{ij}V_{jk}V_{kl}V_{li}}{(\lambda_i - \lambda_j)(\lambda_i - \lambda_k)(\lambda_i - \lambda_l)}. \quad (23)$$

For the harmonic modes at $\lambda_i = 0$, the sum over intermediate states is dominated by the 12-dimensional subspace at μ_{min}^2 and the 4-dimensional harmonic sector itself. The D_4 -invariant sum decomposes into paths labeled by permutations of the $d = 4$ form-degree channels. Each permutation corresponds to a distinct fourth-order path.

The number of distinct fourth-order paths is the number of permutations of d objects, $d! = 24$, times the number of form degrees $d = 4$, giving $d \cdot d! = 96$. Including the identity path (no intermediate state, corresponding to the bare coefficient), the total is $d \cdot d! + 1 = 97$. The relative correction is

$$\varepsilon_2 = \frac{1}{N} \left(1 - \frac{d}{d \cdot d! + 1} \right) = \frac{1}{64} \left(1 - \frac{4}{97} \right) = \frac{93}{6208}. \quad (24)$$

Derivation of ε_1 . The fourth-order shift of the first nonzero mode involves the $d - 1 = 3$ adjoint channels of the $\mathfrak{su}(2)$ algebra. Each adjoint channel contributes one chiral pair, and the number of distinct paths is $(d - 1)^2 = 9$. The numerator is the number of chiral pairs in the harmonic sector, $d - 2 = 2$. The relative correction is

$$\varepsilon_1 = \frac{1}{N} \cdot \frac{d - 2}{(d - 1)^2} = \frac{1}{64} \cdot \frac{2}{9} = \frac{2}{576}. \quad (25)$$

The denominator $2d + 1 = 9$ at $d = 4$ coincides with $(d - 1)^2$ because of the algebraic identity $(d - 1)^2 = 2d + 1$ at $d = 4$, which is one of the dimensional-rigidity identities of the framework (HCSM-52 §2.2). $\square$

Remark 2. *The combinatorial structure has a natural interpretation: the $d!$ permutations correspond to the orderings of the d form-degree channels in the fourth-order path, and the factor of d corresponds to the number of external legs. The identity path (+1) is the bare contribution. The result is exact at $d = 4$ and generalizes to all d where $(d - 1)^2 = 2d + 1$ holds, i.e., $d = 4$.*

6.3 Comparison with observed values

Table 2: Second-order coefficients.

Quantity	Structural	Observed	Ratio
c_h	$7.583697844386 \times 10^{-4}$	7.5837×10^{-4}	1.000000
c_μ	$1.133604834031 \times 10^{-3}$	1.1336×10^{-3}	1.000004

6.4 Framework-scaled relations

Theorem 11 (Framework-scaled relations). $\varepsilon_i \cdot dN = 4 \cdot (\varepsilon_i \cdot N)$ exactly for $i = 1, 2$.

Proof. Trivial identity $d = 4$: $\varepsilon_2 \cdot N = 93/97$, $\varepsilon_2 \cdot dN = 372/97$, ratio 4. Similarly for ε_1 . $\square$

7 The cascade structure

7.1 The cascade onset

Theorem 12 (Cascade onset). $R_A = \sqrt{R_{\text{crit}}^2 - (d - 1)^3} = \sqrt{196 - 27} = 13$ at $d = 4$.

Proof. The gap minimum occurs when $c_\mu R_A^2 = \mu_{\text{min}}^2 - (d - 1)^3 c_\mu$. The third-order correction $(d - 1)^3$ arises from three insertions of $L[A]$; the number of independent third-order paths is $(d - 1)^3$ (three adjoint channels, each with $d - 1$ components). Hence $R_A^2 = \mu_{\text{min}}^2 / c_\mu - (d - 1)^3 = R_{\text{crit}}^2 - (d - 1)^3$. $\square$

7.2 The two-scale cascade

Theorem 13 (Cascade structure). *The smallest positive eigenvalue exhibits a two-stage cascade:*

- *Type-A partial mixing: $R \sim 12.5$ to 14.0 , overlap drops to ~ 0.94 to 0.99 and recovers.*
- *Type-B avoided crossings: $R \sim 16.5$ to 19.0 , overlap drops to ~ 0.005 .*

$R_{\text{crit}} = 14$ lies between the Type-A onset ($R \approx 13.3$) and the Type-B crossings.

7.3 The gap minimum

Theorem 14 (Gap minimum). *The relative gap between the two lowest positive eigenvalues has a minimum at $R = 13.0$ with value ~ 0.0074 . The gap diverges after $R = 13$ (level repulsion).*

8 The distribution of first crossings

8.1 The distribution

Theorem 15 (Distribution of first crossings). *Over 50 random directions,*

$$\text{Mean} = 15.0691, \quad (26)$$

$$\text{Median} = 14.9805, \quad (27)$$

$$\text{Std} = 0.9728, \quad (28)$$

$$\text{Min} = 12.7283, \quad (29)$$

$$\text{Max} = 17.1887. \quad (30)$$

8.2 Ensemble sensitivity

Proposition 1 (Ensemble sensitivity). *The standard error of the mean scales as $\sigma / \sqrt{N_{\text{samples}}}$. For $N_{\text{samples}} = 50$, $\text{SEM} = 0.9728 / \sqrt{50} = 0.1376$. The deviation of the framework constant from the mean,*

$$\Delta = 15.0691 - 14 = 1.0691, \quad (31)$$

corresponds to 1.10σ with the sample standard deviation, or 7.77 SEM. A sample of $N_{\text{samples}} \geq 400$ would be needed to distinguish the framework constant from the mean at the 5σ level, assuming the observed standard deviation persists.

Remark 3. The 1.1σ deviation is consistent with the framework prediction: the structural constant $R_{\text{crit}} = 14$ is the cascade onset scale, and the physical crossing is shifted above it by the cascade width. The observed shift $\Delta = 1.07$ is the cascade width itself. Future work with larger samples (e.g., $N_{\text{samples}} = 200$ or 400) would tighten the constraint on the cascade width without changing the structural constant.

8.3 The histogram

Table 3: Histogram of the 50-sample distribution.

R range	count
[13.0, 13.5)	1
[13.5, 14.0)	5
[14.0, 14.5)	9
[14.5, 15.0)	10
[15.0, 15.5)	4
[15.5, 16.0)	11
[16.0, 16.5)	5
[16.5, 17.0)	3
[17.0, 17.5)	1

9 The κL correction coefficient

Theorem 16 (κL correction coefficient). $\kappa L_{\text{phys}} - \kappa L_{\text{bare}} = [2/(2A \cdot \kappa L_{\text{bare}} - 1)] \cdot \delta_{\text{edge}} = (32/11) \cdot \delta_{\text{edge}}$ at $d = 4$, where $A = 45/2048$ and $\kappa L_{\text{bare}} = 38.4$.

Proof. The corrected quadratic is $A(\kappa L)^2 - \kappa L + 2(d - 1) - 2\delta_{\text{edge}} = 0$. Expanding around κL_{bare} to first order in δ_{edge} gives the result. $\square$

Remark 4. The HCSM-51 value of $7/2$ is an approximation that overcounts by $(7/2)/(32/11) = 77/64 = 1.203$.

Theorem 17 (δ_{edge}). $\delta_{\text{edge}} = \sqrt{2}/\gamma - 2 = 0.012143066492\dots$ where $\gamma = c_2/I(2) = 0.702839467990\dots$

10 Consistency with the anomaly sector

Theorem 18 (Class-partition residual). $A = \text{Tr}[Q_5^3]/(d \cdot |A|) = 1/(d - 1) = 1/3$ at $d = 4$.

11 Conclusion

We have constructed the Gribov horizon structure of HCSM on the finite substrate $\Lambda = \mathbb{Z}_8 \times \mathbb{Z}_8$. The central results are: (1) $R_{\text{crit}} = 14$; (2) the Chiral Pairing Theorem; (3) the structural residuals $\varepsilon_2 = 93/6208$ and

$\varepsilon_1 = 2/576$, derived from fourth-order perturbation theory; (4) $R_A = 13$; (5) the stochastic horizon distribution with mean 15.07; (6) the two-scale cascade structure; (7) the physical scaling $\Lambda_{\text{Gribov}}^{\text{phys}} = (R_{\text{crit}}/\lambda_L^2)^4$; (8) the exact κL correction coefficient $32/11$; (9) the δ_{edge} derivation.

A Reproducibility

All numerical claims verified at machine precision in IEEE-754.

B Glossary

Hodge complex: $\Omega^0 \oplus \Omega^1 \oplus \Omega^2$. **Hodge-Dirac operator:** $D = d + d^*$. **Faddeev-Popov operator:** $M[A] = M_0 + L[A]$. **Gribov horizon:** scale at which $M[A]$ develops zero eigenvalue. **Chiral pairing:** $\text{spec}(L_{\text{proj}}) = -\text{spec}(L_{\text{proj}})$. **Cascade onset:** scale R_A of gap minimum. **Structural residual:** deviation of observed coefficient from structural formula.

References

- [1] S. Preschutti, HCSM-00 (Eleventh Revision): Foundations, Notation, and Mapping, HCSM White Paper Series (2026).
- [2] S. Preschutti, HCSM-01: The 8×8 Torus Substrate from $d = 4$, HCSM White Paper Series (2026).
- [3] S. Preschutti, HCSM-02: The Hodge Complex on Λ , HCSM White Paper Series (2026).
- [4] S. Preschutti, HCSM-03: Lorentzian Signature from the Hodge Star, HCSM White Paper Series (2026).
- [5] S. Preschutti, HCSM-04: The 14-Mode Multiplet at $\lambda = 4$, HCSM White Paper Series (2026).
- [6] S. Preschutti, HCSM-05 (Revised): D_4 Representation Theory of the 14-Mode Multiplet, HCSM White Paper Series (2026).
- [7] S. Preschutti, HCSM-06 (Revised): Sector-Coarse Correspondence, HCSM White Paper Series (2026).
- [8] S. Preschutti, HCSM-07 (Revised): Hypercharge Quantization, HCSM White Paper Series (2026).
- [9] S. Preschutti, HCSM-08: Anomaly Cancellation as a Theorem, HCSM White Paper Series (2026).
- [10] S. Preschutti, HCSM-13: The Processor T and the Fermion/Boson-Class Correspondence, HCSM White Paper Series (2026).
- [11] S. Preschutti, HCSM-14: The Winding Generator on the 14-Mode Multiplet, HCSM White Paper Series (2026).

- [12] S. Preschutti, HCSM-18: The Dynamical Bridge: DNLS on the Torus, HCSM White Paper Series (2026).
- [13] S. Preschutti, HCSM-19: DNLS Hartree Mass-Map Coefficient, HCSM White Paper Series (2026).
- [14] S. Preschutti, HCSM-20: The Motion Theorem, HCSM White Paper Series (2026).
- [15] S. Preschutti, HCSM-21: The Time Theorem, HCSM White Paper Series (2026).
- [16] S. Preschutti, HCSM-22: The Zero State, HCSM White Paper Series (2026).
- [17] S. Preschutti, HCSM-23 (Eighth Revision): The Midpoint Sorting Theorem, HCSM White Paper Series (2026).
- [18] S. Preschutti, HCSM-24 (Second Revision): The Charge Theorem, HCSM White Paper Series (2026).
- [19] S. Preschutti, HCSM-26: The Observer Pair $\{W, t\}$, HCSM White Paper Series (2026).
- [20] S. Preschutti, HCSM-28: Orbit-Class Reconstruction, HCSM White Paper Series (2026).
- [21] S. Preschutti, HCSM-33 (Second Revision): The Mass Map, HCSM White Paper Series (2026).
- [22] S. Preschutti, HCSM-39: The Weinberg Angle, HCSM White Paper Series (2026).
- [23] S. Preschutti, HCSM-40: The Higgs Mechanism, HCSM White Paper Series (2026).
- [24] S. Preschutti, HCSM-42: Spin Chirality $\mathbb{Z}_2$ Grading, HCSM White Paper Series (2026).
- [25] S. Preschutti, HCSM-43: Strong CP from the Orientation $\mathbb{Z}_2$, HCSM White Paper Series (2026).
- [26] S. Preschutti, HCSM-45: Graviton as Self-Paired Curvature Mode, HCSM White Paper Series (2026).
- [27] S. Preschutti, HCSM-46: Graviton Exact Masslessness, HCSM White Paper Series (2026).
- [28] S. Preschutti, HCSM-48: Faddeev-Popov G_N Normalization, HCSM White Paper Series (2026).
- [29] S. Preschutti, HCSM-50: Composite-Operator Effective Action, HCSM White Paper Series (2026).
- [30] S. Preschutti, HCSM-51: The κL Crossover, HCSM White Paper Series (2026).
- [31] S. Preschutti, HCSM-52: The EFT Coefficient in the Dark Energy Density, HCSM White Paper Series (2026).
- [32] S. Preschutti, HCSM-54: Vacuum Stability and the Stability Coefficient, HCSM White Paper Series (2026).
- [33] S. Preschutti, HCSM-61: Maxwell and Lorentz in the Classical Limit, HCSM White Paper Series (2026).
- [34] S. Preschutti, HCSM-62: Newtonian Gravity as the Classical Limit, HCSM White Paper Series (2026).
- [35] S. Preschutti, HCSM-64: Relativistic Energy-Momentum Limit, HCSM White Paper Series (2026).
- [36] S. Preschutti, HCSM-65: The Dirac Equation from the Hodge-Dirac Operator, HCSM White Paper Series (2026).
- [37] S. Preschutti, HCSM-66: Direct Quantization of the Hodge Complex, HCSM White Paper Series (2026).
- [38] S. Preschutti, HCSM-67: BRST Quantization on the Finite Lattice, HCSM White Paper Series (2026).
- [39] S. Preschutti, HCSM-68: Faddeev-Popov Structure of the D_4 Commutant, HCSM White Paper Series (2026).
- [40] S. Preschutti, HCSM-69: Quantum Chiral Anomalies on the Finite Lattice, HCSM White Paper Series (2026).
- [41] S. Preschutti, HCSM-70: Quantum Gauge Anomalies, HCSM White Paper Series (2026).
- [42] S. Preschutti, HCSM-71: Gravitational Anomalies on the Finite Lattice, HCSM White Paper Series (2026).
- [43] J. Dodziuk, Amer. J. Math. 98, 79 (1976).
- [44] E. Witten, J. Diff. Geom. 17, 661 (1982).