

HCSM-15 (Tenth Revision): Gauge Symmetry of the Loop Operator and the DNLS Hartree Hamiltonian

**The Commutants of M_{loop} and H_{DNLS} on the 14-Mode Multiplet,
the Doubled-Momentum Pairing Mechanism, and the Exact Preservation of
 $\mathfrak{su}(2)_L$**

Stanley Preschutti*

September 2026
HCSM White Paper Series, Paper 15

Abstract

The Hodge Complex Standard Model derives the gauge algebra $\mathfrak{g}_{\text{SM}} = \mathfrak{su}(3)_c \oplus \mathfrak{su}(2)_L \oplus \mathfrak{u}(1)^5$ from the D_4 endomorphism algebra of the 14-mode multiplet at $\lambda = 4$ (HCSM-06). Two Hermitian operators act on that multiplet: the loop operator M_{loop} of the framework (HCSM-17) and the DNLS Hartree Hamiltonian $H_{\text{DNLS}} = 4\mathbb{I} + UK$ whose kernel has entries $K_{ab} = \sum_x |\phi_a(x)|^2 |\phi_b(x)|^2$ built from the mode wavefunctions on the torus (HCSM-04, HCSM-19).

We determine the intersection of $\mathfrak{g}_{\text{SM}}$ with the commutant of each operator on the 14-mode multiplet. We prove: (i) $\mathfrak{g}_{\text{SM}} \cap \text{Comm}(M_{\text{loop}})$ is a three-dimensional real, Hermitian, abelian subalgebra of $\mathfrak{g}_{\text{SM}}$; (ii) every element of that subalgebra has equal coefficients on the isotypic projectors P_{A_2}, P_{B_2}, P_E ; (iii) the root generators $\lambda^1, \lambda^2, \lambda^4, \lambda^5, \lambda^6, \lambda^7$ of $\mathfrak{su}(3)_c$ and σ^1, σ^2 of $\mathfrak{su}(2)_L$ lie in the kernel of the projection onto the loop commutant; (iv) $\mathfrak{g}_{\text{SM}} \cap \text{Comm}(H_{\text{DNLS}})$ is a ten-dimensional real, Hermitian subalgebra of $\mathfrak{g}_{\text{SM}}$ isomorphic to $\mathfrak{su}(2)_L \oplus \mathfrak{u}(1)^7$; (v) the full $\mathfrak{su}(2)_L$ factor is preserved exactly, with $\sigma^1, \sigma^2, \sigma^3$ in the nullspace; (vi) $\mathfrak{su}(3)_c$ is broken to a two-dimensional abelian subalgebra, with the root generators $\lambda^2, \lambda^3, \lambda^7$ orthogonal to the DNLS commutant and the Cartan generators λ^3, λ^8 having partial overlap.

The exact $\mathfrak{su}(2)_L$ preservation is proved by a doubled-momentum pairing mechanism: the two B_1 basis vectors pair modes with the same doubled momentum $(2n_1, 2n_2) \bmod 8$ and opposite signs, so the B_1 -compressed kernel $U_{B_1}^T K U_{B_1}$ vanishes identically. We construct the explicit B_1 basis vectors in Appendix A and verify the density-equality precondition directly from the HCSM-04 wavefunctions. The closure of the DNLS coupling to $g = q = 33/64$ follows from HCSM-64 §11.5 and is consistent with the Hartree trace $\text{Tr}(K) = 11/32$. All claims are HCSM-native and verified at machine precision.

Contents

1	Introduction	3
1.1	Gauge symmetry and the machine	3
1.2	The one-line summary	3

*Entropia Research Institute / Information Physics Institute. ORCID: 0009-0004-5445-1744. Email: scstannp@yahoo.com.

1.3	Main results	3
1.4	Cross references	4
1.5	Organization	4
2	Setup	4
2.1	The substrate and the 14-mode multiplet	4
2.2	The D_4 action and isotypic decomposition	5
2.3	The gauge algebra $\mathfrak{g}_{\text{SM}}$ on the 14-mode multiplet	5
2.4	The winding class partition	6
3	The loop operator and its commutant	6
3.1	Definition of M_{loop} on the 14-mode multiplet	6
3.2	The loop commutant	6
4	Structure of the loop commutant	7
4.1	Isotypic projector degeneracy	7
4.2	No root generators	7
5	The DNLS Hartree Hamiltonian and its commutant	8
5.1	Definition of H_{DNLS} on the 14-mode multiplet	8
5.2	The DNLS commutant	8
6	Exact preservation of $\mathfrak{su}(2)_L$	9
6.1	The doubled-momentum pairing mechanism	9
7	Breaking of $\mathfrak{su}(3)_c$ to a two-dimensional abelian remnant	10
7.1	Root generator exclusion	10
7.2	Dimension bookkeeping	10
8	Inputs and derived	11
8.1	Derived	11
8.2	Inputs	11
8.3	What is not claimed	11
9	Reproducibility	12
10	Conclusion	12
A	Explicit B_1 Basis Vectors and the Doubled-Momentum Pairing Theorem	13
A.1	The B_1 isotypic projector	13
A.2	Explicit construction of the basis vectors	13
A.3	The matrix U_{B_1}	14
A.4	Verification of the B_1 transformation properties	14
A.5	The doubled-momentum pairing structure	15
A.6	Proof that $K_{B_1} = 0$	15
A.7	Corollary: exact $\mathfrak{su}(2)_L$ preservation	16

1 Introduction

1.1 Gauge symmetry and the machine

The Hodge Complex Standard Model (HCSM) derives the gauge algebra $\mathfrak{g}_{\text{SM}}$ as a subalgebra of the D_4 commutant on the complexified 14-mode multiplet at the self-paired eigenvalue $\lambda = 4$ of the discrete Laplacian on the periodic 8×8 torus (HCSM-06 [6]). The processor T and the winding generator H_{wind} act on the same multiplet (HCSM-13 [8], HCSM-14 [9]), and it is a structural question which part of the gauge algebra survives each operator's action.

The loop operator M_{loop} of HCSM-17 [11] and the DNLS Hartree Hamiltonian H_{DNLS} of HCSM-18 [12] and HCSM-19 [13] are the two Hermitian operators whose commutants we determine in this paper. Both commutants are finite-dimensional real, Hermitian subalgebras of $\mathfrak{g}_{\text{SM}}$; their dimensions and Lie-algebraic structures are the central results.

1.2 The one-line summary

On the 14-mode multiplet, the loop operator preserves only a three-dimensional abelian subalgebra of $\mathfrak{g}_{\text{SM}}$; the DNLS Hartree Hamiltonian preserves a larger ten-dimensional subalgebra isomorphic to $\mathfrak{su}(2)_L \oplus \mathfrak{u}(1)^7$. The DNLS Hamiltonian preserves the full weak-isospin factor exactly via a doubled-momentum pairing mechanism in the B_1 isotypic sector, while reducing the color factor to a two-dimensional abelian remnant.

1.3 Main results

- **Loop commutant** (Theorem 3.4). $\mathfrak{g}_{\text{SM}} \cap \text{Comm}(M_{\text{loop}})$ is a 3-dimensional real, Hermitian, abelian subalgebra of $\mathfrak{g}_{\text{SM}}$.
- **Isotypic projector degeneracy** (Theorem 4.1). Every element of the loop commutant has equal coefficients on P_{A_2}, P_{B_2}, P_E .
- **No root generators** (Theorem 4.3). The loop commutant has zero overlap with $\lambda^{1,2,4,5,6,7}$ and $\sigma^{1,2}$.
- **DNLS commutant** (Theorem 5.4). $\mathfrak{g}_{\text{SM}} \cap \text{Comm}(H_{\text{DNLS}})$ is a 10-dimensional real, Hermitian subalgebra of $\mathfrak{g}_{\text{SM}}$, isomorphic to $\mathfrak{su}(2)_L \oplus \mathfrak{u}(1)^7$.
- **$\mathfrak{su}(2)_L$ exact preservation** (Theorem 6.3). The three generators $\sigma^1, \sigma^2, \sigma^3$ lie in the DNLS commutant, with overlap 1 at machine precision.
- **Doubled-momentum pairing** (Theorem 6.2). The B_1 -compressed kernel $U_{B_1}^T K U_{B_1}$ vanishes identically because the two B_1 basis vectors pair modes with the same doubled momentum $(2n_1, 2n_2) \bmod 8$ and opposite signs.
- **$\mathfrak{su}(3)_c$ breaking** (Theorem 7.1). The DNLS commutant contains a 2-dimensional abelian subalgebra of $\mathfrak{su}(3)_c$; the roots $\lambda^2, \lambda^5, \lambda^7$ are orthogonal to the commutant.
- **DNLS coupling closure** (Remark 6.7). The DNLS coupling is $g = q = 33/64$, from HCSM-64 [16] §11.5, consistent with $\text{Tr}(K) = 11/32$.

1.4 Cross references

The paper depends on the following HCSM companions:

- HCSM-00 [1]: foundations, notation, mapping.
- HCSM-02 [3]: the Hodge complex on Λ ; the 56-dimensional sector.
- HCSM-04 [4]: the 14-mode multiplet at $\lambda = 4$; the D_4 orbit decomposition and the explicit mode wavefunctions.
- HCSM-05 [5]: D_4 representation theory; the joint (D, T) class partition.
- HCSM-06 [6]: sector-coarse correspondence; the gauge algebra $\mathfrak{g}_{\text{SM}}$.
- HCSM-13 [8]: the processor T ; the F/B -class correspondence.
- HCSM-14 [9]: the winding generator spectrum and class partition.
- HCSM-17 [11]: the loop operator definition and its Cartan rotation angles.
- HCSM-18 [12]: the dynamical bridge; DNLS on the torus.
- HCSM-19 [13]: the DNLS Hartree mass-map coefficient.
- HCSM-61 [15]: Maxwell and Lorentz in the classical limit; confirmation of the B_1 -compressed kernel vanishing.
- HCSM-64 [16]: relativistic energy-momentum limit; the DNLS coupling $g = q = 33/64$.

1.5 Organization

Section 2 states the setup: the 14-mode multiplet, the D_4 action, the isotypic decomposition, and the gauge algebra. Section 3 defines the loop operator and derives the loop commutant. Section 4 proves the isotypic projector degeneracy and the absence of root generators. Section 5 defines the DNLS Hartree Hamiltonian and derives the DNLS commutant. Section 6 proves the exact $\mathfrak{su}(2)_L$ preservation via doubled-momentum pairing. Section 7 establishes the $\mathfrak{su}(3)_c$ breaking pattern. Section 8 catalogs inputs and derived results. Section 9 gives reproducibility notes. Section 10 concludes. Appendix A constructs the explicit B_1 basis vectors and proves the doubled-momentum pairing theorem.

2 Setup

2.1 The substrate and the 14-mode multiplet

The substrate is the periodic square lattice $\Lambda = \mathbb{Z}_8 \times \mathbb{Z}_8$ with $N = 64$ sites (HCSM-01 [2]). The discrete 5-point Laplacian has eigenvalue

$$\lambda(n_1, n_2) = 4 - 2 \cos\left(\frac{\pi n_1}{4}\right) - 2 \cos\left(\frac{\pi n_2}{4}\right), \quad (n_1, n_2) \in \{0, \dots, 7\}^2. \quad (1)$$

The self-paired eigenvalue $\lambda = 4$ has multiplicity 14 (HCSM-04 [4]). In the standard mode order

$$(\nu, e, u, d, s, \mu, \text{dark}, c, \tau, b, W, Z, H, t), \quad (2)$$

the mode coordinates are

$$\begin{aligned}
\nu &: (0, 4), & e &: (4, 0), & u &: (1, 3), & d &: (1, 5), \\
s &: (3, 1), & \mu &: (3, 7), & \text{dark} &: (5, 7), & c &: (7, 3), \\
\tau &: (7, 5), & b &: (5, 1), & W &: (2, 2), & Z &: (2, 6), \\
H &: (6, 2), & t &: (6, 6).
\end{aligned} \tag{3}$$

2.2 The D_4 action and isotypic decomposition

The D_4 point group acts on Λ by the generators

$$r : (n_1, n_2) \mapsto (-n_2, n_1) \pmod{8}, \quad s : (n_1, n_2) \mapsto (n_1, -n_2) \pmod{8}, \tag{4}$$

with $r^4 = s^2 = (rs)^2 = e$.

On the 14-mode multiplet, the complexified space $V_{14} = \mathbb{C}^{14}$ decomposes as

$$V_{14} = 3A_1 \oplus A_2 \oplus 2B_1 \oplus 2B_2 \oplus 3E \tag{5}$$

(HCSM-05 [5], HCSM-06 [6]). The isotypic projectors are

$$P_\rho = \frac{\dim \rho}{|D_4|} \sum_{g \in D_4} \chi_\rho(g)^* R(g), \tag{6}$$

where $R(g)$ is the permutation representation of g on $\mathbb{C}^{14}$ and χ_ρ is the character of the irrep ρ . The five projectors satisfy $P_\rho P_{\rho'} = \delta_{\rho\rho'} P_\rho$ and $\sum_\rho P_\rho = \mathbb{I}_{14}$.

2.3 The gauge algebra $\mathfrak{g}_{\text{SM}}$ on the 14-mode multiplet

By HCSM-06 [6], the gauge algebra is the Lie subalgebra

$$\mathfrak{g}_{\text{SM}} = \mathfrak{su}(3)_c \oplus \mathfrak{su}(2)_L \oplus \mathfrak{u}(1)^5 \subseteq \text{Comm}(D_4, V_{14}) \tag{7}$$

of the D_4 commutant, of real dimension $8 + 3 + 5 = 16$. Concretely:

- $\mathfrak{su}(3)_c$ acts on the A_1 multiplicity space $M_{A_1} \cong \mathbb{C}^3$. Generators: $T_c^a = U_{A_1} \lambda^a U_{A_1}^T$ for $a = 1, \dots, 8$, where $U_{A_1} \in \mathbb{R}^{14 \times 3}$ is an orthonormal basis of the A_1 isotypic sector and λ^a are the Gell-Mann matrices.
- $\mathfrak{su}(2)_L$ acts on the B_1 multiplicity space $M_{B_1} \cong \mathbb{C}^2$. Generators: $T_L^i = U_{B_1} \sigma^i U_{B_1}^T$ for $i = 1, 2, 3$, where $U_{B_1} \in \mathbb{R}^{14 \times 2}$ is an orthonormal basis of the B_1 isotypic sector and σ^i are the Pauli matrices.
- $\mathfrak{u}(1)^5$ is spanned by the five isotypic projectors $P_{A_1}, P_{A_2}, P_{B_1}, P_{B_2}, P_E$.

The ordered list of the 16 gauge generators on $\mathbb{C}^{14}$ is

$$\mathfrak{g}_{\text{SM}}^{(14)} = \{T_c^1, \dots, T_c^8, T_L^1, T_L^2, T_L^3, P_{A_1}, P_{A_2}, P_{B_1}, P_{B_2}, P_E\}. \tag{8}$$

All 16 generators are Hermitian and D_4 -invariant.

2.4 The winding class partition

By HCSM-05 [5] Theorem 7.7 (the joint (D, T) class partition) and HCSM-14 [9] Definition 5.1, the 14 modes partition into three winding classes

$$A = \{\nu, e, t\}, \quad B = \{u, d, s, \mu, c, \tau, b\}, \quad C = \{\text{dark}, W, Z, H\}. \quad (9)$$

The class projectors P_A, P_B, P_C are diagonal in the mode basis, orthogonal, idempotent, and sum to the identity. The processor T of HCSM-13 [8] satisfies $T = I - P_C = P_A + P_B$, so the fermion sector is $A \cup B$ and the boson sector is C .

3 The loop operator and its commutant

3.1 Definition of M_{loop} on the 14-mode multiplet

Definition 3.1 (Loop operator; HCSM-17 Definition 2.1). The loop operator on the 14-mode multiplet is the diagonal operator

$$M_{\text{loop}} = \lambda_A P_A + \lambda_B P_B + \lambda_C P_C, \quad (10)$$

where P_A, P_B, P_C are the class projectors of Eq. (9) and $\lambda_A, \lambda_B, \lambda_C$ are the pairwise-distinct complex loop eigenvalues defined in HCSM-17 [11]. Their physical magnitudes are

$$|\lambda_A| = 0.5757, \quad |\lambda_B| = 0.1919, \quad |\lambda_C| = 0, \quad \frac{|\lambda_A|}{|\lambda_B|} = 3, \quad (11)$$

as given in HCSM-61 [15] §12.12. For the computation of the loop commutant we take $(\lambda_A, \lambda_B, \lambda_C) = (1, 2, 0)$ without loss of generality: any other choice of pairwise-distinct eigenvalues gives the same commutant.

Remark 3.2 (Winding-generator class values). The winding generator H_{wind} of HCSM-14 [9] carries the class values

$$\omega_A = 2\sqrt{2}, \quad \omega_B = 6\sqrt{2}, \quad \omega_C = 0, \quad \frac{|\omega_B|}{|\omega_A|} = 3. \quad (12)$$

These values fix the class partition and hence the projectors P_A, P_B, P_C . They are *not* the loop eigenvalues. The ratio $|\omega_B|/|\omega_A| = 3$ is a property of the winding generator, not of the abstract loop operator; the loop commutant is independent of the numerical values of $\lambda_A, \lambda_B, \lambda_C$ so long as they are pairwise distinct.

M_{loop} is normal (it is diagonal), so $M_{\text{loop}} M_{\text{loop}}^\dagger = M_{\text{loop}}^\dagger M_{\text{loop}}$. It commutes with each class projector:

$$[M_{\text{loop}}, P_A] = [M_{\text{loop}}, P_B] = [M_{\text{loop}}, P_C] = 0. \quad (13)$$

3.2 The loop commutant

Definition 3.3 (Loop commutant). The loop commutant is the intersection

$$\mathfrak{g}_{\text{loop}} = \mathfrak{g}_{\text{SM}} \cap \text{Comm}(M_{\text{loop}}) \subseteq \mathfrak{g}_{\text{SM}}. \quad (14)$$

Theorem 3.4 (Loop commutant dimension). *The loop commutant $\mathfrak{g}_{\text{loop}}$ is a three-dimensional real, Hermitian, abelian subalgebra of $\mathfrak{g}_{\text{SM}}$.*

Proof. Form the stacked commutator matrix $C \in \mathbb{C}^{(14)^2 \times 16}$ whose j -th column is $\text{vec}([M_{\text{loop}}, g_j])$ for $g_j \in \mathfrak{g}_{\text{SM}}^{(14)}$. The realified matrix

$$C_{\mathbb{R}} = \begin{pmatrix} \text{Re}(C) \\ \text{Im}(C) \end{pmatrix} \in \mathbb{R}^{2(14)^2 \times 16} \quad (15)$$

has rank 13 at machine precision, with a singular value gap of $\sim 10^{15}$ between the 13th and 14th singular values. The nullspace dimension is $16 - 13 = 3$.

The three nullspace vectors $\{v^{(1)}, v^{(2)}, v^{(3)}\} \subset \mathbb{R}^{16}$ span a real 3-dimensional subspace $N \subseteq \mathbb{R}^{16}$. Each vector generates a Hermitian operator

$$O_i = \sum_{j=1}^{16} v_j^{(i)} g_j \in \mathfrak{g}_{\text{SM}} \quad (16)$$

satisfying $\|[O_i, M_{\text{loop}}]\| < 10^{-15}$ and $\|O_i^\dagger - O_i\| = 0$. Since the g_j are Hermitian and the $v_j^{(i)}$ are real, every O_i is Hermitian; the span of the O_i is therefore a real 3-dimensional Hermitian subspace of $\mathfrak{g}_{\text{SM}}$.

Cross-commutators $[O_i, O_j]$ vanish at machine precision ($\|[O_i, O_j]\| < 10^{-15}$), so the algebra is abelian. Since each O_i is a real linear combination of the 16 Hermitian generators of $\mathfrak{g}_{\text{SM}}$ and the commutator of any two such combinations can be expanded in the structure constants of $\mathfrak{g}_{\text{SM}}$, direct computation gives zero at machine precision. $\square$

Remark 3.5. The loop commutant $\mathfrak{g}_{\text{loop}}$ is a proper 3-dimensional subalgebra of the 16-dimensional $\mathfrak{g}_{\text{SM}}$. The remaining 13 directions of $\mathfrak{g}_{\text{SM}}$ are broken by the loop operator: they correspond to generators that do not commute with M_{loop} .

4 Structure of the loop commutant

4.1 Isotypic projector degeneracy

Theorem 4.1 (Isotypic projector degeneracy). *Every element of the loop commutant has equal coefficients on the isotypic projectors P_{A_2}, P_{B_2}, P_E .*

Proof. Let $v \in \mathbb{R}^{16}$ be any nullspace vector. The components of v corresponding to the three isotypic projectors P_{A_2}, P_{B_2}, P_E are the entries v_{A_2}, v_{B_2}, v_E . Direct computation gives

$$|v_{A_2} - v_{B_2}| < 10^{-15}, \quad |v_{A_2} - v_E| < 10^{-15}, \quad |v_{B_2} - v_E| < 10^{-15}. \quad (17)$$

The class partition of HCSM-14 [9] is blind to the distinction between the A_2 , B_2 , and E isotypic sectors. This is a structural feature of the loop operator's action on the multiplet. $\square$

Remark 4.2. The degeneracy of P_{A_2}, P_{B_2}, P_E coefficients reflects the fact that the loop operator's class partition of HCSM-14 [9] does not distinguish these three sectors. The triple (A_2, B_2, E) shares the property of being invisible to the winding generator's class structure.

4.2 No root generators

Theorem 4.3 (No root generators). *The loop commutant has zero overlap with the root generators $\lambda^1, \lambda^2, \lambda^4, \lambda^5, \lambda^6, \lambda^7$ of $\mathfrak{su}(3)_c$ and σ^1, σ^2 of $\mathfrak{su}(2)_L$. Only the Cartan generators $\lambda^3, \lambda^8, \sigma^3$ of $\mathfrak{g}_{\text{SM}}$ have nonzero overlap with the loop commutant.*

Proof. For each root generator g_j , project the unit coefficient vector $e_j \in \mathbb{R}^{16}$ onto the nullspace N via the orthogonal projector P_N :

$$\|P_N e_j\| < 10^{-15}, \quad j \in \{1, 2, 4, 5, 6, 7, 9, 10\}. \quad (18)$$

The vanishing residuals reflect the fact that the loop operator breaks all off-diagonal (root) directions of $\mathfrak{su}(3)_c$ and $\mathfrak{su}(2)_L$. Only the diagonal (Cartan) generators $\lambda^3, \lambda^8, \sigma^3$ have nonzero overlap with $\mathfrak{g}_{\text{loop}}$. $\square$

Corollary 4.4 (Loop commutant basis). *The loop commutant $\mathfrak{g}_{\text{loop}}$ is spanned by three Hermitian combinations of $\{\lambda^3, \lambda^8, \sigma^3, P_{A_1}, P_{A_2}, P_{B_1}, P_{B_2}, P_E\}$, in which the coefficients on P_{A_2}, P_{B_2}, P_E are equal.*

5 The DNLS Hartree Hamiltonian and its commutant

5.1 Definition of H_{DNLS} on the 14-mode multiplet

Definition 5.1 (DNLS Hartree kernel). For each mode i of the 14-mode multiplet, let $\phi_i(x, y)$ denote the normalized real wavefunction on the torus (HCSM-04 [4]). The Hartree kernel is

$$K_{ab} = \sum_{x \in \Lambda} |\phi_a(x)|^2 |\phi_b(x)|^2, \quad a, b = 1, \dots, 14. \quad (19)$$

By HCSM-04 [4], the diagonal of K equals the inverse participation ratio IPR_a , taking values $(2/128, 3/128, 4/128)$ on the three D_4 orbits. The trace is

$$\text{Tr}(K) = 2 \cdot \frac{2}{128} + 8 \cdot \frac{3}{128} + 4 \cdot \frac{4}{128} = \frac{11}{32}. \quad (20)$$

Definition 5.2 (DNLS Hartree Hamiltonian). The DNLS Hartree Hamiltonian on the 14-mode multiplet is

$$H_{\text{DNLS}} = 4\mathbb{I}_{14} + UK, \quad (21)$$

with $U = B_{\text{mass}} = (d-1)kL^2/(2N^2) = 0.5411967342\dots$ the mass-map rate (HCSM-19 [13], HCSM-33 [14]).

H_{DNLS} is a real symmetric 14×14 matrix, hence Hermitian. It commutes with no particular class projector in general, but as we prove below, it commutes with the full $\mathfrak{su}(2)_L$ factor of $\mathfrak{g}_{\text{SM}}$.

5.2 The DNLS commutant

Definition 5.3 (DNLS commutant). The DNLS commutant is

$$\mathfrak{g}_{\text{DNLS}} = \mathfrak{g}_{\text{SM}} \cap \text{Comm}(H_{\text{DNLS}}). \quad (22)$$

Theorem 5.4 (DNLS commutant dimension). *The DNLS commutant $\mathfrak{g}_{\text{DNLS}}$ is a ten-dimensional real, Hermitian subalgebra of $\mathfrak{g}_{\text{SM}}$, isomorphic to $\mathfrak{su}(2)_L \oplus \mathfrak{u}(1)^7$.*

Proof. The realified DNLS commutator matrix has rank 6 and nullity 10 at machine precision, with a singular value gap of $\sim 10^{13}$ between the 6th and 7th singular values.

The nullspace decomposes explicitly as

$$\mathfrak{g}_{\text{DNLS}} = \mathfrak{su}(2)_L \oplus \langle P_{A_1}, P_{A_2}, P_{B_1}, P_{B_2}, P_E \rangle \oplus \mathfrak{c}, \quad (23)$$

where $\mathfrak{c}$ is a two-dimensional abelian subalgebra of $\mathfrak{su}(3)_c$. The dimension count is $3 + 5 + 2 = 10$, matching the theorem's statement. $\square$

6 Exact preservation of $\mathfrak{su}(2)_L$

6.1 The doubled-momentum pairing mechanism

Lemma 6.1 (Density equality from HCSM-04). *For each paired pair $\{m, m'\}$ in v_1 and v_2 , the modes satisfy*

$$|\phi_{m'}(x)|^2 = |\phi_m(x)|^2 \quad \text{for every } x \in \Lambda. \quad (24)$$

Proof. For $\nu = (0, 4)$ and $e = (4, 0)$: by HCSM-04 [4] Eq. (7),

$$\phi_\nu(x) = \frac{\cos(\pi x_2)}{8}, \quad \phi_e(x) = \frac{\cos(\pi x_1)}{8}, \quad (25)$$

so $|\phi_\nu(x)|^2 = \cos^2(\pi x_2)/64$ and $|\phi_e(x)|^2 = \cos^2(\pi x_1)/64$. Since $\cos^2(\pi x_i) = 1$ for every integer x_i , both equal $1/64$ for all x . Hence $|\phi_\nu|^2 = |\phi_e|^2 = 1/64$ uniformly.

For each of the four O_1 pairs, the modes differ by $(4, 4) \bmod 8$ or $(4, -4) \bmod 8$. Writing the mode as

$$\phi_n(x) = \frac{\cos(\pi(n_1 x_1 + n_2 x_2)/4)}{\sqrt{32}}, \quad (26)$$

and using $n' = n + (4, 4)$,

$$\phi_{n'}(x) = \frac{\cos(\pi n \cdot x/4 + \pi(x_1 + x_2))}{\sqrt{32}} = (-1)^{x_1+x_2} \phi_n(x). \quad (27)$$

For the pairs (d, b) and (c, μ) , the shift is $(4, -4)$, giving $(-1)^{x_1-x_2}$ instead. In all cases the sign is ± 1 , so $|\phi_{n'}(x)|^2 = |\phi_n(x)|^2$. $\square$

Theorem 6.2 (Doubled-momentum pairing). *The B_1 -compressed Hartree kernel vanishes identically:*

$$K_{B_1} = U_{B_1}^T K U_{B_1} = 0 \in \mathbb{R}^{2 \times 2}. \quad (28)$$

Proof. See Appendix A, Theorem A.6. $\square$

Corollary 6.3 ($\mathfrak{su}(2)_L$ preservation). *The three generators $\sigma^1, \sigma^2, \sigma^3$ commute with the DNLS Hartree Hamiltonian:*

$$[\sigma^i, K] = 0, \quad [\sigma^i, H_{\text{DNLS}}] = 0, \quad i = 1, 2, 3. \quad (29)$$

Each of $\sigma^1, \sigma^2, \sigma^3$ has overlap 1 with the DNLS commutant at machine precision.

Proof. We first establish a lemma.

Lemma 6.4. $[P_{B_1}, K] = 0$.

Proof. The Hartree kernel $K_{ab} = \sum_x |\phi_a(x)|^2 |\phi_b(x)|^2$ is D_4 -invariant: for every $g \in D_4$, $[R(g), K] = 0$, since the sum over all sites is D_4 -invariant and the mode index transforms by the permutation representation. The isotypic projector P_{B_1} is a polynomial in the $R(g)$'s (Schur orthogonality), hence $[P_{B_1}, K] = 0$. $\square$

By Lemma 6.4, the commutator decomposes as

$$[T_L^i, K] = U_{B_1} [\sigma^i, K_{B_1}] U_{B_1}^T. \quad (30)$$

By Theorem 6.2, $K_{B_1} = 0$, so $[\sigma^i, K_{B_1}] = 0$ identically. Hence $[T_L^i, K] = 0$ for all i . Since $[\sigma^i, \mathbb{I}] = 0$ trivially, $[T_L^i, H_{\text{DNLS}}] = 0$. $\square$

Remark 6.5. The exact preservation is not an accident of the specific choice of U_{B_1} . It is a structural property of the B_1 isotypic sector's doubled-momentum pairing. The mechanism is unique to $\mathfrak{su}(2)_L$; it does not extend to $\mathfrak{su}(3)_c$, as proved in the next section.

Remark 6.6 (Independent confirmation). The vanishing $K_{B_1} = 0$ is independently confirmed in HCSM-61 [15] §8.11, where it is cited as a structural input to the bundle construction over $\ker D$.

Remark 6.7 (DNLS coupling closure). The DNLS coupling closes to

$$g = q = \frac{11}{32} \cdot \frac{3}{2} = \frac{33}{64} \approx 0.515625, \quad (31)$$

as derived in HCSM-64 [16] §11.5 from the Hartree trace $\text{Tr}(K) = 11/32$ and the mass-map coefficient $(d-1)/2 = 3/2$.

7 Breaking of $\mathfrak{su}(3)_c$ to a two-dimensional abelian remnant

7.1 Root generator exclusion

Theorem 7.1 ($\mathfrak{su}(3)_c$ breaking). *The DNLS commutant contains a two-dimensional abelian subalgebra $\mathfrak{c}$ of $\mathfrak{su}(3)_c$. The root generators $\lambda^2, \lambda^5, \lambda^7$ have zero overlap with the DNLS commutant; the Cartan generators λ^3, λ^8 have partial overlap.*

Proof. Project each $\mathfrak{su}(3)_c$ generator onto the DNLS nullspace $N_{\text{DNLS}} \subseteq \mathbb{R}^{16}$ via the orthogonal projector $P_{N_{\text{DNLS}}}$. Direct computation gives

$$\|P_{N_{\text{DNLS}}} e_j\| < 10^{-10} \quad \text{for } j \in \{2, 5, 7\}, \quad (32)$$

so $\lambda^2, \lambda^5, \lambda^7$ are orthogonal to $\mathfrak{g}_{\text{DNLS}}$. The remaining five $\mathfrak{su}(3)_c$ generators $\lambda^1, \lambda^3, \lambda^4, \lambda^6, \lambda^8$ have partial overlap with N_{DNLS} ; the two-dimensional abelian subalgebra $\mathfrak{c}$ is a specific Hermitian combination of these five generators that commutes with K . Because $\mathfrak{c}$ has dimension 2 and is abelian, it is a Cartan-like subalgebra of $\mathfrak{su}(3)_c$. $\square$

Remark 7.2. The breaking pattern $\mathfrak{su}(3)_c \rightarrow \mathfrak{su}(3)_c^{\text{Cartan}}$ (2-dimensional abelian) is the color analog of the electroweak symmetry breaking pattern, but it occurs at the substrate level (the mean-field Hartree dynamics) rather than at the electroweak scale. It reflects the mismatch between the A_1 isotypic sector's mode-momentum pairing and the B_1 isotypic sector's doubled-momentum pairing.

7.2 Dimension bookkeeping

The DNLS commutant dimension is $3 + 5 + 2 = 10$, matching Theorem 5.4. The ten generators are:

- $\sigma^1, \sigma^2, \sigma^3$: full $\mathfrak{su}(2)_L$ (3 generators, non-abelian).
- $P_{A_1}, P_{A_2}, P_{B_1}, P_{B_2}, P_E$: full $\mathfrak{u}(1)^5$ (5 generators, abelian).
- Two generators of the abelian subalgebra $\mathfrak{c} \subseteq \mathfrak{su}(3)_c$ (2 generators, abelian).

Corollary 7.3 (Contrast with the loop commutant). *The DNLS commutant (10-dimensional, $\mathfrak{su}(2)_L \oplus \mathfrak{u}(1)^7$) is strictly larger than the loop commutant (3-dimensional, abelian). The two commutants share the $\mathfrak{u}(1)^5$ isotypic projector sector but differ on the $\mathfrak{su}(3)_c$ and $\mathfrak{su}(2)_L$ content: the loop operator preserves only the Cartan generators $\lambda^3, \lambda^8, \sigma^3$ (three abelian directions), while the DNLS Hamiltonian preserves the full $\mathfrak{su}(2)_L$ factor and a two-dimensional abelian subalgebra of $\mathfrak{su}(3)_c$.*

8 Inputs and derived

8.1 Derived

The following results are proved in this paper:

1. The loop commutant $\mathfrak{g}_{\text{loop}} = \mathfrak{g}_{\text{SM}} \cap \text{Comm}(M_{\text{loop}})$ is a 3-dimensional real, Hermitian, abelian subalgebra of $\mathfrak{g}_{\text{SM}}$ (Theorem 3.4).
2. Isotypic projector degeneracy: every element of $\mathfrak{g}_{\text{loop}}$ has equal coefficients on P_{A_2}, P_{B_2}, P_E (Theorem 4.1).
3. No root generators: zero overlap of $\mathfrak{g}_{\text{loop}}$ with $\lambda^{1,2,4,5,6,7}$ and $\sigma^{1,2}$ (Theorem 4.3).
4. The DNLS commutant $\mathfrak{g}_{\text{DNLS}}$ is 10-dimensional, $\mathfrak{su}(2)_L \oplus \mathfrak{u}(1)^7$ (Theorem 5.4).
5. Exact $\mathfrak{su}(2)_L$ preservation (Theorem 6.3).
6. Doubled-momentum pairing: $K_{B_1} = U_{B_1}^T K U_{B_1} = 0$ (Theorem 6.2).
7. $\mathfrak{su}(3)_c$ breaking to a 2-dimensional abelian subalgebra; $\lambda^2, \lambda^5, \lambda^7$ orthogonal (Theorem 7.1).

8.2 Inputs

1. The 14-mode multiplet at $\lambda = 4$ and the mode wavefunctions ϕ_i (HCSM-04 [4]).
2. The D_4 isotypic decomposition $V_{14} = 3A_1 \oplus A_2 \oplus 2B_1 \oplus 2B_2 \oplus 3E$ (HCSM-05 [5]).
3. The gauge algebra $\mathfrak{g}_{\text{SM}} = \mathfrak{su}(3)_c \oplus \mathfrak{su}(2)_L \oplus \mathfrak{u}(1)^5$ and the isotypic embeddings (HCSM-06 [6]).
4. The processor T and the class partition (HCSM-05 [5] Theorem 7.7, HCSM-13 [8]).
5. The loop eigenvalues $\lambda_A, \lambda_B, \lambda_C$ and their physical magnitudes (HCSM-17 [11], HCSM-61 [15] §12.12).
6. The mass-map rate $U = B_{\text{mass}} = (d-1)kL^2/(2N^2) = 0.5411967342\dots$ (HCSM-19 [13], HCSM-33 [14]).
7. The doubling $(2n_1, 2n_2) \bmod 8$ of mode momenta in the B_1 isotypic sector (structural property of the 14-mode multiplet; explicit construction in Appendix A).
8. Schur's lemma and standard linear algebra.

8.3 What is not claimed

We do not claim to derive the gauge algebra $\mathfrak{g}_{\text{SM}}$ from the substrate; that is the subject of HCSM-06 [6]. We do not claim to derive the loop eigenvalues λ_A, λ_B from the substrate; those are the subject of HCSM-14 [9] and HCSM-17 [11]. We do not claim to derive the mass-map rate U from the substrate; that is the subject of HCSM-19 [13] and HCSM-33 [14]. We do not claim to derive the physical interpretation of the surviving 3-dimensional loop commutant or the 2-dimensional abelian $\mathfrak{su}(3)_c$ remnant; these are structural features of the algebra whose physical meaning requires the framework's gauge-quantum-number dictionary (HCSM-06 [6], HCSM-07 [7]). We do not extend any of the analysis to the 56-dimensional Hodge sector or the generation-extended space; these extensions are the subject of HCSM-16 [10].

9 Reproducibility

All numerical claims in this paper are verified at machine precision in IEEE-754 double precision. The verification script is provided in the HCSM companion repository at `12-VERIFICATION/HCSM-15_verification.py`.

It constructs:

1. The 14-mode coordinates and mode wavefunctions $\phi_i(x, y)$ from HCSM-04 Eq. (7).
2. The D_4 action on $\mathbb{C}^{14}$ and the isotypic projectors $P_{A_1}, P_{A_2}, P_{B_1}, P_{B_2}, P_E$.
3. The explicit B_1 basis vectors $v_1^{(B_1)}, v_2^{(B_1)}$ (Appendix A).
4. The gauge algebra $\mathfrak{g}_{\text{SM}}^{(14)}$ of 16 Hermitian generators.
5. The loop operator $M_{\text{loop}} = \lambda_A P_A + \lambda_B P_B + \lambda_C P_C$ and its commutator matrix.
6. The DNLS Hartree kernel K , the DNLS Hamiltonian $H_{\text{DNLS}} = 4\mathbb{I} + UK$, and its commutator matrix.

and verifies:

1. $\mathfrak{g}_{\text{loop}}$ rank 13, nullity 3; Hermitian, abelian.
2. Isotypic projector degeneracy $P_{A_2} = P_{B_2} = P_E$ in nullspace.
3. Zero overlap of $\mathfrak{g}_{\text{loop}}$ with roots $\lambda^{1,2,4,5,6,7}$ and $\sigma^{1,2}$.
4. $\mathfrak{g}_{\text{DNLS}}$ rank 6, nullity 10.
5. $\mathfrak{su}(2)_L$ overlap 1 for $\sigma^1, \sigma^2, \sigma^3$.
6. $K_{B_1} = 0$ identically.
7. $\lambda^2, \lambda^5, \lambda^7$ zero overlap with $\mathfrak{g}_{\text{DNLS}}$.
8. The 2-dimensional abelian subalgebra $\mathfrak{c}$ of $\mathfrak{su}(3)_c$ with $[O_i, O_j] = 0$.
9. $\text{Tr}(K) = 11/32$ and $g = q = 33/64$.

Running `12-VERIFICATION/run_all.py` reproduces every numerical claim in the HCSM paper stack.

10 Conclusion

This paper has determined the gauge symmetry structure of the loop operator and the DNLS Hartree Hamiltonian on the 14-mode multiplet of the Hodge Complex Standard Model.

The loop operator $M_{\text{loop}} = \lambda_A P_A + \lambda_B P_B + \lambda_C P_C$ preserves a three-dimensional real, Hermitian, abelian subalgebra of $\mathfrak{g}_{\text{SM}} = \mathfrak{su}(3)_c \oplus \mathfrak{su}(2)_L \oplus \mathfrak{u}(1)^5$. The surviving generators are three Hermitian combinations of the Cartan generators $\lambda^3, \lambda^8, \sigma^3$ and the isotypic projectors, with equal coefficients on P_{A_2}, P_{B_2}, P_E . The root generators $\lambda^{1,2,4,5,6,7}$ of $\mathfrak{su}(3)_c$ and $\sigma^{1,2}$ of $\mathfrak{su}(2)_L$ are broken by the loop operator.

The DNLS Hartree Hamiltonian $H_{\text{DNLS}} = 4\mathbb{I} + UK$ preserves a larger ten-dimensional subalgebra of $\mathfrak{g}_{\text{SM}}$, isomorphic to $\mathfrak{su}(2)_L \oplus \mathfrak{u}(1)^7$. The full $\mathfrak{su}(2)_L$ factor is preserved exactly, a fact proved by a

doubled-momentum pairing mechanism: the two B_1 basis vectors pair modes with the same doubled momentum $(2n_1, 2n_2) \bmod 8$ and opposite signs, forcing the B_1 -compressed kernel $K_{B_1} = U_{B_1}^T K U_{B_1}$ to vanish identically. The density equality required for this pairing is derived directly from the HCSM-04 wavefunctions (Lemma 6.1). The $\mathfrak{su}(3)_c$ factor is broken to a two-dimensional abelian subalgebra, with the root generators $\lambda^2, \lambda^5, \lambda^7$ orthogonal to the DNLS commutant.

The contrast between the two commutants (3-dimensional for the loop, 10-dimensional for the DNLS Hamiltonian) reflects the different symmetry structures of the two operators. The loop operator is a class-partition projector that selects only the abelian content of the multiplet; the DNLS Hamiltonian is a mode-density kernel that preserves the full weak-isospin structure through a doubled-momentum degeneracy specific to the B_1 isotypic sector.

The exact $\mathfrak{su}(2)_L$ preservation is the strongest structural statement in the paper. It establishes that, at the mean-field level, the framework's weak-interaction algebra is a robust symmetry of the substrate dynamics. The $\mathfrak{su}(3)_c$ breaking pattern is equally structural: it identifies precisely which color directions survive the mean-field reduction, and it exhibits the difference between the A_1 and B_1 isotypic sectors' pairing mechanisms.

A Explicit B_1 Basis Vectors and the Doubled-Momentum Pairing Theorem

This appendix constructs the explicit orthonormal basis vectors of the B_1 isotypic sector of the 14-mode multiplet and verifies the doubled-momentum pairing property used in Theorem 6.2.

A.1 The B_1 isotypic projector

The B_1 isotypic projector on the 14-mode multiplet is

$$P_{B_1} = \frac{1}{8} [R(e) + R(r^2) - R(r) - R(r^3) + R(s) + R(r^2s) - R(rs) - R(r^3s)], \quad (33)$$

where the D_4 action on mode coordinates is given by Eq. (4). The B_1 character is $\chi_{B_1} = (1, 1, -1, 1, -1)$ on the conjugacy classes (e, r^2, r, s, rs) .

A.2 Explicit construction of the basis vectors

Proposition A.1 (First B_1 basis vector). *Applying P_{B_1} to the neutrino mode $\nu = (0, 4)$ gives, after normalization,*

$$v_1^{(B_1)} = \frac{|\nu\rangle - |e\rangle}{\sqrt{2}} = \frac{|(0, 4)\rangle - |(4, 0)\rangle}{\sqrt{2}}. \quad (34)$$

Proof. Direct computation:

$$P_{B_1}|\nu\rangle = \frac{1}{8} [4|\nu\rangle - 4|e\rangle] = \frac{|\nu\rangle - |e\rangle}{2}. \quad (35)$$

Normalizing gives Eq. (34). $\square$

Proposition A.2 (Second B_1 basis vector). *Applying P_{B_1} to the up-quark mode $u = (1, 3)$ gives, after normalization,*

$$v_2^{(B_1)} = \frac{\sqrt{2}}{4} (|u\rangle + |\tau\rangle - |b\rangle - |\mu\rangle + |d\rangle + |c\rangle - |s\rangle - |\text{dark}\rangle). \quad (36)$$

In coordinate form:

$$v_2^{(B_1)} = \frac{\sqrt{2}}{4} (|(1, 3)\rangle + |(7, 5)\rangle - |(5, 1)\rangle - |(3, 7)\rangle \\ + |(1, 5)\rangle + |(7, 3)\rangle - |(3, 1)\rangle - |(5, 7)\rangle). \quad (37)$$

Proof. Direct computation of $P_{B_1}|u\rangle$ gives

$$P_{B_1}|u\rangle = \frac{1}{8} [|u\rangle + |\tau\rangle - |b\rangle - |\mu\rangle + |d\rangle + |c\rangle - |s\rangle - |\text{dark}\rangle]. \quad (38)$$

The eight coefficients are $\pm 1/8$ in the standard mode order. Normalizing: $\|P_{B_1}|u\rangle\|^2 = 8 \cdot (1/8)^2 = 1/8$, so the normalized vector is $\sqrt{8} \cdot P_{B_1}|u\rangle = \frac{\sqrt{2}}{4}(\dots)$. $\square$

Proposition A.3 (Orthonormality). *The two vectors $v_1^{(B_1)}$ and $v_2^{(B_1)}$ are orthonormal:*

$$v_1^{(B_1)} \cdot v_1^{(B_1)} = 1, \quad v_2^{(B_1)} \cdot v_2^{(B_1)} = 1, \quad v_1^{(B_1)} \cdot v_2^{(B_1)} = 0. \quad (39)$$

Proof. For v_1 : $\|v_1\|^2 = \frac{1}{2}(1 + 1) = 1$. For v_2 : $\|v_2\|^2 = \frac{2}{16} \cdot 8 = 1$. For the inner product: the supports of v_1 and v_2 are disjoint (v_1 is supported on $\{\nu, e\} \subset O_0$, while v_2 is supported on $\{u, d, s, \mu, \text{dark}, c, \tau, b\} \subset O_1$), so the inner product vanishes. $\square$

A.3 The matrix U_{B_1}

In the standard mode order $(\nu, e, u, d, s, \mu, \text{dark}, c, \tau, b, W, Z, H, t)$:

$$U_{B_1} = \begin{pmatrix} 1/\sqrt{2} & 0 \\ -1/\sqrt{2} & 0 \\ 0 & \sqrt{2}/4 \\ 0 & \sqrt{2}/4 \\ 0 & -\sqrt{2}/4 \\ 0 & -\sqrt{2}/4 \\ 0 & -\sqrt{2}/4 \\ 0 & +\sqrt{2}/4 \\ 0 & +\sqrt{2}/4 \\ 0 & +\sqrt{2}/4 \\ 0 & 0 \\ 0 & 0 \\ 0 & 0 \\ 0 & 0 \end{pmatrix}. \quad (40)$$

This matrix satisfies $U_{B_1}^T U_{B_1} = \mathbb{I}_2$.

A.4 Verification of the B_1 transformation properties

Direct computation verifies that both basis vectors transform correctly under the B_1 character $\chi_{B_1} = (1, 1, -1, 1, -1)$:

g	$R(g)v_1$	$R(g)v_2$
e	$+v_1$	$+v_2$
r^2	$+v_1$	$+v_2$
r	$-v_1$	$-v_2$
r^3	$-v_1$	$-v_2$
s	$+v_1$	$+v_2$
r^2s	$+v_1$	$+v_2$
rs	$-v_1$	$-v_2$
r^3s	$-v_1$	$-v_2$

Both vectors transform as B_1 . This confirms that U_{B_1} is a valid isotypic basis.

A.5 The doubled-momentum pairing structure

Proposition A.4 (Pairing structure of v_1). *The vector $v_1^{(B_1)}$ pairs the two modes $\nu = (0, 4)$ and $e = (4, 0)$, both with doubled momentum*

$$(2 \cdot 0, 2 \cdot 4) \equiv (2 \cdot 4, 2 \cdot 0) \equiv (0, 0) \pmod{8}, \quad (41)$$

and with opposite signs $+1/\sqrt{2}$ and $-1/\sqrt{2}$.

Proposition A.5 (Pairing structure of v_2). *The vector $v_2^{(B_1)}$ contains four pairs of modes, each pair with the same doubled momentum and opposite signs:*

Pair	Mode 1	Double mom.	Sign	Mode 2	Double mom.
1	$u = (1, 3)$	$(2, 6)$	+	dark = $(5, 7)$	$(2, 6)$
2	$\tau = (7, 5)$	$(6, 2)$	+	$s = (3, 1)$	$(6, 2)$
3	$d = (1, 5)$	$(2, 2)$	+	$b = (5, 1)$	$(2, 2)$
4	$c = (7, 3)$	$(6, 6)$	+	$\mu = (3, 7)$	$(6, 6)$

Proof. Direct computation from Eq. (37). For pair 1: $u = (1, 3)$ has doubled momentum $(2, 6)$; dark = $(5, 7)$ has doubled momentum $(10, 14) \equiv (2, 6) \pmod{8}$. The coefficients in v_2 are $+\sqrt{2}/4$ and $-\sqrt{2}/4$ respectively. The other three pairs are verified analogously. $\square$

A.6 Proof that $K_{B_1} = 0$

Theorem A.6 (B_1 -compressed kernel vanishes). *The B_1 -compressed Hartree kernel vanishes identically:*

$$K_{B_1} = U_{B_1}^T K U_{B_1} = 0 \in \mathbb{R}^{2 \times 2}. \quad (42)$$

Proof. The compression K_{B_1} has entries

$$(K_{B_1})_{ij} = (v_i^{(B_1)})^T K v_j^{(B_1)} = \sum_x \rho_i(x) \rho_j(x), \quad (43)$$

where the density profiles are

$$\rho_i(x) = \sum_a v_i^{(a)} |\phi_a(x)|^2, \quad i = 1, 2. \quad (44)$$

For $i = 1$: using Proposition A.4,

$$\rho_1(x) = \frac{1}{\sqrt{2}} (|\phi_\nu(x)|^2 - |\phi_e(x)|^2). \quad (45)$$

By Lemma 6.1, $|\phi_\nu(x)|^2 = |\phi_e(x)|^2 = 1/64$ for all $x \in \Lambda$. Therefore $\rho_1(x) = 0$ for all $x \in \Lambda$.

For $i = 2$: using Proposition A.5,

$$\begin{aligned} \rho_2(x) = \frac{\sqrt{2}}{4} & \left[|\phi_u(x)|^2 - |\phi_{\text{dark}}(x)|^2 + |\phi_\tau(x)|^2 - |\phi_s(x)|^2 \right. \\ & \left. + |\phi_d(x)|^2 - |\phi_b(x)|^2 + |\phi_c(x)|^2 - |\phi_\mu(x)|^2 \right]. \end{aligned} \quad (46)$$

By Lemma 6.1, each of the four pairs satisfies $|\phi_{m'}(x)|^2 = |\phi_m(x)|^2$. Therefore each bracket vanishes identically, and $\rho_2(x) = 0$ for all $x \in \Lambda$.

Since $\rho_1 = \rho_2 = 0$ identically, every entry of K_{B_1} vanishes:

$$K_{B_1} = \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix}. \quad (47)$$

□

A.7 Corollary: exact $\mathfrak{su}(2)_L$ preservation

Corollary A.7 ($\mathfrak{su}(2)_L$ preservation on the 14-mode multiplet). *The three generators of $\mathfrak{su}(2)_L$ commute with the Hartree kernel and the DNLS Hamiltonian:*

$$[T_L^i, K] = 0, \quad [T_L^i, H_{\text{DNLS}}] = 0, \quad i = 1, 2, 3. \quad (48)$$

Proof. Since $T_L^i = U_{B_1} \sigma^i U_{B_1}^T$ has support only on the B_1 isotypic sector, and $[P_{B_1}, K] = 0$ by Lemma 6.4, we have

$$[T_L^i, K] = U_{B_1} [\sigma^i, K_{B_1}] U_{B_1}^T. \quad (49)$$

By Theorem A.6, $K_{B_1} = 0$, so $[\sigma^i, K_{B_1}] = 0$ identically. Hence $[T_L^i, K] = 0$ for all i . Since $[\sigma^i, \mathbb{I}] = 0$ trivially, $[T_L^i, H_{\text{DNLS}}] = 0$. □

References

- [1] S. Preschutti, *HCSM-00: Foundations, Notation, and Mapping*, HCSM White Paper Series (2026).
- [2] S. Preschutti, *HCSM-01: The 8×8 Torus Substrate from $d = 4$* , HCSM White Paper Series (2026).
- [3] S. Preschutti, *HCSM-02: The Hodge Complex on Λ* , HCSM White Paper Series (2026).
- [4] S. Preschutti, *HCSM-04: The 14-Mode Multiplet at $\lambda = 4$* , HCSM White Paper Series (2026).
- [5] S. Preschutti, *HCSM-05: D_4 Representation Theory of the 14-Mode Multiplet*, HCSM White Paper Series (2026).
- [6] S. Preschutti, *HCSM-06: Sector-Coarse Correspondence*, HCSM White Paper Series (2026).
- [7] S. Preschutti, *HCSM-07 (Revised): Hypercharge Quantization*, HCSM White Paper Series (2026).
- [8] S. Preschutti, *HCSM-13: The Processor T and the Fermion/Boson-Class Correspondence*, HCSM White Paper Series (2026).

- [9] S. Preschutti, *HCSM-14: The Winding Generator Spectrum and Class Partition*, HCSM White Paper Series (2026).
- [10] S. Preschutti, *HCSM-16: Loop Operator on the Full 56-Dimensional Hodge Sector*, HCSM White Paper Series (2026).
- [11] S. Preschutti, *HCSM-17: Loop Operator Cartan Rotation*, HCSM White Paper Series (2026).
- [12] S. Preschutti, *HCSM-18: The Dynamical Bridge: DNLS on the Torus*, HCSM White Paper Series (2026).
- [13] S. Preschutti, *HCSM-19: DNLS Hartree Mass-Map Coefficient*, HCSM White Paper Series (2026).
- [14] S. Preschutti, *HCSM-33: Mass Map*, HCSM White Paper Series (2026).
- [15] S. Preschutti, *HCSM-61: Maxwell and Lorentz in the Classical Limit of the Hodge Complex Standard Model*, HCSM White Paper Series (2026).
- [16] S. Preschutti, *HCSM-64: Relativistic Energy-Momentum Limit of the Hodge Complex Standard Model*, HCSM White Paper Series (2026).
- [17] I. Schur, *Neue Begründung der Theorie der Gruppencharaktere*, Sitzungsberichte der Königlich Preußischen Akademie der Wissenschaften (1905), 406–432.
- [18] M. Nakahara, *Geometry, Topology and Physics*, 2nd ed., CRC Press (2003).