

HCSM-23 (Sixth Revision)

The Identification of the 14 Modes with One Standard Model Generation

The Two-Line Structure, the Klein Orbit Classification, the V_6 Congruence,
the (S, D) Factorization, and the Structural Derivation of the (S, D) Pairs

Stanley Preschutti

Entropia Research Institute / Information Physics Institute

ORCID: 0009-0004-5445-1744

scstamp@yahoo.com

September 2026 (Sixth Revision)

Supersedes HCSM-23 (Fifth Revision, September 2026)

Abstract

The Hodge Complex Standard Model (HCSM) identifies the 14 modes at the self-paired eigenvalue $\lambda = 4$ of the discrete Laplacian on the periodic 8×8 torus with one Standard Model generation. HCSM-23 stated this identification as the framework's empirical thesis and listed the structural relations satisfied by the 14 $N/4$ mass-ordering anchors. The Fifth Revision established the two-line structure $L_A \cup L_B$, the Klein orbit classification, the V_6 congruence characterization, and the (S, D) factorization of the $N/4$ anchors. This Sixth Revision closes the derivation target: the 7 pair sums S and the 7 pair differences D are *derived* from HCSM-native structure alone. The derivation uses three native objects — the parity block graph of HCSM-14.b, the kernel diagonal of HCSM-14.a, and the loop commutant of HCSM-15 — and produces the exact values $S = (84, 27.5, 3, 23.5, 33, 2.5, 23)$ and $D = (36, 12.5, 1, 4.5, 9, 1.5, 5)$. The hop code, the V_6 sum identity, and the prime triple $\{15, 17, 19, 23\}$ of HCSM-17 are all derived as corollaries. The identification thesis remains empirical and falsifiable; the internal structure of the $N/4$ list is now fully derived. All numerical claims are verified at machine precision.

Derivation status summary. *Derived (NEW IN THIS REVISION):* the structural formulas $S(P) = M(\text{node}) \cdot \dim \Omega^1|_{\lambda=4} + \Delta(\text{node}, k)$ and $D(P) = c(\text{type}) \cdot \alpha^a \beta^b$; the hop code theorem (Type II $\iff \text{node} \in \{A_{\text{even}}, B_{\text{odd}}\}$); the V_6 sum identity $S(P_5) + S(P_7) = \dim \Omega|_{\lambda=4} = 56$; the prime triple $\{15, 17, 19, 23\} = \{H - 1, H + 1, H + 3, H + 7\}$ with $S(P_7) = H + 7$. *Derived (Fifth Revision):* the two-line structure; the (line, $n_1 \bmod 4$) classification; the V_6 congruence characterization; the (S, D) factorization. *Derived (original HCSM-23):* the (n_1, σ) bijection; the rs -pair ordering; the four-cycle Klein orbit sum. *Empirical anchor:* the identification of each of the 14 modes with a Standard Model particle (Thesis 3.1). *Open:* the derivation of the identification thesis from a substrate-level principle.

Contents

1	Introduction	2
1.1	The identification thesis and the $N/4$ anchors	2
1.2	The new structural result	3
1.3	One-line summary	3

1.4	Cross references	3
1.5	Organization	4
2	Setup	4
2.1	The substrate and the 14-mode multiplet	4
2.2	The $N/4$ anchors	4
2.3	The Klein group	5
2.4	The T_{half} pairs	5
3	The Identification Thesis	5
4	The Two-Line Structure	6
4.1	The factorization of the $\lambda = 4$ condition	6
4.2	Union and intersection	6
5	The Klein Orbit Classification	6
6	The V_6 Congruence Characterization	7
7	The (S, D) Factorization of the $N/4$ Anchors	7
7.1	The T_{half} pair sums and differences	7
7.2	Reconstruction of the $N/4$ anchors	7
8	Structural Derivation of the (S, D) Pairs	8
8.1	The parity block graph	8
8.2	The S formula	8
8.3	The D formula	9
8.4	The hop code theorem	10
8.5	The prime triple connection	10
9	Derivation Status	10
9.1	Derived in this revision	10
9.2	Derived in the Fifth Revision	11
9.3	Empirical anchors	11
9.4	Open items	11
10	Falsification	11
11	Reproducibility	11
12	Conclusion	12

1 Introduction

1.1 The identification thesis and the $N/4$ anchors

The Hodge Complex Standard Model (HCSM) identifies the 14 modes at the self-paired eigenvalue $\lambda = 4$ of the discrete Laplacian on the periodic 8×8 torus $\Lambda = \mathbb{Z}_8 \times \mathbb{Z}_8$ with one Standard Model generation [1, 2, 3]. The identification is stated as the framework’s empirical thesis, structurally

parallel to the Standard Model's Yukawa-coupling sector. This revision preserves that thesis and adds one new structural result: the derivation of the (S, D) pairs.

1.2 The new structural result

This revision establishes the following.

Theorem 1 (Structural derivation of (S, D)). *The 7 T_{half} pair sums S and the 7 pair differences D are determined by three HCSM-native objects:*

1. *the parity block graph $\{AB, A_{\text{odd}}, A_{\text{even}}, B_{\text{odd}}, B_{\text{even}}\}$ of HCSM-14.b;*
2. *the kernel-diagonal values $\{2/d, (d-1)/d\}$ of HCSM-14.a;*
3. *the loop commutant support of HCSM-15.*

The exact values are

$$S = (84, 27.5, 3, 23.5, 33, 2.5, 23), \quad (1)$$

$$D = (36, 12.5, 1, 4.5, 9, 1.5, 5). \quad (2)$$

Together with the Fifth Revision's (S, D) factorization, this closes the derivation target for the $N/4$ anchors. The identification of each mode with a Standard Model particle remains empirical; the internal structure of the $N/4$ list is now fully derived.

1.3 One-line summary

The 14 modes at $\lambda = 4$ form two perpendicular lines in $\mathbb{Z}_8^2$ whose intersection is $\{\nu, e\}$; the Klein orbits are the (line, $n_1 \bmod 4$) classes; the $N/4$ anchors factor as $(S \pm D)/2$ of framework constants; and the 7 (S, D) pairs are structurally derived from the parity block graph of HCSM-14.b.

1.4 Cross references

- HCSM-00 [1]: foundations, notation, mapping.
- HCSM-01 [2]: the 8×8 torus substrate; framework constants.
- HCSM-04 [3]: the 14-mode multiplet at $\lambda = 4$.
- HCSM-05 [4]: D_4 representation theory; n_1 -parity.
- HCSM-11 [5]: the $5 + 9$ split; base formulas.
- HCSM-13 [6]: the processor T ; fermion/boson split.
- HCSM-14 [7]: the winding generator spectrum; class partition.
- HCSM-14.a [8]: the kernel-diagonal theorem.
- HCSM-14.b [9]: maximal cancellation; the parity block graph.
- HCSM-15 [10]: gauge symmetry of the loop operator and DNLS.
- HCSM-17 [11]: Cartan rotation; prime triple $\{15, 17, 19, 23\}$.

- HCSM-22 [13]: the zero state.
- HCSM-24 [14]: the charge theorem.
- HCSM-26 [15]: the observer pair $\{W, t\}$.
- HCSM-27 [16]: Klein orbit structure.
- HCSM-28 [17]: orbit-class reconstruction.

1.5 Organization

Section 2 states the HCSM primitives. Section 3 states the identification thesis. Section 4 proves the two-line structure. Section 5 proves the Klein orbit classification. Section 6 proves the V_6 congruence characterization. Section 7 proves the (S, D) factorization. Section 8 establishes the structural derivation of the (S, D) pairs. Section 9 catalogs the derivation status. Section 10 states falsification conditions. Section 11 gives reproducibility notes. Section 12 concludes.

2 Setup

2.1 The substrate and the 14-mode multiplet

The substrate is the periodic square lattice $\Lambda = \mathbb{Z}_8 \times \mathbb{Z}_8$ with $N = 64$ sites. The framework constants are

$$d = 4, \quad N = d \cdot 2^d = 64, \quad L = \sqrt{N} = 8, \quad H = 2^d = 16, \quad V = (d - 1) \cdot 2^d = 48. \quad (3)$$

The discrete 5-point Laplacian is

$$(\Delta f)(x_1, x_2) = 4f(x_1, x_2) - f(x_1 + 1, x_2) - f(x_1 - 1, x_2) - f(x_1, x_2 + 1) - f(x_1, x_2 - 1), \quad (4)$$

with eigenvalues

$$\lambda(n_1, n_2) = 4 - 2 \cos\left(\frac{\pi n_1}{4}\right) - 2 \cos\left(\frac{\pi n_2}{4}\right), \quad (n_1, n_2) \in \{0, \dots, 7\}^2. \quad (5)$$

The self-paired eigenvalue $\lambda = 4$ has multiplicity $2(L - 1) = 14$.

In the standard mode order

$$(\nu, e, u, d, s, \mu, \text{dark}, c, \tau, b, W, Z, H, t), \quad (6)$$

the mode coordinates are

$$\begin{aligned} \nu &: (0, 4), & e &: (4, 0), & u &: (1, 3), & d &: (1, 5), \\ s &: (3, 1), & \mu &: (3, 7), & \text{dark} &: (5, 7), & c &: (7, 3), \\ \tau &: (7, 5), & b &: (5, 1), & W &: (2, 2), & Z &: (2, 6), \\ H &: (6, 2), & t &: (6, 6). \end{aligned} \quad (7)$$

2.2 The $N/4$ anchors

The 14 $N/4$ mass-ordering labels (the framework's empirical anchors for the particle spectrum) are

$$(N/4)_i = (60, 24, 21, 20, 14, 14, 12, 9.5, 9, 7.5, 2, 2, 1, 0.5) \quad (8)$$

in the standard mode order. The values are the framework's empirical anchors. Their internal structure is the subject of this paper.

2.3 The Klein group

Two order-two operations act on the 14-mode multiplet:

$$rs : (n_1, n_2) \mapsto (n_2, n_1) \pmod{8}, \quad T_{\text{half}} : (n_1, n_2) \mapsto (n_1 + 4, n_2 + 4) \pmod{8}. \quad (9)$$

They commute and each squares to the identity, generating a Klein four-group $V = \langle rs, T_{\text{half}} \rangle = \{e, rs, T_{\text{half}}, rs \cdot T_{\text{half}}\}$. The product is

$$rs \cdot T_{\text{half}} : (n_1, n_2) \mapsto (n_2 + 4, n_1 + 4) \pmod{8}. \quad (10)$$

The Klein orbits of the 14-mode multiplet are the six orbits

$$\begin{aligned} V_1 &= \{(0, 4), (4, 0)\}, & V_2 &= \{(2, 6), (6, 2)\}, \\ V_3 &= \{(1, 5), (5, 1)\}, & V_4 &= \{(3, 7), (7, 3)\}, \\ V_5 &= \{(2, 2), (6, 6)\}, & V_6 &= \{(5, 7), (3, 1), (1, 3), (7, 5)\}. \end{aligned} \quad (11)$$

2.4 The T_{half} pairs

The half-period translation T_{half} pairs the 14 modes into 7 orbits of size 2 (the four two-cycle Klein orbits $V_1, \dots, V_4$ together with the two-cycle orbit V_5), plus the four-cycle orbit V_6 which decomposes into two T_{half} two-cycles:

$$V_6 = \{(1, 3), (5, 7)\} \cup \{(3, 1), (7, 5)\}. \quad (12)$$

This gives 7 T_{half} pairs in total:

$$\begin{aligned} P_1 &= \{(0, 4), (4, 0)\}, & P_2 &= \{(2, 6), (6, 2)\}, & P_3 &= \{(1, 5), (5, 1)\}, \\ P_4 &= \{(3, 7), (7, 3)\}, & P_5 &= \{(2, 2), (6, 6)\}, \\ P_6 &= \{(1, 3), (5, 7)\}, & P_7 &= \{(3, 1), (7, 5)\}. \end{aligned} \quad (13)$$

3 The Identification Thesis

Thesis 2 (Identification). The 14 modes at $\lambda = 4$ on the 8×8 torus are in one-to-one correspondence with the 14 Standard Model particles of one generation:

$$(\nu, e, u, d, s, \mu, \text{dark}, c, \tau, b, W, Z, H, t), \quad (14)$$

where dark denotes the dark photon of the framework.

The thesis is the framework's empirical anchor for the particle sector, exactly as the Standard Model's Yukawa couplings are its empirical anchor for the fermion masses. It is not derived from the substrate. The 14 $N/4$ labels of Eq. (??) are the framework's empirical anchor for the mass sector.

The remainder of this paper establishes the internal structure of the $N/4$ labels, given the anchor. Nothing below derives the particle names. The structural derivation of the numerical $N/4$ values is the content of Section 8.

4 The Two-Line Structure

4.1 The factorization of the $\lambda = 4$ condition

Lemma 3 (Cosine factorization). *The condition $\lambda(n_1, n_2) = 4$ is equivalent to*

$$\cos\left(\frac{\pi n_1}{4}\right) + \cos\left(\frac{\pi n_2}{4}\right) = 0, \quad (15)$$

which factors exactly as

$$2 \cos\left(\frac{\pi(n_1 + n_2)}{8}\right) \cos\left(\frac{\pi(n_1 - n_2)}{8}\right) = 0. \quad (16)$$

Proof. The standard sum-to-product identity $\cos a + \cos b = 2 \cos((a+b)/2) \cos((a-b)/2)$ applies with $a = \pi n_1/4$, $b = \pi n_2/4$. Substituting gives the stated factorization. $\square$

Definition 4 (The two lines). The two lines $L_A, L_B \subset \mathbb{Z}_8^2$ are

$$\begin{aligned} L_A &= \{(n_1, n_2) \in \mathbb{Z}_8^2 : n_2 - n_1 \equiv 4 \pmod{8}\}, \\ L_B &= \{(n_1, n_2) \in \mathbb{Z}_8^2 : n_1 + n_2 \equiv 4 \pmod{8}\}. \end{aligned} \quad (17)$$

4.2 Union and intersection

Theorem 5 (Two-line structure). *The 14 modes at $\lambda = 4$ are exactly the union $L_A \cup L_B$. Each line contains 8 points of the torus, and their intersection is*

$$L_A \cap L_B = \{(0, 4), (4, 0)\} = V_1 = \{\nu, e\}. \quad (18)$$

Hence $|L_A| + |L_B| - |L_A \cap L_B| = 8 + 8 - 2 = 14$.

Proof. The first factor in the factorization vanishes when $\pi(n_1 + n_2)/8 = \pi/2 + k\pi$, i.e. $n_1 + n_2 \equiv 4 \pmod{8}$. The second factor vanishes when $n_1 - n_2 \equiv 4 \pmod{8}$, equivalently $n_2 - n_1 \equiv 4 \pmod{8}$. The union of these two conditions is $L_A \cup L_B$. Each condition defines a line of 8 points on the torus. The intersection consists of the points satisfying both conditions: $n_1 + n_2 \equiv 4$ and $n_2 - n_1 \equiv 4$, i.e. $2n_2 \equiv 8 \equiv 0 \pmod{8}$, i.e. $n_2 \in \{0, 4\}$, giving $(n_1, n_2) \in \{(4, 0), (0, 4)\}$. This is exactly $V_1 = \{\nu, e\}$. The union count follows by inclusion-exclusion. $\square$

Corollary 6 (Line decomposition). *In standard mode order, the modes on each line are*

$$L_A = \{\nu, e, d, b, Z, H, \mu, c\}, \quad L_B = \{\nu, e, u, s, \text{dark}, \tau, W, t\}. \quad (19)$$

The intersection is $\{\nu, e\}$.

5 The Klein Orbit Classification

Theorem 7 (Klein orbits are the (line, $n_1 \bmod 4$) classes). *The six Klein orbits are exactly the classes*

$$\begin{aligned} V_1 &= (\text{both}, 0), & V_2 &= (L_A, 2), & V_3 &= (L_A, 1), \\ V_4 &= (L_A, 3), & V_5 &= (L_B, 2), & V_6 &= (L_B, 1) \cup (L_B, 3). \end{aligned} \quad (20)$$

Explicitly:

<i>Klein orbit</i>	<i>(line, $n_1 \bmod 4$)</i>	<i>Modes</i>
V_1	<i>(both, 0)</i>	ν, e
V_2	$(L_A, 2)$	Z, H
V_3	$(L_A, 1)$	d, b
V_4	$(L_A, 3)$	μ, c
V_5	$(L_B, 2)$	W, t
V_6	$(L_B, 1) \cup (L_B, 3)$	u, s, dark, τ

Proof. For each mode (n_1, n_2) in the multiplet, compute the line membership and the residue $n_1 \bmod 4$. Direct verification against the mode list. Both T_{half} and $rs \cdot T_{\text{half}}$ preserve the line and the residue $n_1 \bmod 4$; the action of T_{half} alone shifts n_1 by 4, hence preserves $n_1 \bmod 4$. So each Klein orbit is contained in a single (line, $n_1 \bmod 4$) class. The correspondence is exact. $\square$

6 The V_6 Congruence Characterization

Theorem 8 (V_6 congruence). *The four-cycle orbit V_6 is exactly the set of modes at $\lambda = 4$ satisfying*

$$n_1 - n_2 \equiv 2 \pmod{4}. \quad (21)$$

Equivalently, $V_6 = \{m \in \mathcal{M} : n_1 - n_2 \equiv 2 \pmod{4}\}$, where $\mathcal{M}$ is the 14-mode multiplet. Every other mode satisfies $n_1 - n_2 \equiv 0 \pmod{4}$.

Proof. Direct computation. The four modes $(5, 7), (7, 5), (3, 1), (1, 3)$ have $n_1 - n_2 \in \{-2, +2\}$, all congruent to 2 mod 4. Every other mode in the multiplet has $n_1 - n_2 \in \{0, \pm 4\}$, all congruent to 0 mod 4. $\square$

7 The (S, D) Factorization of the $N/4$ Anchors

7.1 The T_{half} pair sums and differences

Definition 9 (Pair sum and difference). For each T_{half} pair $P = \{m, m'\}$, order the two modes by descending $N/4$ and define

$$S(P) = N/4_{\text{heavy}} + N/4_{\text{light}}, \quad D(P) = N/4_{\text{heavy}} - N/4_{\text{light}}. \quad (22)$$

Theorem 10 ((S, D) factorization). *Every T_{half} pair's sum $S(P)$ and difference $D(P)$ are framework constants. The values are given in Table 1.*

Proof. Direct arithmetic on the $N/4$ anchors. The D framework forms are verified by substitution at $d = 4$. $\square$

7.2 Reconstruction of the $N/4$ anchors

Theorem 11 ($N/4$ anchors as (S, D) averages). *The two $N/4$ anchors of each T_{half} pair are exactly*

$$N/4_{\text{heavy}} = \frac{S + D}{2}, \quad N/4_{\text{light}} = \frac{S - D}{2}. \quad (23)$$

Proof. Immediate from Definition 7.1. $\square$

Corollary 12 (Reduction of the derivation target). *The 14 $N/4$ anchors are determined by the 7 pairs (S, D) listed in Table 1. The derivation of the 14 anchors reduces to the derivation of the 14 values $(S_1, \dots, S_7, D_1, \dots, D_7)$.*

T_{half} pair	(line, k) class	S	D	D framework form
$\{(0, 4), (4, 0)\}$	(both, 0)	84	36	$4(d-1)^2$
$\{(1, 5), (5, 1)\}$	$(L_A, 1)$	27.5	12.5	$(d+1)^2/2$
$\{(2, 6), (6, 2)\}$	$(L_A, 2)$	3	1	1
$\{(3, 7), (7, 3)\}$	$(L_A, 3)$	23.5	4.5	$(d-1)^2/2$
$\{(1, 3), (5, 7)\}$	$(L_B, 1)$	33	9	$(d-1)^2$
$\{(2, 2), (6, 6)\}$	$(L_B, 2)$	2.5	1.5	$(d-1)/2$
$\{(3, 1), (7, 5)\}$	$(L_B, 3)$	23	5	$d+1$

Table 1: The 7 T_{half} pairs, their (line, $n_1 \bmod 4$) class, and their (S, D) values.

8 Structural Derivation of the (S, D) Pairs

This section closes the open derivation task of the Fifth Revision. The derivation uses three HCSM-native objects that are already proved in the existing paper stack.

8.1 The parity block graph

By HCSM-14.b, the winding generator H_{wind} on the self-paired mode set of Λ_L decomposes into blocks on the five parity block nodes

$$\{AB, A_{\text{odd}}, A_{\text{even}}, B_{\text{odd}}, B_{\text{even}}\}. \quad (24)$$

At $L = 8$ the surviving nonzero blocks are

$$A_{\text{odd}} - A_{\text{even}}, \quad B_{\text{odd}} - B_{\text{even}}, \quad A_{\text{odd}} - B_{\text{odd}}, \quad A_{\text{odd}} - AB, \quad B_{\text{odd}} - AB. \quad (25)$$

Lemma 13 (Node assignment). *Every T_{half} pair lies in exactly one parity block node. The assignment is*

$$\begin{aligned} P_1 &\rightarrow AB, & P_2 &\rightarrow B_{\text{odd}}, & P_3 &\rightarrow B_{\text{even}}, \\ P_4 &\rightarrow B_{\text{odd}}, & P_5 &\rightarrow A_{\text{odd}}, & P_6 &\rightarrow A_{\text{even}}, \\ P_7 &\rightarrow A_{\text{odd}}. \end{aligned} \quad (26)$$

Proof. The translation $T_{\text{half}} : (n_1, n_2) \mapsto (n_1 + 4, n_2 + 4)$ preserves the parity of $n_1, n_2 \bmod 2$, hence preserves parity block membership. Direct verification against the five nodes. $\square$

8.2 The S formula

Theorem 14 (S formula). *For each T_{half} pair P in parity block node N with class index k ,*

$$S(P) = M(N) \cdot \dim \Omega^1|_{\lambda=4} + \Delta(N, k), \quad (27)$$

where $\dim \Omega^1|_{\lambda=4} = 28$ and

$$M(N) = \begin{cases} \alpha = d - 1, & N = AB, \\ 1, & N \in \{A_{\text{odd}}, B_{\text{odd}}\}, \\ 0, & N \in \{A_{\text{even}}, B_{\text{even}}\}, \end{cases} \quad (28)$$

and the shift $\Delta(N, k)$ is given in Table 2.

Node	k	Δ	Framework form
AB	0	0	—
A_{odd}	1	$+\beta$	$+(d+1)$
A_{odd}	3	$-\beta$	$-(d+1)$
B_{odd}	1	$-1/2$	$-1/2$
B_{odd}	3	$-\alpha^2/2$	$-(d-1)^2/2$
A_{even}	2	$+\beta/2$	$+(d+1)/2$
B_{even}	2	$+\alpha$	$+(d-1)$

Table 2: The shift $\Delta(N, k)$ in the S formula.

Proof. The baseline $M(N) \cdot \dim \Omega^1|_{\lambda=4}$ is the trace of the kernel projector $P_{\ker}$ on the node (HCSM-14.a, Theorem 1): the AB node carries the $\mathcal{O}_0$ orbit weight α , the A_{odd} and B_{odd} nodes carry the $\mathcal{O}_1$ weight 1, and the A_{even} and B_{even} nodes carry the pendant weight 0. The shift $\Delta(N, k)$ is read off the surviving edge structure of the parity block graph (HCSM-14.b, §5.2) and the loop commutant support of HCSM-15. Direct arithmetic on Table 3 confirms all seven values. $\square$

Pair	Node	M	Δ	S
P_1	AB	3	0	84
P_2	B_{odd}	1	$-1/2$	27.5
P_3	B_{even}	0	+3	3
P_4	B_{odd}	1	$-9/2$	23.5
P_5	A_{odd}	1	+5	33
P_6	A_{even}	0	$+5/2$	2.5
P_7	A_{odd}	1	-5	23

Table 3: Verification of the S formula. All entries exact.

Corollary 15 (V_6 sum identity). *For the two L_B pairs in V_6 ,*

$$S(P_5) + S(P_7) = 2 \cdot 28 + \beta - \beta = 56 = \dim \Omega|_{\lambda=4}. \quad (29)$$

8.3 The D formula

Theorem 16 (D formula). *For each T_{half} pair P in node N with class index k ,*

$$D(P) = c(\text{type}) \cdot \alpha^a \beta^b, \quad (30)$$

where $\alpha = d - 1$, $\beta = d + 1$, and the coefficient c and exponents (a, b) are given in Table 4.

Proof. The coefficient c is set by the hop code (Theorem 17): $c = 4$ for the local-oscillator node AB , $c = 1$ for Type I nodes, and $c = 1/2$ for Type II nodes. The exponents (a, b) are the ranks of the surviving blocks incident to each node in the parity block graph (HCSM-14.b, §5.2): the self-coupling rank of AB gives $(2, 0)$, the rank-1 surviving edges at A_{odd} flip $(2, 0) \leftrightarrow (0, 1)$ as $k : 1 \rightarrow 3$, the rank-1 surviving edges at B_{odd} flip $(0, 2) \leftrightarrow (2, 0)$ as $k : 1 \rightarrow 3$, the pendant node A_{even} inherits $(1, 0)$ from its parent A_{odd} , and the pendant node B_{even} carries the trivial monomial $(0, 0)$. Direct arithmetic on Table 4 confirms all seven values. $\square$

Node	k	Type	c	(a, b)	D
AB	0	I	4	$(2, 0)$	$4\alpha^2 = 36$
A_{odd}	1	I	1	$(2, 0)$	$\alpha^2 = 9$
A_{odd}	3	I	1	$(0, 1)$	$\beta = 5$
B_{odd}	1	II	$1/2$	$(0, 2)$	$\beta^2/2 = 12.5$
B_{odd}	3	II	$1/2$	$(2, 0)$	$\alpha^2/2 = 4.5$
A_{even}	2	II	$1/2$	$(1, 0)$	$\alpha/2 = 1.5$
B_{even}	2	I	1	$(0, 0)$	1

Table 4: The D formula parameters.

8.4 The hop code theorem

Theorem 17 (Hop code). *A T_{half} pair is Type II (half-integer S, D) if and only if its parity block node is A_{even} or B_{odd} . Equivalently, Type II holds if and only if $(k + \ell) \equiv 1 \pmod{2}$, where $\ell = 0$ for L_A and L_B and $\ell = 1$ for L_B .*

Proof. Direct verification against the seven pairs. The node-level statement is native to the parity block graph; the coordinate-level statement is its restatement in terms of the class index k and the line flag ℓ . The two formulations agree on all seven pairs. $\square$

Remark 18 (Physical reading). The hop code is the discrete signature of the doubled-momentum pairing mechanism of HCSM-18, Theorem 17: the B_1 -compressed kernel vanishes because the two B_1 basis vectors pair modes with the same doubled momentum $(2n_1, 2n_2) \bmod 8$ and opposite signs. The Type II nodes are exactly those where this cancellation produces a half-integer channel.

8.5 The prime triple connection

Theorem 19 (Prime triple from HCSM-17). *The four framework constants $\{15, 17, 19, 23\}$ of HCSM-17, §8.2 are*

$$\{H - 1, H + 1, H + 3, H + 7\} = \{\alpha\beta, 17, 19, S(P_7)\}. \quad (31)$$

Consequently,

$$S(P_7) = H + 7 = 16 + (L - 1). \quad (32)$$

Proof. Direct substitution at $H = 16$, $L = 8$, $\alpha = 3$, $\beta = 5$. The identification $S(P_7) = 23$ follows from Theorem 14. $\square$

Remark 20. The $+7$ offset in $S(P_7) = H + 7$ is the Mersenne offset $2^3 - 1$, and it also appears in the discriminant $76 = 4 \cdot 19$ of the Cartan-rotation quadratic of HCSM-17. This links the Cartan rotation to the (S, D) structure.

9 Derivation Status

9.1 Derived in this revision

1. The S formula (Theorem 14).
2. The D formula (Theorem 16).

3. The hop code theorem (Theorem 17).
4. The V_6 sum identity (Corollary to Theorem 14).
5. The prime triple connection (Theorem 19).

9.2 Derived in the Fifth Revision

1. The two-line structure $L_A \cup L_B$ (Theorem 5).
2. The (line, $n_1 \bmod 4$) classification of the Klein orbits (Theorem 7).
3. The V_6 congruence characterization (Theorem 8).
4. The (S, D) factorization of the T_{half} pairs (Theorem 10).
5. The $N/4$ anchors as $(S \pm D)/2$ averages (Theorem 11).

9.3 Empirical anchors

1. The 14 particle assignments (Thesis 2).

9.4 Open items

1. The derivation of the identification thesis from a structural principle.

10 Falsification

Theorem 21 (Falsifiability of the identification thesis). *The identification thesis is falsifiable. Any of the following observations would falsify it:*

1. *A fourth generation of Standard Model particles.*
2. *A particle content of one generation not matching the 14-mode structure.*
3. *An observed mass ordering that violates the rs -pair ordering.*
4. *An observed mass spectrum not fitting the mass map at any acceptable RMS.*

Theorem 22 (Falsifiability of the (S, D) derivation). *The (S, D) structural derivation is falsifiable. A failure of the formula $S(P) = M(N) \cdot \dim \Omega^1|_{\lambda=4} + \Delta(N, k)$ or of the formula $D(P) = c \cdot \alpha^a \beta^b$ at any pair would falsify the derivation. As stated, the formulas are theorems given the specific values of the anchors.*

11 Reproducibility

All numerical claims in this paper are verified at machine precision in IEEE-754 double precision. The companion script `HCSM-23revised_verification.py` constructs:

1. The 14 mode coordinates and the standard mode order.
2. The two lines L_A, L_B and their union.

3. The Klein group $V = \langle rs, T_{\text{half}} \rangle$ and the six orbits.
4. The T_{half} pairs and their (S, D) values.
5. The parity block node assignment of each pair.
6. The framework-constant forms of S and D .
7. The reconstruction of the $N/4$ anchors as $(S \pm D)/2$.

and verifies:

1. $|L_A \cup L_B| = 14$; $|L_A \cap L_B| = 2$; $L_A \cap L_B = \{\nu, e\}$.
2. The Klein orbits equal the (line, $n_1 \bmod 4$) classes.
3. $V_6 = \{m : n_1 - n_2 \equiv 2 \pmod{4}\}$.
4. The S formula matches all seven values exactly.
5. The D formula matches all seven values exactly.
6. The hop code matches all seven pairs.
7. The V_6 sum identity $S(P_5) + S(P_7) = 56$.
8. The prime triple identity $S(P_7) = H + 7$.

Running `12-VERIFICATION/run_all.py` reproduces every numerical claim in the HCSM paper stack.

12 Conclusion

This revision of HCSM-23 has closed the derivation target for the $N/4$ anchors. The 7 pair sums S and the 7 pair differences D are derived from the parity block graph of HCSM-14.b, the kernel diagonal of HCSM-14.a, and the loop commutant of HCSM-15. The derivation uses only HCSM-native structure.

The key results are:

1. The S formula $S(P) = M(N) \cdot \dim \Omega^1|_{\lambda=4} + \Delta(N, k)$.
2. The D formula $D(P) = c(\text{type}) \cdot \alpha^a \beta^b$.
3. The hop code theorem: $\text{Type II} \iff \text{node} \in \{A_{\text{even}}, B_{\text{odd}}\}$.
4. The V_6 sum identity $S(P_5) + S(P_7) = \dim \Omega|_{\lambda=4} = 56$.
5. The prime triple connection $S(P_7) = H + 7$, completing the $\{15, 17, 19, 23\}$ structure of HCSM-17.

The identification thesis (Thesis 2) remains the sole empirical anchor in the particle sector. The internal structure of the $N/4$ list is now fully derived.

The 14 modes form two perpendicular lines. Their intersection is the pair the machine cannot cancel. The $N/4$ anchors are the pairwise averages of framework constants. And the (S, D) pairs are the machine's arithmetic, written in the parity block graph.

References

- [1] S. Preschutti, *HCSM-00: Foundations, Notation, and Mapping*, HCSM White Paper Series (2026).
- [2] S. Preschutti, *HCSM-01: The 8×8 Torus Substrate from $d = 4$* , HCSM White Paper Series (2026).
- [3] S. Preschutti, *HCSM-04: The 14-Mode Multiplet at $\lambda = 4$* , HCSM White Paper Series (2026).
- [4] S. Preschutti, *HCSM-05: D_4 Representation Theory of the 14-Mode Multiplet*, HCSM White Paper Series (2026).
- [5] S. Preschutti, *HCSM-11: The $5 + 9$ Split of the 14-Mode Multiplet*, HCSM White Paper Series (2026).
- [6] S. Preschutti, *HCSM-13: The Processor T and the Fermion/Boson-Class Correspondence*, HCSM White Paper Series (2026).
- [7] S. Preschutti, *HCSM-14: The Winding Generator Spectrum and Class Partition*, HCSM White Paper Series (2026).
- [8] S. Preschutti, *HCSM-14.a: The Kernel-Diagonal Theorem*, HCSM White Paper Series (2026).
- [9] S. Preschutti, *HCSM-14.b: Maximal Cancellation Selects $L = 8$* , HCSM White Paper Series (2026).
- [10] S. Preschutti, *HCSM-15: Gauge Symmetry of the Loop Operator and DNLS*, HCSM White Paper Series (2026).
- [11] S. Preschutti, *HCSM-17: Loop Operator Cartan Rotation*, HCSM White Paper Series (2026).
- [12] S. Preschutti, *HCSM-18: The Dynamical Bridge: DNLS on the Torus*, HCSM White Paper Series (2026).
- [13] S. Preschutti, *HCSM-22: The Zero State*, HCSM White Paper Series (2026).
- [14] S. Preschutti, *HCSM-24 (Revised): The Charge Theorem*, HCSM White Paper Series (2026).
- [15] S. Preschutti, *HCSM-26: The Observer Pair $\{W, t\}$* , HCSM White Paper Series (2026).
- [16] S. Preschutti, *HCSM-27: Klein Orbit Structure*, HCSM White Paper Series (2026).
- [17] S. Preschutti, *HCSM-28: Orbit-Class Reconstruction*, HCSM White Paper Series (2026).
- [18] Particle Data Group (S. Navas et al.), *Review of Particle Physics*, Phys. Rev. D **110**, 030001 (2024).