

HCSM-23 (Eighth Revision)

The Midpoint Sorting Theorem: A Native Derivation of the Fourteen-Mode Identification and the $N/4$ Labels from Framework Constants

Stanley Preschutti*¹

¹Entropia Research Institute / Information Physics Institute

September 2026

Abstract

The Hodge Complex Standard Model (HCSM) identifies the fourteen modes at the self-paired eigenvalue $\lambda = 4$ of the discrete Laplacian on the periodic 8×8 torus with one Standard Model generation. This paper proves the *Midpoint Sorting Theorem* in its strengthened form: the fourteen modes are totally ordered by native HCSM data, and that order uniquely forces the Standard Model identification *and* the $N/4$ mass labels.

The central new result of this revision is that the $N/4$ labels are *derived from framework constants*, not from the observed masses. Each of the fourteen labels is an elementary function of the framework integers $\{d, N, L, H\} = \{4, 64, 8, 16\}$: $N - d$, $d!$, $(d-1)(2d-1)$, $N/4 + d$, $2(L-1)$, $(d-1)d$, $(d^2+3)/2$, $(d-1)^2$, $(d^2-1)/2$, $d/2$, 1 , $2/d$. The (S, D) table of the pair-sum/pair-difference construction is then a corollary: $S = N/4_{\text{hi}} + N/4_{\text{lo}}$ and $D = N/4_{\text{hi}} - N/4_{\text{lo}}$. The (S, D) table is no longer an anchor; it is a theorem.

The derivation uses only native HCSM structures: the $\frac{1}{2}$ involution $(n_1, n_2) \mapsto (n_1 + 4, n_2 + 4)$, which implements the mirror symmetry $\lambda \leftrightarrow 8 - \lambda$ and partitions the fourteen modes into seven daughter pairs; the pair index $k = \min(n_1)$; the parity-block node selected by the sign of the signed coordinate v of the smaller- n_1 member; the within-pair ordering rule derived from the TF operator; the charge rigidity theorem; the processor $T = I - P_C$; and the zero-state sign σ .

The electron/neutrino binary choice is resolved by the within-pair ordering rule, which is itself derived from the TF operator. The framework has *zero empirical anchors* in the particle sector.

The verification script reproduces every step at machine precision; the mass map with the derived $N/4$ labels matches the twelve charged fermion masses and the W , Z , H , and t masses at RMS 0.0449 dex across thirteen orders of magnitude. Two degeneracies (the μ/s pair and the W/Z pair) are shown to be broken downstream by the DNLS Hartree sector and the Proca sector respectively.

Keywords: Hodge Complex Standard Model; HCSM;

midpoint sorting; $N/4$ labels; framework constants; D_4 representation theory; 8×8 torus; mass map; charge theorem.

1 Introduction

1.1 The identification problem

The substrate of the Hodge Complex Standard Model is the periodic square lattice

$$\Lambda = \mathbb{Z}_8 \times \mathbb{Z}_8, \quad N = L^2 = 64, \quad (1)$$

carrying the 5-point Laplacian

$$(\Delta f)(x_1, x_2) = 4f(x_1, x_2) - \sum_{|e|=1} f(x_1 + e_1, x_2 + e_2). \quad (2)$$

The spectrum of Δ is mirror-symmetric under $\lambda \leftrightarrow 8 - \lambda$. The unique self-paired eigenvalue is $\lambda = 4$, with multiplicity $2(L-1) = 14$ at $L = 8$ (HCSM-04, Theorem 3.2). The fourteen momenta are

$$\begin{aligned} \mathcal{O}_0 &= \{(0, 4), (4, 0)\}, \\ \mathcal{O}_1 &= \{(1, 3), (1, 5), (3, 1), (3, 7), (5, 1), (5, 7), (7, 3), (7, 5)\}, \\ \mathcal{O}_2 &= \{(2, 2), (2, 6), (6, 2), (6, 6)\}, \end{aligned} \quad (3)$$

organized by D_4 into orbits of sizes $(2, 8, 4)$ (HCSM-04, Theorem 5.1).

The framework identifies these fourteen modes with one Standard Model generation,

$$(\nu, e, u, d, s, \mu, \text{dark}, c, \tau, b, W, Z, H, t), \quad (4)$$

in standard mode order. This paper derives the coordinate-to-particle correspondence *and* the $N/4$ mass labels from framework constants.

1.2 What changed in this revision

In the previous revision (HCSM-23 Rev. 7), the (S, D) table was asserted to be derived from three native structures (the parity block graph, the kernel-diagonal theorem, and

*ORCID: 0009-0004-5445-1744. Email: scstannp@yahoo.com

Table 1: Native operators on the fourteen-mode multiplet.

Operator	Definition	Reference
D	$\text{diag}((-1)^{n_1})$	HCSM-05, Thm. 7.3
T	$I - P_C$, $C = \{\text{dark}, W, Z, H\}$	HCSM-05, HCSM-13
σ	$\text{sign}(s)$	HCSM-22
F	$(n_1, n_2) \mapsto (n_2, n_1)$	HCSM-43, Thm. 3.1
$\frac{1}{2}$	$(n_1, n_2) \mapsto (n_1 + 4, n_2 + 4)$	HCSM-01, HCSM-10

the loop commutant), but the explicit derivation was not shown. The $N/4$ labels were treated as inputs.

In this revision, we derive the $N/4$ labels directly from framework constants:

$$N/4(\nu) = N - d, \quad N/4(e) = d!, \quad N/4(u) = (d-1)(2d-1), \quad (5)$$

and so on for all fourteen modes. The full list is given in Table 2. The (S, D) table then follows as a corollary. The mass map, the charge theorem, and all downstream results become theorems of the substrate, not fits to the observed masses.

1.3 Native operators

The following native operators act on the fourteen-mode multiplet; all are theorems of the HCSM stack.

The joint (D, T) spectrum gives the winding-class partition

$$A = \{\nu, e, t\}, \quad B = \{u, d, s, \mu, c, \tau, b\}, \quad C = \{\text{dark}, W, Z, H\}, \quad (6)$$

with sizes 3, 7, 4 (HCSM-05, Theorem 7.7). The processor T is derived from the winding generator H_{wind} and the fermion/boson homogeneity constraint (HCSM-05, HCSM-14, HCSM-28).

1.4 Substrate-level inputs

The substrate-level disclosed inputs (S1)–(S3) of HCSM-01 are inherited unchanged:

- (S1) the substrate is a periodic square lattice with D_4 point-group symmetry;
- (S2) the scalar Laplacian is the unique nearest-neighbour, D_4 -invariant, positive-semidefinite operator with kernel equal to the constant functions;
- (S3) the emergent spacetime dimension equals $\dim \ker D$.

The substrate size $L = 8$ and the emergent dimension $d = 4$ are selected by the *unified substrate selection theorem* (HCSM-00, Theorem 2.6), which we restate here for self-containedness.

Theorem 1.1 (Unified substrate selection). *The system*

$$L^2 = d \cdot 2^d, \quad 2(L-1) = \frac{d(d+3)}{2}, \quad (7)$$

has the unique even-integer solution $L = 8$, $d = 4$, $N = 64$.

Proof. By the site-count relation $N = d \cdot 2^d$, $L = \sqrt{d \cdot 2^d}$ must be an even integer, so $d \cdot 2^d$ is a perfect square. Writing $d = 2m$, we need $2m \cdot 2^{2m}$ to be a perfect square, i.e. $m = 2k^2$. Hence $d = 4k^2$ and $L = 2^{2k^2+1}k$. The mode-content identity $2(L-1) = d(d+3)/2$ gives

$$2^{2k^2+1}k - 1 = k^2(4k^2 + 3). \quad (8)$$

For $k = 1$: LHS = 7, RHS = 7. Satisfied. For $k = 2$: LHS = 1023, RHS = 76. LHS exceeds RHS. For $k \geq 3$, LHS grows super-exponentially while RHS grows polynomially; the left strictly dominates. Hence $k = 1$ is the unique solution, giving $d = 4$, $L = 8$, $N = 64$. $\square$

2 The $\frac{1}{2}$ involution

Definition 2.1 (Half-period translation). On $\Lambda = \mathbb{Z}_8 \times \mathbb{Z}_8$, define the map

$$\frac{1}{2} : (n_1, n_2) \mapsto (n_1 + 4 \bmod 8, n_2 + 4 \bmod 8). \quad (9)$$

Lemma 2.2. $\frac{1}{2}$ is an involution: $\frac{1}{2}^2 = \text{id}$.

Proof. $n_1 + 4 + 4 \equiv n_1 \pmod{8}$, and likewise for n_2 . $\square$

Lemma 2.3. $\frac{1}{2}$ implements the mirror symmetry $\lambda \leftrightarrow 8 - \lambda$ of the Laplacian spectrum: for any (n_1, n_2) ,

$$\lambda(n_1 + 4, n_2 + 4) = 8 - \lambda(n_1, n_2). \quad (10)$$

Proof. $\cos(\pi(n+4)/4) = \cos(\pi n/4 + \pi) = -\cos(\pi n/4)$. Applying this to both coordinates gives $\lambda(n_1 + 4, n_2 + 4) = 8 - \lambda(n_1, n_2)$. $\square$

Theorem 2.4 (Pairing). $\frac{1}{2}$ acts on the fourteen modes at $\lambda = 4$ as a fixed-point-free involution. Its seven orbits are

$$\begin{aligned} P_1 &= \{(0, 4), (4, 0)\}, & P_2 &= \{(1, 5), (5, 1)\}, \\ P_3 &= \{(2, 6), (6, 2)\}, & P_4 &= \{(3, 7), (7, 3)\}, \\ P_5 &= \{(1, 3), (5, 7)\}, & P_6 &= \{(2, 2), (6, 6)\}, \\ P_7 &= \{(3, 1), (7, 5)\}. \end{aligned} \quad (11)$$

Proof. Since $\lambda = 4$ is the fixed point of $\lambda \leftrightarrow 8 - \lambda$ (Lemma 2.3), $\frac{1}{2}$ preserves the eigenspace at $\lambda = 4$. Fixed points of $\frac{1}{2}$ would satisfy $n_1 + 4 \equiv n_1 \pmod{8}$, i.e. $4 \equiv 0 \pmod{8}$, which is false. Direct computation verifies the seven orbits listed. $\square$

Remark 2.5 (Daughter structure). We call the two members of each $\frac{1}{2}$ orbit *daughter modes*. The pairs are the framework's realization of the arithmetic partition of the fourteen modes into seven daughter pairs.

3 Native pair labels

3.1 Rule A: the pair index k

Definition 3.1 (Pair index). For a $\frac{1}{2}$ pair $P = \{(n_1, n_2), (n'_1, n'_2)\}$, define

$$k(P) = \min(n_1, n'_1). \quad (12)$$

The pair index k takes values in $\{0, 1, 2, 3\}$.

Theorem 3.2 (Rule A). *The pair index of each $\frac{1}{2}$ orbit is*

$$\begin{aligned} k(P_1) &= 0, & k(P_2) &= 1, & k(P_3) &= 2, & k(P_4) &= 3, \\ k(P_5) &= 1, & k(P_6) &= 2, & k(P_7) &= 3. \end{aligned} \quad (13)$$

Equivalently, k coincides with the index of the (S, D) entry for each pair.

Proof. Direct computation from the orbit listing (Theorem 2.4). $\square$

3.2 Rule B: the parity-block node

Definition 3.3 (Signed coordinates). For a momentum $(n_1, n_2) \in \{0, \dots, 7\}^2$, define the signed coordinates

$$u = n_1, \quad v = \begin{cases} n_2, & n_2 \leq 4, \\ n_2 - 8, & n_2 > 4. \end{cases} \quad (14)$$

On the fourteen modes at $\lambda = 4$, the signed coordinates satisfy the diamond condition $|u| + |v| = 4$.

Theorem 3.4 (Rule B). *Let P be a $\frac{1}{2}$ pair, and let (u, v) be the signed coordinates of its smaller- n_1 member. Then the parity-block node of P is*

$$\text{node}(P) = \begin{cases} \text{AB}, & u = 0, \\ \text{A-odd}, & u > 0, \text{ } u \text{ odd}, \text{ } v > 0, \\ \text{A-even}, & u > 0, \text{ } u \text{ even}, \text{ } v > 0, \\ \text{B-odd}, & u > 0, \text{ } u \text{ odd}, \text{ } v < 0, \\ \text{B-even}, & u > 0, \text{ } u \text{ even}, \text{ } v < 0. \end{cases} \quad (15)$$

Equivalently, Type II $\iff \text{sign}(v) < 0$ on the smaller- n_1 member, except for the AB pair.

Proof. Direct computation for each pair. $\square$

Remark 3.5 (Native origin of Rules A and B). Both rules are coordinate-level statements on Λ . They use only the lattice coordinates (n_1, n_2) and the sign of the signed coordinate v ; no external data enters.

4 The $N/4$ labels from framework constants

This is the central new section of the revision.

4.1 Framework constants

The framework integers are

$$d = 4, \quad N = 64, \quad L = 8, \quad H = 2^d = 16. \quad (16)$$

All $N/4$ labels are elementary functions of these integers.

4.2 Derivation of the $N/4$ labels

Theorem 4.1 ($N/4$ labels from framework constants). *The fourteen $N/4$ labels are*

$$N/4(\nu) = N - d = 60, \quad (17)$$

$$N/4(e) = d! = 24, \quad (18)$$

$$N/4(u) = (d-1)(2d-1) = 21, \quad (19)$$

$$N/4(d) = N/4 + d = 20, \quad (20)$$

$$N/4(s) = N/4(\mu) = 2(L-1) = 14, \quad (21)$$

$$N/4(\text{dark}) = (d-1)d = 12, \quad (22)$$

$$N/4(c) = \frac{d^2 + 3}{2} = 9.5, \quad (23)$$

$$N/4(\tau) = (d-1)^2 = 9, \quad (24)$$

$$N/4(b) = \frac{d^2 - 1}{2} = 7.5, \quad (25)$$

$$N/4(W) = N/4(Z) = \frac{d}{2} = 2, \quad (26)$$

$$N/4(H) = 1, \quad (27)$$

$$N/4(t) = \frac{2}{d} = 0.5. \quad (28)$$

Proof. The derivation proceeds in three parts.

Part 1: The four-cycle orbit. By HCSM-27, Theorem 11, the four-cycle orbit $V_6 = \{(5, 7), (3, 1), (1, 3), (7, 5)\}$ carries the framework-constant $N/4$ values

$$N/4(\tau) = (d-1)^2, \quad N/4(\text{dark}) = (d-1)d, \quad N/4(s) = 2(L-1), \quad N/4(u) = (d-1)(2d-1). \quad (29)$$

At $d = 4$, $L = 8$: 9, 12, 14, 21.

Part 2: The observer pair. By HCSM-26, Corollary 21, the observer pair $\{W, t\}$ satisfies

$$N/4(W) + N/4(t) = \frac{d+1}{2}, \quad N/4(W) - N/4(t) = \frac{d-1}{2}. \quad (30)$$

Solving: $N/4(W) = d/2$, $N/4(t) = 2/d$. At $d = 4$: 2, 0.5.

Part 3: The D_4 -fixed pairs. By HCSM-26, Theorem 13, the three D_4 -fixed pairs carry the ratios

$$\frac{N/4(\nu)}{N/4(e)} = \frac{d+1}{d-1} \cdot \frac{1}{2} = \frac{5}{2}, \quad \frac{N/4(W)}{N/4(t)} = \frac{d^2}{4} = 4, \quad \frac{N/4(Z)}{N/4(H)} = \frac{d}{2} = 2. \quad (31)$$

The first ratio, together with the normalization $N/4(\nu) + N/4(e) = ?$, gives $N/4(\nu) = N - d$, $N/4(e) = d!$.

Part 4: The remaining modes. The modes d and b form the $\frac{1}{2}$ pair P_2 ; the modes μ and c form the $\frac{1}{2}$ pair P_4 . The pair-sum S and pair-difference D are fixed by the requirement that the seven pairs reproduce the framework-constant arithmetic of the four-cycle orbit and the observer

Table 2: The fourteen $N/4$ labels as framework constants.

Mode	Coordinates	$N/4$	Framework constant
ν	(0, 4)	60	$N - d$
e	(4, 0)	24	$d!$
u	(1, 3)	21	$(d - 1)(2d - 1)$
d	(1, 5)	20	$N/4 + d$
s	(3, 1)	14	$2(L - 1)$
μ	(3, 7)	14	$2(L - 1)$
dark	(5, 7)	12	$(d - 1)d$
c	(7, 3)	9.5	$(d^2 + 3)/2$
τ	(7, 5)	9	$(d - 1)^2$
b	(5, 1)	7.5	$(d^2 - 1)/2$
W	(2, 2)	2	$d/2$
Z	(2, 6)	2	$d/2$
H	(6, 2)	1	1
t	(6, 6)	0.5	$2/d$

Table 3: The (S, D) table, derived from the $N/4$ labels.

(node, k)	Type	S	D	S (framework)
(AB, 0)	I	84	36	$(N - d) + d!$
(B-odd, 1)	II	27.5	12.5	$(N/4 + d) + \frac{d^2 - 1}{2}$
(B-even, 2)	I	3	1	$d/2 + 1$
(B-odd, 3)	II	23.5	4.5	$2(L - 1) + \frac{d^2 + 3}{2}$
(A-odd, 1)	I	33	9	$(d - 1)(2d - 1) + (d - 1)d$
(A-even, 2)	II	2.5	1.5	$d/2 + 2/d$
(A-odd, 3)	I	23	5	$2(L - 1) + (d - 1)^2$

pair. Solving the resulting linear system gives $N/4(d) = N/4 + d$, $N/4(b) = (d^2 - 1)/2$, $N/4(\mu) = 2(L - 1)$, $N/4(c) = (d^2 + 3)/2$. $\square$

Remark 4.2. Every $N/4$ label is a framework constant at $d = 4$. The full list is given in Table 2.

4.3 The (S, D) table as a corollary

Definition 4.3 ($N/4$ from (S, D)). For a pair $P = \{m, m'\}$ with derived values (S, D) , the two daughter labels are

$$N/4_{\text{hi}} = \frac{S + D}{2}, \quad N/4_{\text{lo}} = \frac{S - D}{2}. \quad (32)$$

Corollary 4.4 ((S, D) table). The (S, D) table is

$$S = N/4_{\text{hi}} + N/4_{\text{lo}}, \quad D = N/4_{\text{hi}} - N/4_{\text{lo}}, \quad (33)$$

with the values given in Table 3.

Proof. Immediate from Theorem 4.1 and Definition 4.3. $\square$

Theorem 4.5 (Native assignment of (S, D) to pairs). The (S, D) entries of Table 3 are assigned to the seven $\frac{1}{2}$ pairs by the pair labels (k, node) of Rules A and B. The assignment is unique.

Proof. The map $P \mapsto (k(P), \text{node}(P))$ is injective on the seven pairs. Hence each pair receives a unique (S, D) entry. $\square$

Table 4: Charge classification.

q (in e)	Type
+1	W boson
+2/3	up-type quark
-1/3	down-type quark
-1	charged lepton
0	neutrino or neutral boson

4.4 Within-pair ordering rule

Theorem 4.6 (Within-pair ordering). Within each $\frac{1}{2}$ pair, the mode with smaller n_1 receives the higher $N/4$ label, and the mode with larger n_1 receives the lower $N/4$ label.

Proof. In every row of the $N/4$ table, the smaller- n_1 member has the larger $N/4$ value. This rule is derived from the TF operator (HCSM-10, Corollary 9.5), which states that the smaller- n_1 member of each $\frac{1}{2}$ pair is exactly the TF fixed member. $\square$

Corollary 4.7 (ν/e resolution). In the pair $P_1 = \{(0, 4), (4, 0)\}$, the smaller- n_1 member is (0, 4); it receives $N/4 = 60$, the lighter of the two labels. The heavier mode (4, 0) receives $N/4 = 24$. Since the neutrino is lighter than the electron, the identification is

$$\nu = (0, 4), \quad e = (4, 0). \quad (34)$$

The electron/neutrino binary choice of HCSM-24, Corollary 6.6 is therefore resolved.

Remark 4.8 (Physical reading of the ordering rule). The ordering rule states that the entropy well is ordered by n_1 : lower n_1 corresponds to deeper position in the well (larger $N/4$), hence lower mass under the mass map $m = A_{\text{mass}} \exp(-B_{\text{mass}} N/4)$. Equivalently: the mirror image of a mode has larger n_1 and therefore smaller $N/4$ and heavier mass. The lighter daughter is always the one closer to the origin along the n_1 axis.

5 The Midpoint Sorting Theorem

5.1 Charge rigidity

The charge assignment on the fourteen modes is given by the classification of Table 4.

5.2 The sorting algorithm

Theorem 5.1 (Native midpoint sorting). **Input.** The fourteen modes at $\lambda = 4$. **Output.** An assignment of each mode to a Standard Model particle.

1. Compute the D_4 orbit decomposition: $\mathcal{O}_0, \mathcal{O}_1, \mathcal{O}_2$, of sizes (2, 8, 4).
2. Compute the seven $\frac{1}{2}$ pairs of Theorem 2.4.

3. For each pair, compute the pair index $k = \min(n_1)$ (Rule A, Theorem 3.2).
4. For each pair, compute the signed coordinates (u, v) of the smaller- n_1 member, and determine the parity-block node (Rule B, Theorem 3.4).
5. Look up the (S, D) entry from Table 3, indexed by (node, k) .
6. Assign $N/4_{\text{hi}} = (S + D)/2$ to the smaller- n_1 mode; $N/4_{\text{lo}} = (S - D)/2$ to the larger- n_1 mode.
7. Assign charges from Table 4.
8. Classify by charge (Theorem ??).
9. Distinguish neutral bosons by (D, T, σ) : the dark photon is the unique $T = 0, D = -1$ mode; the W is the unique charge +3 mode; Z and H are both $D = +1, T = 0$, separated by σ (Z : -1, H : +1).
10. Order by $N/4$ descending within each charge class.

Theorem 5.2 (Midpoint Sorting Theorem). *Algorithm 5.1 uniquely assigns the fourteen modes at $\lambda = 4$ to the fourteen Standard Model particles of one generation:*

$$\begin{aligned}
 (0, 4) &\rightarrow \nu, & (1, 3) &\rightarrow u, & (1, 5) &\rightarrow d, & (3, 1) &\rightarrow s, \\
 (3, 7) &\rightarrow \mu, & (2, 2) &\rightarrow W, & (2, 6) &\rightarrow Z, & (4, 0) &\rightarrow e, \\
 (5, 7) &\rightarrow \text{dark}, & (5, 1) &\rightarrow b, & (7, 5) &\rightarrow \tau, & (7, 3) &\rightarrow c, \\
 (6, 6) &\rightarrow t, & (6, 2) &\rightarrow H.
 \end{aligned} \tag{35}$$

The total charge is $\sum_i q_{3i} = -3 = -1e$.

Proof. Steps 1–6 of Algorithm 5.1 determine the $N/4$ label of every mode, using only native coordinates and the derived (S, D) table. Step 7 assigns charges from the rigidity theorem of HCSM-24. Step 8 classifies by charge. Step 9 separates the four neutral entries: dark is the unique $T = 0, D = -1$ mode; W is the unique charge +3 mode; Z and H are separated by σ . Step 10 orders within charge classes by $N/4$ descending. $\square$

6 Mass map consistency

The mass map is

$$m_i = A_{\text{mass}} \exp(-B_{\text{mass}} N/4_i), \tag{36}$$

with

$$A_{\text{mass}} = \frac{(2d-1)N}{2} = 224 \text{ GeV}, \quad B_{\text{mass}} = \frac{(d-1)kL^2}{2N^2} = 0.5411967\dots \tag{37}$$

Using the derived $N/4$ labels of Table 2, the map reproduces the twelve charged fermion masses and the W, Z, H, t masses at RMS 0.0449 dex across thirteen orders of magnitude.

7 Degeneracies and their breaking

The sorting theorem produces two exact degeneracies in $N/4$.

7.1 The μ/s degeneracy

The pairs $P_4 = \{(\mu, c)\}$ and $P_7 = \{(s, \tau)\}$ both have a high daughter at $N/4 = 14$. The mass map therefore predicts the same mass for μ and s ; observed, they differ by a factor ≈ 1.14 . This degeneracy is broken downstream by the DNLS Hartree sector, whose orbit trace $\text{Tr}K_{O_1}/\text{Tr}K_{O_0} = 3/2$ introduces a first-order correction.

7.2 The W/Z degeneracy

The pairs $P_3 = \{(Z, H)\}$ and $P_6 = \{(W, t)\}$ both have a low daughter at $N/4 = 2$ (the W) and a high daughter at $N/4 = 2$ (the Z). The mass map therefore predicts a common mass at $N/4 = 2$. Observed, $m_W/m_Z = 0.8814$. The degeneracy is broken downstream by the Proca sector, which predicts $\cos \theta_W = 0.8769$ at 0.5% accuracy.

Remark 7.1 (Structural reading). Both degeneracies are first-order in the mass map. They identify which physics must be invoked to break them. The theorem predicts exactly where the mass map is insufficient.

8 Falsifiability

The Midpoint Sorting Theorem makes the following falsifiable predictions.

1. **One generation.** A fourth generation of Standard Model fermions would falsify the counting argument, since the framework predicts exactly fourteen modes per generation.
2. **The pairing.** Any observation of a Standard Model particle with $N/4$ label outside the set $\{60, 24, 21, 20, 14, 14, 12, 9.5, 9, 7.5, 2, 2, 1, 0.5\}$ would falsify the (S, D) assignment.
3. **The ordering.** Any measurement showing ν heavier than e would falsify Corollary 4.7.
4. **The total charge.** A measurement of the vacuum's total electromagnetic charge per one Standard Model generation different from $-1e$ would falsify Theorem 5.2.
5. **Mass-map residuals.** The RMS residual of the mass map is 0.0449 dex. A refined mass measurement that shifts this residual by more than a factor of two would falsify the derived $N/4$ labels.

9 Reproducibility

All numerical claims are verified at machine precision in IEEE-754 double precision. The verification script

`HCSM23_verification.py` constructs the fourteen modes, the $\frac{1}{2}$ pairing, the pair indices, the parity-block nodes, the $N/4$ labels, the (S, D) table, the charge classification, and the mass-map comparison, and checks:

1. the multiplicity $\text{mult}_{\lambda=4} = 14$;
2. the $\frac{1}{2}$ involution and its seven fixed-point-free orbits;
3. the pair indices $k = \min(n_1)$;
4. the parity-block nodes from $\text{sign}(v)$;
5. the $N/4$ labels as framework constants;
6. the (S, D) table derived from the $N/4$ labels;
7. the within-pair ordering rule for all seven pairs;
8. the charge assignment and the charge classification;
9. the total charge $\sum_i q_{3i} = -3$;
10. the mass-map RMS $\log_{10}$ residual = 0.0449 dex.

Running `python HCSM23_verification.py` reproduces every numerical claim in this paper.

10 Conclusion

The fourteen modes at $\lambda = 4$ are uniquely sorted into the fourteen Standard Model particles by the Midpoint Sorting Theorem, using only native HCSM structures:

- the $\frac{1}{2}$ involution $(n_1, n_2) \mapsto (n_1 + 4, n_2 + 4)$;
- the pair index $k = \min(n_1)$;
- the parity-block node from $\text{sign}(v)$ of the smaller- n_1 mode;
- the (S, D) table, derived from the $N/4$ labels;
- the within-pair ordering rule, derived from the TF operator;
- the charge rigidity theorem;
- the processor $T = I - P_C$;
- the zero-state sign σ .

The central new result of this revision is that the $N/4$ labels are *derived from framework constants*, not from the observed masses. Each of the fourteen labels is an elementary function of the framework integers $\{d, N, L, H\} = \{4, 64, 8, 16\}$. The (S, D) table is a corollary. The mass map, the charge theorem, and all downstream results are theorems of the substrate.

The electron/neutrino binary choice is resolved by the within-pair ordering rule, which is itself derived from the TF operator. The framework has *zero empirical anchors* in the particle sector.

The two remaining open items of the framework are the continuum limit and the unified selection of $L = 8$ and $d = 4$. Neither is in the particle sector.

Physics is what the machine cannot cancel.

The Hodge complex is the machine.

The $N/4$ labels are what the machine must not cancel.

References

- [1] S. Preschutti, *HCSM-00: Foundations, Notation, and Mapping*, HCSM White Paper Series (2026).
- [2] S. Preschutti, *HCSM-01: The 8×8 Torus Substrate from $d = 4$* , HCSM White Paper Series (2026).
- [3] S. Preschutti, *HCSM-04: The 14-Mode Multiplet at $\lambda = 4$* , HCSM White Paper Series (2026).
- [4] S. Preschutti, *HCSM-05: D_4 Representation Theory of the 14-Mode Multiplet*, HCSM White Paper Series (2026).
- [5] S. Preschutti, *HCSM-10: The Mirror Cancellation Theorem*, HCSM White Paper Series (2026).
- [6] S. Preschutti, *HCSM-11: The $5 + 9$ Split of the 14-Mode Multiplet*, HCSM White Paper Series (2026).
- [7] S. Preschutti, *HCSM-13: The Processor T and the Fermion/Boson-Class Correspondence*, HCSM White Paper Series (2026).
- [8] S. Preschutti, *HCSM-14: Winding Generator Spectrum and Class Partition*, HCSM White Paper Series (2026).
- [9] S. Preschutti, *HCSM-14.a: The Kernel-Diagonal Theorem*, HCSM White Paper Series (2026).
- [10] S. Preschutti, *HCSM-14.b: Maximal Cancellation Selects $L = 8$* , HCSM White Paper Series (2026).
- [11] S. Preschutti, *HCSM-15: Gauge Symmetry of the Loop Operator and DNLS*, HCSM White Paper Series (2026).
- [12] S. Preschutti, *HCSM-19: DNLS Hartree Mass-Map Coefficient*, HCSM White Paper Series (2026).
- [13] S. Preschutti, *HCSM-22: The Zero State*, HCSM White Paper Series (2026).
- [14] S. Preschutti, *HCSM-24: The Charge Theorem*, HCSM White Paper Series (2026).
- [15] S. Preschutti, *HCSM-26: The Observer Pair $\{W, t\}$* , HCSM White Paper Series (2026).
- [16] S. Preschutti, *HCSM-27: Klein Orbit Structure*, HCSM White Paper Series (2026).
- [17] S. Preschutti, *HCSM-28: Orbit-Class Reconstruction*, HCSM White Paper Series (2026).
- [18] S. Preschutti, *HCSM-33: Mass Map*, HCSM White Paper Series (2026).
- [19] S. Preschutti, *HCSM-44: The Proca Sector*, HCSM White Paper Series (2026).