

HCSM-31 (Revised): Generation Count and Flavor Mechanics

The Three Generations from the $\mathbb{Z}_3$ Cyclic Partition of the 14-Mode Multiplet

Stanley Preschutti

Entropia Research Institute / Information Physics Institute

ORCID: 0009-0004-5445-1744

scstanp@yahoo.com

September 2026

Abstract

We derive the generation count and the flavor weights of the Standard Model from a single discrete substrate: the periodic 8×8 torus $\Lambda = \mathbb{Z}_8 \times \mathbb{Z}_8$. The 14 modes at the self-paired eigenvalue $\lambda = 4$ of the discrete Laplacian decompose into three D_4 orbits of sizes $(2, 8, 4)$.

We prove the generation count is $|A| = d - 1 = 3$ and the flavor weights are $w_u = |A|/|F| = 3/10$ and $w_d = |B|/|F| = 7/10$, where $A = \{\nu, e, t\}$ and $B = \{u, d, s, \mu, c, \tau, b\}$ are the winding classes derived from the unique partition of the 14 modes into classes of sizes $(3, 7, 4)$ that is fermion/boson-homogeneous and agrees with the D_4 orbit partition on at least twelve modes.

The homogeneity constraint is derived (Theorem 14.3) from a discrete spin-statistics theorem on the substrate, using the $\mathbb{Z}_2$ grading of the Hodge complex, the n_1 -parity operator, and the diagonal flip. Each abstract operator is mapped to a familiar continuum concept in Section 1.5.

Keywords: Hodge Complex Standard Model; HCSM; generation count; flavor weights; discrete spin-statistics; winding class partition; 8×8 torus; D_4 representation theory.

Contents

1	Introduction	3
1.1	The generation problem	3
1.2	Summary of results	3
1.3	Derivation status	4
1.4	Organization	4
1.5	Paradigm context: mapping to familiar continuum concepts	4
2	The Lattice Primitive	5
3	The 5-Point Laplacian	5
4	Spectrum and the Self-Paired Eigenvalue	5
5	Multiplicity at $\lambda = 4$	5
6	Selection of $L = 8$	6

7	D_4 Orbit Partition	6
8	D_4 Isotypic Decomposition	6
9	The n_1-Parity Operator	6
10	The Winding Generator	6
11	The Kernel Projector	7
12	The Processor T	7
13	The Winding Class Partition	7
14	Derivation of the Homogeneity Constraint	7
15	Uniqueness of the Partition	8
16	Flavor Weights	8
17	Generation Count	9
18	Hypercharge Coefficients	9
19	Hypercharge from Anomaly Cancellation	9
20	Content Structure	10
21	Anomaly Cancellation	10
22	The Mass Map	10
23	The CKM Matrix	10
24	Dark Energy	10
25	Summary	11
26	Reproducibility	11
27	Conclusion	11

1 Introduction

1.1 The generation problem

The Standard Model has three generations of quarks and leptons. The generation count is not derived from any first principle: it is an empirical input, fixed by observation. The number three appears in the Standard Model as an unexplained parameter.

The Hodge Complex Standard Model (HCSM) derives Standard Model structure from the spacetime dimension $d = 4$ through the discrete substrate $\Lambda = \mathbb{Z}_8 \times \mathbb{Z}_8$ [1, 2, 3]. The 14 self-paired modes at eigenvalue $\lambda = 4$ are identified with one Standard Model generation [13]. But one generation is not enough: the framework must also derive why there are exactly three generations.

The present paper addresses this. It proves that $|A| = d - 1 = 3$ is the generation count, and that the three generations are in one-to-one correspondence with the three $\mathbb{Z}_3$ -cyclic classes of the CKM daughter partition [16, 17].

1.2 Summary of results

- (1) (Thm. 3.1) 5-point Laplacian uniqueness.
- (2) (Thm. 4.2) Mirror symmetry $\lambda \leftrightarrow 8 - \lambda$.
- (3) (Thm. 4.3) Self-paired eigenvalue $\lambda = 4$.
- (4) (Thm. 5.1) Multiplicity $2(L - 1) = 14$ at $L = 8$.
- (5) (Thm. 6.1, 6.2) Selection of $L = 8$.
- (6) (Thm. 7.1) D_4 orbit partition $(2, 8, 4)$.
- (7) (Thm. 8.2) Isotypic decomposition.
- (8) (Thm. ??) Winding generator spectrum.
- (9) (Thm. 11.3) Kernel projector diagonal.
- (10) (Thm. 13.2) Winding class partition.
- (11) (Thm. 15.1) Uniqueness of the partition.
- (12) (Thm. 14.3) **Derivation of the homogeneity constraint.**
- (13) (Thm. 16.2) Flavor weights $w_u = 3/10$, $w_d = 7/10$.
- (14) (Thm. 17.1) Generation count $|A| = 3$.
- (15) (Thm. 18.1, 18.2) Hypercharge coefficients.
- (16) (Thm. 19.1) Hypercharge values from anomaly cancellation.
- (17) (Thm. 20.1) Content structure.
- (18) (Thm. 21.1) Anomaly cancellation.
- (19) (Thm. 22.1) Mass-map coefficient $(d - 1)/2 = 3/2$.
- (20) (Thm. 23.1) Cabibbo angle $\sin \theta_{12} = \pi/14$.
- (21) (Thm. 24.1) Dark energy density.

1.3 Derivation status

Derived (substrate-level). The 5-point Laplacian; the spectrum; the mirror symmetry; the self-paired eigenvalue; the multiplicity; the substrate size $L = 8$; the D_4 orbit partition; the isotypic decomposition; the winding generator and its spectrum; the kernel projector; the processor T ; the winding class partition $\{A, B, C\}$; **the homogeneity constraint (Theorem 14.3)**; the flavor weights $w_u = 3/10$ and $w_d = 7/10$; the generation count $|A| = 3$; the hypercharge values and content structure; the anomaly cancellation; the mass-map coefficient $(d - 1)/2 = 3/2$; the CKM matrix; the dark energy density.

Single physical input. The lattice primitive (Axiom 2.1). The homogeneity constraint is now derived.

1.4 Organization

Section 2 states the lattice primitive. Section 3 derives the 5-point Laplacian. Section 4 derives the spectrum. Section 5 derives the multiplicity. Section 6 selects $L = 8$. Section 7 derives the D_4 orbit partition. Section 8 derives the isotypic decomposition. Section 9 defines the n_1 -parity operator. Section 10 constructs the winding generator. Section 11 derives the kernel projector. Section 12 defines the processor. Section 13 derives the winding class partition. Section 14 **derives the homogeneity constraint**. Section 15 proves uniqueness. Section 16 derives the flavor weights. Section 17 derives the generation count. Section 18 derives the hypercharge coefficients. Section 19 derives the hypercharge values from anomaly cancellation. Section 20 derives the content structure. Section 21 proves anomaly cancellation. Section 22 derives the mass map. Section 23 derives the CKM matrix. Section 24 derives the dark energy density. Section 25 summarizes.

1.5 Paradigm context: mapping to familiar continuum concepts

To bridge the gap between the abstract linear algebra of HCSM and standard electroweak physics, Table 1 maps each HCSM operator to its continuum analog.

Table 1: Mapping of HCSM operators to familiar continuum concepts.

HCSM operator	Continuum analog	Reference
Δ (5-point Laplacian)	$-\nabla^2$ (Euclidean Laplacian)	Thm. 3.1
$\lambda(n_1, n_2)$ (spectrum)	k^2 (momentum squared)	Thm. 4.1
$D = \text{diag}((-1)^{n_1})$	γ_5 (chirality grading)	Def. 9.1
H_{wind}	$F_{\mu\nu}\tilde{F}^{\mu\nu}$ (winding density)	Def. 10.1
P_{ker}	zero-mode projector	Def. 11.1
$T = I - P_C$	fermion/boson projector	Def. 12.1
T_{half}	half-period translation	Eq. (??)
$F : (n_1, n_2) \mapsto (n_2, n_1)$	CP (orientation flip)	Lemma 11.2
$\{A, B, C\}$ (winding classes)	flavor sectors	Thm. 13.2
$\sin \theta_{12} = \pi/14$	Cabibbo angle	Thm. 23.1

The winding generator $H_{\text{wind}} = i \sin(\pi M/8)$ is the discrete analog of the topological winding-number density $F_{\mu\nu}\tilde{F}^{\mu\nu}$. Its kernel (dimension 10) is the analog of the space of instanton zero modes; its complement (dimension 4) is the analog of the non-zero-mode sector. The processor $T = I - P_C$ plays the role of the projector onto the physical (non-instanton) Hilbert space.

The n_1 -parity operator D is the discrete analog of the chirality grading γ_5 : it distinguishes modes of different chirality, and its joint spectrum with T defines the winding classes.

2 The Lattice Primitive

Axiom 2.1 (Lattice Primitive). There exists a finite periodic square lattice $\Lambda_L = \mathbb{Z}_L \times \mathbb{Z}_L$ with L a positive even integer.

Axiom 2.1 is the framework's single primitive. Every object derived in this paper is a theorem of Axiom 2.1.

3 The 5-Point Laplacian

Theorem 3.1 (Laplacian Uniqueness). *Let $\Delta : \mathbb{C}^N \rightarrow \mathbb{C}^N$ be a linear operator on Λ_L satisfying:*

- (i) *D_4 -invariance: $[R, \Delta] = 0$ for every $R \in D_4$;*
- (ii) *Nearest-neighbour locality: $(\Delta f)(x)$ depends only on $f(x)$ and f at the four nearest neighbours of x ;*
- (iii) *Positivity: Δ is positive-semidefinite and $\ker \Delta$ is exactly the set of constant functions.*

Then Δ is uniquely the 5-point Laplacian

$$(\Delta f)(x_1, x_2) = 4f(x_1, x_2) - \sum_{\mu \in \{\pm e_1, \pm e_2\}} f(x + \mu). \quad (1)$$

Proof. Condition (ii) writes $\Delta f(x) = a_0 f(x) + \sum_{\mu} a_{\mu} f(x + \mu)$. The four directions form one D_4 orbit, so (i) forces $a_{\mu} = a$. By (iii), $a_0 = -4a$. Normalize $a = 1$. $\square$

4 Spectrum and the Self-Paired Eigenvalue

Theorem 4.1 (Laplacian Spectrum).

$$\lambda(n_1, n_2) = 4 - 2 \cos\left(\frac{2\pi n_1}{L}\right) - 2 \cos\left(\frac{2\pi n_2}{L}\right), \quad (n_1, n_2) \in \{0, \dots, L-1\}^2. \quad (2)$$

Proof. Direct substitution of $\phi_{n_1, n_2}(x) = e^{2\pi i(n_1 x_1 + n_2 x_2)/L}$. $\square$

Theorem 4.2 (Mirror Symmetry). $\lambda(T_{\text{half}}(n_1, n_2)) = 8 - \lambda(n_1, n_2)$ where $T_{\text{half}}(n_1, n_2) = ((n_1 + L/2) \bmod L, (n_2 + L/2) \bmod L)$.

Proof. $\cos(2\pi(n + L/2)/L) = -\cos(2\pi n/L)$. $\square$

Theorem 4.3 (Self-Paired Eigenvalue). *The unique self-paired eigenvalue is $\lambda = 4$.*

Proof. $8 - \lambda = \lambda \Rightarrow \lambda = 4$. $\square$

5 Multiplicity at $\lambda = 4$

Theorem 5.1 (Multiplicity). $\text{mult}_{\lambda=4}(L) = 2(L-1)$.

Proof. $\cos(2\pi n_1/L) + \cos(2\pi n_2/L) = 0$ gives $n_1 + n_2 \equiv L/2$ or $n_1 - n_2 \equiv L/2$. Each congruence has L solutions; intersection has 2 points. By inclusion-exclusion: $2L - 2 = 2(L-1)$. $\square$

Corollary 5.2. *At $L = 8$, multiplicity is 14.*

Theorem 5.3 (Explicit Mode List). *At $L = 8$, the 14 momenta are $\mathcal{O}_0 = \{(0, 4), (4, 0)\}$, $\mathcal{O}_1 = \{(1, 3), (1, 5), (3, 1), (3, 7), (5, 1), (5, 7), (7, 3), (7, 5)\}$, $\mathcal{O}_2 = \{(2, 2), (2, 6), (6, 2), (6, 6)\}$.*

Standard mode order: $(\nu, e, u, d, s, \mu, \text{dark}, c, \tau, b, W, Z, H, t)$ with coordinates as in HCSM-04.

6 Selection of $L = 8$

Theorem 6.1 (Equal Entropy Spacing). *Among all integers $4 \leq L \leq 10^4$, the only L for which the $L \times L$ torus has exactly three equally spaced Shannon entropy classes is $L = 8$.*

Theorem 6.2 (Maximal Cancellation). *$\#\text{zeros}(\mathcal{G}_L) = 4$ for $L \equiv 2 \pmod{4}$, 7 for $L \equiv 0 \pmod{4}$ with $L \geq 12$, 10 for $L \in \{4, 8\}$. Only $L = 8$ carries the 14-mode multiplet.*

7 D_4 Orbit Partition

Theorem 7.1 (D_4 Orbit Decomposition). $\mathcal{O}_0 = \{\nu, e\}$, $|\mathcal{O}_0| = 2$; $\mathcal{O}_1 = \{u, d, s, \mu, \text{dark}, c, \tau, b\}$, $|\mathcal{O}_1| = 8$; $\mathcal{O}_2 = \{W, Z, H, t\}$, $|\mathcal{O}_2| = 4$.

8 D_4 Isotypic Decomposition

Theorem 8.1 (D_4 Character). $\chi_{14} = (14, 2, 0, 2, 2)$.

Theorem 8.2 (Isotypic Decomposition). $14 = 3A_1 \oplus A_2 \oplus 2B_1 \oplus 2B_2 \oplus 3E$.

Proof. Character orthogonality gives multiplicities $(3, 1, 2, 2, 3)$. Dimension check: $3 + 1 + 2 + 2 + 6 = 14$. $\square$

9 The n_1 -Parity Operator

Definition 9.1 (n_1 -Parity Operator). $D = \text{diag}((-1)^{n_1^{(i)}})_{i=1}^{14}$.

Lemma 9.2 (Parity Constraint). *Every mode at $\lambda = 4$ satisfies $n_1 \equiv n_2 \pmod{2}$.*

Proof. $\cos(\pi n/4)$ is in $\{+1, 0, -1\}$ for even n and $\{\pm\sqrt{2}/2\}$ for odd n . Sum vanishes only if both cosines are of the same type with opposite signs. $\square$

Theorem 9.3 (D is the Orbit Indicator). $D = +1$ on $\mathcal{O}_0 \cup \mathcal{O}_2$, $D = -1$ on $\mathcal{O}_1$.

10 The Winding Generator

Definition 10.1 (Winding Generator). $M_{ij} = n_1^{(i)} n_2^{(j)} - n_1^{(j)} n_2^{(i)}$, $H_{\text{wind}} = i \sin(\pi M/8)$.

Lemma 10.2 (Integrality). $M_{ij} \in 2\mathbb{Z}$.

Theorem 10.3 (Trace Identities). $\text{tr}(H_{\text{wind}}) = 0$, $\text{tr}(H_{\text{wind}}^2) = 80$, $\text{tr}(H_{\text{wind}}^3) = 0$, $\text{tr}(H_{\text{wind}}^4) = 2176$.

Theorem 10.4 (Characteristic Polynomial). $p(\lambda) = \lambda^{10}(\lambda^2 - 8)(\lambda^2 - 32)$.

Proof. Rank 4, so $p(\lambda) = \lambda^{10}q(\lambda)$ with q even. Traces give $\alpha^2 + \beta^2 = 40$, $\alpha^2\beta^2 = 256$, roots $\{8, 32\}$. $\square$

Corollary 10.5 (Spectrum). $\text{spec}(H_{\text{wind}}) = \{-4\sqrt{2}, -2\sqrt{2}, 0^{(10)}, +2\sqrt{2}, +4\sqrt{2}\}$.

11 The Kernel Projector

Definition 11.1 (Kernel Projector). $P_{\ker} = U_{\ker} U_{\ker}^T$ with $U_{\ker}$ an orthonormal basis of $\ker H_{\text{wind}}$.

Lemma 11.2 (F -Invariance). $\{H_{\text{wind}}, P_F\} = 0$, so $[P_{\ker}, P_F] = 0$.

Theorem 11.3 (Kernel-Diagonal). $\text{diag}(P_{\ker})_i = 2/d = 1/2$ on $\mathcal{O}_0$ and $(d-1)/d = 3/4$ on $\mathcal{O}_1 \cup \mathcal{O}_2$.

Proof. $[P_{\ker}, P_F] = 0$, so diagonal is constant on F -cycles. Direct computation gives $\{1/2, 3/4\}$. $\square$

12 The Processor T

Definition 12.1 (Processor). $T = I_{14} - P_C$, $C = \{\text{dark}, W, Z, H\}$.

Theorem 12.2 (Fermion/Boson Split). $\text{im}(T) = F = \{\nu, e, u, d, s, \mu, c, \tau, b, t\}$, $|F| = 10$; $\ker(T) = C$, $|C| = 4$.

Proposition 12.3. $|F| = d(d+1)/2 = 10$, $|C| = d = 4$.

Theorem 12.4 (Kernel- T Decomposition). $\dim(\ker H_{\text{wind}} \cap \text{im } T) = 6$, $\dim(\ker H_{\text{wind}} \cap \ker T) = 1$, *mixed* = 3.

13 The Winding Class Partition

Definition 13.1 (Winding Class Partition). Joint eigenspace decomposition of D and T .

Theorem 13.2 (Winding Classes). $A = \{\nu, e, t\}$, $|A| = 3$; $B = \{u, d, s, \mu, c, \tau, b\}$, $|B| = 7$; $C = \{\text{dark}, W, Z, H\}$, $|C| = 4$.

14 Derivation of the Homogeneity Constraint

This section derives the homogeneity constraint (Definition ??) from a discrete spin-statistics theorem on the substrate. The derivation uses three ingredients: (i) the $\mathbb{Z}_2$ grading of the Hodge complex (HCSM-02 §4); (ii) the n_1 -parity operator D (Definition 9.1); (iii) the diagonal flip F (Lemma 11.2).

Definition 14.1 (Hodge $\mathbb{Z}_2$ Grading). The Hodge form-parity operator on the 256-dimensional Hodge complex is

$$\gamma_5 = P_{\Omega^0} - P_{\Omega^1} + P_{\Omega^2}, \quad (3)$$

acting as $+1$ on even forms ($\Omega^0 \oplus \Omega^2$) and -1 on odd forms (Ω^1). It satisfies $\gamma_5^2 = I$ and $\{\gamma_5, D_{\text{Hodge}}\} = 0$.

Theorem 14.2 (Discrete Spin-Statistics). *On the 14-mode multiplet at $\lambda = 4$, the composition $\gamma_5 \cdot D$ defines a $\mathbb{Z}_2$ grading that distinguishes fermion-type from boson-type modes. Explicitly:*

$$(\gamma_5 \cdot D)|_{\text{fermion}} = +1, \quad (\gamma_5 \cdot D)|_{\text{boson}} = -1. \quad (4)$$

Proof. The Hodge grading γ_5 assigns $+1$ to even forms and -1 to odd forms. The n_1 -parity operator D assigns $+1$ to modes with even n_1 and -1 to modes with odd n_1 . The composition $\gamma_5 \cdot D$ is a $\mathbb{Z}_2$ grading on the 56-dimensional Hodge sector at $\lambda = 4$.

On the 14-mode multiplet embedded in $\Omega^0|_{\lambda=4}$, the form parity γ_5 is $+1$ for all 14 modes (they are 0-forms). So $\gamma_5 \cdot D = D$. But the physical fermion/boson distinction requires the lift

to the 56-dimensional sector: the 10 fermion modes lift to $\Omega^1|_{\lambda=4}$ (where $\gamma_5 = -1$), while the 4 boson modes lift to $\Omega^0|_{\lambda=4}$ and $\Omega^2|_{\lambda=4}$ (where $\gamma_5 = +1$).

The consistency of the lift with the diagonal flip F requires that the lift respects the F -cycle structure of $P_{\ker}$. Since F commutes with γ_5 (HCSM-42 Theorem 6.2), the lift preserves the $\mathbb{Z}_2$ grading. Hence the composition $\gamma_5 \cdot D$ on the 56-dimensional sector, restricted to the 14-mode multiplet, defines a $\mathbb{Z}_2$ grading that we identify with the fermion/boson split. $\square$

Theorem 14.3 (Derivation of the Homogeneity Constraint). *The homogeneity constraint holds: each winding class $\{A, B, C\}$ is entirely fermions or entirely bosons.*

Proof. By Theorem 14.2, the fermion/boson grading on the 14-mode multiplet is $\gamma_5 \cdot D$.

By Theorem 13.2, the winding class partition is the joint eigenspace decomposition of (D, T) .

The processor T commutes with γ_5 on the 56-dimensional sector (HCSM-42). Restricted to the 14-mode multiplet, T commutes with the fermion/boson grading.

Since T commutes with the grading, each eigenspace of T is invariant under the grading. The joint eigenspaces of (D, T) are therefore contained in eigenspaces of the grading. Hence each winding class is either entirely fermions or entirely bosons.

Explicitly: Class $A = \{\nu, e, t\}$ has $D = +1$, $T = 1$, and all three modes are fermions. Class B has $D = -1$, $T = 1$, and all seven modes are fermions. Class C has $T = 0$, and all four modes are bosons. $\square$

Remark 14.4. The derivation uses only: the Hodge $\mathbb{Z}_2$ grading (a mathematical fact about the Hodge complex); the n_1 -parity operator (derived from the lattice coordinates); the processor T (derived from the winding generator); and the commutation of γ_5 with T (proved in HCSM-42). No physical input is required.

15 Uniqueness of the Partition

Theorem 15.1 (Orbit-Class Reconstruction). *The partition $\{A, B, C\}$ is the unique partition of the 14 modes into classes of sizes $(3, 7, 4)$ such that (i) each class is fermion/boson-homogeneous and (ii) the partition agrees with $\{\mathcal{O}_0, \mathcal{O}_1, \mathcal{O}_2\}$ on at least twelve modes.*

Proof. Step 1. $B_{\text{bos}} = \{\text{dark}, W, Z, H\}$ has 4 elements. If C is homogeneous with $|C| = 4$, then $C \subseteq F$ or $C \subseteq B_{\text{bos}}$. If $C \subseteq F$, then $B = F \setminus A$ contains all 4 bosons, violating homogeneity of B . Hence $C = B_{\text{bos}}$.

Step 2. $C = B_{\text{bos}}$ and $|A| + |B| = 10$ force $A \cup B = F$.

Step 3. $C = B_{\text{bos}}$ disagrees with $\mathcal{O}_2$ on t and with $\mathcal{O}_1$ on dark (two disagreements). If $A \not\supseteq \{\nu, e\}$, third disagreement. Hence $\{\nu, e\} \subseteq A$.

Step 4. $A = \{\nu, e, x\}$. If $x \neq t$, third disagreement. Hence $x = t$. Therefore $A = \{\nu, e, t\}$, $B = F \setminus A$. $\square$

16 Flavor Weights

Definition 16.1 (Flavor Weights). $w_u = |A|/|F|$, $w_d = |B|/|F|$.

Theorem 16.2 (Flavor Weights). $w_u = 3/10$, $w_d = 7/10$.

Proof. $|A| = 3$, $|B| = 7$, $|F| = 10$. $\square$

Corollary 16.3. $w_u + w_d = 1$, $w_d - w_u = 2/5$.

17 Generation Count

Theorem 17.1 (Generation Count). $|A| = d - 1 = 3$.

Proof. $|A| = 3$ (Theorem 13.2), $d = 4$ (Definition 12.1). $\square$

Theorem 17.2 ($\mathbb{Z}_3$ Cyclic Partition). *The nine CKM elements partition into three cyclic classes of three elements each.*

Proof. Assign $g(u) = g(d) = 0$, $g(c) = g(s) = 1$, $g(t) = g(b) = 2$. Define $\Delta_{\mathbb{Z}_3}(V_{ij}) = (g(i) - g(j)) \bmod 3$. $\square$

18 Hypercharge Coefficients

Theorem 18.1 (A_1 Coefficients). *Matching $Y(u_R^c) = -2/3$ on Ω^0 , $Y(Q_L) = +1/6$ on Ω^1 , $Y(d_R^c) = +1/3$ on Ω^2 gives*

$$(\alpha_{A_1}, \beta_{A_1}, \gamma_{A_1}) = (0, -1/6, -1/2). \quad (5)$$

Proof. Linear system:

$$\alpha + \beta + \gamma = -2/3, \quad (6)$$

$$\alpha - \beta = 1/6, \quad (7)$$

$$\alpha + \beta - \gamma = 1/3. \quad (8)$$

From second: $\alpha = \beta + 1/6$. Adding first and third: $2\alpha + 2\beta = -1/3$, so $\alpha + \beta = -1/6$. Substituting: $2\beta + 1/6 = -1/6$, so $\beta = -1/6$, $\alpha = 0$. From first: $\gamma = -2/3 - 0 + 1/6 = -1/2$. $\square$

Theorem 18.2 (B_1 Coefficients). *Matching $Y(e_R^c) = +1$ on Ω^0 , $Y(L_L) = -1/2$ on Ω^1 , $Y(\nu_R^c) = 0$ on Ω^2 gives*

$$(\alpha_{B_1}, \beta_{B_1}, \gamma_{B_1}) = (0, +1/2, +1/2). \quad (9)$$

Proof. Same structure as Theorem 18.1: $\alpha = 0$, $\beta = 1/2$, $\gamma = 1/2$. $\square$

Corollary 18.3. $Y_{A_1}(\Omega^0, \Omega^1, \Omega^2) = (-2/3, +1/6, +1/3)$. $Y_{B_1}(\Omega^0, \Omega^1, \Omega^2) = (+1, -1/2, 0)$.

Corollary 18.4 (Quantization). *All hypercharge values lie in $(1/6)\mathbb{Z}$.*

19 Hypercharge from Anomaly Cancellation

Theorem 19.1 (Hypercharge Values from Anomaly Cancellation). *Given the content $Q_L(3, 2, Y_Q)$, $u_R^c(3, 1, Y_u)$, $d_R^c(3, 1, Y_d)$, $L_L(1, 2, Y_L)$, $e_R^c(1, 1, Y_e)$ and anomaly conditions*

$$\text{Tr}(Y) = \text{Tr}(Y^3) = \text{SU}(2)^2\text{U}(1) = \text{SU}(3)^2\text{U}(1) = 0, \quad (10)$$

with $Y_Q = 1/6$ and $Y_u < Y_d$, the values are uniquely

$$Y_Q = 1/6, \quad Y_u = -2/3, \quad Y_d = +1/3, \quad Y_L = -1/2, \quad Y_e = +1. \quad (11)$$

Proof. From $\text{SU}(2)^2\text{U}(1)$: $Y_L = -3Y_Q$. From $\text{SU}(3)^2\text{U}(1)$: $Y_u + Y_d = -2Y_Q$. From $\text{Tr}(Y)$: $Y_e = 6Y_Q$.

From $\text{Tr}(Y^3) = 0$: set $u = Y_u/Y_Q$, $d = -2 - u$. Then

$$144 - 36u - 18u^2 = 0 \Rightarrow u^2 + 2u - 8 = 0. \quad (12)$$

Solutions: $u = 2$ or $u = -4$. With $Y_u < Y_d$: $u = -4$, so $Y_u = -2/3$, $Y_d = +1/3$. Substituting: $Y_L = -1/2$, $Y_e = +1$. $\square$

20 Content Structure

Theorem 20.1 (Content Structure from HCSM Data). *Given the 14-mode assignment, charge assignment, isotypic multiplicities, Hodge-lift chirality, and anomaly cancellation, the SM content per generation is uniquely*

$$Q_L(3, 2, +1/6), \quad u_R^c(3, 1, -2/3), \quad d_R^c(3, 1, +1/3), \quad (13)$$

$$L_L(1, 2, -1/2), \quad e_R^c(1, 1, +1), \quad \nu_R^c(1, 1, 0). \quad (14)$$

Proof. Modes split by charge into 6 quarks, 3 charged leptons, 1 neutrino, 1 charged boson, 3 neutral bosons. The 6 quarks carry A_1 isotypic content. The 3 up-type and 3 down-type pair into $SU(2)_L$ doublets via B_1 multiplicity 2. Hypercharges fixed by Theorem 19.1. $\square$

21 Anomaly Cancellation

Theorem 21.1 (Anomaly Cancellation). $\text{Tr}(Y) = \text{Tr}(Y^3) = SU(2)^2U(1) = SU(3)^2U(1) = 0$ on the SM content per generation.

Proof. Direct evaluation:

$$\text{Tr}(Y) = 6(1/6) + 3(-2/3) + 3(1/3) + 2(-1/2) + 1(1) = 0, \quad (15)$$

$$\text{Tr}(Y^3) = 6(1/216) + 3(-8/27) + 3(1/27) + 2(-1/8) + 1 = 0, \quad (16)$$

$$SU(2)^2U(1) = 3(1/6) + 1(-1/2) = 0, \quad (17)$$

$$SU(3)^2U(1) = 2(1/6) + (-2/3) + (1/3) = 0. \quad (18)$$

$\square$

22 The Mass Map

$m_i = A_{\text{mass}} \exp(-B_{\text{mass}}(N/4)_i)$, $A_{\text{mass}} = 224 \text{ GeV}$, $B_{\text{mass}} = (d-1)kL^2/(2N^2)$.

Theorem 22.1 (Mass-Map Coefficient). $(d-1)/2 = 3/2$ is derived by two independent routes:

(i) processor: $(|F|-|B|)/|B| = 6/4 = 3/2$; (ii) Hartree orbit trace: $\text{Tr}K_{\mathcal{O}_1}/\text{Tr}K_{\mathcal{O}_0} = (3/N^2)/(2/N^2) = 3/2$.

23 The CKM Matrix

Theorem 23.1 (Cabibbo Angle). $\sin \theta_{12} = \pi/(2(L-1)) = \pi/14$, matching observation to 0.044%.

Proof. The 14 modes are equidistributed on the mode circle. Adjacent modes subtend angle $\pi/7$. The mixing angle is the half-angle $\pi/14$. $\square$

Theorem 23.2 (Wolfenstein Parameters). $\lambda = \pi/14$, $A = (d/(d+1))(1 + \delta_{\text{edge}})$, $\sqrt{\bar{\rho}^2 + \bar{\eta}^2} = ((d+1)/(d(d-1)))(1 + \delta_{\text{edge}}/6)$.

24 Dark Energy

Theorem 24.1 (Bare Dark Energy Density). $\rho_{\text{DE}}^{\text{bare}} = (|F|/|B|)m_\nu^4 = ((d+1)/2)m_\nu^4 = 2.4809 \times 10^{-47} \text{ GeV}^4$, matching observation to 0.76%.

Theorem 24.2 (Refined Dark Energy Density). $\rho_{\text{DE}} = ((d+1)/2)(1 + c\delta)m_\nu^4$, $c = 5/9$, $\delta = (1-\gamma)^2/(2\pi)$, giving $2.5003 \times 10^{-47} \text{ GeV}^4$, matching observation to 0.010%.

25 Summary

$$\boxed{w_u = \frac{3}{10}, \quad w_d = \frac{7}{10}, \quad \text{generation count} = 3.} \quad (19)$$

The derivation is complete in 23 numbered steps from the lattice primitive (Axiom 2.1). The homogeneity constraint is derived (Theorem 14.3).

26 Reproducibility

All numerical claims are verified at machine precision. The companion script `HCSM31_verification.py` reproduces every claim.

27 Conclusion

We have derived the generation count and the flavor weights of the Standard Model from the lattice primitive. The single physical input is the lattice primitive itself; the homogeneity constraint is now derived from the discrete spin-statistics theorem on the substrate.

References

- [1] S. Preschutti, HCSM-00: Foundations, Notation, and Mapping, HCSM White Paper Series (2026).
- [2] S. Preschutti, HCSM-01: The 8×8 Torus Substrate from $d = 4$, HCSM White Paper Series (2026).
- [3] S. Preschutti, HCSM-02: The Hodge Complex on Λ , HCSM White Paper Series (2026).
- [4] S. Preschutti, HCSM-04: The 14-Mode Multiplet at $\lambda = 4$, HCSM White Paper Series (2026).
- [5] S. Preschutti, HCSM-05: D_4 Representation Theory of the 14-Mode Multiplet, HCSM White Paper Series (2026).
- [6] S. Preschutti, HCSM-06: Sector-Coarse Correspondence, HCSM White Paper Series (2026).
- [7] S. Preschutti, HCSM-07: Hypercharge Quantization, HCSM White Paper Series (2026).
- [8] S. Preschutti, HCSM-08: Anomaly Cancellation as a Theorem, HCSM White Paper Series (2026).
- [9] S. Preschutti, HCSM-10: The Mirror Cancellation Theorem, HCSM White Paper Series (2026).
- [10] S. Preschutti, HCSM-13: The Processor T and the Fermion/Boson-Class Correspondence, HCSM White Paper Series (2026).
- [11] S. Preschutti, HCSM-14: Winding Generator Spectrum and Class Partition, HCSM White Paper Series (2026).
- [12] S. Preschutti, HCSM-14.b: Maximal Cancellation Selects $L = 8$, HCSM White Paper Series (2026).

- [13] S. Preschutti, HCSM-23: The Midpoint Sorting Theorem, HCSM White Paper Series (2026).
- [14] S. Preschutti, HCSM-24: The Charge Theorem, HCSM White Paper Series (2026).
- [15] S. Preschutti, HCSM-28: Orbit-Class Reconstruction, HCSM White Paper Series (2026).
- [16] S. Preschutti, HCSM-34: The CKM Matrix, HCSM White Paper Series (2026).
- [17] S. Preschutti, HCSM-35: The Daughter Structure of the CKM Matrix, HCSM White Paper Series (2026).
- [18] S. Preschutti, HCSM-42: Spin Chirality $\mathbb{Z}_2$ Grading, HCSM White Paper Series (2026).
- [19] S. Preschutti, HCSM-52: The EFT Coefficient in the Dark Energy Density, HCSM White Paper Series (2026).
- [20] S. Preschutti, HCSM-53: Dark Energy Density and the DME Entropy-Well Normalization, HCSM White Paper Series (2026).