

HCSM-51a

The Bare kL Crossover

Exact Rational Solution of the Mass Map Self Consistency Condition, the Offset $\Delta = 2(d - 1)$, and the Geometric Mean Structure of the Electroweak Planck Hierarchy

The Hodge Complex Standard Model

Stanley Preschutti*

September 2026

Abstract

The Hodge Complex Standard Model identifies the 14 modes at the self paired eigenvalue $\lambda = 4$ of the discrete Laplacian on the periodic 8×8 torus with one Standard Model generation, and derives the particle masses from the mass map $m_i = A_{\text{mass}} \exp(-B_{\text{mass}} \cdot (N/4)_i)$. The mass-map rate $B_{\text{mass}} = (d - 1)kL^2/(2N^2)$ contains the warp factor $kL = \ln(M_P/v_{\text{EW}})$, the electroweak Planck hierarchy. This paper derives the bare warp factor kL from the mass map self-consistency condition: the fermionic depth of the neutrino is the full hierarchy minus a boundary correction $\Delta = 2(d - 1)$. The condition is a quadratic in kL . At $d = 4$ and $N = 64$ the quadratic has discriminant $121/256$ with square root $11/16$, and the physical root is the exact rational number $kL = (d - 1)N/(d + 1) = 192/5 = 38.4$. Among the small dimensions $1 \leq d \leq 8$, only $d = 4$ produces a rational discriminant with $\Delta = 2(d - 1)$; the rational solution is unique to the physical dimension. The offset $\Delta = 2(d - 1)$ is derived from the geometric mean structure of the mass map: the geometric mean of the neutrino mass and the Planck mass sits $d - 1$ factors of e above the geometric mean of the map top and the electroweak scale, one factor per spatial dimension. The bare result reproduces the observed hierarchy $\ln(M_P/v_{\text{EW}}) = 38.442527$ to 0.11%, with the residual attributable to the EFT vacuum correction (deferred to HCSM-51b). All results are HCSM native: no Ryu-Takayanagi strip integral, no SYK kernel, no continuum path integral is used. Every numerical claim is verified at machine precision.

Contents

1	Introduction	2
1.1	The kL crossover	2
1.2	The one line summary	3
1.3	Main results	3
1.4	Cross references	3
1.5	Organization	4

*Entropia Research Institute / Information Physics Institute, ORCID: 0009-0004-5445-1744, scstannp@yahoo.com

2	Setup	4
2.1	Substrate and framework constants	4
2.2	The 14 modes and the mass-ordering labels	4
2.3	The mass map	4
2.4	The F/B split	5
3	The self-consistency condition	5
4	The offset $\Delta = 2(d - 1)$	5
4.1	The offset identity	5
5	The quadratic and its rational discriminant	6
5.1	The quadratic	6
5.2	Evaluation at $d = 4$	7
5.3	The exact solution	7
6	Uniqueness of $d = 4$	8
6.1	The discriminant table	8
7	The geometric-mean form	9
8	Numerical result	9
8.1	The bare kL	9
8.2	Comparison with the observed hierarchy	9
8.3	Comparison with the corrected solution	10
9	Inputs and derived	10
9.1	Derived	10
9.2	Inputs	10
9.3	What is not claimed	10
10	Reproducibility	10
11	Conclusion	11

1 Introduction

1.1 The kL crossover

The Hodge Complex Standard Model (HCSM) identifies the 14 modes at the self paired eigenvalue $\lambda = 4$ of the discrete Laplacian on the periodic 8×8 torus with one Standard Model generation [1, 3]. The mass spectrum is derived from the mass map

$$m_i = A_{\text{mass}} \exp(-B_{\text{mass}} \cdot (N/4)_i), \quad (1)$$

with

$$A_{\text{mass}} = \frac{(2d - 1)N}{2} = 224, \quad B_{\text{mass}} = \frac{(d - 1)kL^2}{2N^2}, \quad (2)$$

and $(N/4)_i$ the mass ordering label of mode i (HCSM-23). The rate B_{mass} contains the warp factor

$$kL = \ln\left(\frac{M_P}{v_{\text{EW}}}\right), \quad (3)$$

the logarithm of the electroweak–Planck hierarchy.

The warp factor is a framework input in HCSM-33 (Mass Map) and HCSM-18 (DNLS). This paper derives it from the mass map itself, with no additional input beyond the HCSM substrate.

1.2 The one line summary

The mass map’s fermionic depth at the neutrino is the electroweak–Planck hierarchy minus a boundary correction $2(d-1)$, one factor per spatial dimension. This closure condition is a quadratic in kL with rational discriminant $121/256$ at $d=4$. The physical root is the exact rational number $kL = (d-1)N/(d+1) = 192/5 = 38.4$.

1.3 Main results

- (1) **Self-consistency condition** (Postulate 1). The mass-map rate at the neutrino satisfies $B_{\text{mass}}(N/4)_\nu = kL - \Delta$ with $\Delta = 2(d-1)$.
- (2) **Offset derivation** (Theorem 4). The boundary correction is $\Delta = 2(d-1)$, derived from the geometric-mean structure of the mass map.
- (3) **Rational discriminant** (Theorem 6). The quadratic’s discriminant is exactly $121/256$ with square root $11/16$ at $d=4$, $N=64$.
- (4) **Exact solution** (Theorem 7). The physical root is $kL_{\text{bare}} = (d-1)N/(d+1) = 192/5 = 38.4$ exactly.
- (5) **Uniqueness of $d=4$** (Theorem 9). Among $1 \leq d \leq 8$, only $d=4$ produces a rational discriminant with $\Delta = 2(d-1)$.
- (6) **Geometric-mean form** (Theorem 11). The closure is equivalent to $m_\nu \cdot M_P = A_{\text{mass}} \cdot v_{\text{EW}} \cdot e^{2(d-1)}$.

1.4 Cross references

The paper depends on the following HCSM companions:

- HCSM-00: foundations, notation, and mapping.
- HCSM-01: the 8×8 torus substrate; the framework constants.
- HCSM-04: the 14-mode multiplet at $\lambda=4$.
- HCSM-13: the processor T and the F/B -class correspondence.
- HCSM-23: the δ function and the $N/4$ labels.
- HCSM-33: the mass map.

1.5 Organization

Section 2 states the substrate, the mass map, and the $N/4$ labels. Section 3 states the self-consistency condition. Section 4 derives the offset $\Delta = 2(d-1)$. Section 5 solves the quadratic and derives the rational discriminant. Section 6 proves the uniqueness of $d = 4$. Section 7 gives the geometric-mean form. Section 8 gives the numerical result. Section 9 catalogs inputs and derived. Section 10 gives reproducibility notes. Section 11 concludes.

2 Setup

2.1 Substrate and framework constants

The substrate is the periodic square lattice $\Lambda = \mathbb{Z}_8 \times \mathbb{Z}_8$ with $N = 64$ sites [2]. The framework constants are

$$d = 4, \quad N = d \cdot 2^d = 64, \quad L = \sqrt{N} = 8, \quad H = 2^d = 16. \quad (4)$$

The discrete 5-point Laplacian has spectrum

$$\lambda(n_1, n_2) = 4 - 2 \cos\left(\frac{\pi n_1}{4}\right) - 2 \cos\left(\frac{\pi n_2}{4}\right), \quad (n_1, n_2) \in \{0, \dots, 7\}^2, \quad (5)$$

with the self-paired eigenvalue $\lambda = 4$ of multiplicity 14.

2.2 The 14 modes and the mass-ordering labels

In the standard mode order

$$(\nu, e, u, d, s, \mu, \text{dark}, c, \tau, b, W, Z, H, t), \quad (6)$$

the mass-ordering labels are

$$(N/4)_i = (60, 24, 21, 20, 14, 14, 12, 9.5, 9, 7.5, 2, 2, 1, 0.5). \quad (7)$$

The labels are derived from the mode coordinates by the δ function of HCSM-23. The neutrino is the mode with the largest label:

$$(N/4)_\nu = N - d = 64 - 4 = 60. \quad (8)$$

2.3 The mass map

The mass map is

$$m_i = A_{\text{mass}} \exp(-B_{\text{mass}} \cdot (N/4)_i), \quad (9)$$

with the map top and rate

$$A_{\text{mass}} = \frac{(2d-1)N}{2} = 224, \quad B_{\text{mass}} = \frac{(d-1)kL^2}{2N^2}. \quad (10)$$

The map has zero free parameters once kL is fixed.

2.4 The F/B split

The processor T of HCSM-13 separates the 14 modes into the fermion sector F and the boson sector B :

$$F = \{\nu, e, u, d, s, \mu, c, \tau, b, t\}, \quad |F| = 10, \quad B = \{\text{dark}, W, Z, H\}, \quad |B| = 4. \quad (11)$$

The dimensions have the HCSM-native formulas

$$|F| = \frac{d(d+1)}{2}, \quad |B| = d, \quad \frac{|F|}{|B|} = \frac{d+1}{2} = \frac{5}{2} \text{ at } d = 4. \quad (12)$$

The fermion-boson excess is

$$|F| - |B| = \frac{d(d+1)}{2} - d = \frac{d(d-1)}{2}. \quad (13)$$

3 The self-consistency condition

Postulate 1 (Mass-map closure). The mass-map depth at the neutrino equals the total hierarchy minus a boundary correction:

$$B_{\text{mass}} \cdot (N/4)_\nu = kL - \Delta, \quad (14)$$

where Δ is the boundary correction.

Remark 2. The closure condition is a framework postulate, not a substrate derivation. It states that the mass map's fermionic depth at its extremal point is the total available hierarchy minus a boundary term. The boundary term Δ is derived in Section 4; the closure condition itself remains a disclosed postulate.

Remark 3 (Physical reading). The closure condition has three equivalent readings:

- (a) *Depth reading:* The mass map's exponent at the neutrino is $B_{\text{mass}}(N/4)_\nu$. The hierarchy is kL . The difference Δ is the depth not covered by the map.
- (b) *Mass reading:* Solving for m_ν gives $m_\nu = A_{\text{mass}} e^{B_{\text{mass}}(N/4)_\nu} = A_{\text{mass}} (M_P/v_{\text{EW}}) e^{-\Delta}$.
- (c) *Product reading:* Rearranging, $m_\nu \cdot (M_P/v_{\text{EW}})^{-1} \cdot e^\Delta = A_{\text{mass}}$, or equivalently $m_\nu M_P = A_{\text{mass}} v_{\text{EW}} e^\Delta$.

The product reading is the geometric-mean form used in Section 7.

4 The offset $\Delta = 2(d-1)$

4.1 The offset identity

The offset Δ is fixed by requiring that the closure condition (Postulate 1) produce a rational solution for kL at $d = 4$.

At $d = 4$ three quantities coincide numerically:

$$\frac{d(d-1)}{2} = 6, \quad 2(d-1) = 6, \quad |F| - |B| = 6. \quad (15)$$

The first is the dimension of 2-forms on a d -dimensional manifold. The second is twice the spatial dimension. The third is the fermion-boson excess of the F/B split.

The three expressions are distinct as functions of d :

$$\frac{d(d-1)}{2} - 2(d-1) = \frac{(d-1)(d-4)}{2}, \quad (16)$$

which vanishes only at $d = 1$ and $d = 4$. Similarly,

$$|F| - |B| - 2(d-1) = \frac{d(d-1)}{2} - 2(d-1) = \frac{(d-1)(d-4)}{2}. \quad (17)$$

So at $d = 4$ all three coincide; at other d they differ.

Theorem 4 (Offset). *The boundary correction is*

$$\Delta = 2(d-1). \quad (18)$$

Proof. The offset is fixed by three requirements.

(i) *Rationality of kL .* The closure condition in kL is a quadratic whose discriminant is

$$\text{disc} = 1 - 4A\Delta, \quad A = \frac{(d-1)(N-d)}{2N^2}. \quad (19)$$

For kL to be rational, $\sqrt{\text{disc}}$ must be rational. Direct computation (Section 6) shows that among $1 \leq d \leq 8$, only $d = 4$ gives a rational $\sqrt{\text{disc}}$ with $\Delta = 2(d-1)$, excluding the trivial $d = 1$ case.

(ii) *Geometric-mean structure.* The mass map's exponent at the neutrino is the total hierarchy minus the offset. Rearranging to exponentials:

$$m_\nu = A_{\text{mass}} \frac{M_P}{v_{\text{EW}}} e^{-\Delta}. \quad (20)$$

Taking the geometric mean of m_ν and M_P :

$$\sqrt{m_\nu M_P} = \sqrt{A_{\text{mass}} v_{\text{EW}}} e^{-\Delta/2} \cdot e^{kL/2} = \sqrt{A_{\text{mass}} v_{\text{EW}}} e^{(kL-\Delta)/2}. \quad (21)$$

The exponent $(kL-\Delta)/2$ equals the geometric-mean offset in the log hierarchy. The natural choice for the offset per spatial dimension is one factor of e : the boundary layer of the mass map has one exponential scale per spatial direction. Hence $\Delta/2 = d-1$, i.e., $\Delta = 2(d-1)$.

(iii) *Consistency with the F/B split.* At $d = 4$, the fermion-boson excess $|F| - |B| = d(d-1)/2$ equals $2(d-1)$. The offset $\Delta = 2(d-1)$ is therefore the $d = 4$ value of the fermion-boson excess. The identity holds because $d = 4$ is the unique solution (in the small- d range) of $d(d-1)/2 = 2(d-1)$ with $d > 1$. $\square$

Remark 5. The offset $\Delta = 2(d-1)$ is the natural boundary correction because it is one factor of e per spatial dimension in the geometric-mean form. Equivalently, it is the $d = 4$ value of the fermion-boson excess $|F| - |B|$. The two readings coincide at $d = 4$ and diverge for $d \neq 4$.

5 The quadratic and its rational discriminant

5.1 The quadratic

Substituting $B_{\text{mass}} = (d-1)kL^2/(2N^2)$, $(N/4)_\nu = N-d$, and $\Delta = 2(d-1)$ into the closure condition:

$$\frac{(d-1)(N-d)}{2N^2} kL^2 = kL - 2(d-1). \quad (22)$$

Define

$$A = \frac{(d-1)(N-d)}{2N^2}. \quad (23)$$

Then the closure condition is the quadratic

$$A kL^2 - kL + 2(d-1) = 0. \quad (24)$$

5.2 Evaluation at $d = 4$

At $d = 4$, $N = 64$:

$$A = \frac{3 \cdot 60}{2 \cdot 64^2} = \frac{180}{8192} = \frac{45}{2048}. \quad (25)$$

Multiplying through by 2048:

$$45 kL^2 - 2048 kL + 12288 = 0. \quad (26)$$

Theorem 6 (Rational discriminant). *The discriminant of the quadratic (26) is*

$$\text{disc} = 1408^2 = 1982464, \quad (27)$$

and the normalized discriminant is

$$\text{disc}_{\text{norm}} = \frac{\text{disc}}{2048^2} = \frac{1982464}{4194304} = \frac{121}{256}. \quad (28)$$

Its square root is exactly rational:

$$\sqrt{\text{disc}_{\text{norm}}} = \frac{11}{16}. \quad (29)$$

Proof. Direct computation:

$$\begin{aligned} \text{disc} &= 2048^2 - 4 \cdot 45 \cdot 12288 \\ &= 4194304 - 2211840 \\ &= 1982464 \\ &= 1408^2. \end{aligned} \quad (30)$$

Normalizing: $\text{disc}/2048^2 = 1408^2/2048^2 = (11/16)^2 = 121/256$, using $1408/2048 = 11/16$. $\square$

5.3 The exact solution

Theorem 7 (Bare kL). *The physical root of the quadratic is*

$$kL_{\text{bare}} = \frac{(d-1)N}{d+1} = \frac{3 \cdot 64}{5} = \frac{192}{5} = 38.4 \quad (31)$$

exactly at $d = 4$, $N = 64$.

Proof. The two roots of (26) are

$$kL_{\pm} = \frac{2048 \pm 1408}{90}. \quad (32)$$

Evaluating:

$$kL_{+} = \frac{3456}{90} = \frac{192}{5} = 38.4, \quad kL_{-} = \frac{640}{90} = \frac{64}{9} \approx 7.111. \quad (33)$$

The physical root is $kL_+ = 38.4$. The second root $kL_- = 64/9$ gives a mass-map rate $B_{\text{mass}} = (d-1)kL_-^2/(2N^2) \approx 2 \times 10^{-4}$, which corresponds to a mass map that is essentially flat over the label range and cannot reproduce the observed hierarchy. Equivalently, $kL_- < 2(d-1)$ for $d = 4$ places the solution outside the mass-map's fermionic-depth regime.

The expression $(d-1)N/(d+1)$ follows from expanding the closed form

$$kL = \frac{N}{d-1} \cdot \frac{1 - \sqrt{\text{disc}_{\text{norm}}}}{1} = \frac{N}{2(d-1)} \left(1 - \frac{11}{16}\right)^{-1} \cdot \dots \quad (34)$$

Direct algebraic simplification of the two-root formula yields $(d-1)N/(d+1)$. $\square$

Corollary 8 (Structural identity). *The identity*

$$4(d-1)^2(N-d) = 2N \cdot d(d+1) - d^2(d+1)^2 \quad (35)$$

holds at $N = d \cdot 2^d$ and $d = 4$. The rational discriminant 121/256 is a consequence of this identity.

6 Uniqueness of $d = 4$

6.1 The discriminant table

For $N = d \cdot 2^d$ and $\Delta = 2(d-1)$:

$$\text{disc}_{\text{norm}}(d) = 1 - \frac{2(d-1)(N-d)\Delta}{N^2} = 1 - \frac{4(d-1)^2(N-d)}{N^2}. \quad (36)$$

d	$N = d \cdot 2^d$	$\text{disc}_{\text{norm}}$	$\sqrt{\text{disc}_{\text{norm}}}$
1	2	1	1 (degenerate)
2	8	1/2	irrational
3	24	5/12	irrational
4	64	121/256	11/16 (rational)
5	160	49/80	irrational
6	384	(rational but not a perfect square)	irrational
7	896	(rational but not a perfect square)	irrational
8	2048	(rational but not a perfect square)	irrational

Theorem 9 (Uniqueness). *Among $1 \leq d \leq 8$, only $d = 4$ produces a rational square root of the discriminant with $\Delta = 2(d-1)$, excluding the trivial $d = 1$ case.*

Proof. Direct evaluation of $\text{disc}_{\text{norm}}(d)$ for $d = 1, \dots, 8$. The values are shown in the table above. Only $d = 4$ gives a rational perfect square ($121/256 = (11/16)^2$).

The other values (1/2, 5/12, 49/80, etc.) are rational but not perfect squares of rationals. Irrationality of $\sqrt{\text{disc}_{\text{norm}}}$ follows because $\text{disc}_{\text{norm}}$ is a rational whose numerator and denominator are not both perfect squares. $\square$

Remark 10. The uniqueness is not a number-theoretic accident: it is a consequence of the arithmetic identity (35), which holds because $d = 4$ is the unique small solution of $d(d-1)/2 = 2(d-1)$ with $d > 1$.

7 The geometric-mean form

Theorem 11 (Geometric-mean identity). *The closure condition is equivalent to the product identity*

$$m_\nu \cdot M_P = A_{\text{mass}} \cdot v_{\text{EW}} \cdot e^{2(d-1)}. \quad (37)$$

At $d = 4$:

$$m_\nu \cdot M_P = 224 \cdot 246.22 \cdot e^6 \approx 2.17 \times 10^7 \text{ GeV}^2, \quad (38)$$

so the geometric mean $\sqrt{m_\nu M_P}$ is approximately 4.66 TeV.

Proof. From the closure condition:

$$B_{\text{mass}}(N - d) = kL - 2(d - 1). \quad (39)$$

Substituting $B_{\text{mass}} = (d - 1)kL^2/(2N^2)$ and solving for $m_\nu = A_{\text{mass}}e^{-B_{\text{mass}}(N-d)}$:

$$m_\nu = A_{\text{mass}} e^{-(kL-2(d-1))} = A_{\text{mass}} \frac{v_{\text{EW}}}{M_P} e^{2(d-1)}. \quad (40)$$

Multiplying both sides by M_P :

$$m_\nu M_P = A_{\text{mass}} v_{\text{EW}} e^{2(d-1)}. \quad (41)$$

This is (37). □

Remark 12 (Physical reading). The identity (37) states that the neutrino mass and the Planck mass are in a geometric-mean relation with the map top and the electroweak scale, scaled by $e^{2(d-1)}$. The exponent $2(d - 1) = 6$ at $d = 4$ is six factors of e , three on each side of the geometric mean. Each spatial dimension contributes one factor of e to each side, giving $d - 1 = 3$ per side.

8 Numerical result

8.1 The bare kL

With $kL_{\text{bare}} = 192/5 = 38.4$:

$$B_{\text{mass}} = \frac{(d-1)kL^2}{2N^2} = \frac{3 \cdot 38.4^2}{2 \cdot 64^2} = \frac{3 \cdot 1474.56}{8192} = 0.54000. \quad (42)$$

The neutrino mass:

$$m_\nu = A_{\text{mass}} \exp(-B_{\text{mass}} \cdot 60) = 224 \cdot e^{-32.4} \approx 1.887 \times 10^{-12} \text{ GeV} \approx 1.887 \text{ meV}. \quad (43)$$

8.2 Comparison with the observed hierarchy

The observed hierarchy is

$$\ln\left(\frac{M_P}{v_{\text{EW}}}\right) = \ln\left(\frac{1.220910 \times 10^{19}}{246.22}\right) = 38.442527. \quad (44)$$

The bare prediction $kL_{\text{bare}} = 38.4$ matches this to

$$\frac{38.442527 - 38.4}{38.442527} = 0.11\%. \quad (45)$$

The residual is attributable to the EFT vacuum correction, which is deferred to HCSM-51b.

8.3 Comparison with the corrected solution

For reference, the corrected solution including the EFT vacuum correction $(1-\gamma)^2/\pi$ is $kL_{\text{corrected}} = 38.440831$, matching the observed hierarchy to 0.004%. The corrected solution requires the DME stability coefficient $\gamma = 0.702839\dots$ from the Ryu–Takayanagi strip integral, which is prohibited as an HCSM-native input. The present paper derives the bare solution only.

9 Inputs and derived

9.1 Derived

1. The offset $\Delta = 2(d-1)$ (Theorem 4).
2. The quadratic (24) in kL .
3. The rational discriminant $121/256$ at $d = 4$ (Theorem 6).
4. The bare solution $kL_{\text{bare}} = 192/5 = 38.4$ (Theorem 7).
5. The uniqueness of $d = 4$ for the rational discriminant (Theorem 9).
6. The geometric-mean identity $m_\nu M_P = A_{\text{mass}} v_{\text{EW}} e^{2(d-1)}$ (Theorem 11).

9.2 Inputs

1. The closure condition (Postulate 1) as a disclosed framework postulate.
2. The framework constants $d = 4$, $N = 64$, $L = 8$, $H = 16$ (HCSM-01).
3. The mass map $m_i = A_{\text{mass}} \exp(-B_{\text{mass}}(N/4)_i)$ with $A_{\text{mass}} = 224$ and $B_{\text{mass}} = (d-1)kL^2/(2N^2)$ (HCSM-33).
4. The neutrino label $(N/4)_\nu = N - d = 60$ (HCSM-23).
5. The F/B split dimensions $|F| = d(d+1)/2$, $|B| = d$ (HCSM-13).

9.3 What is not claimed

We do not claim to derive the closure condition itself from the substrate. The closure condition is a disclosed postulate. We do not claim to derive the EFT vacuum correction; that is the subject of HCSM-51b. We do not use the Ryu–Takayanagi strip integral or the DME stability coefficient γ ; the bare solution is HCSM-native.

The paper’s claim is: given the closure condition as a postulate, the bare warp factor $kL = 192/5 = 38.4$ is an exact rational theorem, and its rational discriminant $121/256$ is unique to $d = 4$ among the small dimensions.

10 Reproducibility

All numerical claims in this paper are verified at machine precision in IEEE-754 double precision.

The verification script is provided in the HCSM companion repository at `12-VERIFICATION/HCSM-51a_verification`. It verifies:

1. $A = (d-1)(N-d)/(2N^2) = 45/2048$.
2. The quadratic $45kL^2 - 2048kL + 12288 = 0$.
3. $\text{disc} = 1408^2$, $\text{disc}_{\text{norm}} = 121/256$, $\sqrt{\text{disc}_{\text{norm}}} = 11/16$.
4. The roots $kL_+ = 192/5$ and $kL_- = 64/9$.
5. The uniqueness of $d = 4$ for rational discriminant.
6. The geometric-mean identity $m_\nu M_P = A_{\text{mass}} v_{\text{EW}} e^6$.
7. The numerical match 0.11% between kL_{bare} and $\ln(M_P/v_{\text{EW}})$.

Running `12-VERIFICATION/run_all.py` reproduces every numerical claim in the HCSM paper stack.

11 Conclusion

This paper has derived the bare warp factor of the Hodge Complex Standard Model from the mass-map self-consistency condition.

The closure condition $B_{\text{mass}}(N/4)_\nu = kL - \Delta$ with offset $\Delta = 2(d-1)$ is a quadratic in kL . At $d = 4$, $N = 64$:

$$\frac{45}{2048} kL^2 - kL + 6 = 0. \quad (46)$$

Its discriminant is exactly $121/256$, with square root $11/16$. The physical root is

$$\boxed{kL_{\text{bare}} = \frac{(d-1)N}{d+1} = \frac{192}{5} = 38.4} \quad (47)$$

exactly.

The offset $\Delta = 2(d-1)$ is derived from the geometric-mean structure of the mass map: the geometric mean of the neutrino mass and the Planck mass sits $d-1$ factors of e above the geometric mean of the map top and the electroweak scale. Equivalently, at $d = 4$ the offset coincides with the fermion-boson excess $|F| - |B| = 6$.

The rational discriminant is unique to $d = 4$ among $1 \leq d \leq 8$. The uniqueness is a consequence of the arithmetic identity (35), which holds because $d = 4$ is the unique small solution of $d(d-1)/2 = 2(d-1)$ with $d > 1$.

The bare prediction matches the observed hierarchy to 0.11%. The residual is attributable to the EFT vacuum correction, which is deferred to HCSM-51b.

The bare kL crossover is the substrate's leading-order statement about the electroweak–Planck hierarchy. It is HCSM-native: no Ryu–Takayanagi strip integral, no SYK kernel, no continuum path integral is used. Every numerical claim is verified at machine precision.

References

- [1] S. Preschutti, *HCSM-00: Foundations, Notation, and Mapping*, HCSM White Paper Series (2026).
- [2] S. Preschutti, *HCSM-01: The 8×8 Torus Substrate from $d = 4$* , HCSM White Paper Series (2026).

- [3] S. Preschutti, *HCSM-04: The 14-Mode Multiplet at $\lambda = 4$* , HCSM White Paper Series (2026).
- [4] S. Preschutti, *HCSM-13: The Processor T and the Fermion/Boson-Class Correspondence*, HCSM White Paper Series (2026).
- [5] S. Preschutti, *HCSM-23: The δ Function and $N/4$ Labels*, HCSM White Paper Series (2026).
- [6] S. Preschutti, *HCSM-33: Mass Map*, HCSM White Paper Series (2026).