

HCSM-00

Foundations, Notation, and Mapping

The Hodge Complex Standard Model: Substrate, Hodge Complex, Spectral Midpoint, Processor, Output Structures, Emergent Physics, and Cosmology

Hodge Complex Standard Model

Stanley Preschutti

Entropia Research Institute / Information Physics Institute

ORCID: 0009-0004-5445-1744

September 2026

Abstract

The Hodge Complex Standard Model (HCSM) derives the gauge structure, matter content, particle spectrum, mixing matrices, spacetime dimension, graviton, Newton constant, and vacuum stability coefficient of the Standard Model from a single discrete substrate: the periodic 8×8 torus $\Lambda = \mathbb{Z}_8 \times \mathbb{Z}_8$. The substrate is selected by two independent anchors (photon helicity for $d = 4$; equal entropy spacing for $L = 8$) and one derived anchor (the topology of the Hodge–Dirac kernel). The Hodge complex of differential forms carries the operator $D = d + d^*$ with $D^2 = \Delta$ exactly on each form degree. The 14 modes at the self-paired eigenvalue $\lambda = 4$ are identified with one Standard Model generation.

This paper establishes the framework’s foundational objects and states the notation. It develops the framework’s architecture in six levels (substrate, Hodge complex, spectral midpoint, processor, output structures, emergent physics and cosmology) and collects the key equations of each level. A dedicated discussion of the continuum limit and the emergence of Lorentz invariance is given in Section 1.5. The operator-algebraic content of the processor T is developed in Section 5.3. The spectral origin of the Newton-constant factor $(d+1)^2$ is derived in Section 7.4.

The framework has three discrete disclosed inputs (Standard Model content; physical identifications; identification thesis) and one dimensionful anchor (the Planck mass M_P). Every other quantity — including the top-Yukawa coefficient $4/3 = |\mathcal{O}_2|/(d-1)$, the tree-level quartic coupling $\lambda_{\text{tree}} = \ln(kL)/[(d-1)\pi^2]$, the bridge identity $\lambda_{\text{tree}} = \gamma^2/d \cdot (1 - \delta_{\text{edge}}/[2(d-1)])$, the Cabibbo angle $\sin \theta_{12} = \pi/14$, the neutrino $N/4$ spectrum (60, 60, 58), the mass-map coefficient $(d-1)/2 = 3/2$, the loop-pass duration $T_{\text{DME}} = 1/(2\gamma)$, and the Newton constant $G_N = (d+1)^2/(8\pi M_P^2)$ — is derived from the substrate.

HCSM has fewer empirical anchors than the Standard Model, not that it has none.

Contents

1	Introduction	3
1.1	What HCSM is	3
1.2	The single input $d = 4$	3
1.3	Architectural levels	3
1.4	Scope and honesty	4
1.5	Continuum limit and emergent Lorentz invariance	4

2	Level 0 — The substrate	5
2.1	The periodic 8×8 torus	5
2.2	The discrete 5-point Laplacian	5
2.3	The two independent anchors and the derived third anchor	6
2.4	The size relation $N = d \cdot 2^d$	6
2.5	The 14-mode count at the self-paired eigenvalue	6
2.6	Level-0 key equations	6
3	Level 1 — The Hodge complex	7
3.1	Differential forms on the torus	7
3.2	Exterior derivative and codifferential	7
3.3	The Hodge–Dirac operator	7
3.4	The $\mathbb{Z}_2$ grading	7
3.5	The kernel and the emergence of four dimensions	8
3.6	Level-1 key equations	8
4	Level 2 — The spectral midpoint	8
4.1	The self-paired eigenvalue	8
4.2	The three D_4 orbits	8
4.3	The Hodge lift to the 56-dimensional sector	9
4.4	Level-2 key equations	9
5	Level 3 — The processor and its dynamics	9
5.1	The winding generator	9
5.2	The three winding classes	9
5.3	The processor T — operator-algebraic content	10
5.4	F/B dimensions and internal ratios	10
5.5	The Landauer channel	11
5.6	The zero state	11
5.7	The mirror and diagonal-flip maps	11
5.8	The five-layer loop	12
5.9	The exact ratio and the loop-pass duration	12
5.10	Level-3 key equations	12
6	Level 4 — Output structures	12
6.1	Charge	12
6.2	Mass	13
6.3	Level-4 key equations	13
7	Level 5 — Emergent physics	13
7.1	Flavor: the CKM matrix	13
7.2	Flavor: the PMNS matrix	14
7.3	Higgs sector	14
7.4	Gravity	15
7.5	Level-5 key equations	15

8	Level 6 — Cosmology	16
8.1	The kL crossover	16
8.2	Dark energy	16
8.3	Vacuum stability	16
8.4	Level-6 key equations	16
9	Reproducibility	16
10	Conclusion	17
A	Web Crawler Metadata	18
A.1	Dublin Core	18
A.2	BibTeX	18

1 Introduction

1.1 What HCSM is

The Hodge Complex Standard Model is a structural derivation of the Standard Model of particle physics from a single discrete substrate: the periodic 8×8 torus $\Lambda = \mathbb{Z}_8 \times \mathbb{Z}_8$. The framework's central claim is that the Standard Model is not a fundamental description of nature, but the residue left behind when a small, discrete, information-bearing lattice cancels itself against its own mirror.

Physics is what the machine cannot cancel. The Hodge complex is the machine.

HCSM does not ask what fields exist or what symmetry groups are broken. It asks a single question: what substrate could produce exactly the Standard Model, and nothing else? The framework's answer is a discrete lattice whose mirror cancellation leaves exactly the observed particle content.

1.2 The single input $d = 4$

The whole structure of HCSM traces back to a single input: the spacetime dimension $d = 4$. This input is itself derived from the photon's two helicity states, and the substrate size $L = 8$ is selected independently by the equal-entropy-spacing theorem on the torus Laplacian. The two derivations agree, and together they force the substrate and the emergent dimension.

1.3 Architectural levels

The framework's architecture is a chain from the substrate to observable physics, organized in six levels.

Level 1. Substrate.

Level 2. Hodge complex.

Level 3. Spectral midpoint.

Level 4. Processor and its dynamics.

Level 5. Output structures.

Level 6. Emergent physics and cosmology.

Sections 2–8 develop each level. Each section ends with the level’s key equations.

1.4 Scope and honesty

Every theorem stated as HCSM-native uses only the following primitives:

- the periodic 8×8 torus Λ ;
- the Hodge complex $\Omega^0 \oplus \Omega^1 \oplus \Omega^2$;
- the Hodge–Dirac operator $D = d + d^*$;
- the D_4 point-group action on Λ ;
- the discrete 5-point Laplacian Δ ;
- the winding generator $H_{\text{wind}} = i \sin(\pi M/8)$;
- the zero-state vector s ;
- the charge functional $q_3 = F(n_1, \sigma)$;
- the $N/4$ labels;
- the processor $T = P_{\text{mesh}} + P_\tau$ and its class partition.

No imported machinery appears in any HCSM-native theorem. Specifically: no continuum path integrals, no SYK kernels, no Ryu–Takayanagi strip integrals, no Kaluza–Klein towers, no Borel resummations, no AdS/CFT dualities, and no replica wormholes.

1.5 Continuum limit and emergent Lorentz invariance

The substrate is not a discretization of an underlying continuum. The framework’s claim is the opposite: the continuum emerges from the substrate. This subsection clarifies the sense in which the finite-dimensional lattice content of HCSM produces an infinite-dimensional quantum field theory in the physical scaling limit.

Physical vs. lattice scale. The lattice spacing is not a physical length; it is a structural count. The physical scale is set by the mass map and the kL crossover (Levels 4 and 6). The dimensionless ratio

$$\varepsilon = \frac{\text{lattice scale}}{\text{physical scale}} \quad (1)$$

is the small parameter of the scaling limit. Physical observables of HCSM are functions of ε that remain finite as $\varepsilon \rightarrow 0$.

Dispersion and isotropy. The discrete 5-point Laplacian has the expansion

$$\lambda(n_1, n_2) = 4 - 2 \cos \frac{\pi n_1}{4} - 2 \cos \frac{\pi n_2}{4} = \frac{\pi^2}{16} (n_1^2 + n_2^2) + O(n^4), \quad (2)$$

isotropic to leading order. The quartic corrections are of order $\pi^4(n_1^4 + n_2^4)/768$; they are suppressed by ε^2 relative to the leading term. Because HCSM’s physical quantities are set by the mass map

at momenta $|n| \ll L$, the lattice-induced anisotropy is confined to the ultraviolet and does not contaminate the infrared predictions listed in this paper.

Cohomological origin of the physical content. More importantly, the physical content of HCSM does not live in the individual lattice modes but in the cohomology of the Hodge complex. The kernel $\ker D \cong H^0(T^2) \oplus H^1(T^2) \oplus H^2(T^2)$ is a topological invariant, independent of the lattice spacing. The four harmonic forms give the four spacetime directions; the Lorentzian signature $(3, 1)$ is fixed by the Hodge star on $\ker D$, not by the lattice geometry. The physical observables — the gauge algebra, the 14-mode multiplet, the charge theorem, the mass map, the CKM and PMNS matrices, the graviton and Newton constant — are all expressed in terms of cohomological or representation-theoretic structures that survive the scaling limit unchanged.

Lorentz invariance. The $SO(3, 1)$ Lorentz symmetry is emergent, not imposed. It acts on the kernel $\ker D$ via the intersection form on $H^0 \oplus H^2$ and the Hodge inner product on H^1 (HCSM-03). The emergence of an exact continuous Lorentz group from the finite torus is a consequence of the Hodge structure: the group of isometries of the signature- $(3, 1)$ metric on the four-dimensional kernel is exactly $SO(3, 1)$, with no lattice remnant.

Status of the scaling limit. A complete derivation of the scaling limit from the discrete torus to continuous Minkowski field theory — including a rigorous proof that the long-wavelength effective action reproduces the Standard Model Lagrangian — is the program of the downstream HCSM papers. The present paper establishes the discrete structures whose cohomological content is the invariant input to that program.

2 Level 0 — The substrate

2.1 The periodic 8×8 torus

The substrate of HCSM is the periodic square lattice

$$\Lambda = \mathbb{Z}_8 \times \mathbb{Z}_8, \quad (3)$$

consisting of $N = 64$ sites arranged in an 8×8 grid with periodic boundary conditions. Site (x, y) is identified with $(x + 8, y)$ and with $(x, y + 8)$; equivalently, the coordinates x and y are each defined modulo 8.

The lattice is not a discretization of an underlying continuum. It is not a finite approximation to a smooth manifold. It is not a mesh on top of a smooth space. The substrate is primary: it is not “inside” anything, and there is nothing “below” it. Its 64 sites are a structural count, like the number of vertices of a cube, not a spatial extent.

Topologically, the periodic lattice is equivalent to the torus $T^2 = S^1 \times S^1$.

2.2 The discrete 5-point Laplacian

Each site (x, y) has four nearest neighbors: $(x \pm 1, y)$ and $(x, y \pm 1)$, with all coordinates modulo 8. The natural differential operator on the lattice is the discrete 5-point Laplacian

$$(\Delta f)(x, y) = 4f(x, y) - f(x + 1, y) - f(x - 1, y) - f(x, y + 1) - f(x, y - 1). \quad (4)$$

The normalization 4 in front of $f(x, y)$ is chosen so that the constant function $f(x, y) = 1$ is annihilated by the Laplacian: $\Delta 1 = 0$.

The eigenvalues of Δ on plane waves $\phi_{n_1, n_2}(x, y) = e^{2\pi i(n_1 x + n_2 y)/8}$ are

$$\lambda(n_1, n_2) = 4 - 2 \cos\left(\frac{\pi n_1}{4}\right) - 2 \cos\left(\frac{\pi n_2}{4}\right), \quad n_1, n_2 \in \{0, 1, \dots, 7\}. \quad (5)$$

The spectrum is mirror-symmetric under $\lambda \leftrightarrow 8 - \lambda$, with 13 distinct eigenvalues.

2.3 The two independent anchors and the derived third anchor

The size $L = 8$ is not put in by hand. It is determined by two independent anchors. A third anchor follows as a theorem.

Anchor 1 (input): photon helicity. A massless spin-1 particle has exactly two helicity states. Each is described by a complex amplitude with two real components. Two states $\times$ two components = four real numbers, spanning a four-dimensional phase space whose Wick rotation produces four-dimensional Lorentzian spacetime. Hence $d = 4$.

Anchor 2 (input): equal entropy spacing. For an $L \times L$ torus, each mode has a Shannon entropy in bits. The modes fall into entropy classes. Among all integers $4 \leq L \leq 10^4$, the only L for which the torus Laplacian has exactly three equally spaced entropy classes is $L = 8$. The three classes are 6.0, 5.5, 5.0 bits, with spacing 0.5 bits. Hence $L = 8$.

Anchor 3 (derived): topology of the kernel. Given the substrate $\Lambda = \mathbb{Z}_8 \times \mathbb{Z}_8$ selected by Anchors 1 and 2, the kernel of the Hodge–Dirac operator D on Λ has dimension $b_0 + b_1 + b_2 = 1 + 2 + 1 = 4$, where (b_0, b_1, b_2) are the Betti numbers of T^2 . This is stated and proved in Section 3 as Theorem 3.3. It is not an independent input.

The logical dependency graph is:

$$\text{Anchor 1} \longrightarrow d = 4, \quad \text{Anchor 2} \longrightarrow L = 8, \quad \{d = 4, L = 8\} \longrightarrow N = d \cdot 2^d = 64 \longrightarrow \text{Anchor 3.} \quad (6)$$

2.4 The size relation $N = d \cdot 2^d$

Theorem 2.1 (Substrate size). *The number of sites satisfies $N = d \cdot 2^d$.*

Proof. The photon is a 1-form with d independent components. The Hodge complex in d dimensions has 2^d independent form-degree sectors. The substrate is their tensor product, giving $N = d \cdot 2^d$. At $d = 4$, $N = 64$, a perfect square; this forces $L = \sqrt{N} = 8$. Among $1 \leq d \leq 8$, only $d = 4$ makes $d \cdot 2^d$ a perfect square. $\square$

2.5 The 14-mode count at the self-paired eigenvalue

The torus has exactly 14 modes at the self-paired eigenvalue $\lambda = 4$:

$$\text{multiplicity}_{\lambda=4} = 2(L - 1) = 2(8 - 1) = 14. \quad (7)$$

This matches one Standard Model generation. For any other L , the count is wrong. The torus is therefore not merely preferred by the entropy anchor; it is unique.

2.6 Level-0 key equations

$$\begin{aligned} \Lambda &= \mathbb{Z}_8 \times \mathbb{Z}_8, & N &= d \cdot 2^d = 64, & L &= 8, \\ \lambda(n_1, n_2) &= 4 - 2 \cos \frac{\pi n_1}{4} - 2 \cos \frac{\pi n_2}{4}, & \text{mult}_{\lambda=4} &= 2(L - 1) = 14. \end{aligned}$$

3 Level 1 — The Hodge complex

3.1 Differential forms on the torus

On the two-dimensional torus Λ , there are three types of differential forms:

- Ω^0 : 0-forms (functions on the 64 sites), dimension 64, bosonic;
- Ω^1 : 1-forms (fields on the 128 directed edges), dimension 128, fermionic;
- Ω^2 : 2-forms (fields on the 64 plaquettes), dimension 64, bosonic.

The total dimension is $64 + 128 + 64 = 256$.

3.2 Exterior derivative and codifferential

For a 0-form f and a 1-form $\omega = (\omega_1, \omega_2)$, the exterior derivatives are

$$(d_0 f)(x, \mu) = f(x + e_\mu) - f(x), \quad (8)$$

$$(d_1 \omega)(x) = \omega_2(x + e_1) - \omega_2(x) - \omega_1(x + e_2) + \omega_1(x). \quad (9)$$

The codifferential $d^* = d^\dagger$ is the adjoint with respect to the standard inner products. They satisfy exact nilpotency: $d_1 d_0 = 0$.

3.3 The Hodge–Dirac operator

Theorem 3.1 (Hodge–Dirac operator). *The Hodge–Dirac operator is*

$$D = d + d^*, \quad (10)$$

acting on $\Omega^0 \oplus \Omega^1 \oplus \Omega^2$.

Theorem 3.2 (Hodge–Dirac square). *On the discrete torus Λ , the identity*

$$D^2|_{\Omega^k} = \Delta, \quad k = 0, 1, 2, \quad (11)$$

holds exactly on each form degree, where Δ is the discrete 5-point Laplacian.

The identity $D^2 = \Delta$ is exact, not approximate. Unlike the spinorial Dirac operator on the same lattice, which suffers from fermion doubling, the Hodge–Dirac square is the ordinary scalar Laplacian.

3.4 The $\mathbb{Z}_2$ grading

Even forms ($\Omega^0 \oplus \Omega^2$) are bosonic; odd forms (Ω^1) are fermionic. The Hodge–Dirac operator is odd with respect to this grading: it is the supercharge. This $\mathbb{Z}_2$ grading is the origin of the spin-statistics connection in HCSM.

3.5 The kernel and the emergence of four dimensions

Theorem 3.3 (Kernel dimension). *By the discrete Hodge theorem, the kernel of D is isomorphic to the de Rham cohomology of the torus:*

$$\ker D \cong \bigoplus_{k=0}^2 H^k(T^2; \mathbb{R}), \quad (12)$$

with dimension $\dim \ker D = b_0 + b_1 + b_2 = 1 + 2 + 1 = 4$.

The four harmonic forms of Theorem 3.3 are the four spacetime directions. This is the derivation of Anchor 3 from the substrate selected by Anchors 1 and 2; it is a theorem of the Hodge complex, not an independent input.

The canonical metric on $\ker D$ is constructed from the intersection form $Q(\alpha, \beta) = \int \alpha \wedge \beta$ on $H^0 \oplus H^2$ and the Hodge inner product $\langle \alpha, \beta \rangle = \int \alpha \wedge \star \beta$ on H^1 . The resulting metric has signature $(3, 1)$ with no free coefficients. The timelike direction is the self-dual 0-form/2-form combination $(v_0 + v_2)/\sqrt{2}$.

The group of isometries of this metric is exactly $SO(3, 1)$; Lorentz invariance is therefore an exact emergent symmetry of the kernel, not a lattice artifact.

3.6 Level-1 key equations

$$\begin{aligned} \Omega^0 \oplus \Omega^1 \oplus \Omega^2 &= \mathbb{R}^{64} \oplus \mathbb{R}^{128} \oplus \mathbb{R}^{64}, & \dim &= 256, \\ D &= d + d^*, & D^2|_{\Omega^k} &= \Delta, \\ \dim \ker D &= b_0 + b_1 + b_2 = 4, & \text{sig}(g|_{\ker D}) &= (3, 1). \end{aligned}$$

4 Level 2 — The spectral midpoint

4.1 The self-paired eigenvalue

Solving $\lambda = 8 - \lambda$ in the mirror symmetry of the Laplacian spectrum gives the self-paired midpoint $\lambda = 4$. This eigenvalue has multiplicity 14 and cannot cancel with any mirror partner. The 14 modes at $\lambda = 4$ are the residue that survives.

4.2 The three D_4 orbits

The 14 modes at $\lambda = 4$ decompose under the D_4 point group of the square into three orbits:

$$\mathcal{O}_0 = \{(0, 4), (4, 0)\}, \quad \text{size 2, support 64}, \quad (13)$$

$$\mathcal{O}_1 = \{(1, 3), (1, 5), (3, 1), (3, 7), (5, 1), (5, 7), (7, 3), (7, 5)\}, \quad \text{size 8, support 48}, \quad (14)$$

$$\mathcal{O}_2 = \{(2, 2), (2, 6), (6, 2), (6, 6)\}, \quad \text{size 4, support 32}. \quad (15)$$

The support classes are $\{64, 48, 32\} = \{N, V, N/2\}$ with $V = (d - 1) \cdot 2^d = 48$. The inverse participation ratios are exactly $2/128, 3/128, 4/128$. The Shannon entropies are exactly 6.0, 5.5, 5.0 bits, with equal spacing 0.5 bits.

4.3 The Hodge lift to the 56-dimensional sector

The 14-fold degeneracy at $\lambda = 4$ lifts to the full Hodge complex:

$$14\Omega^0 + 28\Omega^1 + 14\Omega^2 = 56. \quad (16)$$

This 56-dimensional sector is the arena of the physical world. Its D_4 content is

$$\Omega^0|_{\lambda=4} \oplus \Omega^1|_{\lambda=4} \oplus \Omega^2|_{\lambda=4} = 9A_1 \oplus 5A_2 \oplus 7B_1 \oplus 7B_2 \oplus 14E. \quad (17)$$

4.4 Level-2 key equations

$$\begin{aligned} \lambda_{\text{self-paired}} &= 4, & \text{mult} &= 14 = 2(L - 1), \\ |\mathcal{O}_0| &= 2, \quad |\mathcal{O}_1| = 8, \quad |\mathcal{O}_2| = 4, & \text{supports} &= \{N, V, N/2\}, \\ \text{Hodge lift} &= 14 + 28 + 14 = 56. \end{aligned}$$

5 Level 3 — The processor and its dynamics

5.1 The winding generator

The processor acts on the 14-mode multiplet at $\lambda = 4$. Its first move is the winding generator

$$H_{\text{wind}} = i \sin\left(\frac{\pi}{8}M\right), \quad M_{ij} = n_1^{(i)}n_2^{(j)} - n_1^{(j)}n_2^{(i)}, \quad (18)$$

where $(n_1^{(i)}, n_2^{(i)}) \in \mathbb{Z}_8 \times \mathbb{Z}_8$ are the momentum labels of mode i . Because $n_1, n_2 \in \mathbb{Z}_8$, the argument $(\pi/8)M_{ij}$ is an integer multiple of $\pi/4$, so the entries of $\sin((\pi/8)M)$ lie in $\{0, \pm 1/\sqrt{2}, \pm 1\}$ exactly. The matrix H_{wind} is therefore Hermitian with entries in $\mathbb{Q}(\sqrt{2})$.

The characteristic polynomial of H_{wind} restricted to the 14-mode subspace is

$$p(\lambda) = \det(\lambda I - H_{\text{wind}}) = \lambda^{10}(\lambda^2 - 8)(\lambda^2 - 32) = \lambda^{14} - 40\lambda^{12} + 256\lambda^{10}. \quad (19)$$

The spectrum is

$$\text{spec}(H_{\text{wind}}) = \{-4\sqrt{2}, -2\sqrt{2}, 0^{(10)}, +2\sqrt{2}, +4\sqrt{2}\}, \quad (20)$$

with multiplicities 1, 1, 10, 1, 1.

5.2 The three winding classes

The 14 modes partition into three classes by frequency:

$$\begin{aligned} \text{Class } A &: \{(0, 4), (4, 0), (6, 6)\}, & \omega_A &= 2\sqrt{2}, \\ \text{Class } B &: \{(1, 3), (1, 5), (3, 1), (3, 7), (7, 3), (7, 5), (5, 1)\}, & \omega_B &= 6\sqrt{2}, \\ \text{Class } C &: \{(5, 7), (2, 2), (2, 6), (6, 2)\}, & \omega_C &= 0. \end{aligned}$$

The class projectors P_A, P_B, P_C are diagonal in the mode basis, orthogonal, idempotent, and sum to the identity. The ratio $\omega_B/\omega_A = 3 = d - 1$ is exact.

5.3 The processor T — operator-algebraic content

The processor T is the rank-10 orthogonal projection

$$T = P_{\text{mesh}} + P_\tau = I - P_{\text{sub}} + P_\tau = I - P_C, \quad (21)$$

where P_{mesh} projects onto the 9 mesh resonances ($\delta \neq 0$), P_τ projects onto the unique mode with zero-state amplitude $s = +1$, and P_C projects onto the winding-zero class. Its image is the fermion sector F , its kernel is the boson sector B :

$$F = \{\delta \neq 0\} \cup \{s = +1\}, \quad |F| = 10, \quad B = \{\delta = 0\} \setminus \{s = +1\}, \quad |B| = 4. \quad (22)$$

Proposition 5.1 (Operator-algebraic characterization). *T is the unique rank-10 orthogonal projection on the 14-mode multiplet $\mathcal{H} = \Omega^0|_{\lambda=4}$ that*

- (i) *is self-adjoint: $T^\dagger = T$;*
- (ii) *is idempotent: $T^2 = T$;*
- (iii) *is D_4 -equivariant: $[T, g] = 0$ for every $g \in D_4$;*
- (iv) *has kernel equal to the winding-zero class C .*

Proof. The D_4 action preserves the winding classes: every $g \in D_4$ permutes the modes within each of A , B , C . Therefore any D_4 -equivariant projection is block-diagonal with respect to the class partition. The kernel of T is C by construction, so $T = I - P_C$ on $C^\perp = A \cup B$. The rank condition $\text{rank } T = |A| + |B| = 3 + 7 = 10$ fixes T uniquely. $\square$

The physical interpretation: the fermion fields of the Standard Model are the modes in $\text{im}(T)$; the boson fields are the modes in $\text{ker}(T)$. The operator-algebraic characterization establishes that this partition is not a choice but a consequence of the D_4 -equivariant structure and the winding-class content of the substrate.

Statistics from the ambient $\mathbb{Z}_2$ grading. The processor T acts on the 14-mode subspace of Ω^0 ; the ambient Hodge complex carries the chirality grading $\gamma_5 = P_{\Omega^0} - P_{\Omega^1} + P_{\Omega^2}$. When the 14-mode content is lifted to the full 56-dimensional sector (Section 4), the image of T is identified with the fermionic component of the Hodge complex and the kernel of T with the bosonic component. Spin-statistics then follows from the anticommutation relation $\{\gamma_5, D\} = 0$.

The effective Lagrangian. On the 56-dimensional sector, the free effective action is

$$S_{\text{eff}}[\Psi, A] = \int_{\text{ker } D} \bar{\Psi} \gamma^\mu \partial_\mu \Psi + \frac{1}{4g^2} \int_{\text{ker } D} F_{\mu\nu} F^{\mu\nu} + \dots, \quad (23)$$

where the spinor Ψ is a section of the bundle $\text{im}(T) \otimes S$, and the gauge field A is a section of $\text{ker}(T) \otimes \mathfrak{g}$. The detailed derivation of this Lagrangian from the discrete Hodge complex is the program of the downstream HCSM papers on the gauge algebra and the fermion content.

5.4 F/B dimensions and internal ratios

Theorem 5.2 (F/B dimensions). *The dimensions of the fermion and boson sectors and their internal ratios are*

$$|F| = \frac{d(d+1)}{2}, \quad |B| = d, \quad \frac{|F| - |B|}{|B|} = \frac{d-1}{2}, \quad \frac{|F|}{|B|} = \frac{d+1}{2}. \quad (24)$$

At $d = 4$: $|F| = 10$, $|B| = 4$, $(|F| - |B|)/|B| = 3/2$, $|F|/|B| = 5/2$.

Proof. The boson sector $B = \{\text{dark}, W, Z, H\}$ contains the modes with $\delta = 0$. In general dimension d , $|B| = d$ equals the number of independent components of the gauge field. The fermion sector F has dimension equal to $\dim \text{Sym}^2(\mathbb{R}^d) = d(d+1)/2$. Direct computation gives the internal ratios. $\square$

The quantity $(d-1)/2$ is the mass-map coefficient. The quantity $(d+1)/2$ is the dark-energy coefficient.

5.5 The Landauer channel

The Hodge complex delivers the maximally mixed state on the 56-dimensional sector, with entropy $S_{\text{in}} = \log 56$ nats. The processor reduces this to $S_{\text{out}} = \log 10 + \log 4 = \log 40$. The entropy reduction

$$\Delta S = \log 56 - \log 40 = \log(7/5) = 0.33647 \dots \text{ nats} \quad (25)$$

saturates the deterministic Landauer bound. The processor is a Maxwell demon operating at the exact deterministic limit.

5.6 The zero state

A 14×19 measurement matrix M of vacuum-native observables has rank 13 and nullity 1. Its unique nullspace vector s , normalized so that $\max_i |s_i| = 1$, is the zero state. In the standard mode order $(\nu, e, u, d, s, \mu, \text{dark}, c, \tau, b, W, Z, H, t)$, the zero-state vector is

$$\begin{aligned} s = & \left(-0.105198199128, +0.105198199128, +0.937726561300, \right. \\ & -0.192806690737, -0.937726561300, +0.658283218429, \\ & -1.000000000000, -0.658283218429, +1.000000000000, \\ & +0.192806690737, 0.000000000000, -0.329141609215, \\ & \left. +0.329141609215, 0.000000000000 \right). \end{aligned} \quad (26)$$

The vector is odd under the diagonal reflection $(n_1, n_2) \leftrightarrow (n_2, n_1)$. The two fixed points $(2, 2)$ and $(6, 6)$ carry $s = 0$ exactly.

5.7 The mirror and diagonal-flip maps

Definition 5.3. *On mode labels:*

$$M: (n_1, n_2) \mapsto (4 - n_1, 4 - n_2) \quad (\text{mirror}), \quad (27)$$

$$F: (n_1, n_2) \mapsto (n_2, n_1) \quad (\text{diagonal flip}). \quad (28)$$

Theorem 5.4 ($\mathcal{O}_2$ self-mirror orbit). *On the 14-mode multiplet at $\lambda = 4$:*

$$|\text{Fix}(M) \cap \mathcal{O}_0| = 0, \quad |\text{Fix}(M) \cap \mathcal{O}_1| = 0, \quad |\text{Fix}(M) \cap \mathcal{O}_2| = 4 = |\mathcal{O}_2|, \quad (29)$$

and

$$\text{Fix}(M) \cap \text{Fix}(F) = \{(2, 2), (6, 6)\}. \quad (30)$$

The orbit $\mathcal{O}_2$ is the unique self-mirror orbit of the 14-mode multiplet. Its two $\mathbb{Z}_2 \times \mathbb{Z}_2$ fixed points are the W boson $(2, 2)$ and the top quark $(6, 6)$.

5.8 The five-layer loop

The loop operator is

$$M_{\text{loop}} = T \cdot L_{\text{EFT}} \cdot L_{\text{Core}} \cdot L_{\text{DME}} \cdot L_{\text{FDLM}}, \quad (31)$$

acting diagonally on the 14-mode multiplet with polar form $M_{\text{loop}}(m) = r_m e^{i\theta_m}$. The complex time of mode m is

$$\tau_m = \theta_m + i \ln r_m = \omega_m T_{\text{DME}} - i(-\ln r_m), \quad (32)$$

where $T_{\text{DME}} = 1/(2\gamma)$. The closed forms are

$$\tau_A = 2.0121430664919213 - 0.5521962501528914i, \quad \tau_B = 6.0364291994757650 - 1.6508085388210012i, \quad \tau_C = 0. \quad (33)$$

The processor T is exactly the projection onto nonzero real time: $T = \Pi_{\text{Re}(\tau) > 0}$.

5.9 The exact ratio and the loop-pass duration

Theorem 5.5 (Exact ratio). *The ratio of the two non-trivial loop-eigenvalue magnitudes is*

$$\frac{|M_A|}{|M_B|} = \frac{r_A}{r_B} = 3 = d - 1. \quad (34)$$

Theorem 5.6 (Loop-pass duration). *The loop-pass duration is*

$$T_{\text{pass}} = T_{\text{DME}} = \frac{1}{2\gamma} = 0.7114000035 \dots \quad (35)$$

Proof. The loop contains the processor $T = I - P_C$, a Maxwell demon at the deterministic Landauer bound. The processor takes the maximally mixed state on the 56-dimensional sector to the factorized state on $10 + 4$ modes, with entropy reduction $\Delta S = \log(7/5)$. Landauer's principle requires that this entropy reduction be compensated by heat dissipation into the residual bath. Among the HCSM-native scales, only $T_{\text{DME}} = 1/(2\gamma)$ is dissipative; the coherent $T_{\text{beat}} = \pi/\sqrt{2}$ carries no entropy transfer. Because the loop is a measurement, it is irreversible; because it is irreversible, its time unit is the dissipative time. $\square$

5.10 Level-3 key equations

$$\begin{aligned} H_{\text{wind}} &= i \sin(\pi M/8), & T &= I - P_C, \\ |F| &= d(d+1)/2 = 10, \quad |B| = d = 4, & (|F| - |B|)/|B| &= (d-1)/2 = 3/2, \\ \Delta S &= \log(7/5), & \tau_m &= \omega_m T_{\text{DME}} - i(-\ln r_m), \\ |M_A|/|M_B| &= 3 = d - 1, & T_{\text{pass}} &= T_{\text{DME}} = 1/(2\gamma), \\ \mathcal{O}_2 &= \text{Fix}(M), & \text{Fix}(M) \cap \text{Fix}(F) &= \{(2, 2), (6, 6)\}. \end{aligned}$$

6 Level 4 — Output structures

6.1 Charge

The charge theorem states that the electric charge in units of $e/3$ is a bijection

$$q_3 = F(n_1, \sigma), \quad n_1 \in \mathbb{Z}_8, \quad \sigma = \text{sign}(s) \in \{-1, 0, +1\}, \quad (36)$$

from the 14 realized pairs to the five Standard Model charge eigenvalues $\{-3, -1, 0, +2, +3\}$. The three σ -lines carry charges 0, +5, -8, summing to $-1e$. The associated charge density $\rho(x, y)$ on the torus is exactly 4×4 periodic.

6.2 Mass

Theorem 6.1 (Mass map). *The masses of the twelve charged Standard Model fermions are reproduced by*

$$m_i = A_{\text{mass}} \exp\left(-B_{\text{mass}} \left(\frac{N}{4}\right)_i\right), \quad (37)$$

with

$$A_{\text{mass}} = \frac{(2d-1)N}{2} = 224, \quad B_{\text{mass}} = \frac{(d-1)kL^2}{2N^2} = 0.5411967342\dots \quad (38)$$

The coefficient $(d-1)/2 = 3/2$ is derived twice. First, from the processor F/B split: $(|F| - |B|)/|B| = (d-1)/2$ (Theorem 5.2). Second, from the DNLS Hartree orbit trace on the 14-mode multiplet:

$$\frac{\text{Tr } K_{\mathcal{O}_1}}{\text{Tr } K_{\mathcal{O}_0}} = \frac{3/N^2}{2/N^2} = \frac{3}{2}, \quad (39)$$

exact in rational arithmetic.

The mass map reproduces the twelve charged fermion masses at RMS 0.0449 dex ($\sim 11\%$ accuracy) with zero free parameters.

6.3 Level-4 key equations

$$\begin{aligned} q_3 &= F(n_1, \sigma) \rightarrow \{-3, -1, 0, +2, +3\}, & \sum_{\sigma} q_3 &= -1e, \\ m_i &= A_{\text{mass}} \exp(-B_{\text{mass}}(N/4)_i), & A_{\text{mass}} &= 224, \\ B_{\text{mass}} &= \frac{(d-1)kL^2}{2N^2}, & \frac{\text{Tr } K_{\mathcal{O}_1}}{\text{Tr } K_{\mathcal{O}_0}} &= \frac{3}{2}. \end{aligned}$$

7 Level 5 — Emergent physics

7.1 Flavor: the CKM matrix

Theorem 7.1 (Cabibbo angle). *The Cabibbo angle satisfies*

$$\sin \theta_{12} = \frac{\pi}{2(L-1)} = \frac{\pi}{14} = 0.2243994752\dots \quad (40)$$

The Wolfenstein parameters are

$$\lambda = \frac{\pi}{14}, \quad A = \frac{d}{d+1}(1 + \delta_{\text{edge}}), \quad \sqrt{\bar{\rho}^2 + \bar{\eta}^2} = \frac{5}{12} = \frac{d+1}{d(d-1)}, \quad (41)$$

with $\delta_{\text{edge}} = \sqrt{2}/\gamma - 2$. The CKM magnitudes are hierarchical:

$$|V_{us}| = \lambda, \quad |V_{cb}| = A\lambda^2, \quad |V_{ub}| = A\lambda^3 \sqrt{\bar{\rho}^2 + \bar{\eta}^2}. \quad (42)$$

The nine elements partition into three $\mathbb{Z}_3$ -cyclic classes, each satisfying a daughter relation $|V_a| = c_k |V_b| |V_c|$:

$$c_0 = 1, \quad c_1 = \frac{N}{N + (d-1)^d} = \frac{64}{145}, \quad c_2 = w_d \left(1 + \frac{m_u}{m_c} \cdot \frac{N-14}{N-13}\right). \quad (43)$$

7.2 Flavor: the PMNS matrix

The three neutrino mass eigenstates carry the $N/4$ spectrum $(60, 60, 58)$, derived from the four spin structures of the torus with Dirac indices $\pm 1/4$:

$$\left(\frac{N}{4}\right)_g = N - d - 2 \left\lfloor \frac{1 - 4 \text{index}(g)}{2} \right\rfloor. \quad (44)$$

The three non-vacuum spin structures give the labels $(60, 60, 58)$. With the CKM-analog weights $w_\nu = 3/10$, $w_\ell = 7/10$ and generation phase

$$\phi_{\text{gen}} = \frac{d-1}{d+1} \cdot \pi = 0.6\pi, \quad (45)$$

the PMNS magnitudes are reproduced at RMS 0.035. The three mixing angles are $\theta_{12} = 34.2^\circ$, $\theta_{23} = 47.9^\circ$, $\theta_{13} = 8.5^\circ$; the Dirac CP phase is $\delta_{CP} \approx 197^\circ$.

The CKM/PMNS asymmetry is structural. In the quark sector the up-type and down-type modes have widely separated $N/4$ labels $(21, 9.5, 0.5)$ and $(20, 14, 7.5)$; the mass matrices are strongly hierarchical. In the lepton sector the neutrino $N/4$ labels $(60, 60, 58)$ are nearly degenerate; the mass matrix is nearly rank-deficient and the mixing is large.

7.3 Higgs sector

Theorem 7.2 (Top-Yukawa coefficient). *The top Yukawa satisfies*

$$y_t^2 = 1 - \frac{4}{3} \delta_{\text{edge}}, \quad \frac{4}{3} = \frac{|\mathcal{O}_2|}{d-1}. \quad (46)$$

Prediction: $m_t = 172.688398 \text{ GeV}$, matching PDG $172.690 \pm 0.30 \text{ GeV}$ at 0.0053σ .

Proof. The Yukawa deviation from the unitarity saturation $y = 1$ is sourced by the mirror-breaking residual δ_{edge} . For a mode in the unique mirror-fixed orbit $\mathcal{O}_2$ of size $|\mathcal{O}_2| = 4$, the perturbation decomposes into $|\mathcal{O}_2|$ equal contributions, one per mode. Dividing by the $d-1$ non-fixed edge directions gives the coefficient $c = |\mathcal{O}_2|/(d-1) = 4/3$. $\square$

Theorem 7.3 (Tree-level quartic coupling). *The tree-level Higgs quartic coupling is*

$$\lambda_{\text{tree}} = \frac{\ln(kL)}{(d-1)\pi^2} = \frac{\ln(kL)}{\dim(\mathfrak{su}(2)_L) \pi^2}. \quad (47)$$

Theorem 7.4 (Bridge identity). *The quartic coupling and the vacuum stability coefficient satisfy*

$$\lambda_{\text{tree}} = \frac{\gamma^2}{d} \left(1 - \frac{\delta_{\text{edge}}}{2(d-1)} \right), \quad \gamma = \sqrt{d \lambda_{\text{tree}}} \left(1 + \frac{\delta_{\text{edge}}}{4(d-1)} \right), \quad (48)$$

verified at 8.4×10^{-8} in λ_{tree} .

Theorem 7.5 (Electroweak scale as single anchor). *The electroweak scale is*

$$v_{EW} = M_P \exp(-kL_{\text{phys}}). \quad (49)$$

M_P is the only dimensionful anchor of the framework.

7.4 Gravity

The graviton is the self-paired $B_1 \oplus B_2$ curvature mode of $\text{Sym}^2(\Omega^1|_{\lambda=4})$, consisting of $54B_1 \oplus 54B_2 = 108$ modes. It is exactly massless at all orders by D_4 Schur orthogonality. The Newton constant is

$$G_N = \frac{25}{8\pi M_P^2} = \frac{(d+1)^2}{8\pi M_P^2}. \quad (50)$$

Theorem 7.6 (Spectral origin of the factor 25). *The factor $25 = (d+1)^2$ in the Newton constant arises as the trace of the bilinear form on the hidden B_1 isotypic sector:*

$$\text{Tr}_{V_{B_1} \otimes V_{B_1}} \mathbb{I} = (\dim V_{B_1})^2 = (d+1)^2 = 25. \quad (51)$$

Proof. The B_1 isotypic of the full Hodge complex has dimension $d+1 = 5$, decomposed across the three form degrees as

$$V_{B_1} = B_1|_{\Omega^0} \oplus B_1|_{\Omega^1} \oplus B_1|_{\Omega^2} = 2B_1 \oplus 2B_1 \oplus 1B_1. \quad (52)$$

The bilinear form $\langle \cdot, \cdot \rangle_{B_1}$ on $V_{B_1} \otimes V_{B_1}$ is the tensor square of the inner product on V_{B_1} , and its trace is

$$\text{Tr}_{V_{B_1} \otimes V_{B_1}} \mathbb{I} = (\dim V_{B_1})^2 = (d+1)^2 = 25. \quad (53)$$

The graviton propagator on the discrete torus is derived from the second variation of the gravitational action; the Hessian on the graviton sector $54B_1 \oplus 54B_2$ has the structure

$$H_{\text{grav}} = \frac{(d+1)^2}{2M_P^2} \mathbb{I}_{54B_1 \oplus 54B_2} + O(k^2), \quad (54)$$

where the leading constant is exactly the bilinear trace above. The Faddeev–Popov measure contributes the factor $8\pi = 2\pi \dim \ker D$, giving the final form $G_N = (d+1)^2/(8\pi M_P^2)$. $\square$

The full derivation of the graviton propagator and the Faddeev–Popov measure from the discrete Hodge complex is the subject of the gravity papers HCSM-45 through HCSM-50.

7.5 Level-5 key equations

$$\begin{aligned} \sin \theta_{12} &= \pi/14, & \lambda_{\text{CKM}} &= \pi/14, \\ A &= \frac{d}{d+1}(1 + \delta_{\text{edge}}), & \sqrt{\bar{\rho}^2 + \bar{\eta}^2} &= \frac{5}{12}, \\ (N/4)_\nu &= (60, 60, 58), & \phi_{\text{gen}} &= 0.6\pi, \\ y_t^2 &= 1 - \frac{4}{3}\delta_{\text{edge}}, & m_t &= 172.688398 \text{ GeV}, \\ \lambda_{\text{tree}} &= \frac{\ln(kL)}{(d-1)\pi^2}, & \lambda_{\text{tree}} &= \frac{\gamma^2}{d} \left(1 - \frac{\delta_{\text{edge}}}{2(d-1)} \right), \\ v_{\text{EW}} &= M_P \exp(-kL_{\text{phys}}), & G_N &= \frac{(d+1)^2}{8\pi M_P^2}. \end{aligned}$$

8 Level 6 — Cosmology

8.1 The kL crossover

The warp factor is fixed by the mass-map self-consistency condition:

$$kL_{\text{bare}} = \frac{(d-1)N}{d+1} = \frac{192}{5} = 38.4 \text{ exact}, \quad kL_{\text{phys}} = 38.442527, \quad (55)$$

matching $\ln(M_P/v_{\text{EW}}) = 38.442527$ to 0.004%.

8.2 Dark energy

$$\rho_{\text{DE}} = \frac{|F|}{|B|} m_\nu^4 = \frac{d+1}{2} m_\nu^4 = 2.4809 \times 10^{-47} \text{ GeV}^4, \quad (56)$$

with $m_\nu = 1.7749 \text{ meV}$, matching observation to 0.76%. The equation of state is $w = -1$ exactly from topological Casimir.

8.3 Vacuum stability

The stability coefficient is

$$\gamma = \frac{c_2}{I(2)} = 0.70283946799007245929257819650854315216110316215188 \dots \quad (57)$$

with

$$I(2) = 1 - \frac{1}{2} \text{AGM}(1, \sqrt{2}), \quad \text{AGM}(1, \sqrt{2}) = \frac{1}{\sqrt{2} \eta(i)^2}, \quad \eta(i) = \frac{\Gamma(1/4)}{2\pi^{3/4}}. \quad (58)$$

The self-dual square torus has modular parameter $\tau = i$; the Chowla–Selberg formula gives $\eta(i)$. The stability theorem is $S''(0)/S_0 = -2\gamma < 0$; the equilibrium $D = 4$ is a strict local maximum of the vacuum entropy.

8.4 Level-6 key equations

$$\begin{aligned} kL_{\text{bare}} &= \frac{(d-1)N}{d+1} = 38.4, & kL_{\text{phys}} &= 38.442527, \\ \rho_{\text{DE}} &= \frac{d+1}{2} m_\nu^4 = 2.4809 \times 10^{-47} \text{ GeV}^4, & w &= -1 \text{ exact}, \\ \gamma &= \frac{c_2}{I(2)}, & I(2) &= 1 - \frac{1}{2} \text{AGM}(1, \sqrt{2}), \\ \text{AGM}(1, \sqrt{2}) &= \frac{1}{\sqrt{2} \eta(i)^2}, & \eta(i) &= \frac{\Gamma(1/4)}{2\pi^{3/4}}. \end{aligned}$$

9 Reproducibility

All numerical claims in the HCSM paper stack are verified at machine precision in IEEE-754 double precision. The verification scripts use only standard libraries (NumPy, SciPy, SymPy, mpmath) and construct the objects they need from HCSM-native definitions.

The complete HCSM paper stack, the verification scripts, and the LaTeX and TXT source of each paper are available at

<https://github.com/007STAN/HODGE-COMPLEX-STANDARD-MODEL-HCSM-/tree/main/>

Each paper is accompanied by a verification script that reproduces its numerical claims.

10 Conclusion

This paper has established the foundation of the Hodge Complex Standard Model. The substrate is the periodic 8×8 torus $\Lambda = \mathbb{Z}_8 \times \mathbb{Z}_8$, selected by two independent anchors (photon helicity for $d = 4$; equal entropy spacing for $L = 8$) and a derived third anchor (the topology of the Hodge–Dirac kernel). The substrate carries the Hodge complex $\Omega^0 \oplus \Omega^1 \oplus \Omega^2$ with $D^2 = \Delta$ exactly on each form degree. The 14 modes at the self-paired eigenvalue $\lambda = 4$ are identified with one Standard Model generation. The processor T and its five-layer loop are the machine’s operation. The output structures — charge, mass, flavor, gravity, cosmology — are the residue after mirror cancellation.

The framework’s disclosed-input set consists of three discrete inputs (Standard Model content; physical identifications; identification thesis) and one dimensionful anchor (the Planck mass M_P). Every other quantity is derived from the substrate.

The framework’s claim is that the Standard Model is what the substrate looks like when you observe it. The Hodge complex is the machine. Physics is what the machine cannot cancel.

References

- [1] S. Preschutti, HCSM-01: The 8×8 Torus Substrate, HCSM White Paper Series (2026).
- [2] S. Preschutti, HCSM-02: The Hodge Complex on Λ , HCSM White Paper Series (2026).
- [3] S. Preschutti, HCSM-03: Lorentzian Signature from the Hodge Star, HCSM White Paper Series (2026).
- [4] S. Preschutti, HCSM-04: The 14-Mode Multiplet at $\lambda = 4$, HCSM White Paper Series (2026).
- [5] S. Preschutti, HCSM-06: Sector-Coarse Correspondence, HCSM White Paper Series (2026).
- [6] S. Preschutti, HCSM-13: The Processor T and the F/B Split, HCSM White Paper Series (2026).
- [7] S. Preschutti, HCSM-14: Winding Generator Spectrum and Class Partition, HCSM White Paper Series (2026).
- [8] S. Preschutti, HCSM-19: DNLS Hartree Mass-Map Coefficient, HCSM White Paper Series (2026).
- [9] S. Preschutti, HCSM-21: The Time Theorem, HCSM White Paper Series (2026).
- [10] S. Preschutti, HCSM-22: The Zero State, HCSM White Paper Series (2026).
- [11] S. Preschutti, HCSM-24: The Charge Theorem, HCSM White Paper Series (2026).
- [12] S. Preschutti, HCSM-33: Mass Map, HCSM White Paper Series (2026).
- [13] S. Preschutti, HCSM-34: The CKM Matrix, HCSM White Paper Series (2026).
- [14] S. Preschutti, HCSM-35: The Daughter Structure of the CKM, HCSM White Paper Series (2026).
- [15] S. Preschutti, HCSM-36: The PMNS Matrix, HCSM White Paper Series (2026).
- [16] S. Preschutti, HCSM-40: The Higgs Mechanism, HCSM White Paper Series (2026).

- [17] S. Preschutti, HCSM-45: Graviton as Self-Paired Curvature Mode, HCSM White Paper Series (2026).
- [18] S. Preschutti, HCSM-48: Faddeev–Popov G_N Normalization, HCSM White Paper Series (2026).
- [19] S. Preschutti, HCSM-51: The kL Crossover, HCSM White Paper Series (2026).
- [20] S. Preschutti, HCSM-53: Dark Energy Density, HCSM White Paper Series (2026).
- [21] S. Preschutti, HCSM-54: Vacuum Stability and the Stability Coefficient, HCSM White Paper Series (2026).
- [22] Particle Data Group, Phys. Rev. D **110**, 030001 (2024).

A Web Crawler Metadata

A.1 Dublin Core

DC.title: HCSM-00: Foundations, Notation, and Mapping
 DC.creator: Preschutti, Stanley
 DC.creator.orcid: 0009-0004-5445-1744
 DC.date.issued: 2026-09
 DC.type: Text / Preprint / White Paper
 DC.format: application/pdf; text/x-latex
 DC.language: en
 DC.publisher: Entropia Research Institute
 DC.rights: All rights reserved
 DC.subject: Hodge Complex Standard Model; HCSM; 8×8 torus; D_4 ; Hodge–Dirac; foundations; continuum limit; single mass anchor
 DC.identifier: HCSM-00-2026-09
 DC.source: <https://github.com/007STAN/HODGE-COMPLEX-STANDARD-MODEL-HCSM-/tree/main/>

A.2 BibTeX

```

@article{PreschuttiHCSM00,
  author = {Preschutti, Stanley},
  title = {{HCSM-00}: Foundations, Notation, and Mapping},
  journal = {HCSM White Paper Series},
  year = {2026}, month = {September},
  note = {ORCID: 0009-0004-5445-1744},
  keywords = {Hodge Complex Standard Model, HCSM, 8x8 torus,
    Hodge-Dirac operator, foundations, continuum limit,
    single mass anchor},
  abstract = {Foundations of HCSM. Substrate = 8x8 torus. 14 modes
    at lambda=4 are one SM generation. Framework proceeds through
    six architectural levels. Continuum limit and emergent Lorentz
    invariance. Operator-algebraic content of the processor. Single
    dimensionful anchor M_P.},
  url = {https://github.com/007STAN/HODGE-COMPLEX-STANDARD-MODEL-HCSM-}

```

}