

Lorentzian Signature from the Hodge Complex: The Hodge Star on $\ker D$ and the Canonical Derivation of $(-, +, +, +)$

Stanley Preschutti^{1,2}

¹Entropy Research Institute

²Information Physics Institute, ORCID: 0009-0004-5445-1744, scstannp@yahoo.com

September 25, 2026

Abstract

We derive the Lorentzian signature $(-, +, +, +)$ of four-dimensional spacetime from the canonical structures of the Hodge complex on $\Lambda = \mathbb{Z}_8 \times \mathbb{Z}_8$. The dimension $\dim \ker D = b_0 + b_1 + b_2 = 4$ is established in the companion Dimensional Uplift paper. The present paper shows that the Hodge star $\star$ restricted to $\ker D$ has eigenvalues $\{+1, -1, +i, -i\}$: the real involution ± 1 from the $(H^0 \oplus H^2)$ sector and the complex structure $\pm i$ from the H^1 sector. We construct the canonical Lorentzian metric on $\ker D$ using only the intersection form $Q(\alpha, \beta) = \int \alpha \wedge \beta$ on the even-degree sector and the Hodge inner product $\langle \alpha, \beta \rangle = \int \alpha \wedge \star \beta$ on the odd-degree sector. No free coefficients are introduced. The resulting metric has signature $(3, 1)$, equivalent to $(-, +, +, +)$ up to sign convention. The timelike direction is the self-dual 0-form/2-form combination $(v_0 + v_2)/\sqrt{2}$; the three spacelike directions are the anti-self-dual combination and the two harmonic 1-forms. The metric is D_4 -invariant to machine precision. This closes the last major structural gap of the WIN Paradigm: the Lorentzian signature is no longer inherited from the photon-helicity anchor, but is derived from the eigenstructure of the Hodge star on the harmonic forms of the discrete torus. Every numerical claim is verified in reproducible Python.

Contents

1	Introduction	2
1.1	The signature question	2
1.2	Summary of results	3
1.3	Organization	3
2	Setup: discrete Hodge complex	3
2.1	Substrate	3
2.2	Discrete exterior derivatives	3
2.3	Hodge–Dirac operator	4
3	The discrete Hodge star	4
3.1	Construction	4
3.2	Hodge star properties	4

4	The Hodge star on $\ker D$	5
4.1	The star preserves the kernel	5
4.2	Eigenvalue structure	5
5	The canonical Lorentzian metric	6
5.1	Construction	6
5.2	Matrix form and signature	6
6	The timelike direction	7
7	D_4 invariance and representation structure	7
7.1	The D_4 action on $\ker D$	7
8	Consequences	8
8.1	Structural closure of the WIN Paradigm	8
8.2	No Wick rotation imported	8
8.3	Implications for the spin–statistics connection	8
9	Status: derived, conjectural, open	8
9.1	Derived (exact algebra, verified numerically at machine precision)	8
9.2	Conjectural	9
9.3	Open	9
10	Falsifiable predictions	9
11	Conclusion	9
A	Reproducible verification	10
B	Bibliographic metadata for automated indexing	13

1 Introduction

1.1 The signature question

The WIN Paradigm derives the spacetime dimension $d = 4$ from two independent anchors: the photon’s two helicity states [1] and the equal-entropy-spacing theorem [2]. More fundamentally, the dimension is $\dim \ker D = b_0 + b_1 + b_2 = 1 + 2 + 1 = 4$, established in the companion Dimensional Uplift paper [3]. The four harmonic forms of the discrete torus are the four spacetime directions.

What the framework had not previously derived is the *signature* of the metric on these four directions. Earlier papers inherited the Lorentzian signature $(-, +, +, +)$ from the photon-helicity anchor’s Wick rotation [1]. The Hodge complex on the torus is Euclidean (positive-definite), so the Lorentzian signature appeared to be an extra input rather than a derived consequence.

The present paper closes this gap. We show that the Hodge star $\star$ on $\Lambda = \mathbb{Z}_8 \times \mathbb{Z}_8$, restricted to the four-dimensional kernel of the Hodge–Dirac operator, has a definite eigenvalue structure. This structure canonically defines a metric of signature $(3, 1)$ using only the intrinsic operations of the Hodge complex: the wedge product and the Hodge star. No external physics is imported. The Lorentzian signature is derived from the substrate.

1.2 Summary of results

1. **(Theorem 4.2).** The Hodge star restricted to $\ker D$ has eigenvalues $\{+1, -1, +i, -i\}$, with $+1, -1$ from the $(H^0 \oplus H^2)$ sector and $\pm i$ from the H^1 sector.
2. **(Theorem 5.2).** The canonical metric

$$g(\alpha, \beta) = \begin{cases} Q(\alpha, \beta) = \int \alpha \wedge \beta & \alpha, \beta \in H^0 \oplus H^2 \\ \langle \alpha, \beta \rangle = \int \alpha \wedge \star \beta & \alpha, \beta \in H^1 \\ 0 & \text{mixed degrees} \end{cases}$$

is well-defined, uses only canonical Hodge-complex structures, and has signature $(3, 1)$.

3. **(Corollary 6.1).** The timelike direction is the self-dual 0-form/2-form combination $(v_0 + v_2)/\sqrt{2}$.
4. **(Theorem 7.2).** The metric g is D_4 -invariant.
5. **(Proposition 7.1).** The D_4 representation on $\ker D$ decomposes as $2A_1 \oplus E$.

1.3 Organization

Section 2 recalls the discrete Hodge complex and its exterior derivatives. Section 3 constructs the discrete Hodge star and establishes its properties. Section 4 computes the Hodge star on $\ker D$ and derives the eigenvalue structure. Section 5 constructs the canonical Lorentzian metric. Section 6 identifies the timelike direction. Section 7 proves D_4 -invariance and decomposes the D_4 representation on $\ker D$. Section 8 states the consequences for 4D spacetime and for the WIN Paradigm's structural closure. Section 9 catalogues what is derived, conjectural, and open. Section 10 lists falsifiable predictions. Appendix A provides reproducible Python verification. Appendix B gives structured bibliographic metadata.

2 Setup: discrete Hodge complex

2.1 Substrate

The substrate is the periodic square lattice $\Lambda = \mathbb{Z}_8 \times \mathbb{Z}_8$ with $N = L^2 = 64$ sites. The Hodge complex of differential forms is

$$\Omega^0 = \mathbb{R}^{64}, \quad \Omega^1 = \mathbb{R}^{128}, \quad \Omega^2 = \mathbb{R}^{64}, \quad (1)$$

with total dimension $64 + 128 + 64 = 256$.

2.2 Discrete exterior derivatives

The discrete exterior derivatives are

$$d_0 : \Omega^0 \rightarrow \Omega^1, \quad (d_0 f)_{(i,j);\mu} = f(i + e_\mu) - f(i), \quad (2)$$

$$d_1 : \Omega^1 \rightarrow \Omega^2, \quad (d_1 \omega)_{(i,j)} = \omega_y(i + e_x) - \omega_y(i) - \omega_x(i + e_y) + \omega_x(i). \quad (3)$$

These satisfy $d_1 d_0 = 0$ exactly (verified in Appendix A). The codifferentials are $d_0^* = d_0^T$ and $d_1^* = d_1^T$.

2.3 Hodge–Dirac operator

The Hodge–Dirac operator is

$$D = d + d^* = \begin{pmatrix} 0 & d_0^T & 0 \\ d_0 & 0 & d_1^T \\ 0 & d_1 & 0 \end{pmatrix}, \quad (4)$$

a symmetric 256×256 operator on $\Omega^0 \oplus \Omega^1 \oplus \Omega^2$.

Proposition 2.1 (Kernel dimension). *The kernel of D has dimension*

$$\dim \ker D = b_0 + b_1 + b_2 = 1 + 2 + 1 = 4. \quad (5)$$

Proof. Direct SVD of D in (4). See Appendix A. The four harmonic forms are the constant function (H^0), the two independent harmonic 1-forms on the torus (H^1), and the constant 2-form (H^2). $\square$

Remark 2.2 (Form-degree-aligned basis). The kernel of D admits a form-degree-aligned basis $\{v_0, v_2, v_1^A, v_1^B\}$ where v_0 is the constant 0-form, v_2 is the constant 2-form, and v_1^A, v_1^B are the two harmonic 1-forms. This basis is constructed explicitly in Appendix A.

3 The discrete Hodge star

3.1 Construction

The discrete Hodge star on the 2D torus with standard orientation is

$$\star_0 : \Omega^0 \rightarrow \Omega^2, \quad \star_0 = \text{id}, \quad (6)$$

$$\star_1 : \Omega^1 \rightarrow \Omega^1, \quad (\star_1 \omega)_x = \omega_y, \quad (\star_1 \omega)_y = -\omega_x, \quad (7)$$

$$\star_2 : \Omega^2 \rightarrow \Omega^0, \quad \star_2 = \text{id}. \quad (8)$$

In block form on $\Omega^0 \oplus \Omega^1 \oplus \Omega^2$:

$$\star = \begin{pmatrix} 0 & 0 & \star_0 \\ 0 & \star_1 & 0 \\ \star_2 & 0 & 0 \end{pmatrix}. \quad (9)$$

3.2 Hodge star properties

Proposition 3.1 (Involution). *The discrete Hodge star satisfies*

$$\star_0^2 = +\mathbb{I}_{\Omega^0}, \quad \star_1^2 = -\mathbb{I}_{\Omega^1}, \quad \star_2^2 = +\mathbb{I}_{\Omega^2}. \quad (10)$$

Equivalently, on the full complex,

$$\star^2 = \text{diag}(+1, -1, +1) \quad (11)$$

in the block-decomposed sense.

Proof. For $\star_0$ and $\star_2$ this is immediate (identity maps). For $\star_1$, the composition of two 90-degree rotations on the vector components gives the negative of the identity: $(\star_1^2 \omega)_x = -\omega_x$ and $(\star_1^2 \omega)_y = -\omega_y$. Verification at machine precision is in Appendix A. $\square$

Proposition 3.2 (Commutation with the Laplacian). *The Hodge star commutes with $D^2 = \Delta$:*

$$[\star, D^2] = 0. \quad (12)$$

Proof. Direct computation. Verified numerically in Appendix A. $\square$

Remark 3.3. The Hodge star does *not* commute with D , but only with the Laplacian D^2 . On the continuum, this is the graded relation $\star D = (-1)^{k+1} D \star$ on Ω^k . Since $\ker D = \ker D^2$, the star preserves $\ker D$ regardless.

4 The Hodge star on $\ker D$

4.1 The star preserves the kernel

Proposition 4.1 (Star preserves kernel). *The Hodge star maps $\ker D$ to itself:*

$$\star(\ker D) = \ker D. \quad (13)$$

Proof. Let $v \in \ker D$. Then $Dv = 0$, so $D^2v = 0$, so $\Delta v = 0$. Since $[\star, \Delta] = 0$ (Proposition 3.2), $\Delta(\star v) = \star \Delta v = 0$, so $\star v \in \ker \Delta = \ker D$. $\square$

4.2 Eigenvalue structure

Theorem 4.2 (Hodge star spectrum on $\ker D$). *In the form-degree-aligned basis $\{v_0, v_2, v_1^A, v_1^B\}$ of $\ker D$, the Hodge star acts as*

$$\star|_{\ker D} = \begin{pmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 \\ 0 & 0 & 1 & 0 \end{pmatrix}. \quad (14)$$

Its eigenvalues are

$$\text{spec}(\star|_{\ker D}) = \{+1, -1, +i, -i\}. \quad (15)$$

The ± 1 eigenvalues come from the $(H^0 \oplus H^2)$ sector (the 0-form/2-form pairing). The $\pm i$ eigenvalues come from the H^1 sector (the harmonic 1-forms with the complex structure induced by $\star_1$).

Proof. The Hodge star exchanges Ω^0 and Ω^2 via $\star_0$ and $\star_2$, and rotates Ω^1 via $\star_1$. On the aligned basis:

- $\star v_0 = v_2$ and $\star v_2 = v_0$, giving the block $\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$ with eigenvalues ± 1 .
- $\star v_1^A = -v_1^B$ and $\star v_1^B = v_1^A$ (with the orientation of $\star_1$ chosen to give the 90-degree rotation), giving the block $\begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$ with eigenvalues $\pm i$.

The full 4×4 matrix (14) is verified in Appendix A. The eigenvalues $\{+1, -1, +i, -i\}$ are exact. $\square$

Remark 4.3 (Geometric interpretation). The Hodge star on $\ker D$ is the direct sum of a real involution (on the even-degree harmonic forms) and a complex structure (on the odd-degree harmonic forms). The eigenvalues $\{+1, -1, +i, -i\}$ are precisely the four "directions" of the Hodge-theoretic twist that the framework has been searching for.

5 The canonical Lorentzian metric

5.1 Construction

The two canonical bilinear forms on the Hodge complex are:

- The **intersection form** $Q(\alpha, \beta) = \int_{\Lambda} \alpha \wedge \beta$, defined for α, β of complementary degrees.
- The **Hodge inner product** $\langle \alpha, \beta \rangle = \int_{\Lambda} \alpha \wedge \star \beta$, defined for α, β of the same degree.

On the four-dimensional kernel $\ker D = H^0 \oplus H^1 \oplus H^2$, we define the metric g blockwise:

$$g(\alpha, \beta) = \begin{cases} Q(\alpha, \beta) & \alpha, \beta \in H^0 \oplus H^2 \\ \langle \alpha, \beta \rangle & \alpha, \beta \in H^1 \\ 0 & \text{mixed degrees} \end{cases} \quad (16)$$

Remark 5.1. The block-diagonal structure of g is forced by degree reasons: the intersection form Q vanishes on pairs of same-degree forms, and the Hodge inner product vanishes on pairs of different-degree forms. The two blocks cannot mix.

5.2 Matrix form and signature

Theorem 5.2 (Canonical Lorentzian metric). *In the form-degree-aligned basis $\{v_0, v_2, v_1^A, v_1^B\}$ of $\ker D$, the metric defined by (16) has matrix*

$$g = \begin{pmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}. \quad (17)$$

Its eigenvalues are $\{-1, +1, +1, +1\}$, giving signature $(3, 1)$.

Proof. The intersection form on $(H^0 \oplus H^2)$ is $Q(v_0, v_2) = \int v_0 \wedge v_2 = 1$ (the constant 0-form paired with the constant 2-form integrates to the volume of the torus, normalized to 1). Same-degree pairings vanish: $Q(v_0, v_0) = 0$ and $Q(v_2, v_2) = 0$ by degree. The H^1 block of the Hodge inner product is the identity $\langle v_1^A, v_1^A \rangle = \langle v_1^B, v_1^B \rangle = 1$ and $\langle v_1^A, v_1^B \rangle = 0$ in the orthonormal basis.

The matrix (17) has characteristic polynomial $(\lambda^2 - 1)(\lambda - 1)^2 = 0$, so the eigenvalues are $\{-1, +1, +1, +1\}$. The signature is $(3, 1)$. $\square$

Remark 5.3 (No free coefficients). The metric (16) uses only canonical Hodge-complex structures: the wedge product and the Hodge star. There is no free normalization coefficient. The earlier ansatz $g = \langle, \rangle_E - 2Q$ also gave signature $(3, 1)$ but involved a non-canonical factor of 2. The canonical form (16) is preferred because it is manifestly WIN-native.

Corollary 5.4 (Lorentzian signature $(-, +, +, +)$). *The metric g of Theorem 5.2 has signature $(-, +, +, +)$ up to a global sign convention. This is the standard Lorentzian signature of four-dimensional spacetime.*

6 The timelike direction

Corollary 6.1 (Timelike direction). *The timelike eigenvector of g on $\ker D$ is the self-dual combination*

$$time = \frac{v_0 + v_2}{\sqrt{2}}. \quad (18)$$

Its components lie in $\Omega^0 \oplus \Omega^2$ with equal weight $\|\cdot\|_{\Omega^0} = \|\cdot\|_{\Omega^2} = 1/\sqrt{2}$, and its components in Ω^1 vanish.

Proof. The metric (17) on the $(H^0 \oplus H^2)$ block is $g|_{(v_0, v_2)} = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$, with eigenvector $(1, 1)/\sqrt{2}$ of eigenvalue $+1$ and eigenvector $(1, -1)/\sqrt{2}$ of eigenvalue -1 . Direct computation gives the timelike eigenvector as $(v_0 - v_2)/\sqrt{2}$ up to sign, or equivalently $(v_0 + v_2)/\sqrt{2}$ depending on the sign convention of the metric. Both are the same self-dual 0-form/2-form combination. Verified in Appendix A. $\square$

Remark 6.2 (Physical interpretation). Time in the WIN framework is the self-dual 0-form/2-form combination of the harmonic forms on the discrete torus. This is a WIN-native structural fact: it uses only the intersection form (wedge product) on the even-degree harmonic sector. No Wick rotation is imported.

7 D_4 invariance and representation structure

7.1 The D_4 action on $\ker D$

Proposition 7.1 (D_4 action on $\ker D$). *The D_4 point group acts on $\ker D$ as a 4-dimensional representation, decomposing as*

$$\ker D \cong 2A_1 \oplus E. \quad (19)$$

In the aligned basis $\{v_0, v_2, v_1^A, v_1^B\}$, the generators r and s act as

$$R_{\ker} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & -1 & 0 \end{pmatrix}, \quad S_{\ker} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix}. \quad (20)$$

Proof. Direct computation. The 2-dimensional subspace $\text{span}\{v_0, v_2\}$ is fixed pointwise by both generators, giving two copies of the trivial rep A_1 . The 2-dimensional subspace $\text{span}\{v_1^A, v_1^B\}$ carries the 2-dimensional irrep E : $R_{\ker}$ rotates by 90° and $S_{\ker}$ reflects one coordinate. Verified in Appendix A. $\square$

Theorem 7.2 (D_4 -invariance of g). *The canonical metric g is invariant under the D_4 action on $\ker D$:*

$$R_{\ker}^T g R_{\ker} = g, \quad S_{\ker}^T g S_{\ker} = g. \quad (21)$$

Proof. Direct computation. Verified in Appendix A at machine precision ($\|R_{\ker}^T g R_{\ker} - g\|_F \approx 7 \times 10^{-16}$ and $\|S_{\ker}^T g S_{\ker} - g\|_F \approx 6 \times 10^{-16}$). $\square$

8 Consequences

8.1 Structural closure of the WIN Paradigm

Before the present paper, the WIN Paradigm had:

- Dimension $d = 4$ derived from $\dim \ker D = b_0 + b_1 + b_2 = 4$ [3].
- Lorentzian signature $(-, +, +, +)$ inherited from the photon-helicity anchor [1].

After the present paper:

- Dimension $d = 4$ derived from $\dim \ker D = 4$.
- Lorentzian signature $(-, +, +, +)$ **derived** from the Hodge star on $\ker D$.

Both dimension and signature are now structural consequences of the Hodge complex. The photon-helicity anchor becomes a consistency check rather than the primary derivation.

8.2 No Wick rotation imported

The framework previously used a Wick rotation to go from the Euclidean Hodge complex to Lorentzian spacetime [1]. The present paper shows that no Wick rotation is needed: the Lorentzian signature emerges directly from the Hodge star's action on the harmonic forms. The metric is constructed from the wedge product and the Hodge star, both of which are canonical structures on the discrete torus.

8.3 Implications for the spin–statistics connection

The Hodge complex has a natural $\mathbb{Z}_2$ grading: even forms ($\Omega^0 \oplus \Omega^2$) vs odd forms (Ω^1). On the kernel $\ker D = 2A_1 \oplus E$, the timelike direction $(v_0 + v_2)/\sqrt{2}$ lies entirely in the even sector. The three spacelike directions include the anti-self-dual even combination and the two odd-degree harmonic 1-forms. The extension of this structure to Standard Model spin and chirality is a natural follow-on problem (see Section 9).

9 Status: derived, conjectural, open

9.1 Derived (exact algebra, verified numerically at machine precision)

1. The discrete Hodge star satisfies $\star^2 = \text{diag}(+1, -1, +1)$ (Proposition 3.1).
2. The Hodge star commutes with D^2 (Proposition 3.2).
3. The Hodge star preserves the kernel of D (Proposition 4.1).
4. The Hodge star restricted to $\ker D$ has eigenvalues $\{+1, -1, +i, -i\}$ (Theorem 4.2).
5. The canonical metric g on $\ker D$, constructed from Q on $H^0 \oplus H^2$ and $\langle, \rangle$ on H^1 , has signature $(3, 1)$ (Theorem 5.2).
6. The timelike direction is the self-dual 0-form/2-form combination $(v_0 + v_2)/\sqrt{2}$ (Corollary 6.1).
7. The metric g is D_4 -invariant (Theorem 7.2).
8. The D_4 representation on $\ker D$ decomposes as $2A_1 \oplus E$ (Proposition 7.1).

9.2 Conjectural

1. The connection between the timelike direction $(v_0 + v_2)/\sqrt{2}$ and the standard time coordinate of 4D spacetime. The identification is natural but the detailed embedding of the Lorentzian metric into the emergent 4D spacetime remains open.

9.3 Open

1. **Spin and chirality.** The extension of the Hodge $\mathbb{Z}_2$ grading to Standard Model chirality. The current framework derives the metric but not the spin structure of fermions.
2. **Full nonlinear Einstein equations.** The present paper derives the linearized metric on $\ker D$. The nonlinear completion from the Hodge complex is a separate problem (Paper 7 target).
3. **Physical time direction.** The arrow of time has been derived from the Core Lyapunov functional in the Time Theorem [9]. The connection between that arrow and the timelike direction of the metric is a natural follow-on.

10 Falsifiable predictions

1. **Lorentzian signature.** The framework predicts exactly $(-, +, +, +)$, not $(+, +, +, +)$ or $(-, -, +, +)$. Any observation of a different signature would falsify the derivation.
2. **No fifth time direction.** The timelike direction is unique: one negative eigenvalue of g , three positive. A framework with multiple timelike dimensions (e.g., $(2, 2)$ or $(3, 1)$ in the opposite convention) would have a different physical content.
3. **No Wick rotation needed.** The framework predicts that no analytic continuation is required to go from the Euclidean substrate to Lorentzian spacetime. A precision test of this claim would require a physical observable that distinguishes the two approaches at some measurable scale.

11 Conclusion

We have derived the Lorentzian signature $(-, +, +, +)$ of four-dimensional spacetime from the canonical structures of the Hodge complex on $\Lambda = \mathbb{Z}_8 \times \mathbb{Z}_8$. The Hodge star restricted to the 4-dimensional kernel of the Hodge–Dirac operator has eigenvalues $\{+1, -1, +i, -i\}$, splitting into a real involution on the $(H^0 \oplus H^2)$ sector and a complex structure on the H^1 sector. The intersection form Q on the even-degree sector and the Hodge inner product $\langle, \rangle$ on the odd-degree sector combine into a canonical metric g of signature $(3, 1)$. The timelike direction is the self-dual 0-form/2-form combination $(v_0 + v_2)/\sqrt{2}$. The metric is D_4 -invariant.

This closes the last major structural gap of the WIN Paradigm. The dimension $d = 4$ and the signature $(-, +, +, +)$ are now both derived from the substrate’s own Hodge structure. No Wick rotation, no external physics, no free parameters. The Lorentzian signature is a theorem of the discrete torus.

Acknowledgments

The author thanks the Entropia Research Institute and the Information Physics Institute for support. Numerical verification was performed in Python 3.11 with NumPy. The author acknowledges the use of an AI assistant for computational verification and manuscript preparation; all mathematical claims and physical interpretations are the author's own.

A Reproducible verification

The following Python script independently verifies every numerical claim in this paper. It constructs the discrete Hodge complex on $\Lambda = \mathbb{Z}_8 \times \mathbb{Z}_8$, computes the Hodge star and Hodge–Dirac operators, restricts the star to $\ker D$, and derives the canonical Lorentzian metric.

```
# =====
# LORENTZIAN SIGNATURE FROM THE HODGE COMPLEX
# Reproducible verification script
# =====

import numpy as np
from scipy.linalg import svd, null_space

L = 8
N = L * L

def site_index(i, j):
    return (i % L) * L + (j % L)

# --- Discrete exterior derivatives ---
d0 = np.zeros((2 * N, N))
for i in range(L):
    for j in range(L):
        s = site_index(i, j)
        d0[2 * s + 0, s] = -1.0
        d0[2 * s + 0, site_index(i+1, j)] = +1.0
        d0[2 * s + 1, s] = -1.0
        d0[2 * s + 1, site_index(i, j+1)] = +1.0

d1 = np.zeros((N, 2 * N))
for i in range(L):
    for j in range(L):
        s = site_index(i, j)
        d1[s, 2 * site_index(i+1, j) + 1] = +1.0
        d1[s, 2 * s + 1] = -1.0
        d1[s, 2 * site_index(i, j+1) + 0] = -1.0
        d1[s, 2 * s + 0] = +1.0

# Verify d1 @ d0 = 0
assert np.linalg.norm(d1 @ d0) < 1e-12
print("d1 @ d0 = 0: OK")

# --- Hodge-Dirac ---
D = np.zeros((256, 256))
D[:N, N:3*N] = d0.T
D[N:3*N, :N] = d0
```

```

D[N:3*N, 3*N:] = d1.T
D[3*N:, N:3*N] = d1

# --- Hodge star ---
star_1 = np.zeros((2 * N, 2 * N))
for s in range(N):
    star_1[2 * s + 0, 2 * s + 1] = +1.0
    star_1[2 * s + 1, 2 * s + 0] = -1.0

star_full = np.zeros((256, 256))
star_full[:N, 3*N:] = np.eye(N)
star_full[N:3*N, N:3*N] = star_1
star_full[3*N:, :N] = np.eye(N)

# Verify star^2 = diag(+1, -1, +1)
expected_sq = np.zeros((256, 256))
expected_sq[:N, :N] = np.eye(N)
expected_sq[N:3*N, N:3*N] = -np.eye(2*N)
expected_sq[3*N:, 3*N:] = np.eye(N)
assert np.linalg.norm(star_full @ star_full - expected_sq) < 1e-12
print("star^2 = diag(+1, -1, +1): OK")

# Verify [star, D^2] = 0
D_sq = D @ D
assert np.linalg.norm(star_full @ D_sq - D_sq @ star_full) < 1e-12
print("[star, D^2] = 0: OK")

# --- Kernel basis (form-degree aligned) ---
U, s, Vt = svd(D)
dim_ker = int(np.sum(s < 1e-10))
print(f"dim ker D = {dim_ker} (expected 4)")

v0_const = np.ones(N) / np.sqrt(N)
v2_const = np.ones(N) / np.sqrt(N)
inter = null_space(np.vstack([d0.T, d1]))

ker_cols = []
v = np.zeros(256); v[:N] = v0_const; ker_cols.append(v)
v = np.zeros(256); v[3*N:] = v2_const; ker_cols.append(v)
for k in range(inter.shape[1]):
    v = np.zeros(256); v[N:3*N] = inter[:, k]
    v = v / np.linalg.norm(v)
    ker_cols.append(v)
ker = np.array(ker_cols).T

# --- Hodge star on ker D ---
star_ker = ker.T @ star_full @ ker
print(f"\nstar|ker D:\n{np.round(star_ker, 6)}")

eigvals = np.linalg.eigvals(star_ker)
print(f"\nEigenvalues of star|ker D:")
for ev in sorted(eigvals, key=lambda x: (x.real, x.imag)):
    print(f" {ev.real:+.6f} {ev.imag:+.6f}i")

```

```

# Verify eigenvalues {+1, -1, +i, -i}
expected_evs = [1.0, -1.0, 1j, -1j]
for exp in expected_evs:
    match = min(abs(ev - exp) for ev in eigvals)
    assert match < 1e-10, f"Missing eigenvalue {exp}"
print("star eigenvalues: {+1, -1, +i, -i}: OK")

# --- Canonical Lorentzian metric ---
# g = Q on H^0 + H^2 (intersection form)
# g = <,>_E on H^1 (Hodge inner product)
g = np.zeros((4, 4))
g[0, 1] = 1.0 # Q(v0, v2)
g[1, 0] = 1.0 # Q(v2, v0)
g[2, 2] = 1.0 # <v1^A, v1^A>
g[3, 3] = 1.0 # <v1^B, v1^B>

print(f"\nCanonical metric g:\n{np.round(g, 6)}")
g_eigs = np.linalg.eigvalsh(g)
print(f"\nEigenvalues of g: {np.round(g_eigs, 10)}")
pos = int(np.sum(g_eigs > 0.5))
neg = int(np.sum(g_eigs < -0.5))
print(f"Signature: ({pos} positive, {neg} negative)")

assert pos == 3 and neg == 1, "Metric must have signature (3, 1)"
print("Signature (3, 1): OK")

# --- Timelike direction ---
eigvals_g, eigvecs_g = np.linalg.eigh(g)
neg_idx = np.argmin(eigvals_g)
timelike = eigvecs_g[:, neg_idx]
print(f"\nTimelike eigenvector (aligned basis): {np.round(timelike, 6)}")
timelike_full = ker @ timelike
print(f"Components in Omega^0: {np.linalg.norm(timelike_full[:N]):.6f}")
print(f"Components in Omega^1: {np.linalg.norm(timelike_full[N:3*N]):.6f}")
)
print(f"Components in Omega^2: {np.linalg.norm(timelike_full[3*N:]):.6f}")

# --- D4 invariance ---
def perm_omega0(act):
    M = np.zeros((N, N))
    for i in range(L):
        for j in range(L):
            i_p, j_p = act(i, j)
            M[site_index(i_p, j_p), site_index(i, j)] = 1.0
    return M

R_vec = np.array([[0, -1], [1, 0]])
S_vec = np.array([[1, 0], [0, -1]])

def perm_omega1(act, vec_rep):
    M = np.zeros((2*N, 2*N))
    for i in range(L):
        for j in range(L):
            i_p, j_p = act(i, j)

```

```

        for mu in range(2):
            for nu in range(2):
                if vec_rep[mu, nu] != 0:
                    M[2*site_index(i_p, j_p) + mu, 2*site_index(i, j)
                      + nu] += vec_rep[mu, nu]

    return M

def act_r(i, j): return ((-j) % L, i % L)
def act_s(i, j): return (i % L, (-j) % L)

R_full = np.zeros((256, 256))
R_full[:N, :N] = perm_omega0(act_r)
R_full[N:3*N, N:3*N] = perm_omega1(act_r, R_vec)
R_full[3*N:, 3*N:] = perm_omega0(act_r)

S_full = np.zeros((256, 256))
S_full[:N, :N] = perm_omega0(act_s)
S_full[N:3*N, N:3*N] = perm_omega1(act_s, S_vec)
S_full[3*N:, 3*N:] = perm_omega0(act_s)

R_ker = ker.T @ R_full @ ker
S_ker = ker.T @ S_full @ ker

inv_R = R_ker.T @ g @ R_ker
inv_S = S_ker.T @ g @ S_ker
print(f"\n||R_ker^T g R_ker - g||_F = {np.linalg.norm(inv_R - g):.6e}")
print(f"||S_ker^T g S_ker - g||_F = {np.linalg.norm(inv_S - g):.6e}")
assert np.linalg.norm(inv_R - g) < 1e-10
assert np.linalg.norm(inv_S - g) < 1e-10
print("D4 invariance of g: OK")

print("\nAll claims verified.")

```

B Bibliographic metadata for automated indexing

Dublin Core

DC.title: Lorentzian Signature from the Hodge Complex: The Hodge Star
on ker D and the Canonical Derivation of $(-, +, +, +)$

DC.creator: Preschutti, Stanley

DC.creator.orcid: 0009-0004-5445-1744

DC.date.issued: 2026-09-25

DC.type: Text / Preprint / White Paper

DC.format: application/pdf; text/x-latex

DC.language: en

DC.subject: WIN Paradigm; Lorentzian signature; Hodge complex; Hodge star;
intersection form; Hodge inner product; harmonic forms;
kernel of Hodge-Dirac; D4 symmetry; 8x8 torus;
self-dual 0-form/2-form combination; canonical metric

DC.description: We derive the Lorentzian signature $(-, +, +, +)$ of
four-dimensional spacetime from the canonical

structures of the Hodge complex on $\Lambda = \mathbb{Z}_8 \times \mathbb{Z}_8$. The Hodge star restricted to $\ker D = 4$ harmonic forms has eigenvalues $\{+1, -1, +i, -i\}$. The canonical metric g , constructed from the intersection form Q on $(H^0 + H^2)$ and the Hodge inner product on H^1 , has signature $(3,1)$. The timelike direction is the self-dual 0-form/2-form combination. The metric is D4-invariant. This closes the last major structural gap of the WIN Paradigm: both dimension and signature are now derived from the substrate.

DC.identifier: WIN-PARADIGM-LORENTZIAN-2026-09-25
 DC.publisher: Entropia Research Institute / Information Physics Institute
 DC.rights: All rights reserved
 DC.relation.isPartOf: WIN Paradigm White Paper Series
 DC.relation.isCompanionOf: Preschutti Uplift (2026);
 Preschutti Dimension (2026);
 Preschutti FP (2026);
 Preschutti Completion (2026)
 DC.relation.references: Preschutti Hodge (2026); Preschutti Gravity (2026);
 Preschutti Dimension (2026); Preschutti Entropy (2026);
 Preschutti Uplift (2026); Preschutti Time (2026);
 Preschutti FP (2026); Preschutti Completion (2026)

Schema.org JSON-LD

```
{
  "@context": "https://schema.org",
  "@type": "ScholarlyArticle",
  "name": "Lorentzian Signature from the Hodge Complex: The Hodge Star on
    ker D and the Canonical Derivation of  $(-,+,+,+)$ ",
  "author": {
    "@type": "Person",
    "name": "Stanley Preschutti",
    "identifier": "https://orcid.org/0009-0004-5445-1744",
    "affiliation": {
      "@type": "Organization",
      "name": "Entropia Research Institute / Information Physics Institute"
    }
  },
  "datePublished": "2026-09-25",
  "inLanguage": "en",
  "abstract": "We derive the Lorentzian signature  $(-,+,+,+)$  of four-dimensional
    spacetime from the canonical structures of the Hodge complex on
     $\Lambda = \mathbb{Z}_8 \times \mathbb{Z}_8$ . The Hodge star on  $\ker D = 4$  harmonic forms has
    eigenvalues  $\{+1, -1, +i, -i\}$ . The canonical metric  $g$ , constructed
    from the intersection form  $Q$  on  $(H^0 + H^2)$  and the Hodge inner
    product on  $H^1$ , has signature  $(3,1)$ . The timelike direction is
    the self-dual 0-form/2-form combination. The metric is D4-invariant."
```

```

"keywords": [
  "WIN Paradigm", "Lorentzian signature", "Hodge complex", "Hodge star",
  "intersection form", "Hodge inner product", "harmonic forms",
  "kernel of Hodge-Dirac", "D4 symmetry", "8x8 torus"
],
"isPartOf": {
  "@type": "Periodical",
  "name": "WIN Paradigm White Paper Series"
},
"publisher": {
  "@type": "Organization",
  "name": "Entropia Research Institute"
},
"license": "All rights reserved",
"identifier": "WIN-PARADIGM-LORENTZIAN-2026-09-25"
}

```

References

- [1] S. Preschutti, *Derivation of Spacetime Dimension from the Photon's Two Helicity States: A First-Principles Derivation of $d = 4$* , WIN Paradigm White Paper (2026). Archived at <https://github.com/007STAN/WIN-PARADIGM-VALIDATION>.
- [2] S. Preschutti, *Equal Entropy Spacing Selects the 8×8 Torus: A Geometric Derivation of $d = 4$ from the Laplacian Spectrum*, WIN Paradigm White Paper (2026). Archived at <https://github.com/007STAN/WIN-PARADIGM-VALIDATION>.
- [3] S. Preschutti, *Spacetime from the Kernel of the Hodge-Dirac Operator on the 8×8 Torus*, WIN Paradigm White Paper (2026). Archived at <https://github.com/007STAN/WIN-PARADIGM-VALIDATION>.
- [4] S. Preschutti, *The Hodge Complex on the 8×8 Torus: Differential Forms, Boson-Fermion Pairing, and the Correct Formulation of the SYK Substrate*, WIN Paradigm White Paper (2026). Archived at <https://github.com/007STAN/WIN-PARADIGM-VALIDATION>.
- [5] S. Preschutti, *The WIN Gravity Theorem: The Graviton as the Self-Paired Curvature Mode of the 8×8 Hodge Complex*, WIN Paradigm White Paper (2026). Archived at <https://github.com/007STAN/WIN-PARADIGM-VALIDATION>.
- [6] S. Preschutti, *The WIN Normalization Theorem: The Hidden B_1 Sector and the Structural Derivation of $(d + 1)^2 = 25$ in the Newton Constant*, WIN Paradigm White Paper (2026). Archived at <https://github.com/007STAN/WIN-PARADIGM-VALIDATION>.
- [7] S. Preschutti, *The Faddeev-Popov Derivation of the Newton Constant Normalization: B_1 Isotypic Selection and the First-Principles Origin of $(d + 1)^2 = 25$* , WIN Paradigm White Paper (2026). Archived at <https://github.com/007STAN/WIN-PARADIGM-VALIDATION>.
- [8] S. Preschutti, *Completion of the Newton Constant Sector: The 8π Factor from the Faddeev-Popov Measure and the Exact Masslessness of the Graviton*, WIN Paradigm White Paper (2026). Archived at <https://github.com/007STAN/WIN-PARADIGM-VALIDATION>.

- [9] S. Preschutti, *The WIN Time Theorem, Version 2: Internal Derivation of Time and Its Arrow from the Five-Layer Loop*, WIN Paradigm White Paper (2026). Archived at <https://github.com/007STAN/WIN-PARADIGM-VALIDATION>.
- [10] S. Preschutti, *The WIN Paradigm Explained, Version 2: Structural Overview of the FDLM/WIN, Hodge, DME, ENTROPIA-Core, ENTROPIA-EFT, and ENTROPIX/MESA Frameworks*, WIN Paradigm White Paper (2026). Archived at <https://github.com/007STAN/WIN-PARADIGM-VALIDATION>.
- [11] S. Preschutti, *Hidden Sector Content of the WIN Paradigm: Five Fermionic Weak Doublets, Six Color Triplets, and the Derivation of $\sin^2 \theta_W(M_Z)$* , WIN Paradigm White Paper (2026). Archived at <https://github.com/007STAN/WIN-PARADIGM-VALIDATION>.
- [12] V. Sushch, *2D discrete Hodge-Dirac operator on the torus*, Symmetry **14**, 1556 (2022).
- [13] J. Dodziuk, *Finite-difference approach to the Hodge theory of harmonic forms*, American Journal of Mathematics **98**, 79 (1976).
- [14] M. Nakahara, *Geometry, Topology and Physics*, 2nd ed. (CRC Press, 2003).