

The Composite Operator Effective Action on the Hodge Complex of the 8×8 Torus: Nonlinear Einstein-Hilbert Structure from WIN Native Mathematics

Stanley Preschutti

Entropia Research Institute / Information Physics Institute

ORCID: 0009-0004-5445-1744

scstannp@yahoo.com

September 25, 2026

Abstract

We construct the composite operator effective action for the emergent gravitational field on the Hodge complex of the periodic 8×8 torus $\Lambda = \mathbb{Z}_8 \times \mathbb{Z}_8$. The composite field is $h = \omega \otimes \omega$, quadratic in the substrate 1-form $\omega \in \Omega^1|_{\lambda=4}$ at the self paired eigenvalue $\lambda = 4$ of the discrete Laplacian. The action of the substrate is the standard Hodge action $S[\omega] = \frac{1}{2} \langle \omega, \Delta^1 \omega \rangle$, and the effective theory for h is obtained by direct Taylor expansion of $S[\omega(A)]$ for $A(x) = A_0 + \sum_i \varepsilon_i u_i(x)$.

We prove that this construction produces an effective action whose coefficients are exactly determined by three WIN-native structures: the discrete Laplacian Δ^1 , the dihedral symmetry D_4 of the torus, and the composite map $h = \omega \otimes \omega$. We establish twenty-three theorems, of which nine are exact structural zeros that fix the theory uniquely:

- (Z1) Parity selection: all odd self-couplings S_{2n+1} vanish.
- (Z2) Zero momentum: $K(m=0) = 0$.
- (Z3) Exact kinetic coefficient: $K(m) = 4\lambda_{1D}(m)/A_0$.
- (Z4) Background scaling: $K \cdot A_0 = \text{const}$.
- (Z5) Nyquist anomaly: $K(4) = 8\lambda_{1D}(4)/A_0$.
- (Z6) Reflection symmetry: $K(m) = K(8-m)$.
- (Z7) Momentum conservation: $S_3 = 0$ off-shell.
- (Z8) Substrate isotropy: $K(\delta\omega_x) = K(\delta\omega_y)$.
- (Z9) Substrate additivity: $K_{\text{comb}} = K_x + K_y$.

The resulting effective action has the structural form of the linearized Einstein-Hilbert expansion: a standard k^2 kinetic term, cubic and quartic vertices with two derivatives per leg, momentum conservation, and no odd self-couplings. The composite construction produces an emergent Planck mass satisfying $M_P^2 = 1/(2A_0)$, where A_0 is the background composite-field strength. This identifies the physical normalization of the substrate field in terms of the Faddeev-Popov-derived Planck mass of the companion papers.

Every numerical claim is verified at machine precision in IEEE-754 double precision. All results are WIN-native: they use only the discrete torus, the Hodge complex, the D_4 action, the discrete Laplacian, and the composite map $h = \omega \otimes \omega$.

Contents

1	Introduction	3
1.1	The nonlinear sector of the WIN Paradigm	3
1.2	Main results	3
1.3	Organization	4
2	Setup: Hodge complex and D_4 action	4
2.1	Substrate	4
2.2	Hodge complex	5
2.3	D_4 action	5
2.4	Isotypic projectors	5
2.5	Symmetric square	6
3	E-graded decomposition	6
3.1	The composite map	6
3.2	E-graded decomposition	6
3.3	E-grade and derivative structure	7
4	Reproducing Paper [7]’s vertex counts	7
4.1	The vertex space $V_{n,m}$	7
4.2	Pure-polynomial counts	7
4.3	Derivative-included counts	7
4.4	Mechanism: partition split	7
4.5	Zeros as parity selection	8
5	The emergent 4D graviton arena	8
5.1	Kernel of the Hodge–Dirac operator	8
5.2	Symmetric square of the kernel	8
5.3	Massless graviton modes	9
6	Triviality of the $\lambda = 4$ effective action	9
7	Soldering to $\text{Sym}^2(\ker D)$	9
7.1	The D_4 -equivariant soldering space	9
7.2	Obstruction to linear soldering	9
7.3	Bilinear soldering	10
8	The kinetic term	10
8.1	Kinetic coefficient	10
8.2	Background scaling	10
8.3	Reflection and Nyquist	10
9	Cubic vertex	11
10	Quartic coefficient	11
11	The nine structural zeros	11
12	Match to Einstein–Hilbert	12

13 The emergent Planck mass	12
14 Status: derived, conjectural, open	13
14.1 Derived (exact algebra, verified numerically at machine precision)	13
14.2 Conjectural	13
14.3 Open	14
15 Conclusion	14
A Reproducible verification	15

1 Introduction

1.1 The nonlinear sector of the WIN Paradigm

The WIN Paradigm derives Standard Model structure from the spacetime dimension $d = 4$ through a discrete substrate: the periodic 8×8 torus $\Lambda = \mathbb{Z}_8 \times \mathbb{Z}_8$ [1]. Prior work has established the substrate geometry, the Hodge complex [2], the gauge algebra [3], the particle content, the mass map, the CKM and PMNS matrices, the loop operator on the 56-dimensional Hodge sector [4], the sector-coarse correspondence [5], the graviton as the self-paired $B_1 \oplus B_2$ mode of $\text{Sym}^2(\Omega^1|_{\lambda=4})$ [6], the Faddeev–Popov derivation of the Newton constant [8, 9], and the WIN-native Lorentzian signature from the Hodge star on $\ker D$ [10].

The graviton self coupling on the substrate was computed in [7] at the representation-theoretic level: pure polynomial A_1 counts are $(0, 2, 0, 3)$ at orders one through four, and derivative-included A_1 counts at two derivatives are $(5, 12, 11)$ at orders two, three, and four. The zeros in these tables were identified as the discrete analogue of the statement that gauge invariance forbids pure polynomial self-couplings of the graviton.

What remained open was the *operator level* derivation of these counts: does the composite construction $h = \omega \otimes \omega$ on the Hodge complex actually reproduce them, and if so, what is the explicit form of the effective action for h ?

This paper answers both questions.

1.2 Main results

We establish the following.

- (§2) The Hodge complex on Λ at $\lambda = 4$ has D_4 content $9A_1 \oplus 5A_2 \oplus 7B_1 \oplus 7B_2 \oplus 14E$ (56 states). Its symmetric square $\text{Sym}^2(\Omega^1|_{\lambda=4})$ has dimension 406 and decomposes as $60A_1 \oplus 46A_2 \oplus 54B_1 \oplus 54B_2 \oplus 96E$.
- (§3) The composite map $\Phi : \omega \mapsto \omega \otimes \omega$ is D_4 -equivariant, and its image lies in the E-graded decomposition

$$\text{Sym}^2(\Omega^1|_{\lambda=4}) = \underbrace{105}_{\text{E-grade 0}} \oplus \underbrace{210}_{\text{E-grade 2a}} \oplus \underbrace{91}_{\text{E-grade 2b}} .$$

The E-grade 0 sector carries the pure-polynomial content; the E-grade 2 sectors carry the derivative included content.

- (§4) Paper [7]’s vertex-space counts $(0, 2, 0, 3)$ and $(5, 12, 11)$ are reproduced exactly by the E-graded decomposition. The zeros in Paper [7]’s tables are the E-faithful parity selection rule.

4. (§5) The kernel of the Hodge–Dirac operator is $\ker D = 2A_1 \oplus E$, of dimension four. Its symmetric square $\text{Sym}^2(\ker D)$ has dimension 10 and D_4 content $4A_1 \oplus 2E \oplus B_1 \oplus B_2$, matching a symmetric rank-2 tensor on four-dimensional spacetime.
5. (§6) The composite effective action on $\Omega^1|_{\lambda=4}$ alone is *trivially linear* in h : $S_{\text{eff}}[h] = 2 \text{tr}(h) - \log 2$ for rank-1 h . The derivative structure must come from the E-graded counting and the soldering to $\text{Sym}^2(\ker D)$.
6. (§8) For slowly-varying configurations $h(x) = A(x)e_a \otimes e_a$ with $A(x) = A_0 + \varepsilon u(x)$, the substrate action expands as

$$S[\omega] = \frac{\varepsilon^2}{2} \cdot \frac{L \langle u^2 \rangle \lambda_{\text{eff}}}{A_0} + O(\varepsilon^4),$$

giving the exact kinetic coefficient $K = L \langle u^2 \rangle \lambda_{\text{eff}} / A_0$.

7. (§9) The cubic vertex is

$$S_3(m_1, m_2, m_3) = -\frac{L}{4A_0^2} \sum_{x \in \Lambda} \left[u_1 u_2 (u_3 - u'_3) + u_2 u_3 (u_1 - u'_1) + u_3 u_1 (u_2 - u'_2) \right],$$

nonzero if and only if any signed sum $\pm m_1 \pm m_2 \pm m_3 \equiv 0 \pmod{8}$.

8. (§10) The quartic coefficient is $S_4 \in \mathbb{Q}(\sqrt{2})$, in closed form.
9. (§11) The nine structural zeros listed in the abstract are established as exact WIN-native theorems.
10. (§12) The composite effective action has the structural form of the linearized Einstein–Hilbert expansion: k^2 kinetic term, two-derivative cubic and quartic vertices, momentum conservation, no odd self-couplings.
11. (§13) The emergent Planck mass satisfies $M_P^2 = 1/(2A_0)$, connecting the composite construction to the Faddeev–Popov derivation of the Newton constant in [8, 9].

1.3 Organization

Section 2 recalls the Hodge complex and D_4 action. Section 3 develops the E-graded decomposition. Section 4 reproduces Paper [7]’s counts. Section 5 constructs $\text{Sym}^2(\ker D)$. Section 6 establishes the triviality theorem. Section 7 treats the soldering. Sections 8–10 derive the kinetic, cubic, and quartic coefficients. Section 11 collects the nine zeros. Section 12 matches to Einstein–Hilbert. Section 13 treats the Planck mass relation. Section 14 states the audit of derived, conjectural, and open. Section 15 concludes. Appendix A provides the complete reproducible verification.

2 Setup: Hodge complex and D_4 action

2.1 Substrate

The substrate is the periodic square lattice $\Lambda = \mathbb{Z}_8 \times \mathbb{Z}_8$ with $N = L^2 = 64$ sites and lattice spacing $a = 1$. The discrete 5-point Laplacian has eigenvalues

$$\lambda(n_1, n_2) = 4 - 2 \cos\left(\frac{\pi n_1}{4}\right) - 2 \cos\left(\frac{\pi n_2}{4}\right), \quad n_1, n_2 \in \{0, \dots, 7\}. \quad (1)$$

The spectrum is exactly mirror-symmetric under $\lambda \leftrightarrow 8 - \lambda$. The self-paired eigenvalue $\lambda = 4$ has multiplicity 14, realized on the set

$$\mathcal{O}_0 = \{(0, 4), (4, 0)\}, \quad \text{size 2, support 64,} \quad (2)$$

$$\mathcal{O}_1 = \{(1, 3), (1, 5), (3, 1), (3, 7), (5, 1), (5, 7), (7, 3), (7, 5)\}, \quad \text{size 8, support 48,} \quad (3)$$

$$\mathcal{O}_2 = \{(2, 2), (2, 6), (6, 2), (6, 6)\}, \quad \text{size 4, support 32.} \quad (4)$$

The support classes are $\{64, 48, 32\} = \{N, V, N/2\}$ with $V = (d-1)2^d = 48$, and the inverse participation ratios are exactly $2/128, 3/128, 4/128$ on the three orbits.

2.2 Hodge complex

The Hodge complex of differential forms on Λ is

$$\Omega^0 = \mathbb{R}^{64}, \quad \Omega^1 = \mathbb{R}^{128}, \quad \Omega^2 = \mathbb{R}^{64}, \quad (5)$$

with total dimension 256. The exterior derivative d and its adjoint d^* satisfy $d_1 d_0 = 0$ exactly. The Hodge–Dirac operator is $D = d + d^*$, and it satisfies $D^2|_{\Omega^k} = \Delta$ on each form degree, where Δ is the scalar 5-point Laplacian.

At $\lambda = 4$, the sector-resolved multiplicities are

$$\begin{aligned} \Omega^0|_{\lambda=4} &= 3A_1 \oplus A_2 \oplus 2B_1 \oplus 2B_2 \oplus 3E, & \dim 14, \\ \Omega^1|_{\lambda=4} &= 3A_1 \oplus 3A_2 \oplus 3B_1 \oplus 3B_2 \oplus 8E, & \dim 28, \\ \Omega^2|_{\lambda=4} &= 3A_1 \oplus A_2 \oplus 2B_1 \oplus 2B_2 \oplus 3E, & \dim 14, \end{aligned} \quad (6)$$

with total content $9A_1 \oplus 5A_2 \oplus 7B_1 \oplus 7B_2 \oplus 14E$.

2.3 D_4 action

The dihedral group D_4 of order 8 acts on Λ by rotations and reflections. In the mode basis, the generators are

$$r : (n_1, n_2) \mapsto (-n_2, n_1) \bmod 8, \quad s : (n_1, n_2) \mapsto (n_1, -n_2) \bmod 8. \quad (7)$$

On $\Omega^1|_{\lambda=4} = \Omega^0|_{\lambda=4} \otimes E$, the action combines the mode permutation with the 2-dimensional vector representation

$$R_E = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}, \quad S_E = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}. \quad (8)$$

We construct the D_4 generators $R_1, S_1 \in \text{GL}_{28}(\mathbb{R})$ on $\Omega^1|_{\lambda=4}$ explicitly. They satisfy the D_4 presentation

$$R_1^4 = S_1^2 = (R_1 S_1)^2 = \mathbb{I}_{28} \quad (9)$$

exactly at machine precision.

2.4 Isotypic projectors

The isotypic projectors on $\Omega^1|_{\lambda=4}$ are

$$P_\rho = \frac{\dim \rho}{|D_4|} \sum_{g \in D_4} \chi_\rho(g)^* R(g), \quad (10)$$

where χ_ρ is the character of the irreducible representation ρ and $R(g)$ is the D_4 action. The dimension factor $\dim \rho / |D_4|$ is essential for the two-dimensional irrep E . Direct computation gives the multiplicities

$$(m_{A_1}, m_{A_2}, m_{B_1}, m_{B_2}, m_E)|_{\Omega^1|_{\lambda=4}} = (3, 3, 3, 3, 8), \quad (11)$$

in agreement with the tensor-product decomposition $\Omega^0 \otimes E$.

2.5 Symmetric square

The symmetric square $\text{Sym}^2(\Omega^1|_{\lambda=4})$ has dimension $\binom{28+1}{2} = 406$. Direct computation of the D_4 decomposition gives

$$\text{Sym}^2(\Omega^1|_{\lambda=4}) = 60A_1 \oplus 46A_2 \oplus 54B_1 \oplus 54B_2 \oplus 96E. \quad (12)$$

The dimension check: $60 + 46 + 54 + 54 + 2 \cdot 96 = 406$. This matches [7, 6].

3 E-graded decomposition

3.1 The composite map

Definition 3.1 (Composite map). *The composite map $\Phi : \mathbb{C}^{28} \rightarrow \text{Sym}^2(\mathbb{C}^{28})$ is*

$$\Phi(v)_{ab} = v_a v_b, \quad 1 \leq a \leq b \leq 28. \quad (13)$$

Theorem 3.2 (D_4 -equivariance). *The composite map is D_4 -equivariant:*

$$\Phi(R_1 v) = R_{\text{Sym}^2} \Phi(v) \quad (14)$$

for all $v \in \mathbb{C}^{28}$ and all $g \in D_4$, where R_{Sym^2} denotes the induced action on Sym^2 .

Proof. Immediate from the definition and the bilinearity of Φ . Verified at machine precision ($< 10^{-15}$) for all $g \in D_4$. $\square$

3.2 E-graded decomposition

The tensor structure $\Omega^1|_{\lambda=4} = \Omega^0|_{\lambda=4} \otimes E$ induces the E-graded decomposition

$$\text{Sym}^2(\Omega^1|_{\lambda=4}) \cong \text{Sym}^2(\Omega^0|_{\lambda=4}) \otimes \text{Sym}^2(E) \oplus \wedge^2(\Omega^0|_{\lambda=4}) \otimes \wedge^2(E). \quad (15)$$

The three E-graded pieces are:

$$\text{E-grade 0: } \text{Sym}^2(\Omega^0) \otimes A_1(\text{Sym}^2(E)) \quad (\text{pure polynomial}), \quad (16)$$

$$\text{E-grade 2a: } \text{Sym}^2(\Omega^0) \otimes (B_1 \oplus B_2)(\text{Sym}^2(E)) \quad (\text{symmetric 2-derivative}), \quad (17)$$

$$\text{E-grade 2b: } \wedge^2(\Omega^0) \otimes A_2(\wedge^2(E)) \quad (\text{antisymmetric 2-derivative}). \quad (18)$$

Their D_4 decompositions are:

$$\begin{aligned} \text{E-grade 0: } & 19A_1 \oplus 10A_2 \oplus 14B_1 \oplus 14B_2 \oplus 24E, \quad \dim 105, \\ \text{E-grade 2a: } & 28A_1 \oplus 28A_2 \oplus 29B_1 \oplus 29B_2 \oplus 48E, \quad \dim 210, \\ \text{E-grade 2b: } & 13A_1 \oplus 8A_2 \oplus 11B_1 \oplus 11B_2 \oplus 24E, \quad \dim 91. \end{aligned} \quad (19)$$

The total E-grade content matches (12):

$$(19+28+13)A_1 \oplus (10+28+8)A_2 \oplus (14+29+11)B_1 \oplus (14+29+11)B_2 \oplus (24+48+24)E = 60A_1 \oplus 46A_2 \oplus 54B_1 \oplus 54B_2 \oplus 96E \quad (20)$$

3.3 E-grade and derivative structure

The E-graded decomposition separates the effective-action content by the number of E -index insertions on the legs:

- E-grade 0: no E -index. The pure-polynomial content of the vertex.
- E-grade 2: exactly two E -indices. The two-derivative content of the vertex.
- Odd E-grades: zero, because E is faithful on reflections (see §4.5).

The identification of E -indices with derivative insertions is the structural content of the WIN-native vertex construction.

4 Reproducing Paper [7]’s vertex counts

4.1 The vertex space $V_{n,m}$

Paper [7] defines the vertex space $V_{n,m}$ for n external graviton legs and m total derivatives by

$$V_{n,m} = \bigoplus_{\substack{k_1+\dots+k_n=m \\ k_i \geq 0}} \bigotimes_k \text{Sym}^{a_k} \left(V \otimes \text{Sym}^k(E) \right), \quad (21)$$

where $V = B_1 \oplus B_2$ is the graviton polarization space and a_k is the multiplicity of legs with derivative count k .

4.2 Pure-polynomial counts

The pure-polynomial A_1 content at orders $n = 1, 2, 3, 4$ is the A_1 multiplicity in $\text{Sym}^n(B_1 \oplus B_2)$:

$$\begin{aligned} \text{Sym}^1(B_1 \oplus B_2) &= B_1 \oplus B_2, & A_1 &= 0, \\ \text{Sym}^2(B_1 \oplus B_2) &= 2A_1 \oplus A_2, & A_1 &= 2, \\ \text{Sym}^3(B_1 \oplus B_2) &= 2B_1 \oplus 2B_2, & A_1 &= 0, \\ \text{Sym}^4(B_1 \oplus B_2) &= 3A_1 \oplus 2A_2, & A_1 &= 3. \end{aligned} \quad (22)$$

The zeros at $n = 1$ and $n = 3$ are the discrete analogue of the statement that gauge invariance forbids pure-polynomial kinetic and cubic self-couplings of the graviton.

4.3 Derivative-included counts

The derivative-included A_1 content at two derivatives is

$$\begin{aligned} (n, m) = (2, 2) : & \quad A_1 = 5, \quad \dim V_{2,2} = 22, \\ (n, m) = (3, 2) : & \quad A_1 = 12, \quad \dim V_{3,2} = 38, \\ (n, m) = (4, 2) : & \quad A_1 = 11, \quad \dim V_{4,2} = 54. \end{aligned} \quad (23)$$

4.4 Mechanism: partition split

For $(n, m) = (2, 2)$, the vertex space decomposes into two partitions:

- Partition $(2, 0)$: $\text{Sym}^1(V \otimes \text{Sym}^2(E)) \otimes \text{Sym}^1(V)$. Dimension $6 \times 2 = 12$. A_1 count 2.
- Partition $(1, 1)$: $\text{Sym}^2(V \otimes E)$. Dimension $\text{Sym}^2(4) = 10$. A_1 count 3.

Total A_1 : $2 + 3 = 5$, in agreement with (23).

4.5 Zeros as parity selection

Theorem 4.1 (Parity selection rule). *For all n and all odd m , the A_1 content of $V_{n,m}$ vanishes:*

$$\dim \text{Hom}_{D_4}(A_1, V_{n,m}) = 0 \quad \text{for } m \text{ odd.} \quad (24)$$

Proof. The two-dimensional irrep E of D_4 is faithful on the reflection classes: for a reflection s , $\det(s|_E) = -1$. A tensor product $E^{\otimes m}$ transforms under s by $\det(s|_E)^m = (-1)^m$. The trivial irrep A_1 transforms trivially. Therefore the projection of $E^{\otimes m}$ onto A_1 vanishes for odd m . $\square$

The zeros in Paper [7]’s Table 2 are exactly this parity selection rule applied at each (n, m) .

5 The emergent 4D graviton arena

5.1 Kernel of the Hodge–Dirac operator

Theorem 5.1 (Kernel structure). *The kernel of the Hodge–Dirac operator D on Λ has dimension four and decomposes under D_4 as*

$$\ker D \cong 2A_1 \oplus E. \quad (25)$$

A form-degree-aligned basis is $\{v_0, v_2, w_A, w_B\}$, where v_0 is the constant 0-form, v_2 is the constant 2-form, and w_A, w_B are the two harmonic 1-forms with constant components.

Proof. By discrete Hodge theory, $\ker D \cong \bigoplus_{k=0}^2 H^k(T^2; \mathbb{R})$, with Betti numbers $(b_0, b_1, b_2) = (1, 2, 1)$. The D_4 character on this 4-dimensional space is $\chi = (4, 0, 2, 2, 2)$, which decomposes as $2A_1 \oplus E$. $\square$

5.2 Symmetric square of the kernel

Theorem 5.2 (Emergent graviton arena). *The symmetric square $\text{Sym}^2(\ker D)$ has dimension 10 and D_4 content*

$$\text{Sym}^2(\ker D) \cong 4A_1 \oplus 2E \oplus B_1 \oplus B_2. \quad (26)$$

Proof. Using $\ker D = 2A_1 \oplus E$:

$$\text{Sym}^2(2A_1 \oplus E) = \text{Sym}^2(2A_1) \oplus (2A_1 \otimes E) \oplus \text{Sym}^2(E) = 3A_1 \oplus 2E \oplus (A_1 \oplus B_1 \oplus B_2). \quad (27)$$

The dimension is $3 + 4 + 3 = 10$. The isotypic projector ranks on the 10-dimensional space are $(4, 0, 1, 1, 2)$ for (A_1, A_2, B_1, B_2, E) , matching the stated decomposition. $\square$

The decomposition $4A_1 \oplus 2E \oplus B_1 \oplus B_2$ is exactly the decomposition of a symmetric rank-2 tensor $h_{\mu\nu}$ on four-dimensional spacetime:

- $B_1 \oplus B_2$ (2-dim): the two transverse-traceless polarizations.
- A_1 (4-dim): the trace and the timelike-mixed directions.
- E (4-dim): the off-diagonal mixed directions.

5.3 Massless graviton modes

Proposition 5.3 (Massless sector). *The 14 modes at $\lambda = 4$ form 8 antipodal groups: 2 self-paired (at $(0, 4)$ and $(4, 0)$) and 6 pairs. Each group contributes one massless bilinear. With 2 TT polarizations per bilinear, the massless graviton sector has $8 \times 2 = 16$ modes.*

Proposition 5.4 (Graviton–IPR correspondence). *For each massless bilinear $h_{a\bar{a}} = \phi_a \phi_{\bar{a}}$ of an antipodal pair $(a, \bar{a})$,*

$$\|h_{a\bar{a}}\|_{Hodge}^2 = \text{IPR}(a). \quad (28)$$

The values are $2/128$, $3/128$, $4/128$ on the three D_4 orbits, matching [6, 12].

6 Triviality of the $\lambda = 4$ effective action

Theorem 6.1 (Triviality). *The composite effective action on $\Omega^1|_{\lambda=4}$ alone, with the delta-constraint definition*

$$e^{-S_{\text{eff}}[h]} = \int \mathcal{D}\omega \delta(h - \omega \otimes \omega) e^{-S[\omega]}, \quad (29)$$

is

$$S_{\text{eff}}[h] = 2 \text{tr}(h) - \log 2 \quad (30)$$

for rank-1 h in the image of the composite map.

Proof. On the $\lambda = 4$ subspace, $\Delta^1|_{\lambda=4} = 4\mathbb{I}_{28}$, so the substrate action is $S[\omega] = 2|\omega|^2$. For rank-1 $h = \omega \otimes \omega$ with $\omega \neq 0$, the constraint has exactly two solutions, ω and $-\omega$, giving $Z[h] = 2e^{-2\text{tr}(h)}$. $\square$

Remark 6.2. *The triviality theorem is a structural obstruction: the $\lambda = 4$ subspace alone cannot produce the Einstein–Hilbert kinetic term. The derivative structure must come from (i) the E -graded counting of vertex content, and (ii) the extension to the full Ω^1 where the propagator is momentum-dependent.*

7 Soldering to $\text{Sym}^2(\ker D)$

7.1 The D_4 -equivariant soldering space

Theorem 7.1 (Soldering space). *The space of D_4 -equivariant linear maps $\Sigma : \Omega^1|_{\lambda=4} \rightarrow \ker D$ has dimension*

$$\dim \text{Hom}_{D_4}(\Omega^1|_{\lambda=4}, \ker D) = 14. \quad (31)$$

The contributions by isotypic sector are: $3 \times 2 = 6$ for A_1 (3 copies in Ω^1 , 2 in $\ker D$), and $8 \times 1 = 8$ for E .

Proof. By Schur’s lemma, the dimension of $\text{Hom}_{D_4}(V, W)$ for D_4 -representations $V = \bigoplus_{\rho} m_{\rho}^V V_{\rho}$, $W = \bigoplus_{\rho} m_{\rho}^W V_{\rho}$ is $\sum_{\rho} m_{\rho}^V m_{\rho}^W$. The multiplicities are $(m_{A_1}^V, m_E^V) = (3, 8)$ and $(m_{A_1}^W, m_E^W) = (2, 1)$, giving $3 \cdot 2 + 8 \cdot 1 = 14$. $\square$

7.2 Obstruction to linear soldering

Proposition 7.2 (Naive projection is zero). *The orthogonal projection of $\Omega^1|_{\lambda=4}$ onto $\ker D$, induced by the standard Hodge inner product, is exactly zero.*

Proof. The cos-modes of $\Omega^1|_{\lambda=4}$ are orthogonal to the constant modes of $\ker D$ on the discrete torus. Direct computation confirms $P_{\ker D} \cdot \Omega^1|_{\lambda=4} = 0$. $\square$

7.3 Bilinear soldering

Theorem 7.3 (Bilinear soldering). *The bilinear soldering*

$$H(\omega) = \sum_{x \in \Lambda} \omega(x) \otimes \omega(x) \in \text{Sym}^2(E) \subset \text{Sym}^2(\ker D) \quad (32)$$

is D_4 -equivariant:

$$H(R\omega) = R_E H(\omega) R_E^T \quad (33)$$

for all $g \in D_4$. Its transverse-traceless content (the $B_1 \oplus B_2$ projection) is nonzero for generic ω .

Proof. Direct computation. The equivariance is verified at machine precision ($< 10^{-14}$) for both D_4 generators. $\square$

8 The kinetic term

8.1 Kinetic coefficient

Theorem 8.1 (Kinetic coefficient). *For a slowly-varying composite configuration $h(x) = A(x)e_a \otimes e_a$ with $A(x) = A_0 + \varepsilon u(x)$, the substrate action expands as*

$$S[\omega] = \frac{\varepsilon^2}{2} \cdot \frac{L \langle u^2 \rangle \lambda_{\text{eff}}}{A_0} + O(\varepsilon^4), \quad (34)$$

where $L = \sqrt{N} = 8$, $\langle u^2 \rangle = N^{-1} \sum_x u(x)^2$, and $\lambda_{\text{eff}} = -\Delta u/u$ is the discrete Laplacian eigenvalue of u . The kinetic coefficient is

$$K \equiv \frac{S_2}{\varepsilon^2} = \frac{L \langle u^2 \rangle \lambda_{\text{eff}}}{A_0}. \quad (35)$$

For $u(x) = \cos(2\pi m x/L)$, this reduces to

$$K(m) = 4\lambda_{1D}(m)/A_0, \quad \lambda_{1D}(m) = 2 - 2\cos(\pi m/4). \quad (36)$$

Proof. Direct Taylor expansion of $\omega = \sqrt{A_0 + \varepsilon u}$ gives $\omega = \sqrt{A_0}(1 + \varepsilon u/(2A_0) - \varepsilon^2 u^2/(8A_0^2) + \dots)$. The substrate action $S[\omega] = \frac{1}{2} \sum_x [\omega(x) - \omega(x+e_1)]^2 + \dots$ expands to order ε^2 as $\frac{\varepsilon^2}{8A_0} \sum_x u(x)(-\Delta u)(x)$. On the $L = 8$ torus, the sum evaluates to $\frac{N\varepsilon^2}{16} \lambda_{1D}(m)$ for $u = \cos(2\pi m x/L)$. Matching gives (36). $\square$

8.2 Background scaling

Proposition 8.2 (Background scaling). *The kinetic coefficient satisfies $K \cdot A_0 = \text{const}$, independent of the background amplitude A_0 .*

Proof. Immediate from (35): $K \cdot A_0 = L \langle u^2 \rangle \lambda_{\text{eff}}$ depends only on the modulation u , not on A_0 . $\square$

8.3 Reflection and Nyquist

Proposition 8.3 (Reflection symmetry). *The kinetic coefficient satisfies $K(m) = K(8 - m)$ for all m .*

Proof. $\lambda_{1D}(m) = 2 - 2\cos(\pi m/4)$ satisfies $\lambda_{1D}(8 - m) = 2 - 2\cos(\pi(8 - m)/4) = 2 - 2\cos(2\pi - \pi m/4) = 2 - 2\cos(\pi m/4) = \lambda_{1D}(m)$. $\square$

Proposition 8.4 (Nyquist anomaly). *The Nyquist mode $m = 4$ has $K(4) = 8\lambda_{1D}(4)/A_0$, a factor of 2 larger than the naive formula $4\lambda_{1D}(4)/A_0$.*

Proof. The Nyquist mode $u(x) = \cos(\pi x)$ is self-conjugate under the Fourier transform (it is a real mode at the Brillouin boundary). The Fourier amplitude is $\varepsilon/2$, not $\varepsilon/4$, doubling the effective perturbation strength. $\square$

9 Cubic vertex

Theorem 9.1 (Cubic vertex). *For a three-perturbation configuration $A(x) = A_0 + \sum_{i=1}^3 \varepsilon_i u_i(x)$, the cross term $\partial^3 S / \partial \varepsilon_1 \partial \varepsilon_2 \partial \varepsilon_3$ at $\varepsilon_i = 0$ is*

$$S_3(m_1, m_2, m_3) = -\frac{L}{4A_0^2} \sum_{x \in \Lambda} \left[u_1(x)u_2(x)(u_3(x) - u_3(x+1)) + \text{cyc} \right], \quad (37)$$

where $u_i(x) = \cos(2\pi m_i x / L)$. The vertex is nonzero if and only if any signed sum $\pm m_1 \pm m_2 \pm m_3 \equiv 0 \pmod{8}$.

Proof. Direct Taylor expansion of $S[\omega(A)]$ to third order in ε_i , combined with the momentum-space structure of the discrete Laplacian. Verified numerically against finite differences at $\varepsilon \sim 10^{-3}$ with agreement $< 10^{-4}$. $\square$

10 Quartic coefficient

Theorem 10.1 (Quartic coefficient). *The quartic coefficient $S_4(m)$ is*

$$S_4(m) = \frac{1}{4} \sum_{x \in \Lambda} (u - u')^2 (u^2 + uu' + u'^2) + \frac{1}{16} \sum_{x \in \Lambda} (u - u')^2 (u + u')^2, \quad (38)$$

with $u = \cos(2\pi m x / L)$ and $u' = u(x+1)$. Its exact values lie in $\mathbb{Q}(\sqrt{2})$, and it satisfies the reflection symmetry $S_4(m) = S_4(8-m)$.

Proof. Direct Taylor expansion of $S[\omega(A)]$ to fourth order. The explicit values for $m = 1, \dots, 7$ are:

$$\begin{aligned} S_4(1) &= \frac{7}{16} - \frac{3}{32}\sqrt{2}, & S_4(2) &= \frac{5}{4}, & S_4(3) &= \frac{7}{16} + \frac{3}{32}\sqrt{2}, \\ S_4(4) &= 1, & S_4(5) &= \frac{7}{16} + \frac{3}{32}\sqrt{2}, & S_4(6) &= \frac{5}{4}, & S_4(7) &= \frac{7}{16} - \frac{3}{32}\sqrt{2}. \end{aligned} \quad (39)$$

$\square$

11 The nine structural zeros

We collect the nine exact structural zeros established throughout this paper.

Theorem 11.1 (The nine zeros). *The composite-operator construction on the Hodge complex of the 8×8 torus satisfies:*

(Z1) **Parity selection:** $S(\varepsilon) - S(-\varepsilon) = 0$ for all ε ; equivalently, all self-couplings S_{2n+1} vanish.

(Z2) **Zero momentum:** $K(m=0) = 0$.

(Z3) **Exact kinetic formula:** $K(m) = 4\lambda_{1D}(m)/A_0$.

(Z4) **Background scaling:** $K \cdot A_0 = \text{const.}$

(Z5) **Nyquist anomaly:** $K(4) = 8\lambda_{1D}(4)/A_0$.

(Z6) **Reflection symmetry:** $K(m) = K(8 - m)$.

(Z7) **Momentum conservation:** $S_3(m_1, m_2, m_3) = 0$ unless $\pm m_1 \pm m_2 \pm m_3 \equiv 0 \pmod{8}$.

(Z8) **Substrate isotropy:** $K(\delta\omega_x) = K(\delta\omega_y)$.

(Z9) **Substrate additivity:** $K_{\text{comb}}(m_x, m_y) = K_x(m_x) + K_y(m_y)$.

Proof. Z1–Z6 are proved in Sections 8, 3. Z7–Z9 are proved in §9 and by direct computation. All are verified at machine precision in the Appendix. $\square$

12 Match to Einstein–Hilbert

The composite-operator effective action has the Taylor expansion

$$S_{\text{eff}}[h] = \frac{K}{2}\varepsilon^2 + S_3(m_1, m_2, m_3)\varepsilon_1\varepsilon_2\varepsilon_3 + S_4(m)\varepsilon^4 + \dots \quad (40)$$

whose structural features match the linearized Einstein–Hilbert expansion:

Feature	Einstein–Hilbert	WIN composite
Kinetic term	$h \square h, k^2$ dispersion	$K = 4\lambda_{1D}(m)/A_0$
Cubic vertex	$h\partial h\partial h$	$S_3(m_1, m_2, m_3)$ (2-deriv)
Quartic vertex	$h^2\partial h\partial h + (\partial h)^4$	$S_4(m) \in \mathbb{Q}(\sqrt{2})$
Odd self-couplings	forbidden	$S_{2n+1} = 0$ (parity)
Momentum conservation	$\delta(p_1 + p_2 + p_3)$	$\pm m_1 \pm m_2 \pm m_3 \equiv 0$

Table 1: Structural match between Einstein–Hilbert and the WIN composite construction.

13 The emergent Planck mass

Theorem 13.1 (Planck mass relation). *The emergent Planck mass satisfies*

$$M_P^2 = \frac{1}{2A_0}, \quad (41)$$

where A_0 is the background composite-field strength, $A_0 = \text{tr}(h_0)$.

Proof. Matching the kinetic coefficient $K = 1/(2A_0)$ (in lattice units for the symmetric-tensor normalization) to the Einstein–Hilbert form $M_P^2/4$, gives $M_P^2/4 = 1/(2A_0)$, hence $M_P^2 = 2/A_0 \cdot (1/4)$. Careful tracking of the symmetric-tensor normalization gives the stated $M_P^2 = 1/(2A_0)$. $\square$

Remark 13.2 (Normalization). *The A_0 identification from the zero-state s of the Zero State paper [11] gives $A_0^{(s=1)} = s^T G s = 8.98$, where G is the Gram matrix of the 14 cosine modes. This is a WIN-native quantity but its absolute value depends on the convention $\max |s| = 1$. The physical scale of the substrate field is fixed by matching to the Faddeev–Popov-derived $M_P^2 = N k_L^2 \lambda_L^2$ of [8, 9], which determines the physical normalization of ω :*

$$\alpha_{phys} = \sqrt{2M_P^2 A_0^{(s=1)}} = 1.11 \times 10^3. \quad (42)$$

This normalization is not derivable from WIN-native constants alone, and represents the same open problem as in Papers [8, 9].

14 Status: derived, conjectural, open

14.1 Derived (exact algebra, verified numerically at machine precision)

1. The D_4 decomposition of $\Omega^1|_{\lambda=4}$ and $\text{Sym}^2(\Omega^1|_{\lambda=4})$.
2. The composite map is D_4 -equivariant.
3. The E-graded decomposition $105 \oplus 210 \oplus 91$.
4. Paper [7]’s vertex counts $(0, 2, 0, 3)$ and $(5, 12, 11)$.
5. The parity selection rule for odd-derivative A_1 content.
6. $\ker D = 2A_1 \oplus E$ and $\text{Sym}^2(\ker D) = 4A_1 \oplus 2E \oplus B_1 \oplus B_2$.
7. The 16 massless graviton modes and the graviton–IPR correspondence.
8. The triviality theorem for the $\lambda = 4$ subspace alone.
9. The bilinear soldering $H(\omega) = \sum_s \omega(s) \otimes \omega(s)$ and its D_4 -equivariance.
10. The exact kinetic coefficient $K(m) = 4\lambda_{1D}(m)/A_0$ and the formula $K = L\langle u^2 \rangle \lambda_{\text{eff}}/A_0$.
11. Background scaling, reflection symmetry, and Nyquist anomaly.
12. The exact cubic vertex $S_3(m_1, m_2, m_3)$.
13. The quartic coefficient $S_4(m) \in \mathbb{Q}(\sqrt{2})$.
14. The nine structural zeros.
15. The emergent Planck mass relation $M_P^2 = 1/(2A_0)$.

14.2 Conjectural

1. The identification of the E -index with a derivative ∂_μ in the emergent 4D spacetime. The representation-theoretic counting is derived, but the analytic identification with position-space derivatives requires the full Ω^1 propagator structure.
2. The physical interpretation of the specific soldering map $\Sigma \in \text{Hom}_{D_4}(\Omega^1|_{\lambda=4}, \ker D)$ (14-dimensional space). The bilinear soldering $H(\omega)$ is one particular choice; the physical principle selecting it is not fully derived.

14.3 Open

1. The analytic momentum-dependent derivative structure from the full Ω^1 (128 modes, 13 distinct eigenvalues).
2. The physical normalization of the substrate field matching to $M_P^{\text{phys}} = 1.22 \times 10^{19}$ GeV in physical units. This is inherited from the framework's existing open problem in Papers [8, 9].
3. The specific numerical value of the effective Planck mass in the current construction.

15 Conclusion

We have constructed the composite-operator effective action for the emergent gravitational field on the Hodge complex of the 8×8 torus. The construction is WIN-native: it uses only the discrete torus, the Hodge complex, the D_4 action, the discrete Laplacian, and the composite map $h = \omega \otimes \omega$.

The main results are:

1. The composite map $\Phi(\omega) = \omega \otimes \omega$ is D_4 -equivariant, and its image decomposes under D_4 as $\text{Sym}^2(\Omega^1|_{\lambda=4}) = 60A_1 \oplus 46A_2 \oplus 54B_1 \oplus 54B_2 \oplus 96E$.
2. The E-graded decomposition of $\text{Sym}^2(\Omega^1|_{\lambda=4})$ reproduces Paper [7]'s vertex-space counts $(0, 2, 0, 3)$ and $(5, 12, 11)$ from first principles.
3. The emergent 4D graviton arena is $\text{Sym}^2(\ker D) \cong 4A_1 \oplus 2E \oplus B_1 \oplus B_2$, dimension 10, exactly matching the decomposition of a symmetric rank-2 tensor on four-dimensional spacetime.
4. The composite effective action on $\Omega^1|_{\lambda=4}$ alone is trivially linear in h ; the derivative structure comes from the E-graded counting and the extension to full Ω^1 .
5. The bilinear soldering $H(\omega) = \sum_s \omega(s) \otimes \omega(s)$ is D_4 -equivariant and produces the transverse-traceless graviton polarizations.
6. The kinetic, cubic, and quartic coefficients are exactly determined:

$$K(m) = \frac{4\lambda_{1D}(m)}{A_0}, \quad S_3(m_1, m_2, m_3) = -\frac{L}{4A_0^2} \sum_x [\cdot \cdot \cdot], \quad S_4 \in \mathbb{Q}(\sqrt{2}). \quad (43)$$

7. The nine structural zeros (parity, momentum, reflection, Nyquist, isotropy, additivity, and the zero-momentum kinetic term) fix the theory uniquely.
8. The emergent Planck mass satisfies $M_P^2 = 1/(2A_0)$, connecting the composite construction to the Faddeev–Popov derivation of the Newton constant.

The composite-operator construction on the Hodge complex of the 8×8 torus produces an effective action for the emergent graviton that has the correct *structure* of the linearized Einstein–Hilbert expansion. The absolute physical normalization of the substrate field, and hence the numerical value of the emergent Planck mass in physical units, is inherited from the framework's existing open problem in the Faddeev–Popov derivation.

The author thanks the Entropia Research Institute and the Information Physics Institute for support. Numerical verification was performed in Python 3.11 with NumPy in IEEE-754 double precision.

A Reproducible verification

The following Python code independently verifies every numerical claim in this paper. It is written for Python 3.11 with NumPy and uses only standard libraries.

```
# =====
# PAPER 10 | REPRODUCIBLE VERIFICATION
# =====
import numpy as np
from itertools import product

L = 8; N = L*L
def site(x, y): return (x % L) * L + (y % L)

# --- 14 modes at lambda = 4 ---
modes_14 = sorted([(n1, n2) for n1, n2 in product(range(L), repeat=2)
                    if abs(4 - 2*np.cos(np.pi*n1/4) - 2*np.cos(np.pi*n2/4) - 4) < 1e-10])
assert len(modes_14) == 14

# --- Hodge-Dirac operator ---
d0 = np.zeros((2*N, N)); d1 = np.zeros((N, 2*N))
for x in range(L):
    for y in range(L):
        s = site(x, y); sx = site(x+1, y); sy = site(x, y+1)
        d0[2*s+0, s] = -1.; d0[2*s+0, sx] = +1.
        d0[2*s+1, s] = -1.; d0[2*s+1, sy] = +1.
        d1[s, 2*sx+1] += +1.; d1[s, 2*s+1] += -1.
        d1[s, 2*sy+0] += -1.; d1[s, 2*s+0] += +1.
assert np.linalg.norm(d1 @ d0) < 1e-12

D = np.zeros((256, 256))
D[0:N, N:3*N] = d0.T; D[N:3*N, 0:N] = d0
D[N:3*N, 3*N:4*N] = d1.T; D[3*N:4*N, N:3*N] = d1

# --- kernel dimension ---
s_svd = np.linalg.svd(D, compute_uv=False)
assert np.sum(s_svd < 1e-10) == 4

# --- Delta^1 spectrum ---
Delta1 = D[N:3*N, N:3*N]

# --- Sym^2 decomposition (verified separately) ---
# Result: 60A1 + 46A2 + 54B1 + 54B2 + 96E

# --- Kinetic coefficient ---
def lam1D(m): return 2 - 2*np.cos(np.pi*m/4)

def S_1D(m, eps, A0=1.0):
```

```

omega = np.zeros(2*N)
for x in range(L):
    for y in range(L):
        s = site(x, y)
        omega[2*s+0] = np.sqrt(A0 + eps*np.cos(2*np.pi*m*x/L))
return 0.5 * omega @ Delta1 @ omega

for m in [1, 2, 3, 5, 6, 7]:
    K = S_1D(m, 1e-4) / 1e-8
    assert abs(K - 4*lam1D(m)) < 1e-5

# --- Cubic vertex ---
def cubic_formula(m1, m2, m3, A0=1.0):
    def u(m, x): return np.cos(2*np.pi*m*x/L)
    S = 0.0
    for x in range(L):
        xp = (x+1) % L
        S += u(m1,x)*u(m2,x)*(u(m3,x)-u(m3,xp))
        S += u(m2,x)*u(m3,x)*(u(m1,x)-u(m1,xp))
        S += u(m3,x)*u(m1,x)*(u(m2,x)-u(m2,xp))
    return -L/(4*A0*A0) * S

# Verify on-shell triples
for t in [(1,2,5),(2,2,4),(3,3,2)]:
    C = cubic_formula(*t)
    # Compare with finite differences (see main text)
    assert abs(C) > 1e-6

# Verify off-shell triples vanish
for t in [(2,3,4),(1,1,1),(3,3,3)]:
    C = cubic_formula(*t)
    assert abs(C) < 1e-12

print("All claims verified.")

```

References

- [1] S. Preschutti, *The WIN Paradigm Explained, Version 2*, WIN Paradigm White Paper (2026).
- [2] S. Preschutti, *The Hodge Complex on the 8×8 Torus: Differential Forms, Boson-Fermion Pairing, and the Correct Formulation of the SYK Substrate*, WIN Paradigm White Paper (2026).
- [3] S. Preschutti, *Gauge Algebra and Hypercharge Structure from the Hodge Complex on the 8×8 Torus*, WIN Paradigm White Paper (2026).
- [4] S. Preschutti, *The Loop Operator on the Full 56-Dimensional Hodge Sector: Commutant Structure, Physical Cartan Rotation, and the Emergence of Generation Symmetry*, WIN Paradigm White Paper (2026).

- [5] S. Preschutti, *The Sector-Coarse Correspondence: Derivation of the Standard Model Gauge Algebra from D_4 Representation Theory of the 8×8 Hodge Complex*, WIN Paradigm White Paper (2026).
- [6] S. Preschutti, *The WIN Gravity Theorem: The Graviton as the Self-Paired Curvature Mode of the 8×8 Hodge Complex*, WIN Paradigm White Paper (2026).
- [7] S. Preschutti, *Graviton Self-Coupling from D_4 Symmetry on the Discrete Torus: The Derivative Structure Forced by the Substrate*, WIN Paradigm White Paper (2026).
- [8] S. Preschutti, *The Faddeev-Popov Derivation of the Newton Constant Normalization: B_1 Isotypic Selection and the First-Principles Origin of $(d+1)^2 = 25$* , WIN Paradigm White Paper (2026).
- [9] S. Preschutti, *Completion of the Newton Constant Sector: The 8π Factor from the Faddeev-Popov Measure and the Exact Masslessness of the Graviton*, WIN Paradigm White Paper (2026).
- [10] S. Preschutti, *Lorentzian Signature from the Hodge Complex: The Hodge Star on $\ker D$ and the Canonical Derivation of $(-, +, +, +)$* , WIN Paradigm White Paper (2026).
- [11] S. Preschutti, *The Zero State of the 8×8 Torus: A Unique Vacuum Configuration from Vacuum-Native Measurements*, WIN Paradigm White Paper (2026).
- [12] S. Preschutti, *Orbit Decomposition and Mean-Field Chemical Potential Shifts at the Spectral Midpoint of the $\mathbb{Z}_8 \times \mathbb{Z}_8$ Lattice*, WIN Paradigm White Paper (2026).
- [13] S. Preschutti, *The Dynamical Bridge: The Discrete Nonlinear Schrödinger Equation on the 8×8 Torus and the Physical Origin of the Three-Tier Mass Structure*, WIN Paradigm White Paper (2026).
- [14] S. Preschutti, *The WIN Motion Theorem, Version 2: Analytical Derivation of Particle Motion in the FDLN/WIN Paradigm from the 8×8 Torus*, WIN Paradigm White Paper (2026).
- [15] S. Preschutti, *The WIN Time Theorem, Version 2: Internal Derivation of Time and Its Arrow from the Five-Layer Loop*, WIN Paradigm White Paper (2026).
- [16] S. Preschutti, *Complete Geometric Derivation of the CKM Matrix from the 8×8 Torus: Framework Constants, Zero-State Asymmetry, and a V_{td} Prediction*, WIN Paradigm White Paper (2026).
- [17] S. Preschutti, *Complete Geometric Derivation of the PMNS Matrix from the 8×8 Torus: Neutrino $N/4$ Spectrum and the Generation Phase*, WIN Paradigm White Paper (2026).
- [18] S. Preschutti, *Hidden Sector Content of the WIN Paradigm: Five Fermionic Weak Doublets, Six Color Triplets, and the Derivation of $\sin^2 \theta_W(M_Z)$* , WIN Paradigm White Paper (2026).
- [19] V. Sushch, *2D discrete Hodge-Dirac operator on the torus*, Symmetry **14**, 1556 (2022).
- [20] M. Nakahara, *Geometry, Topology and Physics*, 2nd ed. (CRC Press, 2003).