

## - Lesson 5.3-5.4

8/8 points (100%)

Quiz, 8 questions

 **Congratulations! You passed!**

Next Item

1 / 1  
points

1.

We use the continuous version of Bayes' theorem if:

 $\theta$  is continuous**Correct**If  $\theta$  is continuous, we use a probability density for the prior. $Y$  is continuous $f(y | \theta)$  is continuous

All of the above



None of the above

1 / 1  
points

2.

Consider the coin-flipping example from the lesson. Recall that the likelihood for this experiment was Bernoulli with unknown probability of heads, i.e.,  $f(y | \theta) = \theta^y(1 - \theta)^{1-y}I_{\{0 \leq \theta \leq 1\}}$ , and we started with a uniform prior on the interval  $[0, 1]$ .

After the first flip resulted in heads ( $Y_1 = 1$ ), the posterior for  $\theta$  became  $f(\theta | Y_1 = 1) = 2\theta I_{\{0 \leq \theta \leq 1\}}$ .

Now use this posterior as your prior for  $\theta$  before the next (second) flip. Which of the following represents the posterior PDF for  $\theta$  after the second flip also results in heads ( $Y_2 = 1$ )?



$$f(\theta | Y_2 = 1) = \frac{(1-\theta) \cdot 2\theta}{\int_0^1 (1-\theta) \cdot 2\theta d\theta} I_{\{0 \leq \theta \leq 1\}}$$



$$f(\theta | Y_2 = 1) = \frac{\theta(1-\theta) \cdot 2\theta}{\int_0^1 \theta(1-\theta) \cdot 2\theta d\theta} I_{\{0 \leq \theta \leq 1\}}$$



$$f(\theta | Y_2 = 1) = \frac{\theta \cdot 2\theta}{\int_0^1 \theta \cdot 2\theta d\theta} I_{\{0 \leq \theta \leq 1\}}$$

**Correct**

Lesson 5.3 5.4 This simplifies to the posterior PDF  $f(\theta | Y_2 = 1) = 3\theta^2 I_{\{0 \leq \theta \leq 1\}}$ .

**8/8 points (100%)**

Quiz, 8 questions. Incidentally, if we assume that the two coin flips are independent, we would have arrived at the same posterior if we had again started with a uniform prior and performed a single update using  $Y_1 = 1$  and  $Y_2 = 1$ .

1 / 1  
points

3.

Consider again the coin-flipping example from the lesson. Recall that we used a  $\text{Uniform}(0,1)$  prior for  $\theta$ . Which of the following is a correct interpretation of  $P(0.3 < \theta < 0.9) = 0.6$ ?



(0.3, 0.9) is a 60% credible interval for  $\theta$  before observing any data.

**Correct**

The probability statement came from our prior, so the prior probability that  $\theta$  is in this interval is 0.6.



(0.3, 0.9) is a 60% credible interval for  $\theta$  after observing  $Y = 1$ .



(0.3, 0.9) is a 60% confidence interval for  $\theta$ .



The posterior probability that  $\theta \in (0.3, 0.9)$  is 0.6.

1 / 1  
points

4.

Consider again the coin-flipping example from the lesson. Recall that the posterior PDF for  $\theta$ , after observing  $Y = 1$ , was  $f(\theta | Y = 1) = 2\theta I_{\{0 \leq \theta \leq 1\}}$ . Which of the following is a correct interpretation of  $P(0.3 < \theta < 0.9 | Y = 1) = \int_{0.3}^{0.9} 2\theta d\theta = 0.72$ ?



(0.3, 0.9) is a 72% credible interval for  $\theta$  before observing any data.



(0.3, 0.9) is a 72% credible interval for  $\theta$  after observing  $Y = 1$ .

**Correct**

The probability statement came from the posterior, so the posterior probability that  $\theta$  is in this interval is 0.72.



(0.3, 0.9) is a 72% confidence interval for  $\theta$ .



The prior probability that  $\theta \in (0.3, 0.9)$  is 0.72.

# Lesson 5.3-5.4

1 / 1  
points

8/8 points (100%)

Quiz, 8 questions

Which two quantiles are required to capture the middle 90% of a distribution (thus producing a 90% equal-tailed interval)?

☐ .10 and .90

☒ .05 and .95

## Correct

90% of the probability mass is contained between the .05 and .95 quantiles (or equivalently, the 5th and 95th percentiles). 5% of the probability lies on either side of this interval.

☐ .025 and .975

☐ 0 and .9
1 / 1  
points

6.

Suppose you collect measurements to perform inference about a population mean  $\theta$ . Your posterior distribution after observing data is  $\theta | \mathbf{y} \sim N(0, 1)$ .

Report the upper end of a 95% equal-tailed interval for  $\theta$ . Round your answer to two decimal places.

## Correct Response

The 95% equal-tailed interval for a standard normal distribution is (-1.96, 1.96).

Because the normal distribution is symmetric and unimodal (has only one peak), the equal-tailed interval is also the highest posterior density (HPD) interval.

In R:

```
1 qnorm(p=0.975, mean=0, sd=1)
```

In Excel:

```
1 = NORM.INV(0.975, 0, 1)
2 |
```

where probability=0.975, mean=0, standard\_dev=1.

## Lesson 5.3-5.4 1 / 1

Quiz, 8 questions

points

8/8 points (100%)

7.

What does "HPD interval" stand for?


- ☐ Highest partial density interval
- ☐ Highest precision density interval
- ☐ Highest point distance interval
- ☒ Highest posterior density interval

**Correct**1 / 1  
points


8.

Each of the following graphs depicts a 50% credible interval from a posterior distribution. Which of the intervals represents the HPD interval?

- ☐ 50% interval:  $\theta \in (0.400, 0.756)$



- ☒ 50% interval:  $\theta \in (0.326, 0.674)$



Correct


## Lesson 5.3-5.4

This is the 50% credible interval with the highest posterior density values. It is the shortest possible interval containing 50% of the probability under this posterior distribution.

8/8 points (100%)

Quiz, 8 questions

☐ 50% interval:  $\theta \in (0.500, 1.000)$



☐ 50% interval:  $\theta \in (0.196, 0.567)$

