

Mathematical Biostatistics Bootcamp: Lecture 10, T Confidence Intervals

Brian Caffo

Department of Biostatistics
Johns Hopkins Bloomberg School of Public Health
Johns Hopkins University

September 19, 2012

Table of contents

- 1 Table of contents
- 2 Independent group t intervals
- 3 Likelihood method
- 4 Unequal variances

Independent group t confidence intervals

- Suppose that we want to compare the mean blood pressure between two groups in a randomized trial; those who received the treatment to those who received a placebo
- We cannot use the paired t test because the groups are independent and may have different sample sizes
- We now present methods for comparing independent groups

Notation

- Let X_1, \dots, X_{n_x} be iid $N(\mu_x, \sigma^2)$
- Let Y_1, \dots, Y_{n_y} be iid $N(\mu_y, \sigma^2)$
- Let \bar{X} , \bar{Y} , S_x , S_y be the means and standard deviations
- Using the fact that linear combinations of normals are again normal, we know that $\bar{Y} - \bar{X}$ is also normal with mean $\mu_y - \mu_x$ and variance $\sigma^2(\frac{1}{n_x} + \frac{1}{n_y})$
- The pooled variance estimator

$$S_p^2 = \{(n_x - 1)S_x^2 + (n_y - 1)S_y^2\} / (n_x + n_y - 2)$$

is a good estimator of σ^2

Note

- The pooled estimator is a mixture of the group variances, placing greater weight on whichever has a larger sample size
- If the sample sizes are the same the pooled variance estimate is the average of the group variances
- The pooled estimator is unbiased

$$\begin{aligned} E[S_p^2] &= \frac{(n_x - 1)E[S_x^2] + (n_y - 1)E[S_y^2]}{n_x + n_y - 2} \\ &= \frac{(n_x - 1)\sigma^2 + (n_y - 1)\sigma^2}{n_x + n_y - 2} \end{aligned}$$

- The pooled variance estimate is independent of $\bar{Y} - \bar{X}$ since S_x is independent of \bar{X} and S_y is independent of \bar{Y} and the groups are independent

Result

- The sum of two independent Chi-squared random variables is Chi-squared with degrees of freedom equal to the sum of the degrees of freedom of the summands
- Therefore

$$\begin{aligned}(n_x + n_y - 2)S_p^2/\sigma^2 &= (n_x - 1)S_x^2/\sigma^2 + (n_y - 1)S_y^2/\sigma^2 \\&= \chi_{n_x-1}^2 + \chi_{n_y-1}^2 \\&= \chi_{n_x+n_y-2}^2\end{aligned}$$

Putting this all together

- The statistic

$$\frac{\bar{Y} - \bar{X} - (\mu_y - \mu_x)}{\sigma \left(\frac{1}{n_x} + \frac{1}{n_y} \right)^{1/2}} = \frac{\bar{Y} - \bar{X} - (\mu_y - \mu_x)}{S_p \left(\frac{1}{n_x} + \frac{1}{n_y} \right)^{1/2}}$$

$$\frac{\bar{Y} - \bar{X} - (\mu_y - \mu_x)}{\sqrt{\frac{(n_x + n_y - 2)S_p^2}{(n_x + n_y - 2)\sigma^2}}}$$

is a standard normal divided by the square root of an independent Chi-squared divided by its degrees of freedom

- Therefore this statistic follows Gosset's t distribution with $n_x + n_y - 2$ degrees of freedom
- Notice the form is (estimator - true value) / SE

Confidence interval

- Therefore a $(1 - \alpha) \times 100\%$ confidence interval for $\mu_y - \mu_x$ is

$$\bar{Y} - \bar{X} \pm t_{n_x+n_y-2, 1-\alpha/2} S_p \left(\frac{1}{n_x} + \frac{1}{n_y} \right)^{1/2}$$

- Remember this interval is assuming a constant variance across the two groups
- If there is some doubt, assume a different variance per group, which we will discuss later

Likelihood method

- Exactly as before,

$$\frac{\bar{Y} - \bar{X}}{S_p \left(\frac{1}{n_x} + \frac{1}{n_y} \right)^{1/2}}$$

follows a non-central t distribution with non-centrality parameter $\frac{\mu_y - \mu_x}{\sigma \left(\frac{1}{n_x} + \frac{1}{n_y} \right)^{1/2}}$

- Therefore, we can use this statistic to create a likelihood for $(\mu_y - \mu_x)/\sigma$, a standardized measure of the change in group means

Example

Example from Rosner Fundamentals of Biostatistics, Page 304

- Comparing SBP for 8 oral contraceptive users versus 21 controls
- $\bar{X}_{OC} = 132.86$ mmHg with $s_{OC} = 15.34$ mmHg
- $\bar{X}_C = 127.44$ mmHg with $s_C = 18.23$ mmHg
- Pooled variance estimate

$$s_p^2 = \frac{7(15.34)^2 + 20(18.23)^2}{8 + 21 - 2} = 307.8$$


- $t_{27,.975} = 2.052$ (in R, `qt(.975, df = 27)`)
- Interval

$$132.86 - 127.44 \pm 2.052 \left\{ 307.8 \left(\frac{1}{8} + \frac{1}{21} \right)^{1/2} \right\} = [-9.52, 20.36]$$

Likelihood plot for the effect size

Reasonable values for the effect size from the confidence interval

$$[-9.52, 20.36]/sp = [-.54, 1.16]$$



Unequal variances

- Note that under unequal variances

$$\bar{Y} - \bar{X} \sim N\left(\mu_y - \mu_x, \frac{\sigma_x^2}{n_x} + \frac{\sigma_y^2}{n_y}\right)$$

- The statistic

$$\frac{\bar{Y} - \bar{X} - (\mu_y - \mu_x)}{\left(\frac{\sigma_x^2}{n_x} + \frac{\sigma_y^2}{n_y}\right)^{1/2}}$$

approximately follows Gosset's t distribution with degrees of freedom equal to

$$\frac{(S_x^2/n_x + S_y^2/n_y)^2}{\left(\frac{S_x^2}{n_x}\right)^2 / (n_x - 1) + \left(\frac{S_y^2}{n_y}\right)^2 / (n_y - 1)}$$

Example

- Comparing SBP for 8 oral contraceptive users versus 21 controls
- $\bar{X}_{OC} = 132.86$ mmHg with $s_{OC} = 15.34$ mmHg
- $\bar{X}_C = 127.44$ mmHg with $s_C = 18.23$ mmHg
- $df = 15.04$, $t_{15.04,.975} = 2.13$
- Interval

$$132.86 - 127.44 \pm 2.13 \left(\frac{15.34^2}{8} + \frac{18.23^2}{21} \right)^{1/2} = [-8.91, 19.75]$$