


On Schemes for Exponential Decay

Hans Petter Langtangen^{1,2}

¹Center for Biomedical Computing, Simula Research Laboratory

²Department of Informatics, University of Oslo

Jan 8, 2014



1 Goal

The primary goal of this demo talk is to demonstrate how to write talks with [Doconce](#) and get them rendered in numerous HTML formats.

Layout.

This version utilizes latex document slides with the theme `no theme`.


The talk investigates the accuracy of three finite difference schemes for the ordinary differential equation $u' = -au$ with the aid of numerical experiments. Numerical artifacts are in particular demonstrated.

2 Mathematical problem

$$u'(t) = -au(t), \tag{1}$$

$$u(0) = I, \tag{2}$$

- $t \in (0, T]$
- a , I , and T are prescribed parameters
- $u(t)$ is the unknown function



3 Numerical solution method

- Mesh in time: $0 = t_0 < t_1 < \dots < t_N = T$
- Assume constant $\Delta t = t_n - t_{n-1}$
- u^n : numerical approx to the exact solution at t_n

Numerical scheme:

$$u^{n+1} = \frac{1 - (1 - \theta)a\Delta t}{1 + \theta a\Delta t} u^n, \quad n = 0, 1, \dots, N - 1$$


4 Implementation

The numerical method is implemented in a Python function:

```
def solver(I, a, T, dt, theta):
    """Solve u'=-a*u, u(0)=I, for t in (0,T] with steps of dt."""
    dt = float(dt)          # avoid integer division
    N = int(round(T/dt))     # no of time intervals
    T = N*dt                # adjust T to fit time step dt
    u = zeros(N+1)          # array of u[n] values
    t = linspace(0, T, N+1) # time mesh

    u[0] = I                # assign initial condition
    for n in range(0, N):   # n=0,1,...,N-1
        u[n+1] = (1 - (1-theta)*a*dt)/(1 + theta*dt*a)*u[n]
    return u, t
```

4.1 The Crank-Nicolson method



4.2 The artifacts can be explained by some theory

Exact solution of the scheme:

$$u^n = A^n, \quad A = \frac{1 - (1 - \theta)a\Delta t}{1 + \theta a\Delta t}.$$

- Stability: $|A| < 1$
- No oscillations: $A > 0$
- Always for Backward Euler ($\theta = 1$)
- $\Delta t < 1/a$ for Forward Euler ($\theta = 0$)
- $\Delta t < 2/a$ for Crank-Nicolson ($\theta = 1/2$)

Concluding remarks:

Only the Backward Euler scheme is guaranteed to always give qualitatively correct results.