

11 Quadratic Residues

Introduction: The concept of Quadratic Residues is a fundamental tool which has ramifications in lots of other number theory places: Cryptography, Factoring, etc...

11.1 Quadratic Residues & Nonresidues

1. **Introduction:** Suppose we asked the following, given a modulus m : Which numbers are perfect squares mod m ?

Ex. Let $m = 7$. What are the perfect squares? We could of course work backwards, squaring each value:

$$0^2 \equiv 0 \pmod{7}$$

$$1^2 \equiv 1 \pmod{7}$$

$$2^2 \equiv 4 \pmod{7}$$

$$3^2 \equiv 2 \pmod{7}$$

$$4^2 \equiv 2 \pmod{7}$$

$$5^2 \equiv 4 \pmod{7}$$

$$6^2 \equiv 1 \pmod{7}$$

Then the perfect squares are 0, 1, 2, 4 and 3, 5, 6 are not.

2. Quadratic Residues & Nonresidues - Counting

- (a) **Definition:** Let m be a modulus and $a \in \mathbb{Z}$ with $\gcd(a, m) = 1$. We say a is a *quadratic residue mod m* if $\exists x \in \mathbb{Z}$ such that $x^2 \equiv a \pmod{m}$. Otherwise, we say a is a *quadratic nonresidue mod m* if $\nexists x \in \mathbb{Z}$ such that $x^2 \equiv a \pmod{m}$.

Ex. If $m = 7$ then QR:1, 2, 4, QNR:3, 5, 6, and Neither:0.

- (b) **Theorem:** If p is an odd prime and $a \in \mathbb{Z}$ with $p \nmid a \implies \gcd(p, a) = 1$, then $x^2 \equiv a \pmod{p}$ has either no solutions or exactly two solutions mod p .

Proof. If there are none, we are done. Suppose x is one solution to $x^2 \equiv a \pmod{p}$. Claim $-x$ is also a solution. Then $2x \equiv 0 \pmod{p}$. Since p is odd we can do $x \equiv 0 \pmod{p}$ which implies $p \mid x \implies p \mid x^2$. Then, $x^2 \equiv 0 \pmod{p} \implies a \equiv 0 \pmod{p}$ which contradicts $p \nmid a$.

Let's show that for any two solutions, they are negative of one another. Suppose $x_1^2 \equiv a \pmod{p}$ and $x_2^2 \equiv a \pmod{p}$. Then $x_1^2 - x_2^2 \equiv 0 \pmod{p}$ so

$p \mid (x_1^2 - x_2^2)$ so $p \mid (x_1 - x_2)(x_1 + x_2)$ so $p \mid (x_1 - x_2)$ or $p \mid (x_1 + x_2)$.
If $p \mid (x_1 - x_2)$ then $x_1 \equiv x_2 \pmod{p}$. If $p \mid (x_1 + x_2)$ then $x_1 \equiv -x_2 \pmod{p}$.
Thus, there can only be the two which are negatives of one another \square

- (c) **Theorem:** Suppose p is an odd prime. Then $\exists \frac{p-1}{2}$ QR and $\exists \frac{p-1}{2}$ QNR.

Proof. If we square all of $1, 2, 3, \dots, p-1$ the results will be in pairs (two of every result) the $\frac{p-1}{2}$ we do get are the QR. We miss $\frac{p-1}{2}$ results, those are the QNR. \square

- (d) **Theorem:** Let p be an odd prime and r a primitive root mod p . Suppose $p \nmid a$, then a is a QR mod p if and only if $\text{ind}_r a$ is even.

Proof.

\rightarrow Suppose a is a quadratic residue mod p , $\exists x$ such that $x^2 \equiv a \pmod{p}$.
Then take the index of both sides to get $\text{ind}_r x^2 \equiv \text{ind}_r a \pmod{p-1}$ and
so $2\text{ind}_r x \equiv \text{ind}_r a \pmod{p-1}$. From here we see $\text{ind}_r a = 2\text{ind}_r x + k(p-1)$
for some $k \in \mathbb{Z}$ and so since $p-1$ is even we know $\text{ind}_r a$ is even.

\leftarrow Suppose $\text{ind}_r a$ is even. Say $\text{ind}_r a = 2k$ for $k \in \mathbb{Z}$ so $r^{2k} \equiv a \pmod{p}$ so
 $(r^k)^2 \equiv a \pmod{p}$. Then, a is a quadratic residue mod p . \square

To illustrate: $r = 3$ is a primitive root mod 17.

$a \pmod{17}$	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16
$\text{ind}_3 a$	<u>16</u>	<u>14</u>	1	<u>12</u>	5	15	11	<u>10</u>	<u>2</u>	3	7	13	<u>4</u>	9	<u>6</u>	<u>8</u>

So what this theorem tells us is that $a = 1, 2, 4, 8, 9, 13, 15, 16$ are the quadratic residues

3. The Legendre Symbol and Properties

- (a) **Definition:** Given an odd prime p and $a \in \mathbb{Z}$ with $\gcd(a, p) = 1$, define the Legendre Symbol:

$$\left(\frac{a}{p}\right) = \begin{cases} 1 & \text{if } a \text{ is a quadratic residue mod } p \\ -1 & \text{if } a \text{ is a quadratic nonresidue mod } p \end{cases}$$

Ex. If $p = 7$ we have:

$$\left(\frac{1}{7}\right) = \left(\frac{2}{7}\right) = \left(\frac{4}{7}\right) = 1$$

$$\left(\frac{3}{7}\right) = \left(\frac{5}{7}\right) = \left(\frac{6}{7}\right) = -1$$

Since 1, 2, 4 are QR mod 7 and 3, 5, 6 are QNR mod 7.

- (b) **Euler's Criterion:** If p is an odd prime and $a \in \mathbb{Z}$ with $\gcd(a, p) = 1$ then:

$$\left(\frac{a}{p}\right) \equiv a^{(p-1)/2} \pmod{p}$$

Proof. Suppose $\left(\frac{a}{p}\right) = 1$ then $\exists x$ such that $x^2 \equiv a \pmod{p}$. Then observe, $a^{(p-1)/2} \equiv (x^2)^{(p-1)/2} = x^{p-1} \equiv 1 \pmod{p}$ by Euler's Theorem/Fermat's Little Theorem they are equal.

Suppose $\left(\frac{a}{p}\right) = -1$. Consider the list $\{1, 2, \dots, p-1\}$, each is coprime to p and there are an even number of them because p is odd. Suppose $b \in \{1, 2, \dots, p-1\}$, then consider the equation $bx \equiv a \pmod{p}$. Since $\gcd(b, p) = 1 \mid a$, $\exists!$ solution. Could $x \equiv b \pmod{p}$? No because if $b \cdot b \equiv a \pmod{p} \implies b^2 \equiv a \pmod{p}$ but then a would be a QR mod p . Since the solution is not b it is another element in the set $\{1, 2, \dots, p-1\}$. Thus all of $\{1, 2, \dots, p-1\}$ pair up to give pairs whose products are a . Thus,

$$\underbrace{(1)(2) \cdots (p-1)}_{\text{Wilson's Theorem}} \equiv a^{(p-1)/2} \pmod{p}$$

$$a^{(p-1)/2} \equiv -1 \pmod{p}$$

□

Ex. $\left(\frac{6}{11}\right) = 6^{(11-1)/2} = 6^5 \equiv 10 \equiv -1 \pmod{11}$. So 6 is a QNR mod 11. i.e. $x^2 \equiv 6 \pmod{11}$ has no solution.

- (c) **Theorem:** If p is an odd prime and $a \in \mathbb{Z}$ with $\gcd(a, p) = \gcd(b, p) = 1$ then:

- i. If $a \equiv b \pmod{p}$ then $\left(\frac{a}{p}\right) = \left(\frac{b}{p}\right)$. This statements that we can reduce the numerator mod the denominator.

Proof. Clear because $x^2 \equiv a \pmod{p}$ if and only if $x^2 \equiv b \pmod{p}$ because $a \equiv b \pmod{p}$. □

- ii. $\left(\frac{ab}{p}\right) = \left(\frac{a}{p}\right) \left(\frac{b}{p}\right)$

Proof. Well,

$$\left(\frac{ab}{p}\right) \equiv (ab)^{(p-1)/2} \equiv a^{(p-1)/2} b^{(p-1)/2} \equiv \left(\frac{a}{p}\right) \left(\frac{b}{p}\right) \pmod{p}$$

So $\left(\frac{ab}{p}\right) \equiv \left(\frac{a}{p}\right) \left(\frac{b}{p}\right) \pmod{p}$ so $p \mid \left[\left(\frac{ab}{p}\right) - \left(\frac{a}{p}\right) \left(\frac{b}{p}\right)\right]$ but $p \geq 3$ Since $\left(\frac{ab}{p}\right) - \left(\frac{a}{p}\right) \left(\frac{b}{p}\right)$ is between -2 and 2 and p divides it, we know that it must be 0. Therefore, $\left(\frac{ab}{p}\right) = \left(\frac{a}{p}\right) \left(\frac{b}{p}\right)$. □

iii. $\left(\frac{a^2}{p}\right) = 1$

Proof. Obvious. \square

- (d) **Gauss' Lemma:** Suppose p is an odd prime and $a \in \mathbb{Z}$ with $\gcd(a, p) = 1$. Let s be the number of least nonnegative residues in the set

$$\{a, 2a, \dots, ((p-1)/2)a\}$$

which are $> p/2$. Then $\left(\frac{a}{p}\right) = (-1)^s$.

Ex. Consider $\left(\frac{8}{13}\right)$. Note that $\left(\frac{p-1}{2}\right) = \frac{12}{2} = 6$ so look at

$$\{8, 2 \cdot 8, 3 \cdot 8, \dots, 6 \cdot 8\} \equiv \{8, 3, 11, 6, 1, 9\} \pmod{13}$$

Since only three of these are greater than $p/2 = 6.5$ we have $\left(\frac{8}{13}\right) = (-1)^3 = -1$. Thus, 8 is a quadratic nonresidue mod 13.

4. Two Special Cases

These will turn out to be really useful after 11.2 and 11.3 .

- (a) **Theorem:** Suppose p is an odd prime, then

$$\left(\frac{-1}{p}\right) = \begin{cases} 1 & \text{if } p \equiv 1 \pmod{4} \\ -1 & \text{if } p \equiv 3 \pmod{4} \end{cases}$$

Proof. By Euler's Criterion we have,

$$\left(\frac{-1}{p}\right) \equiv (-1)^{(p-1)/2} \pmod{p}$$

If $p \equiv 1 \pmod{4}$ then $p = 4k + 1$ for some $k \in \mathbb{Z}$ so:

$$(-1)^{(p-1)/2} = (-1)^{(4k+1-1)/2} = (-1)^{2k} = 1$$

If $p \equiv 3 \pmod{4}$ then $p = 4k + 3$ for some $k \in \mathbb{Z}$ so:

$$(-1)^{(p-1)/2} = (-1)^{(4k+3-1)/2} = (-1)^{2k+1} = -1$$

\square

- (b) **Theorem:** Suppose p is an odd prime, then

$$\left(\frac{2}{p}\right) = \begin{cases} 1 & \text{if } p \equiv 1, 7 \pmod{8} \\ -1 & \text{if } p \equiv 3, 5 \pmod{8} \end{cases}$$

Proof. Not obvious as it uses Gauss' Lemma and is lengthy. \square

Note: This is equivalent to

$$\left(\frac{2}{p}\right) = (-1)^{(p^2-1)/8}$$

11.2 Quadratic Reciprocity and Calculation Examples

1. **Introduction:** The Law of Quadratic reciprocity establishes that for odd primes p and q there is a connection between when p is a quadratic residue mod q when q is a quadratic residue mod p .

2. **Theorem:** If p, q are odd primes then

$$\left(\frac{p}{q}\right) \left(\frac{q}{p}\right) = (-1)^{\left(\frac{p-1}{2}\right)\left(\frac{q-1}{2}\right)}$$

Proof. Omitted due to length. □

Use for Calculation: Under what circumstances will $\left(\frac{p}{q}\right)$ and $\left(\frac{q}{p}\right)$ be identical? We would need $\left(\frac{p-1}{2}\right)\left(\frac{q-1}{2}\right)$ to be even. This happens if and only if one of the two is even, say $\frac{p-1}{2}$ is even. That is, $\frac{p-1}{2} = 2k$ for some $k \in \mathbb{Z}$, so $p - 1 = 4k$ so $p \equiv 1 \pmod{4}$. Thus, for calculation, we get:

$$\left(\frac{p}{q}\right) = \begin{cases} \left(\frac{p}{q}\right) & \text{if either } p \equiv 1 \pmod{4} \text{ or } q \equiv 1 \pmod{4} \text{ (or both).} \\ -\left(\frac{q}{p}\right) & \text{if both } p \equiv 3 \pmod{4} \text{ and } q \equiv 3 \pmod{4}. \end{cases}$$

3. **Theorem:**

- (a) If $a \equiv b \pmod{p}$ then $\left(\frac{a}{p}\right) = \left(\frac{b}{p}\right)$. Call this "reducing".
- (b) $\left(\frac{ab}{p}\right) = \left(\frac{a}{p}\right) \left(\frac{b}{p}\right)$. Call this "splitting".
- (c) $\left(\frac{a^2}{p}\right) = 1$. Call this the "square rule".
- (d) $\left(\frac{-1}{p}\right) = \begin{cases} 1 & \text{if } n \equiv 1 \pmod{4} \\ -1 & \text{if } n \equiv 3 \pmod{4} \end{cases}$. Call this the "-1 rule".
- (e) $\left(\frac{2}{p}\right) = \begin{cases} 1 & \text{if } n \equiv 1, 7 \pmod{8} \\ -1 & \text{if } n \equiv 3, 5 \pmod{8} \end{cases}$. Call this the "2 rule".

4. **Examples:**

Ex. Calculate $\left(\frac{48}{29}\right)$:

$$\begin{aligned} \left(\frac{48}{29}\right) &= \left(\frac{19}{29}\right) \text{ by reducing} \\ \left(\frac{19}{29}\right) &= \left(\frac{29}{19}\right) \text{ by LoQR since } 19 \equiv 1 \pmod{4}. \\ \left(\frac{29}{19}\right) &= \left(\frac{10}{19}\right) \text{ by reducing.} \\ \left(\frac{10}{19}\right) &= \left(\frac{2}{19}\right) \left(\frac{5}{19}\right) \text{ by splitting.} \end{aligned}$$

Then, we calculate these separately. First $\left(\frac{2}{19}\right) = -1$ by the "2 rule" because $19 \equiv 2 \pmod{8}$. Then second,

$$\left(\frac{5}{19}\right) = \left(\frac{19}{5}\right) \text{ by LoQR since } 5 \equiv 1 \pmod{4}.$$

$$\left(\frac{19}{5}\right) = \left(\frac{4}{5}\right) \text{ by reducing.}$$

$$\left(\frac{4}{5}\right) = 1 \text{ by square rule.}$$

Thus $\left(\frac{48}{29}\right) = (-1)(1) = -1$.

Ex. Calculate $\left(\frac{105}{1009}\right)$. Note that 105 is not prime so we cannot use the LoQR immediately.

$$\left(\frac{105}{1009}\right) = \left(\frac{3}{1009}\right) \left(\frac{5}{1009}\right) \left(\frac{7}{1009}\right) \text{ by splitting.}$$

Then we calculate these separately. First,

$$\left(\frac{3}{1009}\right) = \left(\frac{1009}{3}\right) \text{ by LoQR since } 1009 \equiv 1 \pmod{4}.$$

$$\left(\frac{1009}{3}\right) = \left(\frac{1}{3}\right) \text{ by reducing}$$

$$\left(\frac{1}{3}\right) = 1$$

Second,

$$\left(\frac{5}{1009}\right) = \left(\frac{1009}{5}\right) \text{ by LoQR since } 1009 \equiv 1 \pmod{4}.$$

$$\left(\frac{1009}{5}\right) = \left(\frac{4}{5}\right) \text{ by reducing}$$

$$\left(\frac{4}{5}\right) = 1 \text{ by the square rule}$$

Third,

$$\left(\frac{7}{1009}\right) = \left(\frac{1009}{7}\right) \text{ by LoQR since } 1009 \equiv 1 \pmod{4}.$$

$$\left(\frac{1009}{7}\right) = \left(\frac{1}{7}\right) \text{ by reducing}$$

$$\left(\frac{1}{7}\right) = 1$$

Thus, $\left(\frac{105}{1009}\right) = (1)(1)(1) = 1$.

11.3 The Jacobi Symbol

1. **Introduction:** The Jacobi symbol is a generalization of the Legendre symbol for when the denominator is odd but not necessarily prime. It preserves many of the same useful properties and almost the same meaning.
2. **Definition:** Let n be an odd positive integer with prime factorization $n = p_1^{\alpha_1} \cdots p_k^{\alpha_k}$ and let $a \in \mathbb{Z}$ be coprime to n . Define:

$$\left(\frac{a}{n}\right) = \left(\frac{a}{p_1}\right)^{\alpha_1} \cdots \left(\frac{a}{p_k}\right)^{\alpha_k}$$

Thus the Jacobi symbol is defined in terms of the Legendre symbol.

3. **Theorem:** Assume $\gcd(a, n) = \gcd(b, n) = 1$.
 - (a) If $a \equiv b \pmod{n}$ then $\left(\frac{a}{n}\right) = \left(\frac{b}{n}\right)$.
 - (b) $\left(\frac{a^2}{n}\right) = 1$
 - (c) $\left(\frac{ab}{n}\right) = \left(\frac{a}{n}\right) \left(\frac{b}{n}\right)$
 - (d) $\left(\frac{-1}{n}\right) = \begin{cases} 1 & \text{if } n \equiv 1 \pmod{4} \\ -1 & \text{if } n \equiv 3 \pmod{4} \end{cases}$
 - (e) $\left(\frac{2}{n}\right) = \begin{cases} 1 & \text{if } n \equiv 1, 7 \pmod{8} \\ -1 & \text{if } n \equiv 3, 5 \pmod{8} \end{cases}$
 - (f) $\left(\frac{m}{n}\right) = \begin{cases} \left(\frac{m}{n}\right) & \text{if either } m \equiv 1 \pmod{4}, n \equiv 1 \pmod{4} \\ -\left(\frac{n}{m}\right) & \text{if both } m \equiv 3 \pmod{4} \text{ and } n \equiv 3 \pmod{4} \end{cases}$

Proof. Lots of calculation. □

4. **Question:** We know $\left(\frac{a}{p}\right)$ tells us if a is a QR or QNR mod p . Does $\left(\frac{a}{n}\right)$ tell us if a is a QR or QNR mod n ? Well, half-yes.

Theorem: Suppose $\gcd(a, n) = 1$ and n is an odd prime.

- (a) If a is a QR mod n then $\left(\frac{a}{n}\right) = 1$.
- (b) If $\left(\frac{a}{n}\right) = 1$ then we cannot conclude a is a QR mod n .

11.4 Problems

1. Determine, by squaring, which of $1, \dots, 16$ are quadratic residues of $p = 17$.
2. Calculate $\left(\frac{3}{17}\right)$ by
 - (a) Euler's Criterion
 - (b) Gauss's Lemma
3. Prove that if p and $q = 2p + 1$ are both odd primes then -4 is a primitive root of q .
4. Prove that if $p \equiv 1 \pmod{4}$ is a prime then -4 and $(p-1)/4$ are both quadratic residues of p .
5. Calculate each of the following:
 - (a) $\left(\frac{21}{59}\right)$
 - (b) $\left(\frac{1463}{89}\right)$
 - (c) $\left(\frac{1547}{1913}\right)$
6. Using the Law of Quadratic Reciprocity, show that if p is an odd prime that

$$\left(\frac{3}{p}\right) = \begin{cases} 1 & \text{if } p \equiv \pm 1 \pmod{12} \\ -1 & \text{if } p \equiv \pm 5 \pmod{12} \end{cases}$$

7. Classify all primes p with $\left(\frac{5}{p}\right) = 1$
8. Calculate each of the following using properties of the Jacobi Symbol, not by raw calculation.
 - (a) $\left(\frac{5}{21}\right)$
 - (b) $\left(\frac{1009}{2307}\right)$
 - (c) $\left(\frac{27}{101}\right)$
9. Categorize all positive integers n which are relatively prime to 15 and for which $\left(\frac{15}{n}\right) = 1$.
10. Show that if $a > 0$ is not a perfect square then there exists a positive integer n such that $\left(\frac{a}{n}\right) = -1$.