
Viscosity by averaging window

$\dot{\gamma} = 100\ s^{-1}$; each average ends at its own file\'s final strain



Final strain by attraction level
0%: 16.2; 10%: 23.55; 25%: 26.3; 50%: 29; 100%: 27.6


Final strain by attraction level
3%: 20.45; 6%: 23; 9%: 22.95; 12%: 22.15; 15%: 25.5

$\overline{\eta} = \overline{\sigma}_{xz}/\dot{\gamma}$. Interval means are weighted by covered strain; all fluctuations are retained.

Logarithmic y axes with panel-specific ranges. Dashed lines connect sampled cases; they are not fitted models.

Relative viscosity by averaging window

$\dot{\gamma} = 100\ s^{-1}$; each average ends at its own file\'s final strain



Final strain by attraction level
0%: 16.2; 10%: 23.55; 25%: 26.3; 50%: 29; 100%: 27.6


Final strain by attraction level
3%: 20.45; 6%: 23; 9%: 22.95; 12%: 22.15; 15%: 25.5

$\overline{\eta}_r = \overline{\eta}/\eta_m$; $\eta_m = 0.000890$ Pa s (pure_Gr), 0.0020427132 Pa s (CMC 4 g/L).

Logarithmic y axes with panel-specific ranges. Dashed lines connect sampled cases; they are not fitted models.

Sensitivity to the averaging start strain

Each series is divided by its value over $[5, \gamma_{end}]$. All three windows share the same endpoint.



Contact counts are weighted means of recorded endpoint counts; viscosity uses solver interval-averaged stress.

pure_Gr 0% has only 1.20 strain units after strain 15. Every requested mean is shown, including this short window.

Overlapping windows are correlated. Agreement is a convergence diagnostic, not an independent-replicate uncertainty estimate.