Clinical Protocol RAG

Clinical Assessment Report

Clinical Assessment

Report ID: CR-20261022-200130

22 Oct 2026, 08:01 PM

Patient Information

PATIENT NAME		PATIENT ID	DATE
Saif Ahmed		**P-001**	**22 Oct 2026, 08:01 PM**
AGE	SEX	CONTACT	DURATION
24	**Male**	**01334173316**	**toa**
ADDRESS		ALLERGIES	
Dhaka		**—**	

Vital Signs

BLOOD PRESSURE	PULSE	TEMPERATURE	SPO_2
120/80 mmHg	**80 bpm**	**99 °F**	**98 %**
WEIGHT	HEIGHT		
62 kg	**168 cm**		

Clinical Presentation

CHIEF COMPLAINTS

Acute watery diarrhoea

ADDITIONAL FINDINGS

Very weak pulse, sunken eyes, unable to drink properly, severe dehydration

Assessment

PRIMARY CONTEXT	DIFFERENTIAL
Cholera	**—**

Probable Diagnosis

Probable Cholera. Consider suspected cholera according to outbreak context. Outside a declared cholera outbreak, a patient aged 2 years or older with acute watery diarrhoea and severe dehydration, or dying from acute watery diarrhoea, meets the suspected cholera case definition. During a declared cholera outbreak, any person presenting with or dying from acute watery diarrhoea meets the suspected case definition.

Investigations

• Classify dehydration severity before selecting treatment. Severe dehydration is present with one or more danger signs such as lethargy or unconsciousness, absent or weak pulse, or respiratory distress, or with at least two severe dehydration findings such as sunken eyes, inability to drink or drinking poorly, and a skin pinch that returns very slowly. Some dehydration is considered when there are no danger signs and at least two findings such as irritability or restlessness, sunken eyes, rapid pulse, thirst with eager drinking, or a skin pinch that returns slowly. If the patient is awake and alert with normal pulse, normal thirst, eyes not sunken and normal skin pinch, classify as no dehydration.

Medication Planning

• Doxycycline; dose: 200 mg if under 12 years; 300 mg for older children and adults; route: oral; frequency: single dose

• Azithromycin; dose: 20 mg/kg if under 12 years; 1 g for older children and adults; route: oral; frequency: single dose; instructions: alternative to doxycycline
• Ringer's lactate; dose: 100 mL/kg total; route: intravenous; frequency: 30 mL/kg initially then 70 mL/kg; duration: age-dependent infusion schedule; instructions: under 12 months: 30 mL/kg in 1 hour then 70 mL/kg over 5 hours; 12 months or older: 30 mL/kg in 30 minutes then 70 mL/kg over 2.5 hours

Advice

No additional patient advice was explicitly supported by the currently applicable cards.

Evidence

• CHOLERA_ASSESS_001: Clinical tools for cholera treatment facilities (WHO 2024), page 6
• CHOLERA_DIAG_001: Clinical tools for cholera treatment facilities (WHO 2024), page 5
• CHOLERA_ABX_001: Clinical tools for cholera treatment facilities (WHO 2024), page 5
• CHOLERA_MGMT_001: Clinical tools for cholera treatment facilities (WHO 2024), page 7

Prepared / Reviewed By	Clinician Signature	Seal / Date