











Ashima Garg

Course: GATE

Computer Science Engineering(CS)

-  HOME
-  MY TEST
-  BOOKMARKS
-  MY PROFILE
- REPORTS
-  BUY PACKAGE
-  ASK AN EXPERT
-  OFFER
-  EXCLUSIVE OFFER FOR OTS STUDENTS ONLY ON BOOK PACKAGES

TOPICWISE : ENGINEERING MATHEMATICS-2 (GATE - 2019) - REPORTS

OVERALL ANALYSIS COMPARISON REPORT SOLUTION REPORT

ALL(17) CORRECT(10) INCORRECT(5) SKIPPED(2)

Q. 1

Consider the rank of matrix 'A' of size $(m \times n)$ is " $m - 1$ ". Then, which of the following is true?

[Have any Doubt ?](#)

A

A A^T will be invertible.

B

A have " $m - 1$ " linearly independent rows and " $m - 1$ " linearly independent column.

Correct Option

Solution :
(b)

Rank of matrix is " $m - 1$ ", so it must have " $m - 1$ " linearly independent rows as well as " m independent columns.

C

A will have " m " linearly independent rows and " n " linearly independent columns.

D

A will have " $m - 1$ " linearly independent rows and " $n - 1$ " independent columns.

QUESTION ANALYTICS

Q. 2

For function $f(x) = 4x^3 - 6x^2$, the maximum occurs in interval $[-1, 2]$ when x is equal to

[Have any Doubt ?](#)

A

0

B

- 1

C

1

D

2

Your answer is **Correct**

Solution :
(d)


$$\begin{aligned} f(x) &= 4x^3 - 6x^2 \\ \frac{d f(x)}{dx} &= 12x^2 - 12x \\ 12x^2 - 12x &= 0 \\ 12x [x - 1] &= 0 \\ x &= 0, 1 \end{aligned}$$




Ashima Garg

Course: GATE

Computer Science Engineering(CS)

HOME

MY TEST


BOOKMARKS


MY PROFILE

REPORTS

BUY PACKAGE

ASK AN EXPERT

OFFER

EXCLUSIVE OFFER FOR OTS
STUDENTS ONLY ON BOOK
PACKAGES

At $x = 1$, $24 \times 1 - 12 = 12 > 0$ minima

So, at

$$x = -1, f(-1) = 4(-1)^3 - 6(-1)^2 = -4 - 6 = -10$$

$$x = 0, f(0) = 4(0)^3 - 6(0)^2 = 0$$

$$x = 1, f(1) = 4(1)^3 - 6(1)^2 = 4 - 6 = -2$$

$$x = 2, f(2) = 4(2)^3 - 6(2)^2 = 32 - 24 = 8$$

So maximum value occurs at $x = 2$.

QUESTION ANALYTICS

Q. 3

Find the limit?

$$\lim_{x \rightarrow \infty} \left[1 + \frac{3}{2x} \right]^{5x}$$

Have any Doubt ?

A
 e^{15}

B
 e^3

C
 $e^{15/2}$

Your answer is **Correct**

Solution :

(c)

$$\lim_{x \rightarrow \infty} \left[1 + \frac{3}{2x} \right]^{5x}$$

Put limit $x \rightarrow \infty$

1^∞ form create,

So, we know, for form 1^∞

$$\lim_{x \rightarrow \infty} f(x)^{g(x)} = e^{\left(\lim_{x \rightarrow \infty} (f(x)-1) g(x) \right)}$$

Apply in given function:

$$= e^{\lim_{x \rightarrow \infty} \left[1 + \frac{3}{2x} - 1 \right] 5x}$$

$$= e^{\lim_{x \rightarrow \infty} \left[\frac{3}{2x} \right] 5x}$$

$$= e^{15/2}$$

D
 $e^{5/3}$

QUESTION ANALYTICS

Q. 4

Consider the following function:

$$f(x) = \begin{cases} -1.5x^2, & x \leq -2 \\ 6x - 5, & x > -2 \end{cases}$$

Which of the following is true at $x = -2$?

Have any Doubt ?

A


Ashima Garg

Course: GATE

Computer Science Engineering(CS)

HOME

MY TEST

BOOKMARKS

MY PROFILE

REPORTS

BUY PACKAGE

ASK AN EXPERT

OFFER

 EXCLUSIVE OFFER FOR OTS
STUDENTS ONLY ON BOOK
PACKAGES

Differentiable and continuous both

C

Differentiable but not continuous

 Your answer is **Wrong**

D

neither continuous nor differentiable

Correct Option

Solution :

(d)

Check for continuous:

$$f(-2) = -1.5 \times (-2)^2 = -6$$

$$f(-2^+) = 6(-2) - 5 = -17$$

$$f(-2^-) = -1.5 \times (-2)^2 = -6$$

$$f(-2^-) \neq f(-2^+)$$

Function is not continuous, hence cannot be differentiable i.e. differentiable \rightarrow continuous.

QUESTION ANALYTICS

Q. 5

Consider a man is known to speak truth 3 out of 5 times, he throw a die and reports the number obtained is 2. What is the probability that the number obtained is actually 2?

Have any Doubt ?

A


$$\frac{13}{30}$$

B

$$\frac{3}{13}$$

 Your answer is **Correct**
Solution :

(b)

Applying Bayes Theorem:


So,

$$P(\text{spoke truth/reports 2}) = \frac{P(\text{spoke truth} \cap \text{reports 2})}{P(\text{reports 2})}$$

$$= \frac{\frac{3}{5} \times \frac{1}{6}}{\frac{3}{5} \times \frac{1}{6} + \frac{2}{5} \times \frac{5}{6}} = \frac{3}{13}$$

C

$$\frac{1}{10}$$

D

None of the above



Ashima Garg

Course: GATE

Computer Science Engineering(CS)

HOME

MY TEST

BOOKMARKS

MY PROFILE

REPORTS

BUY PACKAGE

ASK AN EXPERT

OFFER

 EXCLUSIVE OFFER FOR OTS
STUDENTS ONLY ON BOOK
PACKAGES

Q. 6

The value of

$$\int_0^2 \frac{1}{(3+2x)^2} dx = \text{_____}. \text{ (Upto 3 decimal places)}$$

[Have any Doubt ?](#)

0.095 (0.095 - 0.096)

 Your answer is **Correct**.095

Solution :

0.095 (0.095 - 0.096)

Consider,

$$u = 3 + 2x$$

$$\frac{du}{dx} = 2$$

$$dx = \frac{du}{2}$$

Calculate new limits:

$$x = 0, u = 3 + 2 \cdot x = 3 + 0 = 3$$

$$x = 2, u = 3 + 2 \cdot x = 3 + 2 \times 2 = 7$$

By substitution:

$$= \int_3^7 \frac{1}{u^2} \cdot \frac{1}{2} du$$

$$= \frac{1}{2} [-u^{-1}]_3^7$$

$$= \frac{1}{2} \left[\frac{1}{3} - \frac{1}{7} \right]$$

$$= \frac{1}{2} \left[\frac{4}{21} \right] = \frac{2}{21}$$

$$= 0.095$$

QUESTION ANALYTICS

Q. 7

The maximum value of the function:

$$f(x) = x^3 - 9x^2 + 24x + 5$$

 in interval of $[-3 \text{ to } 3]$ is _____.

[FAQ](#) | [Have any Doubt ?](#)

25

 Your answer is **Correct**25

Solution :

25

$$f(x) = x^3 - 9x^2 + 24x + 5$$

$$\frac{df}{dx} = 3x^2 - 18x + 24$$

Function attains local minimum or maximum at critical points.

 Critical points are those where $f'(x) = 0$

$$3x^2 - 18x + 24 = 0$$

$$x^2 - 6x + 8 = 0$$

$$x^2 - 4x - 2x + 8 = 0$$

$$x(x - 4) - 2(x - 4) = 0$$

$$(x - 2)(x - 4) = 0$$

$$x = 2, 4$$

$$\frac{df'(x)}{dx} = 6x - 18$$



Ashima Garg

Course: GATE

Computer Science Engineering(CS)

- HOME
- MY TEST
- BOOKMARKS
- MY PROFILE
- REPORTS
- BUY PACKAGE
- ASK AN EXPERT
- OFFER
- EXCLUSIVE OFFER FOR OTS
STUDENTS ONLY ON BOOK
PACKAGES

x	$f(x)$
-3	Some value in negative
2	25
3	23

Hence maximum value is 25 at $x = 2$.

QUESTION ANALYTICS

Q. 8

Consider

$$f(x) = \begin{cases} -x, & x \leq 1 \\ 1+x, & x \geq 1 \end{cases} \text{ and } g(x) = \begin{cases} 1-x, & x \leq 0 \\ x^2, & x > 0 \end{cases}$$

The composition of f and g i.e. $g \circ f(x) = g(f(x))$. Then out of $f(x)$, $g(x)$ and $g \circ f(x)$ in the interval $(-\infty, 0)$, how many are discontinuous _____.

Have any Doubt ?

0

Correct Option

Solution :

0

For interval $(-\infty, 0)$

$$f(x) = -x; x < 0$$

$$g(x) = 1 - x; x \leq 0$$

Both are continuous for $x < 0$ and we know composition of two continuous function is continuous. So, $g \circ f(x)$ is also continuous.

Hence no function is discontinuous.

Your Answer is 1

QUESTION ANALYTICS

Q. 9

Consider a 3×3 matrix 'A' having $\det(A) = -5$. The value of $\det(4A)$ is _____.

Have any Doubt ?

-320

Your answer is Correct-320

Solution :

-320

We know that,

$$A = \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix}$$

$$4A = \begin{bmatrix} 4a & 4b & 4c \\ 4d & 4e & 4f \\ 4g & 4h & 4i \end{bmatrix}$$

$$\det(4A) = 4^3 \begin{vmatrix} a & b & c \\ d & e & f \\ g & h & i \end{vmatrix}$$



Ashima Garg

Course: GATE

Computer Science Engineering(CS)

HOME

MY TEST

BOOKMARKS

MY PROFILE

REPORTS

BUY PACKAGE

ASK AN EXPERT

OFFER

EXCLUSIVE OFFER FOR OTS
STUDENTS ONLY ON BOOK
PACKAGES

QUESTION ANALYTICS

Q. 10

Consider the following function:

$$f(x) = \begin{cases} \frac{x-c}{1+c}, & \text{if } x \leq 0 \\ x^2 + c, & \text{if } x > 0 \end{cases}$$

Which of the following value of c , for which function is continuous for every ' x '?

Have any Doubt ?

A
2

B
-2

C
0

D
Both (b) and (c)

Your answer is **Correct**

Solution :
(d)

function $f(x)$ is continuous for every $x \neq 0$ (since $\frac{x-c}{1+c}$ and $x^2 + c$ are polynomials, and polynomials are continuous).

$$\begin{aligned} f(0) &= \frac{0-c}{1+c} = \frac{-c}{1+c} \\ \lim_{x \rightarrow 0^-} \frac{0-c}{1+c} &= \frac{-c}{1+c} \\ \lim_{x \rightarrow 0^+} 0^2 + c &= c \end{aligned}$$

Since $f(x)$ is continuous for every x , hence continuous for $x = 0$.

$$\begin{aligned} \Rightarrow f(0) &= \lim_{x \rightarrow 0^-} f(x) = \lim_{x \rightarrow 0^+} f(x) \\ \Rightarrow \frac{-c}{1+c} &= c \\ \Rightarrow -c &= c(1+c) \\ c^2 + 2c &= 0 \\ c &= -2 \text{ or } c = 0 \end{aligned}$$

So option (d) is correct answer

QUESTION ANALYTICS

Q. 11

Which of the following matrix is LU decomposable?

Have any Doubt ?

A

$$\begin{bmatrix} 1 & 2 & 3 \\ 2 & 4 & 5 \\ 1 & 3 & 4 \end{bmatrix}$$

B

$$\begin{bmatrix} 3 & 2 \end{bmatrix}$$



Ashima Garg

Course: GATE
Computer Science Engineering(CS)

HOME

MY TEST

BOOKMARKS

MY PROFILE

REPORTS

BUY PACKAGE

ASK AN EXPERT

OFFER

EXCLUSIVE OFFER FOR OTS
STUDENTS ONLY ON BOOK
PACKAGES

Solution :

(b)

To check matrix is LU decomposable by checking if principal minors have non-zero determinants.

Check (a):

$$|A_1| = |1| = 1 \neq 0$$

Now

$$|A_2| = \begin{vmatrix} 1 & 2 \\ 2 & 4 \end{vmatrix} = 0$$

So option (a) is not LU decomposable.

Check (b):

$$\begin{bmatrix} 3 & 2 \\ 0 & 1 \end{bmatrix} \text{ here } |A_1| = 3, |A_2| = \begin{vmatrix} 3 & 2 \\ 0 & 1 \end{vmatrix} = 3 - 0 = 3$$

So LU decomposable.

Check (c):

$$\begin{bmatrix} 0 & 1 \\ 3 & 2 \end{bmatrix} \text{ here } |A_1| = 0$$

So not LU decomposable.

Check (d):

$$\begin{bmatrix} 1 & -3 & 7 \\ -2 & 6 & 1 \\ 0 & 3 & -2 \end{bmatrix} \text{ here } |A_1| = 1 \neq 0 \text{ but}$$

$$|A_2| = \begin{vmatrix} 1 & -3 \\ -2 & 6 \end{vmatrix} = |6 - 6| = 0$$

So not LU decomposable.

C

$$\begin{bmatrix} 0 & 1 \\ 3 & 2 \end{bmatrix}$$

D

$$\begin{bmatrix} 1 & -3 & 7 \\ -2 & 6 & 1 \\ 0 & 3 & -2 \end{bmatrix}$$

QUESTION ANALYTICS

Q. 12

Consider the following table with data recorded over a month with 30 days:

Weather		
Mood	Sunny	Not sunny
	12	9
Not Good	4	5

If Rahul recorded on each day, whether it was sunny or not sunny and whether Rahul's mood was good or not good. If given day is sunny, then what is the probability that on given day Rahul's mood is good?

Have any Doubt ?

A

$$\frac{1}{4}$$

B

$$\frac{3}{4}$$

Your answer is Correct



Ashima Garg

Course: GATE

Computer Science Engineering(CS)

- HOME
- MY TEST
- BOOKMARKS
- MY PROFILE
- REPORTS
- BUY PACKAGE
- ASK AN EXPERT
- OFFER
- EXCLUSIVE OFFER FOR OTS STUDENTS ONLY ON BOOK PACKAGES

P(S) represent given day is sunny.

So,

$$P(G | S) = \frac{P(G \cap S)}{P(S)}$$
$$P(G \cap S) = \frac{12}{30}$$
$$P(S) = \frac{16}{30}$$
$$P(G | S) = \frac{\frac{12}{30}}{\frac{16}{30}} = \frac{12}{16}$$
$$= \frac{3}{4}$$

Mood

Good	Sunny	Not sunny	
	12	9	21
Not Good	4	5	9
	16	14	30

C

$\frac{5}{16}$

D

$\frac{16}{30}$

QUESTION ANALYTICS

Q. 13

The value of the integral given below is:

$$\int_{\pi/6}^{\pi/3} \frac{\operatorname{cosec}^2 x}{\cot^2 x} dx$$

Have any Doubt ?

A

$\frac{2}{3}$

B

$\frac{2}{\sqrt{3}}$

Your answer is **Correct**

Solution :

(b)

Consider,

$u = \cot x$

$\frac{du}{dx} = -\operatorname{cosec}^2 x$

$du = -\operatorname{cosec}^2 x \, dx$

$-du = \operatorname{cosec}^2 x \, dx$

Now new limits:

$x = \frac{\pi}{6} \rightarrow u = \cot \frac{\pi}{6} = \sqrt{3}$

$x = \frac{\pi}{3} \rightarrow u = \cot \frac{\pi}{3} = \frac{1}{\sqrt{3}}$

Substitute new limits and $\operatorname{cosec}^2 x \, dx$

$$\int_{\sqrt{3}}^{1/\sqrt{3}} \frac{-du}{u^2} = \left[\frac{u^{-2+1}}{-2+1} \right]_{\sqrt{3}}^{1/\sqrt{3}}$$
$$= \left[u^{-1} \right]_{\sqrt{3}}^{1/\sqrt{3}}$$


Ashima Garg

Course: GATE

Computer Science Engineering(CS)

HOME

MY TEST

BOOKMARKS

MY PROFILE

REPORTS

BUY PACKAGE

ASK AN EXPERT

OFFER

 EXCLUSIVE OFFER FOR OTS
STUDENTS ONLY ON BOOK
PACKAGES

 $\sqrt{3}$

C

 $\frac{3}{2}$

D

 $2\sqrt{3}$

QUESTION ANALYTICS

Q. 14

If the determinant of matrix:

$$A = \begin{bmatrix} 0 & 4 & 2 & 1 \\ 3 & -1 & 0 & 2 \\ 5 & 2 & x & 4 \\ 6 & 1 & -1 & 0 \end{bmatrix}$$

is 245, then which of the following represents the value of 'x'?

Have any Doubt ?

A

-6

B

4

C

-4

D

6

 Your answer is **Correct**
Solution :

(d)

$$A = \begin{bmatrix} 0 & 4 & 2 & 1 \\ 3 & -1 & 0 & 2 \\ 5 & 2 & x & 4 \\ 6 & 1 & -1 & 0 \end{bmatrix} = 245$$

$$\Rightarrow 5 \begin{bmatrix} 4 & 2 & 1 \\ -1 & 0 & 2 \\ 1 & -1 & 0 \end{bmatrix} - 2 \begin{bmatrix} 0 & 2 & 1 \\ 3 & 0 & 2 \\ 6 & -1 & 0 \end{bmatrix} + x \begin{bmatrix} 0 & 4 & 1 \\ 3 & -1 & 2 \\ 6 & 1 & 0 \end{bmatrix} - 4 \begin{bmatrix} 0 & 4 & 2 \\ 3 & -1 & 0 \\ 6 & 1 & -1 \end{bmatrix} = 245$$

$$\Rightarrow 5[1[4] + 1[8 + 1]] - 2[6[4] + 1[-3]] + x[6[8 + 1] - 1[-3]] - 4[-4[-3] + 2[3 + 6]] = 245$$

$$\Rightarrow 5[4 + 9] - 2[24 - 3] + x[54 + 3] - 4[12 + 18] = 245$$

$$\Rightarrow 65 - 42 + 57x - 120 = 245$$

$$57x = 245 + 120 + 42 - 65$$

$$57x = 342$$

$$x = 6$$

Alternate method:

For a shorter method, kindly refer to the video solution corresponding to this question.

QUESTION ANALYTICS

Q. 15



Ashima Garg

Course: GATE

Computer Science Engineering(CS)

HOME

MY TEST

BOOKMARKS

MY PROFILE

REPORTS

BUY PACKAGE

ASK AN EXPERT

OFFER

 EXCLUSIVE OFFER FOR OTS
STUDENTS ONLY ON BOOK
PACKAGES

[Have any Doubt ?](#)

5832

Correct Option

 Solution :
 5832

$$\begin{aligned}
 &\Rightarrow \lim_{x \rightarrow 0} \left[2 + \left(\frac{\log \cos x}{\log \cos(x/2)} \right)^2 \right]^3 \\
 &\Rightarrow \left[2 + \left(\lim_{x \rightarrow 0} \frac{\log \cos x}{\log \cos(x/2)} \right)^2 \right]^3 \\
 &\Rightarrow \text{Apply L'Hospital rule since } \cos 0 = 1 \text{ and } \log(1) = 0 \text{ which form indeterminant form i.} \\
 &\Rightarrow \left[2 + \left(\lim_{x \rightarrow 0} \frac{\frac{\sin x}{\cos x}}{\frac{1}{2} \frac{\sin(x/2)}{\cos(x/2)}} \right)^2 \right]^3 \\
 &\Rightarrow \left[2 + \left(2 \times \lim_{x \rightarrow 0} \frac{\tan x}{\tan(x/2)} \right)^2 \right]^3 \\
 &\Rightarrow \text{Apply L'Hospital rule again make 0/0 form.} \\
 &\Rightarrow \left[2 + \left(4 \times \lim_{x \rightarrow 0} \frac{\sec^2 x}{\sec^2(x/2)} \right)^2 \right]^3 \\
 &\Rightarrow \left[2 + \left(4 \times \frac{\sec^2 0}{\sec^2 0} \right)^2 \right]^3 \\
 &\Rightarrow \left[2 + \left(4 \times \frac{1}{1} \right)^2 \right]^3 \\
 &\Rightarrow [2 + 4^2]^3 \\
 &\Rightarrow [2 + 16]^3 = [18]^3 \\
 &\Rightarrow 5832
 \end{aligned}$$

Your Answer is 216

QUESTION ANALYTICS

Q. 16

Consider Kuldeep purchase a product of company X. The manual on it states that the lifetime T of product is defined as the amount of time (in years) the product works properly until it breaks down, satisfy following equation:

$$P(T \geq t) = e^{-t/4}, \text{ for all } t \geq 0$$

The probability that it breaks down in 3rd year is _____. (Upto 2 decimal places)

[FAQ](#) [Have any Doubt ?](#)

0.13 (0.11 - 0.16)

Correct Option

Solution :

0.13 (0.11 - 0.16)

Consider 'A' be an event that product break down in 3rd year and

So,

$$\begin{aligned}
 P(B) &= P(T \geq 2) \\
 &= e^{-2/4}
 \end{aligned}$$

$$P(A) = P(2 < T < 3)$$


Ashima Garg

Course: GATE

Computer Science Engineering(CS)

HOME

MY TEST

BOOKMARKS

MY PROFILE

REPORTS

BUY PACKAGE

ASK AN EXPERT

OFFER

 EXCLUSIVE OFFER FOR OTS
STUDENTS ONLY ON BOOK
PACKAGES

 $= 0.134 \text{ (approx.)}$
Your Answer is .88

QUESTION ANALYTICS

Q. 17


Consider there are 3 true coins and 1 false coin with tail on both sides. A coin is chosen at random and tosses 4 times. If tail occurs all the 4 times, then the probability that false coin is chosen is _____. (Upto 2 decimal places)

[Have any Doubt ?](#)

0.84 (0.84 - 0.85)

Correct Option
Solution :

0.84 (0.84 - 0.85)

According to Bayes theorem:


$$\text{So, probability of obtaining tail} = \frac{1}{4} \times 1 + \frac{3}{4} \times \frac{1}{2} \times \frac{1}{2} \times \frac{1}{2} \times \frac{1}{2}$$

$$= \frac{1}{4} + \frac{3}{4} \times \frac{1}{16}$$

$$= \frac{1}{4} + \frac{3}{64}$$

$$= \frac{16+3}{64} = \frac{19}{64}$$

$$\text{So, P(False coin/Tail on 4 tosses)} = \frac{\frac{1}{4} \times 1}{\frac{19}{64}} = \frac{\frac{1}{4}}{\frac{19}{64}}$$

$$= \frac{64}{19 \times 4} = \frac{16}{19}$$

$$= 0.842$$

Your Answer is .02

QUESTION ANALYTICS