



Ashima Garg

 Course: GATE
 Computer Science Engineering(CS)

HOME

MY TEST

BOOKMARKS

MY PROFILE

REPORTS

BUY PACKAGE

ASK AN EXPERT

OFFER

 EXCLUSIVE OFFER FOR OTS
 STUDENTS ONLY ON BOOK
 PACKAGES

SINGLE SUBJECT : ENGINEERING MATHEMATICS (GATE - 2019) - REPORTS

[OVERALL ANALYSIS](#)
[COMPARISON REPORT](#)
[SOLUTION REPORT](#)

ALL(33) CORRECT(17) INCORRECT(6) SKIPPED(10)

Q. 1

 At the point $x = 2$, the function:

$$f(x) = \begin{cases} x^3 - 8 & 2 < x < \infty \\ x - 2 & -\infty < x \leq 2 \end{cases} \text{ is}$$

[Solution Video](#) | [Have any Doubt ?](#) |

A

continuous and differentiable

B

continuous and not differentiable

Your answer is **Correct****Solution :**

(b)

$$\lim_{x \rightarrow 2^-} f(x) = \lim_{x \rightarrow 2^-} (x - 2) = (2 - 2) = 0$$

$$\lim_{x \rightarrow 2^+} f(x) = \lim_{x \rightarrow 2^+} (x^3 - 8) = (2^3 - 8) = 0$$

Also $f(2) = 2 - 2 = 0$

Thus $\lim_{x \rightarrow 2^-} f(x) = \lim_{x \rightarrow 2^+} f(x) = f(2)$

 $\therefore f$ is continuous at $x = 2$

$$f'(x) = \begin{cases} 3x^2 & 2 < x < \infty \\ 1 & -\infty < x \leq 2 \end{cases}$$

and $Lf'(2) = 1$ and $Rf'(2) = 12$

 $\therefore f$ is not differentiable at $x = 2$.

C

discontinuous and differentiable

D

discontinuous and not differentiable

QUESTION ANALYTICS

Q. 2

 Given that the determinant of the matrix $\begin{bmatrix} 2 & 6 & 0 \\ 4 & 12 & 8 \\ -2 & 0 & 4 \end{bmatrix}$ is -96, the determinant of the

 matrix $\begin{bmatrix} 4 & 12 & 0 \\ 8 & 24 & 16 \\ -4 & 0 & 8 \end{bmatrix}$ is

[Solution Video](#) | [Have any Doubt ?](#) |

A

192



Ashima Garg

 Course: GATE
 Computer Science Engineering(CS)

- HOME
- MY TEST
- BOOKMARKS
- MY PROFILE
- REPORTS
- BUY PACKAGE
- ASK AN EXPERT
- OFFER

EXCLUSIVE OFFER FOR OTS
 STUDENTS ONLY ON BOOK
 PACKAGES

 C
 -384

 D
 -768

Your answer is Correct

Solution :
 (d)
 $D = -96$ for the given matrix

$$|A| = \begin{vmatrix} 4 & 12 & 0 \\ 8 & 24 & 16 \\ -4 & 0 & 8 \end{vmatrix} = 2^3 \begin{vmatrix} 2 & 6 & 0 \\ 4 & 12 & 8 \\ -2 & 0 & 4 \end{vmatrix}$$

(Taking 2 common from each row)

$$\therefore \text{Det}(A) = (2)^3 \times D \\ = 8 \times (-96) \\ = -768$$

QUESTION ANALYTICS

Q. 3
 The value of $\lim_{x \rightarrow 4} \frac{(2x)^{1/3} - 2}{2x - 8}$ is

[Solution Video](#) | [Have any Doubt ?](#) |

A

$$\frac{1}{3}$$

B

$$\frac{1}{6}$$

C

$$\frac{1}{24}$$

D

$$\frac{1}{12}$$

Your answer is Correct

Solution :
 (d)

$$\lim_{x \rightarrow 4} \frac{(2x)^{1/3} - 2}{2x - 8}$$

 Above form is $\left(\frac{0}{0}\right)$ by putting the value $x = 4$

Applying L' Hospital rule

$$\begin{aligned} &= \lim_{x \rightarrow 4} \frac{\frac{1}{3}(2x)^{\left(\frac{1}{3}-1\right)} \times 2}{2} \\ &= \lim_{x \rightarrow 4} \frac{1}{3} (2x)^{-\frac{2}{3}} \\ &= \frac{1}{3} (8)^{-2/3} = \frac{1}{12} \end{aligned}$$



Ashima Garg

 Course: GATE
 Computer Science Engineering(CS)

HOME

MY TEST

BOOKMARKS

MY PROFILE

REPORTS

BUY PACKAGE

ASK AN EXPERT

OFFER

 EXCLUSIVE OFFER FOR OTS
 STUDENTS ONLY ON BOOK
 PACKAGES
Q. 4

Which one of the following functions is strictly bounded?

[Solution Video](#) [Have any Doubt ?](#)

A

$$\frac{3x^2}{2}$$





B

$$2e^{-x^2}$$

Your answer is Correct

Solution :

(b)

From the graphs below, we can see that only $2e^{-x^2}$ is strictly bounded.

C

$$e^{3x}$$

D

$$x^3$$

QUESTION ANALYTICS

Q. 5
 A continuous random variable X has a probability density function $f(x) = e^{-2x}$, $0 < x < \infty$. Then $P[X > 1]$ is

[Solution Video](#) [Have any Doubt ?](#)

A

0.270

B

0.067

Your answer is Correct



Ashima Garg

Course: GATE
Computer Science Engineering(CS)

HOME

MY TEST

BOOKMARKS

MY PROFILE

REPORTS

BUY PACKAGE

ASK AN EXPERT

OFFER

EXCLUSIVE OFFER FOR OTS
STUDENTS ONLY ON BOOK
PACKAGES

$$\begin{aligned}
 P[X > 1] &= \int_1^{\infty} e^{-ux} dx = \left[e^{-ux} \right]_1^{\infty} \\
 &= -\left(\frac{e^{-2\infty}}{2} - \frac{e^{-2}}{2} \right) \quad [e^{-\infty} = 0] \\
 &= \frac{e^{-2}}{2} = 0.067
 \end{aligned}$$

C
0.034D
0.135

QUESTION ANALYTICS

Q. 6

What are the eigen values of matrix given below:

$$A = \begin{bmatrix} 0 & 2 & 0 \\ 0 & 0 & 2 \\ 0 & -6 & -8 \end{bmatrix}_{3 \times 3}$$

Solution Video | Have any Doubt ?

A
(0, -2, 6)B
(0, 2, 6)C
(0, -2, -6)

Your answer is Correct

Solution :
(c)

$$A = \begin{bmatrix} 0 & 2 & 0 \\ 0 & 0 & 2 \\ 0 & -6 & -8 \end{bmatrix}$$

$$|A - \lambda I| = 0$$

$$\begin{vmatrix} -\lambda & 2 & 0 \\ 0 & -\lambda & 2 \\ 0 & -6 & -8 - \lambda \end{vmatrix} = 0$$

$$-\lambda(8\lambda + \lambda^2 + 12) + 2(0 - 0) = 0$$

$$\text{Either } \lambda = 0 \text{ or } (\lambda^2 + 8\lambda + 12) = 0$$

$$\lambda = 0 \text{ or } (\lambda + 2)(\lambda + 6) = 0$$

$$\text{Therefore } \lambda = 0, -2, -6$$

So option (c) is correct answer.

D
(0, 1, 3)

QUESTION ANALYTICS

Q. 7

A diagnostic test has a probability 95% of giving a positive result when applied to a person suffering from a certain disease and probability 10% of giving a positive when applied to a nonsufferer. It is estimated that 5%



Ashima Garg

 Course: GATE
 Computer Science Engineering(CS)

HOME

MY TEST

BOOKMARKS

MY PROFILE

REPORTS

BUY PACKAGE

ASK AN EXPERT


OFFER

 EXCLUSIVE OFFER FOR OTS
 STUDENTS ONLY ON BOOK
 PACKAGES

 A
 0.045

 B
 0.333

Correct Option

Solution :
 (b)


$$\begin{aligned}
 P\left(\frac{\text{Sufferer}}{\text{Positive}}\right) &= \frac{P(\text{Sufferer}) \times P(\text{Positive})}{P(\text{Sufferer}) \times P(\text{Positive}) + P(\text{Non-sufferer}) \times P(\text{Positive})} \\
 &= \frac{0.95 \times 0.05}{(0.95 \times 0.05) + (0.95 \times 0.1)} \\
 &= \frac{0.0475}{0.0475 + 0.095} = \frac{0.0475}{0.1425} = 0.333
 \end{aligned}$$

 C
 0.099

 D
 0.435

QUESTION ANALYTICS

Q. 8
 What is the result of $\int_0^1 \log(1+x) dx$?

A

$$\log\left(\frac{2}{e}\right)$$

B

$$\log\left(\frac{4}{e}\right)$$

Your answer is Correct

Solution :
 (b)

$$\int_0^1 \log(1+x) dx$$

Using integration by parts:

$$\begin{aligned}
 &= \int_0^1 \log(1+x) \cdot 1 dx \\
 &= \left| \log(1+x) \cdot x \right|_0^1 - \int_0^1 \frac{1}{1+x} \cdot x dx
 \end{aligned}$$



Ashima Garg

 Course: GATE
 Computer Science Engineering(CS)

HOME

MY TEST

BOOKMARKS

MY PROFILE

REPORTS

BUY PACKAGE

ASK AN EXPERT

OFFER

 EXCLUSIVE OFFER FOR OTS
 STUDENTS ONLY ON BOOK
 PACKAGES

$$\begin{aligned}
 &= \log(2) - \int_0^1 \frac{ax}{1+x} dx \\
 &= \log(2) - \int_0^1 1 dx + \int_0^1 \frac{dx}{1+x} \\
 &= \log(2) - \left| x \right|_0^1 + \left| \log(1+x) \right|_0^1 \\
 &= \log(2) - 1 + \log(2) \\
 &= \log(2^2) - 1 \\
 &= \log(2^2) - \log_e(e) \\
 &= \log\left(\frac{4}{e}\right)
 \end{aligned}$$

C

$$\log\left(\frac{e}{2}\right)$$

D

$$\log\left(\frac{e}{4}\right)$$

QUESTION ANALYTICS

Q. 9

Consider Vamshi decides to toss a fair coin repeatedly until he gets a tail. He makes atmost 4 tosses. The value of variance (T) is _____
 (variable T denotes the number of tosses).

[Solution Video](#) | [Have any Doubt ?](#)

A

$$\frac{15}{16}$$

B

$$\frac{1}{16}$$

C

$$\frac{252}{256}$$

Correct Option

Solution :
 (c)

x	1	2	3	4
$P(x)$	$\frac{1}{2}$	$\frac{1}{4}$	$\frac{1}{8}$	$\frac{1}{16}$

$$V(x) = E(x^2) - (E(x))^2$$

$$E(x^2) = 1^2 \times \frac{1}{2} + 2^2 \times \frac{1}{4} + 3^2 \times \frac{1}{8} + 4^2 \times \frac{1}{16} = \frac{58}{16}$$

$$E(x) = 1 \times \frac{1}{2} + 2 \times \frac{1}{4} + 3 \times \frac{1}{8} + 4 \times \frac{1}{16} = \frac{26}{16}$$

$$V(x) = E(x^2) - (E(x))^2$$

$$= \frac{58}{16} - \left(\frac{26}{16} \right)^2 = \frac{252}{256}$$

D



Ashima Garg

 Course: GATE
 Computer Science Engineering(CS)

HOME

MY TEST

BOOKMARKS

MY PROFILE

REPORTS

BUY PACKAGE

ASK AN EXPERT

OFFER

 EXCLUSIVE OFFER FOR OTS
 STUDENTS ONLY ON BOOK
 PACKAGES

QUESTION ANALYTICS

Q. 10

If A is 3×3 matrix and Trace A = 9, $|A| = 24$ and one of the eigen values is 3, then sum of other eigen values is _____.

[Solution Video](#) | [Have any Doubt ?](#) |

A

5

B

8

C

6

Your answer is Correct

Solution :

(c)

Given,

$$\text{Trace } A = 9$$

$$|A| = 24$$

$$\lambda_1 = 3$$

$$\lambda_1 + \lambda_2 + \lambda_3 = 9$$

$$3 + \lambda_2 + \lambda_3 = 9$$

$$\Rightarrow \lambda_2 + \lambda_3 = 6$$

D

9

QUESTION ANALYTICS

Q. 11

If the sum of diagonal elements of a 2×2 symmetric matrix is -8, then the maximum possible value of determinant of the matrix is _____.

[Solution Video](#) | [Have any Doubt ?](#) |

16

Your answer is Correct 16

Solution :

16

Consider a symmetric matrix $A = \begin{bmatrix} a & b \\ b & d \end{bmatrix}$

$$\text{Given } a + d = -8$$

$$|A| = ad - b^2$$

Now since b^2 is always non-negative, maximum determinant will come when $b^2 = 0$.

So we need to maximize

$$\begin{aligned} |A| &= ad \\ &= ad = a \times (-8 + a) = -a^2 - 8a \end{aligned}$$

$$\frac{d|A|}{da} = -2a - 8 = 0$$

$\Rightarrow a = -4$ is the only stationary point

Since, $\left[\frac{d^2|A|}{da^2} \right]_{a=-4} = -2 < 0$, we have a maximum at $a = -4$

Since, $a + d = -8$, $d = -4$. Now maximum value of determinant is $|A| = 16$



Ashima Garg

 Course: GATE
 Computer Science Engineering(CS)

HOME

MY TEST

BOOKMARKS

MY PROFILE

REPORTS

BUY PACKAGE

ASK AN EXPERT

OFFER

 EXCLUSIVE OFFER FOR OTS
 STUDENTS ONLY ON BOOK
 PACKAGES
Q. 12

The probability density function of random variable X is

$$f_x(x) = \begin{cases} \left(\frac{1}{2}\right)e^{-x/2} & n \geq 0 \\ 0 & \text{otherwise} \end{cases}$$

 The expectation $E(x)$ is _____.

[Solution Video](#) | [Have any Doubt ?](#) |

2

Correct Option

Solution :

(2)

$$\begin{aligned} E(x) &= \int_0^{\infty} \left(\frac{1}{2}\right) xe^{-x/2} dx \\ &= \frac{1}{2} \left[\int_0^{\infty} xe^{-x/2} dx \right] \\ &= \frac{1}{2} \left\{ \left[\frac{xe^{-x/2}}{-\frac{1}{2}} \right]_0^{\infty} - \left[\int_0^{\infty} \frac{e^{-x/2}}{-\frac{1}{2}} dx \right] \right\} \\ &= \frac{1}{2} \left\{ - \left[\frac{e^{-x/2}}{-\frac{1}{2} \times -\frac{1}{2}} \right]_0^{\infty} \right\} \\ &= \frac{1}{2} \left[-0 + \frac{1}{\frac{1}{2} \times \frac{1}{2}} \right] = 2 \end{aligned}$$

Your Answer is .735

QUESTION ANALYTICS

Q. 13
 Let A and B be 3×3 matrices such that $A' = -A$ and $B' = B$. Then matrix $3AB + \lambda BA$ is a skew symmetric matrix for λ equal to _____.

[FAQ](#) | [Solution Video](#) | [Have any Doubt ?](#) |

3

Correct Option

Solution :

3

 Since it is skew symmetric, $(3AB + \lambda BA)' = -(3AB + \lambda BA)$

$$\begin{aligned} LHS &= 3(AB)' + \lambda(BA)' \\ &= 3B'A' + \lambda A'B' \end{aligned}$$

Now

 $B' = B$ and $A' = -A$

$$= -3BA - \lambda AB = -(\lambda AB + 3BA)$$

$$\begin{aligned} RHS &= -(3AB + \lambda BA) \\ &= 3BA + \lambda AB = 3AB + \lambda BA \end{aligned}$$

$$\begin{aligned} (\lambda - 3)AB &= (\lambda - 3)BA \\ \lambda &= 3 \end{aligned}$$

QUESTION ANALYTICS



Ashima Garg

 Course: GATE
 Computer Science Engineering(CS)

- HOME
- MY TEST
- BOOKMARKS
- MY PROFILE
- REPORTS
- BUY PACKAGE
- ASK AN EXPERT
- OFFER
- EXCLUSIVE OFFER FOR OTS STUDENTS ONLY ON BOOK PACKAGES

Let $A_n = 1 + 2 + 3 + \dots + n$ and $T_n = \frac{A_2}{A_2 - 1} \times \frac{A_3}{A_3 - 1} \times \frac{A_4}{A_4 - 1} \dots \frac{A_n}{A_n - 1}$, where $n \in N (n \geq 2)$.

Then the value of $\lim_{n \rightarrow \infty} T_n$ is _____.

[Solution Video](#) | [Have any Doubt ?](#) |

3

Correct Option

Solution :

3

$$\begin{aligned}
 A_n &= \frac{n(n+1)}{2} \\
 A_n - 1 &= \frac{n(n+1)}{2} - 1 = \frac{n^2 + n - 2}{2} \\
 &= \frac{(n+2)(n-1)}{2} \\
 \frac{A_n}{A_n - 1} &= \frac{n(n+1)}{(n+2)(n-1)} = \left(\frac{n}{n-1} \right) \left(\frac{n+1}{n+2} \right) \\
 T_n &= \left(\frac{2}{1} \times \frac{3}{2} \times \frac{4}{3} \dots \frac{n}{n-1} \right) \left(\frac{3}{4} \times \frac{4}{5} \dots \frac{n+1}{n+2} \right) \\
 \frac{n}{1} \times \frac{3}{n+2} &= \frac{3n}{n+2} \\
 \lim_{n \rightarrow \infty} T_n &= \lim_{n \rightarrow \infty} \frac{3n}{n+2} = 3
 \end{aligned}$$

QUESTION ANALYTICS

Q. 15

A fair coin is tossed till a head appears for the first time. The probability that the number of required tosses is odd will be _____. (Upto 2 decimal places)

[Solution Video](#) | [Have any Doubt ?](#) |

0.67 [0.65 - 0.68]

Your answer is **Correct.66****Solution :**

0.67 [0.65 - 0.68]

 $P(E) = \text{Probability of head appearing in odd number of tosses}$
 $= P(H) + P(TTH) + \dots$

$$\begin{aligned}
 &= \frac{1}{2} + \left(\frac{1}{2} \right)^2 \left(\frac{1}{2} \right) + \dots \\
 &= \frac{1/2}{1 - \frac{1}{4}} = \frac{2}{3} = 0.67
 \end{aligned}$$

QUESTION ANALYTICS

Q. 16

Let M be a matrix:

$$\begin{bmatrix}
 1 & 0 & 0 & 1 & 0 & 0 \\
 0 & 1 & 0 & 0 & 1 & 0 \\
 0 & 0 & 1 & 0 & 0 & 1 \\
 0 & 0 & 0 & 1 & 0 & 0 \\
 0 & 0 & 0 & 0 & 1 & 0 \\
 0 & 0 & 0 & 0 & 0 & 1
 \end{bmatrix}$$



Ashima Garg

Course: GATE
Computer Science Engineering(CS)

HOME

MY TEST

BOOKMARKS

MY PROFILE

REPORTS

BUY PACKAGE

ASK AN EXPERT

OFFER

 EXCLUSIVE OFFER FOR OTS
STUDENTS ONLY ON BOOK
PACKAGES

60

Correct Option

Solution :

60

$$M = \begin{bmatrix} I & I \\ 0 & I \end{bmatrix}$$

$$M^2 = \begin{bmatrix} I & I \\ 0 & I \end{bmatrix} \begin{bmatrix} I & I \\ 0 & I \end{bmatrix}$$

$$= \begin{bmatrix} I^2 + 0 \times I & I^2 + I^2 \\ 0 \times I + I \times 0 & 0 \times I + I \times I \end{bmatrix}$$

$$= \begin{bmatrix} I^2 & 2I^2 \\ 0 & I^2 \end{bmatrix} \text{ since } [I^2 = I]$$

$$= \begin{bmatrix} I & 2I \\ 0 & I \end{bmatrix}$$

$$M^3 = \begin{bmatrix} I & 2I \\ 0 & I \end{bmatrix} \begin{bmatrix} I & I \\ 0 & I \end{bmatrix} = \begin{bmatrix} I^2 & 2I^2 + I^2 \\ 0 & I \times I + I \times 0 \end{bmatrix}$$

$$= \begin{bmatrix} I^2 & 3I^2 \\ 0 & I^2 \end{bmatrix}$$

$$M^3 = \begin{bmatrix} I & 3I \\ 0 & I \end{bmatrix}$$

$$M^K = \begin{bmatrix} I & KI \\ 0 & I \end{bmatrix}$$

$$M^{18} = \begin{bmatrix} I & 18I \\ 0 & I \end{bmatrix}$$

$$= 3 \times 1 + 3 \times 1 + 18 \times 3 \times 1 \\ = 6 + 54 = 60$$

Your Answer is 9

QUESTION ANALYTICS

Q. 17What is the value of $\lim_{x \rightarrow 0} \left(\frac{a^x + b^x + c^x}{3} \right)^{1/x}$?

Solution Video | Have any Doubt ?

A

abc

Your answer is Wrong

B

 $\sqrt[3]{abc}$

C

 $\sqrt[3]{abc}$

Correct Option

Solution :

(c)

$$\lim_{x \rightarrow 0} \left(\frac{a^x + b^x + c^x}{3} \right)^{1/x}$$



Ashima Garg

 Course: GATE
 Computer Science Engineering(CS)

HOME

MY TEST

BOOKMARKS

MY PROFILE

REPORTS

BUY PACKAGE

ASK AN EXPERT

OFFER

 EXCLUSIVE OFFER FOR OTS
 STUDENTS ONLY ON BOOK
 PACKAGES

$$\begin{aligned}
 &= \lim_{x \rightarrow 0} \left(1 + \frac{a^x + b^x + c^x - 3}{3} \right)^{1/x} \\
 &= \lim_{x \rightarrow 0} \left(1 + \frac{(a^x - 1) + (b^x - 1) + (c^x - 1)}{3} \right)^{1/x}
 \end{aligned}$$

We know that:

$$\begin{aligned}
 \lim_{x \rightarrow 0} (1 + \lambda x)^{1/x} &= e^\lambda \\
 &= e^{\lim_{x \rightarrow 0} \frac{(a^x - 1)}{3x} + \frac{(b^x - 1)}{3x} + \frac{(c^x - 1)}{3x}} \\
 &= e^{\lim_{x \rightarrow 0} \frac{1}{3} \left(\frac{a^x - 1}{x} + \frac{b^x - 1}{x} + \frac{c^x - 1}{x} \right)} \\
 &= e^{1/3 (\log a + \log b + \log c)} \quad \left[\because \lim_{x \rightarrow 0} \frac{a^x - 1}{x} = \log a \right] \\
 &= e^{1/3 \log (abc)} = e^{\log(abc)^{1/3}} = (abc)^{1/3} \\
 &= \sqrt[3]{abc}
 \end{aligned}$$

D

 $(abc)^3$

QUESTION ANALYTICS

Q. 18

An artillery target may be either at point 1 with probability $\frac{8}{9}$ or at point 2 with probability $\frac{1}{9}$. We have 21 shells, each of which can be fired at point 1 or point 2. Each shell may hit the target, independently of other shells, with probability $\frac{1}{2}$. If 12 shells are fired at point 1 and 9 shells are fired at point 2, what is the probability that the target is hit?

[Solution Video](#) [Have any Doubt ?](#)

A

$$\frac{8}{9}2^{12} + \frac{1}{9}2^9$$

B


$$\frac{8}{9} \left(\frac{1}{2^{12}} \right) + \frac{1}{9} \left(\frac{1}{2^9} \right)$$

Your answer is Wrong

C

$$\frac{8}{9} \left(1 - \frac{1}{2^{12}} \right) + \frac{1}{9} \left(1 - \frac{1}{2^9} \right)$$

Correct Option

Solution :
 (c)




Ashima Garg

 Course: GATE
 Computer Science Engineering(CS)

HOME

MY TEST

BOOKMARKS

MY PROFILE

REPORTS

BUY PACKAGE

ASK AN EXPERT

OFFER

 EXCLUSIVE OFFER FOR OTS
 STUDENTS ONLY ON BOOK
 PACKAGES

- D
None of these

QUESTION ANALYTICS

Q. 19

 Which of the following is result of $\int_0^{\pi/3} \sin^3 \theta d\theta$?

 [Solution Video](#) | [Have any Doubt ?](#) |

A

0

B

1

C

 $\frac{1}{3}$

D

 $\frac{5}{24}$

Correct Option

Solution :

(d)

$$I = \int_0^{\pi/3} (1 - \cos^2 \theta) \sin \theta d\theta$$

Let,

$$\cos \theta = t$$

$$-\sin \theta \frac{d\theta}{dt} = 1$$

$$-\sin \theta d\theta = dt$$

$$\text{at } \theta = 0, \cos 0 = 1$$

$$\text{at } \theta = \frac{\pi}{3}, \cos \frac{\pi}{3} = \frac{1}{2}$$

$$I = \int_1^{1/2} -(1-t^2) dt$$

$$= -\left[t - \frac{t^3}{3} \right]_1^{1/2}$$

$$= -\left(\frac{11}{24} \right) - \left(-\frac{2}{3} \right)$$

$$= \frac{5}{24}$$

QUESTION ANALYTICS

Q. 20

Cayley Hamilton theorem states that a square matrix satisfies its own characteristic equation. Consider a

 matrix, $A = \begin{bmatrix} 3 & 1 \\ -2 & 0 \end{bmatrix}$.



Ashima Garg

 Course: GATE
 Computer Science Engineering(CS)

HOME

MY TEST

BOOKMARKS

MY PROFILE

REPORTS

BUY PACKAGE

ASK AN EXPERT

OFFER

 EXCLUSIVE OFFER FOR OTS
 STUDENTS ONLY ON BOOK
 PACKAGES

A $A + 3I + 2A^{-1} = 0$

B

(A - I) (A + 2I) = 0

C

A - 3I - 2A^{-1} = 0

D

A - 3I + 2A^{-1} = 0

Your answer is Correct

Solution :

(d)

$$\begin{aligned} |A - \lambda I| &= 0 \\ \begin{bmatrix} 3 & 1 \\ -2 & 0 \end{bmatrix} - \begin{bmatrix} \lambda & 0 \\ 0 & \lambda \end{bmatrix} &= 0 \\ \begin{vmatrix} 3-\lambda & 1 \\ -2 & -\lambda \end{vmatrix} &= 0 \\ -3\lambda + \lambda^2 + 2 &= 0 \\ \lambda^2 - 3\lambda + 2 &= 0 \\ A^2 - 3A + 2 &= 0 \\ A - 3I + 2A^{-1} &= 0 \quad (\times \text{ by } A^{-1} \text{ on both sides}) \end{aligned}$$

QUESTION ANALYTICS

Q. 21

A matrix has eigen values -6 and -3 , the corresponding eigen vectors are $\begin{bmatrix} 3 \\ -6 \end{bmatrix}$ and $\begin{bmatrix} 3 \\ -3 \end{bmatrix}$ respectively.

The matrix is

 [Solution Video](#) [Have any Doubt ?](#)

A

$$\begin{bmatrix} -3 & 0 \\ 0 & -6 \end{bmatrix}$$

Your answer is Wrong

B

$$\begin{bmatrix} 3 & 3 \\ -3 & -6 \end{bmatrix}$$

C

$$\begin{bmatrix} 0 & 3 \\ -6 & -9 \end{bmatrix}$$

Correct Option

Solution :

(c)

AX = λX

$$\begin{bmatrix} a & b \\ c & d \end{bmatrix} \begin{bmatrix} 3 \\ -6 \end{bmatrix} = (-6) \begin{bmatrix} 3 \\ -6 \end{bmatrix}$$

3a - 6b = -18

3c - 6d = 36

$$\begin{bmatrix} a & b \\ c & d \end{bmatrix} \begin{bmatrix} 3 \\ -3 \end{bmatrix} = (-3) \begin{bmatrix} 3 \\ -3 \end{bmatrix}$$



Ashima Garg

 Course: GATE
 Computer Science Engineering(CS)

HOME

MY TEST

BOOKMARKS

MY PROFILE

REPORTS

BUY PACKAGE

ASK AN EXPERT

OFFER

 EXCLUSIVE OFFER FOR OTS
 STUDENTS ONLY ON BOOK
 PACKAGES

 from equation (ii) and (iv), $c = -6$ and $a = -9$.

$$\therefore A = \begin{bmatrix} a & b \\ c & d \end{bmatrix} = \begin{bmatrix} 0 & 3 \\ -6 & -9 \end{bmatrix}$$

D

$$\begin{bmatrix} 3 & 6 \\ -6 & -12 \end{bmatrix}$$

QUESTION ANALYTICS

Q. 22

 A function $y = 7x^2 + 12x$ is defined over an open interval $x = (1, 3)$. At least at one point in this interval,

 $\frac{dy}{dx}$ is exactly

[Solution Video](#) | [Have any Doubt ?](#)

A

26

B

40

Your answer is Correct

Solution :

(b)

$$y = 7x^2 + 12x$$

Using Lagrange's mean value theorem:

 At $x = 1, y = 7 + 12 = 19$
 $x = 3, y = 63 + 36 = 99$

$$f'(x) = \frac{f(b) - f(a)}{b - a}$$

$$= \frac{99 - 19}{3 - 1} = 40$$

So option (b) is correct answer.

C

62

D

54

QUESTION ANALYTICS

Q. 23

 A random variable x has the following probability distribution.

x	0	1	2	3	4
$P(x)$	c	$2c$	$2c$	c^2	$5c^2$

The mean and variance of x is
[Solution Video](#) | [Have any Doubt ?](#)

A

1.638, 1.45

Your answer is Correct

Solution :

(a)



Ashima Garg

 Course: GATE
 Computer Science Engineering(CS)

HOME

MY TEST

BOOKMARKS

MY PROFILE

REPORTS

BUY PACKAGE

ASK AN EXPERT

OFFER

 EXCLUSIVE OFFER FOR OTS
STUDENTS ONLY ON BOOK
PACKAGES

$$6c^2 + 5c - 1 = 0$$

$$c = \frac{1}{6}, -1$$

 Since $P(x) \geq 0$, the possible value of

$$c = \frac{1}{6}$$

x	0	1	2	3	4
P(x)	$\frac{1}{6}$	$\frac{2}{6}$	$\frac{2}{6}$	$\frac{1}{36}$	$\frac{5}{36}$
xP(x)	0	$\frac{2}{6}$	$\frac{4}{6}$	$\frac{3}{36}$	$\frac{20}{36}$

$$\text{Mean} = \sum_{x=0}^4 xP(x) = 0 + \frac{2}{6} + \frac{4}{6} + \frac{3}{36} + \frac{20}{36}$$

$$= \frac{59}{36} = 1.638$$

$$\text{Variance} = \sigma^2 = E(x^2) - [E(x)]^2$$

$$= \left[0\left(\frac{1}{6}\right) + 1\left(\frac{2}{6}\right) + 4\left(\frac{2}{6}\right) + 9\left(\frac{1}{36}\right) + 16\left(\frac{5}{36}\right) - \left(\frac{59}{36}\right)^2 \right] = 1.45$$

B

1.638, 1.204

C

1.204, 1.45

D

1.45, 1.638

QUESTION ANALYTICS

Q. 24
 Consider function $f(x) = (x^2 - 4)^2$ where x is a real number. Which of the following is true about given function?

[Solution Video](#) | [Have any Doubt ?](#)

A

Has only one minima

B

Has only two minima

Your answer is Correct

Solution :

(b)

$$f(x) = (x^2 - 4)^2$$

$$f'(x) = 2(x^2 - 4) \times 2x$$

$$= 4x(x^2 - 4) = 0$$

 $x = 0, x = 2$ and $x = -2$ are the stationary points

$$f''(x) = 4[x(2x) + (x^2 - 4) \times 1]$$

$$= 4[(2x^2 + x^2 - 4) = 4[3x^2 - 4]$$

$$= 12x^2 - 16$$

$$f''(0) = -16 < 0 \quad (\text{So maxima at } x = 0)$$

$$f''(2) = (12)2^2 - 16 = 32 > 0 \quad (\text{So minima at } x = 2)$$

$$f''(-2) = 12(-2)^2 - 16 = 32 > 0 \quad (\text{So minima at } x = -2)$$

 \therefore There is only one maxima and only two minima for this function.

C

Has three minima

D

Has three maxima



Ashima Garg

 Course: GATE
 Computer Science Engineering(CS)

HOME

MY TEST

BOOKMARKS

MY PROFILE

REPORTS

BUY PACKAGE

ASK AN EXPERT

OFFER

 EXCLUSIVE OFFER FOR OTS
 STUDENTS ONLY ON BOOK
 PACKAGES

Consider the following linear equations of system :

$$x + y + z = 6$$

$$x + 2y + 5z = 10$$

$$2x + 3y + \lambda z = \mu$$

Which of the following is correct about system?

[Solution Video](#) | [Have any Doubt ?](#) |

A

System has unique solution for $\lambda \neq 6$

B

System has no solution for $\lambda = 6$ and $\mu \neq 16$

C

System has infinite solution for $\lambda = 6$ and $\mu = 16$

D

All of the above

Your answer is **Correct****Solution :**

(d)

$$AX = B$$

$$\begin{bmatrix} 1 & 1 & 1 \\ 1 & 2 & 5 \\ 2 & 3 & \lambda \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 6 \\ 10 \\ \mu \end{bmatrix}$$

$$C = (A, B) = \begin{bmatrix} 1 & 1 & 1 & : & 6 \\ 1 & 2 & 5 & : & 10 \\ 2 & 3 & \lambda & : & \mu \end{bmatrix}$$

After performing $R_2 \leftarrow R_2 - R_1$ and $R_3 \leftarrow R_3 - 2R_1$

$$C = \begin{bmatrix} 1 & 1 & 1 & : & 6 \\ 0 & 1 & 4 & : & 4 \\ 0 & 1 & \lambda - 2 & : & \mu - 12 \end{bmatrix}$$

After performing $R_3 \leftarrow R_3 - R_2$

$$C = \begin{bmatrix} 1 & 1 & 1 & : & 6 \\ 0 & 1 & 4 & : & 4 \\ 0 & 0 & \lambda - 6 & : & \mu - 16 \end{bmatrix}$$

Since $R(A) = R(C)$ for unique solutionSo $\lambda - 6 \neq 0, \lambda \neq 6$.For no solution $R(A) \neq R(C)$ then $R(A) = 2$ and $R(C) = 3$

$$\lambda - 6 = 0$$

$$\Rightarrow \lambda = 6 \text{ and } \mu - 16 \neq 0 \Rightarrow \mu \neq 16$$

For infinite solution $R(A) = R(C) = 2$

$$\text{then } \lambda - 6 = 0 \text{ and } \mu - 16 = 0$$

$$\lambda = 6 \text{ and } \mu = 16$$

So all of options are true.

QUESTION ANALYTICS

Q. 26Probability density function of a random variable X is distributed uniformly between 0 and 10. The probability that X lies between 2.5 to 7.5 and the mean square value of X are respectively
[Solution Video](#) | [Have any Doubt ?](#) |



Ashima Garg

 Course: GATE
 Computer Science Engineering(CS)

HOME

MY TEST

BOOKMARKS

MY PROFILE

REPORTS

BUY PACKAGE

ASK AN EXPERT

OFFER

 EXCLUSIVE OFFER FOR OTS
 STUDENTS ONLY ON BOOK
 PACKAGES

2 3

Correct Option

Solution :

(a)

In uniform distribution $[a, b]$

$$k = \frac{1}{b-a}$$

$$= \frac{1}{10-0} = \frac{1}{10}$$

$$P(2.5 \leq X \leq 7.5) = \int_{2.5}^{7.5} \frac{1}{10} dx = \frac{1}{10} x \Big|_{2.5}^{7.5} = \frac{1}{10}(7.5 - 2.5) = \frac{1}{2}$$

$$E(x^2) = \text{Mean square value} = \int_0^{10} x^2 f(x) dx$$

$$\int_0^{10} \frac{1}{10} x^2 dx = \frac{1}{10} \frac{x^3}{3} \Big|_0^{10} = \frac{10^3 - 0^3}{30} = \frac{1000}{30} = \frac{100}{3}$$

B

5 and 100

C

5 and $\frac{100}{3}$

D

 $\frac{1}{2}$ and 100

QUESTION ANALYTICS

Q. 27Assume A and B are matrix of size $n \times n$, which of the following is true?
[Solution Video](#) | [Have any Doubt ?](#) |

A

If A is invertible, then $ABA^{-1} = B$.

B

If A is an idempotent non-singular matrix, then A must be the identity matrix.

Correct Option

Solution :

(b)

- $ABA^{-1} = B$ given,
- $\Rightarrow AB = BA$ since matrix multiplication is not commutative. So false even if A is invertible.
- A is idempotent, so $A^2 = A$, since A is non-singular, so it is invertible i.e. A^{-1} exist.
 $I = A^{-1} \cdot A = A^{-1} \cdot A^2 = IA = A$
 So A must be identity matrix. So true.
- If coefficient matrix A is invertible for $Ax = b$ then $x = A^{-1}$ unique solution exist. So false
- If B is zero matrix, then also $AB = B =$ zero matrix. So false

C

If the coefficient matrix A of the system $Ax = b$ is invertible, then the system has infinitely many solution

D

If $AB = B$ then B is identity matrix.



Ashima Garg

 Course: GATE
 Computer Science Engineering(CS)

HOME

MY TEST

BOOKMARKS

MY PROFILE

REPORTS

BUY PACKAGE

ASK AN EXPERT

OFFER

 EXCLUSIVE OFFER FOR OTS
 STUDENTS ONLY ON BOOK
 PACKAGES
Q. 28Consider a matrix P :

$$P = \begin{bmatrix} 4 & -6 \\ -2 & 8 \end{bmatrix}$$

 If $P \times Q = \begin{bmatrix} 5 \\ 0 \end{bmatrix}$, then which of the following represent the matrix Q ?

[Solution Video](#) | [Have any Doubt ?](#)

A

$$\begin{bmatrix} 2 \\ 4 \end{bmatrix}$$

B

$$\begin{bmatrix} 4 \\ 2 \end{bmatrix}$$

C

$$\begin{bmatrix} 2 \\ 0.5 \end{bmatrix}$$

Correct Option

Solution :

(c)

$$\begin{aligned}
 P \times Q &\Rightarrow \begin{bmatrix} 4 & -6 \\ -2 & 8 \end{bmatrix} \times Q = \begin{bmatrix} 5 \\ 0 \end{bmatrix} \\
 &\Rightarrow Q = \begin{bmatrix} 4 & -6 \\ -2 & 8 \end{bmatrix}^{-1} \times \begin{bmatrix} 5 \\ 0 \end{bmatrix} \\
 &\Rightarrow Q = \frac{1}{20} \begin{bmatrix} 8 & 6 \\ 2 & 4 \end{bmatrix} \times \begin{bmatrix} 5 \\ 0 \end{bmatrix} \\
 &= \begin{bmatrix} \frac{8}{20} & \frac{6}{20} \\ \frac{2}{20} & \frac{4}{20} \end{bmatrix} \times \begin{bmatrix} 5 \\ 0 \end{bmatrix} \\
 &= \begin{bmatrix} \frac{8}{20} \times 5 \\ \frac{2}{20} \times 5 \end{bmatrix} = \begin{bmatrix} 2 \\ 0.5 \end{bmatrix}
 \end{aligned}$$

D

$$\begin{bmatrix} 0.5 \\ 2 \end{bmatrix}$$

QUESTION ANALYTICS

Q. 29Consider $f(x)$ be a function defined by

$$f(x) = \begin{cases} 4x - 5, & \text{if } x \leq 2 \\ x - \lambda, & \text{if } x > 2 \end{cases}$$

In $\lim_{x \rightarrow 2} f(x)$ exist, then the value of λ is _____.
[Solution Video](#) | [Have any Doubt ?](#)



Ashima Garg

Course: GATE
Computer Science Engineering(CS)

HOME

MY TEST

BOOKMARKS

MY PROFILE

REPORTS

BUY PACKAGE

ASK AN EXPERT

OFFER

EXCLUSIVE OFFER FOR OTS STUDENTS ONLY ON BOOK PACKAGES

SOLUTION .
 (-1)

$$f(x) = \begin{cases} 4x - 5, & \text{if } x \leq 2 \\ x - \lambda, & \text{if } x > 2 \end{cases}$$

$$\begin{aligned} \therefore \lim_{x \rightarrow 2^-} f(x) &= \lim_{h \rightarrow 0} f(2-h) \\ &= \lim_{h \rightarrow 0} [4(2-h) - 5] \\ &= \lim_{h \rightarrow 0} [8 - 4h - 5] \\ &= 3 \\ \lim_{x \rightarrow 2^+} f(x) &= \lim_{h \rightarrow 0} f(2+h) \\ &= \lim_{h \rightarrow 0} [2+h - \lambda] \\ &= 2 - \lambda \end{aligned}$$

Since limit exist, so,

$$\begin{aligned} \lim_{x \rightarrow 2^-} f(x) &= \lim_{x \rightarrow 2^+} f(x) \\ 2 - \lambda &= 3 \\ \lambda &= -1 \end{aligned}$$

QUESTION ANALYTICS

Q. 30

A manufacturer makes condensers which on an average are 1% defective. He packs them in boxes of 100. The probability that a box picked at random will contain 3 or more faulty condensers is _____. (Upto 2 decimal places)

[Solution Video](#) | [Have any Doubt ?](#)

Correct Option

0.08 (0.08 - 0.09)

Solution :
 0.08 (0.08 - 0.09)

$$\begin{aligned} P &= 1\% = 0.01 \\ n &= 100 \\ \lambda &= nP = 100 \times 0.01 = 1 \end{aligned}$$

$$P(r) = \frac{e^{-\lambda} \lambda^r}{r!}$$

P(3 or more faulty condensers)

$$\begin{aligned} &= P(3) + P(4) + \dots + P(100) \\ &= 1 - [P(0) + P(1) + P(2)] \\ &= 1 - \left[\frac{e^{-1} 1^0}{0!} + \frac{e^{-1} 1^1}{1!} + \frac{e^{-1} 1^2}{2!} \right] = 1 - \left[e^{-1} + e^{-1} + \frac{e^{-1}}{2} \right] \\ &= 1 - e^{-1} \left[\frac{5}{2} \right] = 0.0803 \end{aligned}$$

QUESTION ANALYTICS

Q. 31

Consider two matrices given below:

$$P = \begin{bmatrix} a & b \\ c & d \end{bmatrix}; Q = \begin{bmatrix} a^2 + b^2 & ac + bd \\ ac + bd & c^2 + d^2 \end{bmatrix}$$

If the rank of matrix P is 2, then the rank of matrix Q will be _____.
[Solution Video](#) | [Have any Doubt ?](#)



Ashima Garg

 Course: GATE
 Computer Science Engineering(CS)

HOME

MY TEST

BOOKMARKS

MY PROFILE

REPORTS

BUY PACKAGE

ASK AN EXPERT

OFFER

 EXCLUSIVE OFFER FOR OTS
STUDENTS ONLY ON BOOK
PACKAGES

(2)

$$\begin{aligned}
 P &= \begin{bmatrix} a & b \\ c & d \end{bmatrix} \\
 Q &= \begin{bmatrix} a^2 + b^2 & ac + bd \\ ac + bd & c^2 + d^2 \end{bmatrix} \\
 &= P \times P^T = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \begin{bmatrix} a & c \\ b & d \end{bmatrix} \\
 &= \begin{bmatrix} a^2 + b^2 & ac + bd \\ ac + bd & c^2 + d^2 \end{bmatrix} = Q
 \end{aligned}$$

i.e. $P \cdot P^T = Q$
 So, rank of matrix Q is same as rank of P .

QUESTION ANALYTICS

Q. 32

A system matrix is given as follows:

$$A = \begin{bmatrix} 0 & 2 & -2 \\ -12 & -22 & 12 \\ -12 & -22 & 10 \end{bmatrix}$$

 The value of the ratio of the absolute maximum eigen value to the absolute minimum eigen value is _____.
 (It is known that one of the eigen value is -6)

[Solution Video](#) | [Have any Doubt ?](#) |

3

Correct Option

Solution :

3

Characteristic equation is $|A - \lambda I| = 0$

$$\begin{vmatrix} -\lambda & 2 & -2 \\ -12 & -22 - \lambda & 12 \\ -12 & -22 & 10 - \lambda \end{vmatrix} = 0$$

$$\Rightarrow -\lambda[-220 + 22\lambda - 10\lambda + \lambda^2 + 264] - 2[-120 + 12\lambda + 144] - 2[264 - 264 - 12\lambda] = 0$$

$$\Rightarrow -\lambda(\lambda^2 + 12\lambda + 44) - 2(12\lambda + 24) + 24\lambda = 0$$

$$\Rightarrow -\lambda^3 - 12\lambda^2 - 44\lambda - 24\lambda - 48 + 24\lambda = 0$$

$$\Rightarrow -\lambda^3 - 12\lambda^2 - 44\lambda - 48 = 0$$

$$\Rightarrow \lambda^3 + 12\lambda^2 + 44\lambda + 48 = 0$$

$$\Rightarrow (\lambda + 2)(\lambda + 4)(\lambda + 6) = 0$$

(one eigen value is -6, so $\lambda + 6$ is factor of this equation so we can obtain the other roots by polynomial division)

$$\Rightarrow \lambda = -2, -4, -6$$

$$\text{Ratio} = \frac{6}{2} = 3$$

QUESTION ANALYTICS

Q. 33
 Consider $f(x) = 3x^3 - 7x^2 + 5x + 6$. The minimum value of $f(x)$ over the interval $[0, 2]$ is _____.
 (Upto 1 decimal place)

[Solution Video](#) | [Have any Doubt ?](#) |



Ashima Garg

 Course: GATE
 Computer Science Engineering(CS)

HOME

MY TEST

BOOKMARKS

MY PROFILE

REPORTS

BUY PACKAGE

ASK AN EXPERT

OFFER

 EXCLUSIVE OFFER FOR OTS
 STUDENTS ONLY ON BOOK
 PACKAGES

$$\begin{aligned}
 f(x) &= 3x^3 - 7x^2 + 5x + 6 \\
 f'(x) &= 9x^2 - 14x + 5 \\
 f'(x) &= 0 \\
 &= 9x^2 - 14x + 5 \\
 &= 9x^2 - 9x - 5x + 5 \\
 &= 9x[x - 1] - 5[x - 1] \\
 &= (9x - 5)(x - 1)
 \end{aligned}$$

$$x = 1 \text{ and } x = \frac{5}{9} = 0.55$$

$$\begin{aligned}
 f''(x) &= 18x - 14 \\
 f''(1) &= 18 - 14 \\
 &= 4 > 0 \text{ (minima)}
 \end{aligned}$$

$$\begin{aligned}
 \text{Minimum } \{f(0), f(1), f(2)\} &= \text{minimum } \{6, 7, 12\} \\
 &= 6
 \end{aligned}$$

Your Answer is 4.6

QUESTION ANALYTICS