A Continuous Life-years Gained Priority Score for Ventilator Allocation

William Parker, Siva Bhavani, Susan Han, Dwight Miller, Monica Malec, Mark Siegler

- 1 Theory
- 2 Simulation using CDC data

Theory

In military triage situations, a utilitarian framework is employed to save the greatest number of wounded soliders in a mass casualty event

Identify patients who will survive without critical care (Green)

In military triage situations, a utilitarian framework is employed to save the greatest number of wounded soliders in a mass casualty event

- Identify patients who will survive without critical care (Green)
- Exclude patients who obviously will not survive with critical care (Blue)

In military triage situations, a utilitarian framework is employed to save the greatest number of wounded soliders in a mass casualty event

- Identify patients who will survive without critical care (Green)
- Exclude patients who obviously will not survive with critical care (Blue)
- 3 Rank order patients who will die without critical care by P(ICUSurvival) (Red > Yellow)

In military triage situations, a utilitarian framework is employed to save the greatest number of wounded soliders in a mass casualty event

- Identify patients who will survive without critical care (Green)
- Exclude patients who obviously will not survive with critical care (Blue)
- 3 Rank order patients who will die without critical care by P(ICUSurvival) (Red > Yellow)
- 4 Treat as many patients as possible in order of P(ICUSurvival)

Problems with military triage approach in the COVID-19 Pandemic

Three patients with COVID-19



28 year old female
• SOFA: 30% survival



80 year old male
• SOFA: 75% survival



60 year old male
• SOFA: 50% survival

Who gets the one remaining ventilator?

New York ventilator allocation policy

Step 2 – Mortality Risk Assessment Using SOFA ¹		
Color Code and Level of Access	Assessment of Mortality Risk/ Organ Failure	
Blue No ventilator provided. Use alternative forms of medical intervention and/or palliative care or discharge. Reassess if ventilators become available.	Exclusion criterion OR SOFA > 11	
Red Highest Use ventilators as available	SOFA < 7 OR Single organ failure ²	
Yellow Intermediate Use ventilators as available	SOFA 8 – 11	
Green Use alternative forms of medical intervention or defer or discharge. Reassess as needed.	No significant organ failure AND/OR No requirement for lifesaving resources	

William Parker, Siva Bhavani, Susan Han, Dwight Miller, Monica Malec, Mark Siegler

Priority rankings under NY triage system



Goes against "youngest first" allocation principles and does not maximize life-years saved

Multiprinciple approach

Principle Specificat	Specification	Point System*			
		1	2	3	4
Save the most lives	Prognosis for short-term survival (SOFA score)	SOFA score <6	SOFA score, 6-9	SOFA score, 10–12	SOFA score >12
Save the most life-years	Prognosis for long-term survival (medical assessment of comorbid conditions)	No comorbid conditions that limit long-term survival	Minor comorbid conditions with small impact on long-term survival	Major comorbid conditions with substantial impact on long-term survival	Severe comorbid conditions; death likely within 1 year
Life-cycle principle†	Prioritize those who have had the least chance to live through life's stages (age in years)	Age 12–40 y	Age 41-60 y	Age 61–74 y	Age ≥75 y

SOFA = Sequential Organ Failure Assessment.

White

et al, Ann Internal Medicine, 2009 What justification for relative weight of each category? Why categorical?

^{*} Persons with the lowest cumulative score would be given the highest priority to receive mechanical ventilation and critical care services.

[†] Pediatric patients may need to be considered separately, because their small size may require the use of different mechanical ventilators and personnel.

Maximizing life-years gained

An alternative utilitarian approach is to maximize life-years gained

Priority Score that maximizes life-years gained

$$PriorityScore = P(ICUSurvival) * (100 - age)$$

Example: Maximizing life-years

Life-years gained allocation



28 year old female

- · SOFA: 30% survival
- 100 28 = 72 years of life left
- · 22 life-years gained with vent



- SOFA: 75% survival
- 100 80 = 20 years of life left
- 15 life-years gained with vent



60 year old male


- SOFA: 50% survival
- 100 60 = 40 years of life left
 - 20 life-years gained with vent

3


2

1

Priority Score vs. Patient Age, by Probability of ICU Survival




Range of possible priority scores by patient age




Simulation using CDC data

Data sources


We took data from the CDC report Severe Outcomes Among Patients with Coronavirus Disease 2019 — United States, February 12–March 16, 2020



Simulated ICU population from CDC data distribution




Simulated SOFA score distribution



Currently drawn from $f(SOFA|age) = N(8 + \frac{age-65}{30}, 3.5)$, need to replace with a distribution estimated from real data.

Calibration of the SOFA score


The Sequential Organ Failure Assesment (SOFA) score is a validated bedside predictor of ICU mortality. The calibration of SOFA scores is drawn from *Raith et al. JAMA*, 2017




SOFA Score by Age

Age	Mean SOFA	Survival with Ventilator
20–44	6.9	77%
45–54	7.0	76%
55–64	8.2	71%
65–74	7.7	73%
75–84	8.5	70%
85	8.7	68%


Simulated Hypotehtical Outcomes by Age



Lottery allocation




lottery allocation - ICU outcomes




A random allocation of 500 ventilators would save 365 out of 1000 who were in need of mechanical ventilation. A lottery saves 12,268 (36%) out of a total of possible 33,927 life years.

William Parker, Siva Bhavani, Susan Han, Dwight Miller, Monica Malec, Mark Siegler

Lottery- age vs. SOFA




Youngest first allocation




Youngest first allocation 500 ventilators would save 369 out of 1000 who were in need of mechanical ventilation. Youngest first saves 17.846 (53%) out of a total of possible 33.927 life years.

William Parker, Siva Bhavani, Susan Han, Dwight Miller, Monica Malec, Mark Siegler


Youngest first - ICU allocation



Youngest first allocation- age vs. SOFA




Maximizing ICU survival




<<<<<< HEAD A P(ICUsurvival) triage system of 500 ventilators would save 434 out of 1000 patients in need. Max ICU surival saves 16.295 out of a total of possible 33.927 (48%)

William Parker, Siva Bhavani, Susan Han, Dwight Miller, Monica Malec, Mark Siegler


Maximizing ICU survival- ICU Outcomes



Max ICU survival- age vs. SOFA




Maximizing Life-years gained




Prioritizing life-years for 500 ventilators would save 405 out of 1000 patients admitted to the ICU. Maximizing life-years gained saves 18.601 out of a total of possible 33.927 (55%) life-years.

William Parker, Siva Bhavani, Susan Han, Dwight Miller, Monica Malec, Mark Siegler

Max Life Years- ICU Outcomes by Age



Max life years- age vs. SOFA



Maximizing life-years vs. ICU survival

Prioritizing young sick patients over old healthy patients leads to more ICU deaths in exchange for more life-years gained.

The Tradeoff

Prioritizing life-years gained over ICU survival saves an additional 2,306 life-years for this 1000 patient sample, at a cost of 29 more deaths in the ICU.