

# Absolute rate constants for hydrocarbon autoxidation. VI. Alkyl aromatic and olefinic hydrocarbons<sup>1</sup>

J. A. HOWARD AND K. U. INGOLD

Division of Applied Chemistry, National Research Council, Ottawa, Canada


Received September 14, 1966

Absolute rate constants have been measured for the autoxidation of a large number of hydrocarbons at 30 °C. The chain-propagating and chain-terminating rate constants depend on the structure of the hydrocarbon and also on the structure of the chain-carrying peroxy radical. With certain notable exceptions which are mainly due to steric hindrance, the rate constants for hydrogen-atom abstraction increase in the order primary < secondary < tertiary; and, for compounds losing a secondary hydrogen atom, the rate constants increase in the order unactivated < acyclic activated by a single  $\pi$ -electron system < cyclic activated by a single  $\pi$ -system < acyclic activated by two  $\pi$ -systems < cyclic activated by two  $\pi$ -systems. The rate constants for chain termination by the self-reaction of two peroxy radicals generally increase in the order tertiary peroxy radicals < acyclic allylic secondary < cyclic secondary < acyclic benzylic secondary < primary peroxy radicals < hydroperoxy radicals.


Canadian Journal of Chemistry. Volume 45, 793 (1967)

## INTRODUCTION

Early work on the liquid phase autoxidation of hydrocarbons suggested that the rate constant for the chain-terminating interaction of two peroxy radicals ( $2k_t$ ),



was insensitive to the structure of R, the hydrocarbon portion in the radical (1). It was therefore assumed that variations in oxidation rates reflected changes in the rate constants for chain propagation,



Bolland (2) utilized an "average" termination constant of  $6 \times 10^5 \text{ M}^{-1} \text{ s}^{-1}$  to develop a set of rules relating olefinic structure to the ease of hydrogen-atom abstraction in chain propagation. These empirical rules have been successfully applied to the prediction of the gross reactivities of olefins containing *primary* or *secondary* allylic hydrogens but they predict too great a reactivity for olefins with *tertiary* allylic hydrogen (3, 4).

This simple picture of hydrocarbon autoxidation which was based on an invariant termination constant became unlikely when Melville and Richards (5) showed that the rate constant for the interaction of the tertiary peroxy radicals from

cumene was only about  $3 \times 10^4 \text{ M}^{-1} \text{ s}^{-1}$ . Russell (6) utilized the difference in the termination constants of *secondary* and *tertiary* peroxy radicals to explain the minimum that occurred in plots of the oxidation rate against composition for mixtures of cumene with a number of hydrocarbons giving secondary peroxy radicals. However, in the absence of evidence to the contrary, Russell was forced to assume that all *secondary* and all *primary* peroxy radicals terminate at the same rate. We have recently presented some rather limited evidence (Part IV of this series (7)) that not only do *primary* peroxy radicals terminate more rapidly than *secondary* peroxy radicals but that there is also a fairly wide variation in the rate constants for the termination of *secondary* peroxy radicals. The present paper extends the measurement of termination rate constants to a much wider variety of hydrocarbons. In addition, the propagation rate constants have been used to compare the relative rates of hydrogen abstraction by peroxy radicals with the relative rates reported for abstraction by some other free radicals.

## EXPERIMENTAL

### Reagents

None of the hydrocarbons and esters used in this work had purities of less than 99.5% as determined by gas-liquid chromatography (g.l.c.). All compounds except neopentylbenzene were passed through alumina immediately before use.

<sup>1</sup>Issued as N.R.C. No. 9318.

### Experimental Procedure

The experimental technique and theory of the rotating sector method have been described in previous papers in this series (7-10). None of the compounds studied showed any variation of the rate with oxygen pressure (200-730 mm). Unreactive liquid substrates were used in the pure state. The solid hydrocarbons and the more reactive liquids were used as solutions in chlorobenzene, their concentrations ranging as low as 0.1 mole/l for the very reactive substrates. In calculations of rate constants for 1,3- and 1,4-cyclohexadiene, 1,2- and 1,4-dihydronaphthalene, and 9,10-dihydroanthracene in chlorobenzene it was assumed that the rate of chain initiation was equal to the value found under similar conditions with less reactive hydrocarbons (see preceding paper (11)). The hydroperoxide yield was determined by the iodine-thiosulfate method so that the rate of hydrogen-atom abstraction ( $k_{abstr}$ ) could be separated from the other chain-propagating steps such as addition ( $k_{add}$ ) (10).

Although it is nearly impossible to assign specific errors to absolute rate constants, the sources of such errors were discussed in some detail in earlier papers (8, 10). In the present work, particular attention was paid to the chain lengths ( $\nu$ ) of the reactions. All the results represent averages of reasonably consistent data which were obtained at not less than two chain lengths. Most of the compounds oxidized quite readily and gave no particular difficulty. For these compounds it is probably safe to assume that  $k_t$  should be accurate to within a factor of 2 and  $k_p$  should be accurate to within  $\pm 50\%$ . Certain compounds exhibited specific problems which are outlined below. The calculation of absolute rate constants from measurements at short chain lengths is also described below.

### Compounds Presenting Special Problems

#### (i) Toluene and *m*-Xylene

Toluene and *m*-xylene both oxidized at rather too low a rate for satisfactory rotating sector measurements at our standard temperature of 30 °C. Fortunately, all the available evidence indicates that  $k_t$  is probably not significantly affected by polar factors (6, 9, 12). It is therefore fairly safe to assume that  $k_t$  for these two hydrocarbons will be close to the value found for *p*-xylene, i.e.  $1.5 \times 10^8 M^{-1} s^{-1}$ . Propagation rate constants for toluene and *m*-xylene were calculated using this assumed termination constant.

#### (ii) Methyl Linolenate

The rate of oxygen absorption decreased sharply as the reaction proceeded. This was not observed with the simple monoolefins (see Fig. 1). The decreased rate was attributed to a decrease in the rate of chain initiation ( $R_i$ ) resulting from the formation of a conjugated triene (these compounds absorb strongly at the initiating wavelength, 365 m $\mu$ ). The results for methyl linolenate are therefore much less reliable than those obtained with most other compounds. A much smaller autoretardation was observed with methyl linoleate, indene, and

2,5-dimethyl hexene-3. In contrast, diphenylmethane, octene-1, 1,3-cyclohexadiene, decyne-5, and 1,2-dihydronaphthalene were slightly autocatalytic. For the remaining hydrocarbons, the rates of oxygen absorption were constant for the duration of the experiments.




FIG. 1. Relative rates of oxygen absorption at 30 °C vs. amount of oxygen absorbed for some unsaturated compounds. (○) Heptene-3; (□) methyl oleate; (△) methyl linoleate; (×) methyl linolenate.

#### (iii) 9,10-Dihydroanthracene, Decyne-5, and Octyne-1

9,10-Dihydroanthracene, decyne-5, and octyne-1 underwent oxidation in the dark. For the first two compounds the relative importance of this thermally initiated reaction could be reduced sufficiently to permit rotating sector studies by oxidizing them as 1.0 *M* solutions in chlorobenzene. For octyne-1 the thermal rate, even in the absence of hydroperoxide, was 30-40% of the photochemical rate under the standard experimental conditions and absolute rate constants could not be obtained.

#### (iv) 1,2-Dihydronaphthalene

1,2-Dihydronaphthalene forms both polyperoxides and hydroperoxides (cf. indene (10, 13)). It also behaves like other hydrocarbons which contain a styrene skeleton with a hydrogen atom in the  $\alpha$ -position in that there is a kinetically first-order chain-termination process (rate constant  $k_x$ ) as well as the usual bimolecular termination process (8-10). This renders the accurate determination of  $k_p$  and  $k_t$  quite difficult. The first-order process is a fairly important source of chain termination and its importance increases during the oxidation (cf.  $\beta$ -methylstyrene (10)). The first-order rate constant in chlorobenzene at 30 °C is about  $0.9 s^{-1}$  which can be compared with values of 0.24, 0.61, and  $1.24 s^{-1}$  for styrene,  $\beta$ -methylstyrene, and indene, respectively, under the same conditions (10). The first-order termination process could be effectively eliminated by the addition of  $D_2O$  (8, 10), the overall rate of oxidation increasing by about 35%. In the presence of  $D_2O$  the measured propagation and termination rate constants were  $415 M^{-1} s^{-1}$  and

$1.2 \times 10^8 M^{-1} s^{-1}$  respectively, whereas in the absence of  $D_2O$  the measured values were  $190 M^{-1} s^{-1}$  and  $3.5 \times 10^7 M^{-1} s^{-1}$ . The former values are considered the more accurate because the first-order correction has been eliminated and so they are quoted in Table III.

(v) *9,10-Dihydrophenanthrene*

9,10-Dihydrophenanthrene had a kinetically first-order chain-termination process which increased in importance as the reaction proceeded. In the initial stages of the reaction  $k_x$  was about  $0.3 s^{-1}$ . Since the addition of  $D_2O$  reduced, but did not eliminate, this process, it is possible that only part of the first-order termination was due to the formation of a phenolic inhibitor (9<sup>2</sup>). The increase in  $k_x$  during the reaction may result from the slow accumulation of the weak inhibitor phenanthrene.

*Calculation of Rate Constants at Short Chain Lengths*

At short chain lengths the usual assumption that the measured rate of gas absorption is proportional to the peroxy radical concentration is no longer valid. This is because the quantities of gas absorbed and evolved in the initiation and termination processes become significant compared with the amount absorbed in the propagation process. It was assumed that for each molecule of initiator (azo-bis-cyclohexylnitrile) decomposed, one molecule of nitrogen was evolved,  $2e$  molecules of oxygen were absorbed by the initiating radicals, and  $e$  molecules of oxygen were finally evolved in the chain-ending step. By using this correction<sup>3</sup> it was found that absolute rate constants obtained at chain lengths as low as 3.0 were not significantly different from those obtained at long chain lengths.

*Reactions with *t*-Butoxy Radicals*

In order to extend the comparison of hydrocarbon reactivity towards peroxy and alkoxy radicals we measured the relative reactivity of a few hydrocarbons towards *t*-butoxy radicals. The radicals were generated photolytically from *t*-butyl hypochlorite using the general techniques and competitive procedures developed by Walling and co-workers (16–18). The reactions were carried out in dilute solutions in  $CCl_4$  at  $40^\circ C$  as we have described previously (19). Analyses were by gas chromatography and selective solvolysis (19). Where comparison is possible, the present results are generally

<sup>2</sup>The oxidation products gave a positive response to the phosphomolybdic acid spot test for phenols (14). The parent hydrocarbon and phenanthrene both gave a negative response to this test.

<sup>3</sup>Under intermittent illumination this correction is independent of the speed of rotation of the sector disc. That is, for our experimental arrangement with a light/dark ratio of 3.0, the correction under intermittent illumination is 1/4 of the correction under steady illumination. In similar circumstances, Tsepalov and Shlyapintokh (15) have used a correction which varies with the sector's rotation speed. We believe our method of correcting the measured rates is more precise.

in quite good agreement with the data of Walling (see Table IV).

In Part IV (7), deuterium isotope effects were reported for the oxidation of  $\alpha, \alpha$ - $d_2$ -diphenylmethane and perdeuterotetralin at  $30^\circ C$ . The measured values of  $(k_p)_H/(k_p)_D$  were 5.1 for diphenylmethane, which is of about the expected magnitude, but  $> 50$  for tetralin. It was suggested that this high value might be due to an impurity in the deuterated compound which retarded its oxidation. Such an impurity should not affect a relative reactivity determined by a competitive procedure. In competition with toluene, the isotope effects for the reaction with *t*-butoxy radicals were 1.7 for diphenylmethane and 2.4 for tetralin, both values being in good agreement with our previous value of 2.4 for toluene itself (19).

## RESULTS

Rate constants for the autoxidation of alkyl aromatic hydrocarbons, olefins, and alkenyl aromatic hydrocarbons at  $30^\circ C$  are given in Tables I, II, and III respectively. The tables include data from the previous papers in this series. Values of  $k_p/(2k_t)^{1/2}$  are in reasonable agreement, after allowing for the  $15^\circ$  temperature difference, with Bolland's results at  $45^\circ C$  (2). The agreement is good for 2,3-dimethyl butene-2, methyl oleate, heptene-3, octene-1, and allylbenzene, but only fair for tetralin, cyclohexene, linoleate, linolenate, and crotylbenzene. Rate constants for hydrogen-atom abstraction are given on an available hydrogen basis, e.g.  $k_{abstr}/H = k_p/3$  for toluene,  $k_p/2$  for ethylbenzene, etc. Compounds which did not contain benzylic or allylic hydrogens, e.g. *t*-butylbenzene, cyclohexane, *n*-decane, and methyl palmitate, did not oxidize under the present conditions. The error involved in ignoring the unactivated hydrogen atoms when calculating  $k_{abstr}/H$  is therefore negligible. Although several alkenes did not give 100% yields of hydroperoxide, the values of  $k_{abstr}/H$  were only corrected when the measured yields fell below 75%. The hydroperoxide yields were generally not measured for those compounds for which yields of over 80% at higher temperatures have been reported. All rate constants are in units of  $\text{mole}^{-1} s^{-1}$ .

In the tables the compounds from which a *primary* hydrogen atom is abstracted are listed ahead of those from which a *secondary*

TABLE I  
Rate constants for the autoxidation of alkyl aromatic hydrocarbons at 30 °C\*

Hydrocarbon	$k_p/(2k_t)^{1/2} \times 10^5$	$k_{abstr}/H$	$k_t \times 10^{-6}$
Primary			
Toluene	1.4	0.08	150†
<i>m</i> -Xylene	2.8	0.08	150†
<i>o</i> -Xylene	3.3	0.07	77
<i>p</i> -Xylene	4.9	0.14	150
Secondary			
Neopentylbenzene‡	2.0	—	—
<i>n</i> -Butylbenzene	8.1	0.28	25
Bibenzyl	13	0.14	10
Ethylbenzene (ref. 7)	21	0.65	20
Indan	170	1.2	4.1
Tetralin (ref. 7)	230	1.6	3.8
9,10-Dihydrophenanthrene	720	14	30
Diphenylmethane (ref. 7)	38	2.4	80
9,10-Dihydroanthracene	5 850	44	4.6
Tertiary			
<i>sec</i> -Amylbenzene	11	0.07	0.22
Phenylcyclohexane	15	0.06	0.08
<i>sec</i> -Butylbenzene	18	0.08	0.09
1,1-Diphenylethane	110	0.34	0.047
Isopropylbenzene	150	0.18	0.0075

\*In mole<sup>-1</sup> s<sup>-1</sup> units.

†Assumed, see text.

‡Oxidized too slowly for a rotating sector study.

TABLE II  
Rate constants for the autoxidation of alkenes at 30 °C†

Olefins‡	Hydroperoxide yield (%)	$k_p/(2k_t)^{1/2} \times 10^5$	$k_{abstr}/H$	$k_t \times 10^{-6}$
Primary				
2,3-Me <sub>2</sub> -butene-2*( <i>p,t</i> )	67	320	0.14	0.32
Secondary ( <i>s,s</i> )				
Methyl oleate	>81§	89	0.22	0.53
Heptene-3	100	54	0.35	3.2
Octene-1*( <i>s,p</i> )	80	6.2	0.50	130
Cyclopentene	86	280	1.7	3.1
Cyclohexene (ref. 7)	88	230	1.5	2.8
1,3-Cyclohexadiene	27	10 000	55	33
1,4-Pentadiene*( <i>s,p</i> )	100	42	7.0	540
Methyl linoleate*	98§	2 100	31	4.4
Methyl linolenate*	~100¶	3 900	59	18
1,4-Cyclohexadiene* (ref. 11)	97	3 900	370	630
Tertiary				
2,5-Me <sub>2</sub> -hexene-3( <i>t,s</i> )	100	540	1.2	0.09
Acetylenes				
Octyne-1	—	12	—	—
Decyne-5	80	74	0.7	7.0

\*A rearranged peroxy radical is probably involved in the reactions. See text.

†In mole<sup>-1</sup> s<sup>-1</sup> units.‡(*p,t*) refers to a reaction in which a primary hydrogen is abstracted to give an allylic radical in which the odd electron is delocalized between a primary and a tertiary carbon, (*s,s*) a secondary hydrogen abstracted and the electron delocalized between two secondary carbon atoms, etc.

§Reference 20.

||Reference 21.

¶Assumed to be ~100% by analogy with methyl linoleate. Titration actually gives only 48%, probably because the hydroperoxide is rather unstable under our conditions.

TABLE III  
Rate constants for the autoxidation of alkenyl aromatic hydrocarbons at 30 °C†

Hydrocarbon	$k_p/(2k_t)^{1/2} \times 10^5$	$k_{abstr}/H$	$k_{add}$	$k_t \times 10^{-6}$
Secondary				
Styrene (ref. 9)	890	—	41	21
$\beta$ -Methyl styrene (ref. 10)	1 300	—	51	16
Allylbenzene* (ref. 7)	49	5.0	—	220
Crotylbenzene*,‡	420	4.1	—	1.9
Indene (ref. 10)	2 840	7.0	128	25
1,2-Dihydronaphthalene	2 750	26§	291	115
1,4-Dihydronaphthalene* (ref. 11)	3 500	225	—	350
Tertiary				
$\alpha$ -Methyl styrene (ref. 10)	1 300	—	10	0.3

\*A rearranged peroxy radical is probably involved in the reactions. See text.

†In mole<sup>-1</sup> s<sup>-1</sup> units.


‡C<sub>6</sub>H<sub>5</sub>CH<sub>2</sub>CH=CHCH<sub>3</sub>.

§Assumed to have four active hydrogen atoms.

hydrogen is abstracted and these precede the *tertiary* hydrogen donors. Those compounds losing *secondary* hydrogens are arranged in the order: acyclic activated by a single  $\pi$ -electron system (i.e.  $\text{>C}-\text{CH}_2-\text{R}$ ); cyclic activated by a single  $\pi$ -system; acyclic activated by two  $\pi$ -systems (i.e.  $\text{>C}-\text{CH}_2-\text{C}<$ ); and cyclic activated by two  $\pi$ -systems. This order corresponds approximately with the order of increasing ease of hydrogen-atom abstraction. It must be emphasized that classification as *primary*, *secondary*, or *tertiary* does not imply that the same *primary*, *secondary*, or *tertiary* peroxy radical is produced and is involved in the abstraction and (or) termination processes. That is, oxygen may add to either end of an allylic radical formed by hydrogen abstraction from an alkene. This can lead, for example, to the production of mainly *primary* peroxy radicals after the abstraction of a *secondary* hydrogen from allylbenzene (22) and 1-octene (23).



Rearrangement of the allylic radical also occurs for methyl linoleate and linolenate (24), although in these cases the peroxy radical remains a *secondary* radical.



In a similar way, 1,4-pentadiene should give a *primary* peroxy radical. All the obvious cases where oxygen does not add at

the position from which hydrogen was abstracted because of the migration of a double bond are marked with an asterisk in the tables. 1,4-Cyclohexadiene and 1,4-dihydronaphthalene give mainly the HOO $\cdot$  radical under the conditions employed (11). In addition to these changes in the monomeric peroxy radicals, compounds containing the styrene skeleton and compounds with conjugated double bonds form polymeric peroxy radicals, R(OOR)<sub>n</sub>OO $\cdot$ . These polymeric radicals are certainly the main hydrogen-atom-abstracting species from indene, 1,2-dihydronaphthalene, and 1,3-cyclohexadiene (2).

The rate of the photochemically initiated autoxidation of *m*-xylene was measured from 30 to 65 °C. The results could be represented by

$$k_p/(2k_t)^{1/2} = 1.7 \times 10^2 \exp(-9400/RT) (M^{-1} \text{ s}^{-1})^{1/2}.$$

Since  $k_t$  for this hydrocarbon is probably quite large (Table I) the chain-termination process cannot be associated with a big activation energy. If we assume it has a value of 1.2 kcal/mole<sup>4</sup> then

$$k_{abstr}/H = 1.3 \times 10^6 \exp(-10000/RT) M^{-1} \text{ s}^{-1}$$

for *m*-xylene and also for toluene.

<sup>4</sup>This is purposely chosen to be slightly smaller than the value of 1.8 kcal/mole found previously for the autoxidation of styrene (8) because  $k_t$  is probably larger for *m*-xylene than for styrene.

The rate of the azo-bis-isobutyronitrile thermally initiated oxidation of cyclohexane was measured at 65 °C. By using a low concentration of AIBN ( $5.2 \times 10^{-4} M$ ) a chain length of 2.4 was attained. The measured rate gave  $k_p/(2k_t)^{1/2} = 2.75 \times 10^{-5} (M^{-1} s^{-1})^{1/2}$ . Taking McCarthy and MacLachlan's (25) value of  $1.4 \times 10^6 M^{-1} s^{-1}$  for  $k_t$  at 25 °C and assuming  $E_t \sim 4.5$  kcal/mole (cf. tetralin (7)) gives  $k_t = 3.5 \times 10^6 M^{-1} s^{-1}$  at 65 °C. Therefore, at 65 °C  $(k_{abstr}/H)_{\text{cyclohexane}} = 6.1 \times 10^{-3} M^{-1} s^{-1}$  and hence  $(k_{abstr}/H)_{\text{cyclohexane}} / (k_{abstr}/H)_{\text{toluene}} = 0.014$ . This small value indicates that peroxy radicals have a high selectivity in hydrogen-atom abstraction reactions (cf. Table IV and the results for a number of additional radicals collected in ref. 26).

#### DISCUSSION

The compounds listed in Tables I, II, and III exhibit wide differences in the ease with which their activated hydrogen atoms can be abstracted by the chain-carrying peroxy radicals and in the rates at which the peroxy radicals interact with one another to terminate reaction chains. These differences must arise from a variety of mechanistic, steric, polar, and radical stabilization effects. Small differences produced by small changes in reactant structure could often be attributed to several of these factors. For this reason, an attempt has been made to restrict the following discussion of chain termination and hydrogen abstraction to those results for which the relation between structure and rate constant appears to be fairly straightforward.

##### Chain Termination


Secondary and, presumably, primary peroxy radicals terminate reaction chains by transfer of an  $\alpha$ -hydrogen atom (27). The ease with which this atom is transferred plays a very important role in determining the rate constant for termination.

##### (i) Primary Peroxy Radicals

Primary peroxy radicals derived from alkylbenzenes and 1-olefins (allylbenzene (22) and octene-1 (23)) give  $k_t$  values in the range  $0.8\text{--}1.5 \times 10^8 M^{-1} s^{-1}$ . As we had predicted (7), previously reported

values of  $k_t$  for octene-1 (i.e.  $0.8\text{--}1.5 \times 10^8 M^{-1} s^{-1}$  (1), and  $4.6 \times 10^6 M^{-1} s^{-1}$  (25)) were too small. The larger termination constant for 1,4-pentadiene (which can be assumed to give mainly *primary* peroxy radicals) indicates that a peroxy radical adjacent to a conjugated system of double bonds terminates more readily than one adjacent to a single double bond, possibly because the  $\alpha$ -hydrogen is more loosely bound and thus more readily transferred.

The low termination constant for 2,3-dimethyl butene-2 ( $3.2 \times 10^5 M^{-1} s^{-1}$ ) suggested that tertiary rather than primary peroxy radicals are involved in the oxidation, i.e.



The allylic radical should have a greater electron density at the *tertiary* carbon (structure II) than at the *primary* carbon (structure I).<sup>5</sup> Oxygen should therefore add mainly to the tertiary position. It has been shown previously that the hydroperoxide derived from III (1,1,2-trimethylallylhydroperoxide) is the main product of the Rose Bengal photosensitized autoxidation (28) and the singlet oxygen oxidation (29) of 2,3-dimethyl butene-2. The same hydroperoxide is produced under our experimental conditions and the low termination rate constant for this olefin is thus explained.

##### (ii) Secondary Peroxy Radicals

Secondary peroxy radicals show wide variations in  $k_t$  for comparatively small changes in structure. Most secondary benzylic peroxy radicals ( $\text{C}_6\text{H}_5\text{C}(\text{H},\text{R})\text{OO}\cdot$ )

<sup>5</sup>This is due to the greater stability of tertiary radicals. Moreover, repulsion between the methyl groups will be reduced by an increase in the central CC bond length, i.e. by an increase in its single bond character.

have values in the range  $1.0\text{--}3.0 \times 10^7 M^{-1} s^{-1}$  which is about an order of magnitude smaller than the values found with the xylenes. This difference can be partly attributed to the greater steric hindrance to termination for secondary than for primary peroxy radicals. A number of *m*- and *p*-substituted styrenes have  $k_t$  values in the range  $2.2\text{--}5.4 \times 10^7 M^{-1} s^{-1}$  (9). Diphenylmethane has the highest  $k_t$  value ( $8.0 \times 10^7 M^{-1} s^{-1}$ ) of all the secondary benzylic compounds examined, possibly because the  $\alpha$ -hydrogen is activated by a second phenyl group.

The difference in termination rate constants for methyl oleate and heptene-3 can presumably be attributed to steric effects. The fact that  $k_t$  for heptene-3 is smaller than for secondary benzylic hydrocarbons may be partly related to steric effects and partly to the rapid isomerization of acyclic allylic peroxy radicals (30). Steric effects must be nearly equal for methyl oleate and methyl linoleate. Comparison of their termination constants again suggests that a peroxy radical adjacent to a conjugated system of double bonds terminates more readily than one adjacent to a single double bond.


It is rather more difficult to interpret the termination constants for cyclic hydrocarbons. We have previously attributed the comparatively low  $k_t$  values for cyclohexene and tetralin ( $\sim 3 \times 10^6 M^{-1} s^{-1}$ ) to the fact that these compounds have non-planar, six-membered rings (7). This leads to steric hindrance in termination because one radical must have its peroxy group in the unfavorable axial position in order for its  $\alpha$ -hydrogen to be in the most favored position for transfer. Rather surprisingly cyclopentene and indan also have small termination rate constants. Since these five-membered rings will be virtually planar, these low values suggest that there is an increase in ring strain in the transition state for termination. The  $HOO\cdot$  radical is responsible for the very high termination constants observed in the oxidation of dilute solutions of 1,4-cyclohexadiene and 1,4-dihydronaphthalene (11). Steric hindrance to termination is probably more

important for 9,10-dihydroanthracene ( $k_t = 4.6 \times 10^6 M^{-1} s^{-1}$ ) than for the remaining cyclic compounds 1,3-cyclohexadiene, indene, 1,2-dihydronaphthalene, and 9,10-dihydrophenanthrene, which have  $k_t$  values in the range  $2.5\text{--}11.5 \times 10^7 M^{-1} s^{-1}$ .



The termination and propagation rate constants for decyne-5 are slightly higher than the values found for comparable olefins.

### (iii) Tertiary Peroxy Radicals

Tertiary peroxy radicals, having no  $\alpha$ -hydrogens, generally terminate chains more slowly than secondary peroxy radicals. The self-reaction of *t*-peroxy radicals has received considerable attention in recent years (see e.g. refs. 31-33). Oxygen is evolved in both a chain-terminating and a non-terminating interaction.



Both processes involve significant activation energies (33, 34). The alkoxy radicals may undergo  $\beta$ -scission to give a ketone and a new radical which, if it contains an  $\alpha$ -hydrogen atom, may become important in the overall termination process, e.g. (32)



The relative importance of the direct termination (reaction [1]) and the indirect termination (via  $\beta$ -scission) cannot be obtained by rotating sector measurements.

Rather surprisingly the measured termination constants decrease as the size of the groups on the *tertiary* carbon decrease (see, *sec*-amyl-, *sec*-butyl-, and isopropylbenzene). Unfortunately, it was not possible to measure  $k_t$  for triphenylmethane because its rate of oxidation depends on the oxygen pressure (6). The differences in the  $k_t$  values for the various tertiary hydrocarbons may be associated with differences in the relative importance of the two termination processes.

The  $k_t$  value for cumene is only about half the value of  $1.8 \times 10^4 \text{ M}^{-1} \text{ s}^{-1}$  obtained as the rate constant for reaction [1] at this temperature by electron spin resonance (34), which suggests that the indirect termination process, via  $\text{CH}_3\text{OO}^\cdot$ , is unimportant under our experimental conditions. This was confirmed by showing that added cumene hydroperoxide had a comparatively small effect on the rate constants for the oxidation of cumene at  $30^\circ\text{C}$ . The presence of  $1.6 \text{ M}$  cumene hydroperoxide in cumene gave  $k_p/(2k_t)^{1/2} = 1.05 \times 10^{-3} \text{ M}^{-1/2} \text{ s}^{-1/2}$ ,  $k_p = 0.14 \text{ M}^{-1} \text{ s}^{-1}$ , and  $k_t = 8.3 \times 10^3 \text{ M}^{-1} \text{ s}^{-1}$ . The differences in the absolute rate constants from the values found in the absence of added hydroperoxide are not significant. The fact that  $k_t$  is unchanged under conditions designed to suppress termination via  $\beta$ -scission (32) again implies that this process is not important at  $30^\circ\text{C}$ . The small decrease in  $k_p/(2k_t)^{1/2}$  produced by added cumene hydroperoxide contrasts with its accelerating effect at  $60^\circ\text{C}$  (32).

#### Hydrogen Abstraction

The rate constants in Tables I, II, and III indicate that singly activated benzylic and allylic hydrogen atoms have similar reactivities in compounds of similar structure. This point is brought out more clearly in Table IV which compares the rates of hydrogen abstraction (per activated hydrogen) from a number of compounds with the rate of abstraction from toluene. The similarity in the reactivity of indan and cyclopentene and of tetralin and cyclohexene is remarkable. In the analogous straight chain compounds the allylic hydrogens seem to be slightly more reactive than the benzylic. This might be due to a lower steric hindrance for the abstraction of an allylic hydrogen or to the greater stability of the resultant allylic radical. The greater reactivity of allylic positions is most clearly apparent for hydrogen atoms activated by two  $\pi$ -systems in the series diphenylmethane, allylbenzene, and 1,4-pentadiene.

We have recently discussed some of the factors which may contribute to the greater reactivity of cyclic hydrocarbons compared

with their straight chain analogues (34), cf. tetralin and indan vs. ethylbenzene, cyclohexene and cyclopentene vs. heptene-3. The difference appears to be due to a combination of radical stabilization, steric effects, and polar factors rather than to one overriding process. The rather high reactivities of 1,4-cyclohexadiene and 1,4-dihydronaphthalene compared with indene and 9,10-dihydroanthracene suggest that the  $\text{HOO}^\cdot$  radical is more active than the secondary peroxy radicals derived from the indene and 9,10-dihydroanthracene. This is probably due more to steric effects than to differences in the  $\text{HOO}-\text{H}$  and  $\text{ROO}-\text{H}$  bond strengths (35).












The polar and radical stabilizing effects of the alkyl groups not directly involved in the reaction are probably mainly responsible for the greater reactivities of 2,3-dimethyl butene-2 and 2,5-dimethyl hexene-3 compared with toluene and cumene respectively (cf. toluene and *p*-xylene in Table I). A similar explanation probably applies to the difference in reactivity of methyl linoleate and 1,4-pentadiene.

Steric effects can be invoked to explain the lower reactivity of methyl oleate compared with heptene-3 and of *sec*-amyl- and *sec*-butylbenzene compared with cumene. The *tertiary* hydrogen in cumene is itself remarkably unreactive when compared with the *secondary* hydrogens of ethylbenzene. When cumene and ethylbenzene are attacked by the same free radical, the reactivity of cumene (per benzylic hydrogen) is *always* greater than the reactivity of ethylbenzene (cf. Table IV and the literature data collected in ref. 34). The low reactivity of cumene in the present work must therefore be attributed to steric factors associated with the attacking *tertiary* peroxy radical. That is, the *tertiary* cumylperoxy radical is a less reactive radical than the *secondary*  $\alpha$ -phenethylperoxy radical. This difference in the reactivities of *secondary* and *tertiary* peroxy radicals appears to be a fairly general phenomenon (34, 36). It has an important bearing on the relative rates of consumption of different hydrocarbons during the autoxidation of hydrocarbon mixtures.



TABLE IV

Reactivities of  $\alpha$ -hydrogen atoms toward abstraction by some free radicals  
(reactivity per active hydrogen relative to toluene)

Compound	$\text{RO}_2^\cdot$ , 30 °C	$(\text{CH}_3)_3\text{CO}^\cdot$ , 40 °C	$\text{C}_6\text{H}_5^\cdot$ , 60 °C
	1.00 <sup>c</sup>	1.00 <sup>c</sup>	1.00 <sup>c</sup>
$(\text{CH}_3)_2\text{C}=\text{C}(\text{CH}_3)_2$	2.6	2.0 <sup>d</sup>	2.1
	8.1	3.2	4.6
	2.0	1.8 <sup>e,f</sup>	2.8 <sup>g</sup>
$\text{CH}_2=\text{CH}(\text{CH}_2)_3\text{CH}_3$	6.3	6.1 <sup>h</sup>	3.4
$\text{R}'\text{CH}=\text{CHCH}_2\text{R}$	4.4 <sup>i</sup>	9.2 <sup>j</sup>	3.3 <sup>k</sup>
	15	19 <sup>l</sup>	8.0
	21	38	9.7
	20	16 <sup>e,l</sup>	12
	18	37	11
	30	3.5 <sup>e,m</sup>	7.7
	63	8.3 <sup>e,n</sup>	10
$\text{CH}_2=\text{CHCH}_2\text{CH}=\text{CH}_2$	88	—	6.6 <sup>o</sup>
	2.3	6.8	9.7
$\text{R}'\text{CH}=\text{CHCH}(\text{CH}_3)_2$	15	18 <sup>o</sup>	16 <sup>o</sup>
	0.014 <sup>p</sup>	1.5	1.0

<sup>a</sup>References 16 and 17.<sup>b</sup>Reference 26.<sup>c</sup>By definition.<sup>d</sup>*cis*-Butene-2 and *trans*-butene-2.<sup>e</sup>This work.<sup>f</sup>A value of 4.1 was obtained for *n*-butylbenzene.<sup>g</sup>G. A. Russell. Private communication.<sup>h</sup>Butene-1.<sup>i</sup>Heptene-3.<sup>j</sup>*cis*-Pentene-2 and *trans*-pentene-2.<sup>k</sup>An average value for a secondary allylic hydrogen derived from several 1- and 2-olefins.<sup>l</sup>A value of 9.2 is given in ref. 18.<sup>m</sup>A value of 4.7 is given in ref. 16.<sup>n</sup>A value of 5.4 is given in ref. 18.<sup>o</sup>3-Methyl-butene-1.<sup>p</sup>At 65 °C.

In Table IV the reactivities of the hydrocarbons towards their own peroxy radicals are compared with their reactivities towards two more active radicals, *t*-butoxy (16–18) and phenyl (26). The most remarkable aspect of this comparison is the general similarity (with the exception of cumene) in the relative reactivities of singly activated hydrogen atoms towards the three radicals. This similarity does not extend to hydrogen atoms activated by two  $\pi$ -systems, the peroxy radicals showing a greater preference for these more weakly bound hydrogens. In a similar way, peroxy radicals are much more reluctant to abstract an unactivated hydrogen atom from cyclohexane than are the *t*-butoxy or phenyl radicals. Since only small differences are observed in the relative reactivities of hydrocarbons containing hydrogens activated by a single  $\pi$ -system for radicals of widely different absolute reactivity, some caution must be employed when relative reactivities are used to identify abstracting radicals.<sup>6</sup> When this method is employed the structures of the reactants and their relative reactivities should cover a wide range. The present results indicate that secondary peroxy radicals are more selective than chlorine, *t*-butoxy, phenyl, and methyl, and less selective than bromine, trichloromethyl, and polystyryl (see Table IV and refs. 26, 38, and 40).

## REFERENCES

1. L. BATEMAN and G. GEE. Proc. Roy. Soc. London, Ser. A, **195**, 376, 391 (1948); Trans. Faraday Soc. **47**, 155 (1951).
2. J. L. BOLLAND. Quart. Rev. London, **3**, 1 (1949); Trans. Faraday Soc. **46**, 358 (1950).
3. W. F. BRILL and B. J. BARONE. J. Org. Chem. **29**, 140 (1964).
4. W. F. BRILL. Advan. Chem. Ser. **51**, 70 (1965).
5. H. W. MELVILLE and S. RICHARDS. J. Chem. Soc. 944 (1954).
6. G. A. RUSSELL. J. Am. Chem. Soc. **77**, 4583 (1955); **78**, 1047 (1956).
7. J. A. HOWARD and K. U. INGOLD. Can. J. Chem. **44**, 1119 (1966).
8. J. A. HOWARD and K. U. INGOLD. Can. J. Chem. **43**, 2729 (1965).
9. J. A. HOWARD and K. U. INGOLD. Can. J. Chem. **43**, 2737 (1965).
10. J. A. HOWARD and K. U. INGOLD. Can. J. Chem. **44**, 1113 (1966).
11. J. A. HOWARD and K. U. INGOLD. Can. J. Chem. This issue.
12. G. A. RUSSELL and R. C. WILLIAMSON, JR. J. Am. Chem. Soc. **86**, 2357, 2364 (1964).
13. G. A. RUSSELL. J. Am. Chem. Soc. **78**, 1035, 1041 (1956).
14. V. M. PLATKOVSKAYA and S. G. VATKINA. Zh. Prikl. Khim. S.S.S.R. **10**, 202 (1937).
15. V. F. TSEPALOV and V. YA. SHLYAPINTOKH. Kinetika i Kataliz, **3**, 870 (1962).
16. C. WALLING and B. B. JACKNOW. J. Am. Chem. Soc. **82**, 6108 (1960).
17. C. WALLING and W. THALER. J. Am. Chem. Soc. **83**, 3877 (1961).
18. P. WAGNER and C. WALLING. J. Am. Chem. Soc. **87**, 5179 (1965).
19. B. R. KENNEDY and K. U. INGOLD. Can. J. Chem. **44**, 2381 (1966).
20. K. R. HARGRAVE and A. L. MORRIS. Trans. Faraday Soc. **52**, 89 (1956).
21. D. E. VAN SICKLE, F. R. MAYO, and R. M. ARLUCK. J. Am. Chem. Soc. **87**, 4824, 4832 (1965).
22. L. BATEMAN. Quart. Rev. London, **8**, 147 (1954).
23. S. J. MOSS and H. STEINER. J. Chem. Soc. 2372 (1965).
24. N. A. KHAN, W. O. LUNDBERG, and R. T. HOLMAN. J. Am. Chem. Soc. **76**, 1779 (1954).
25. R. L. MCCARTHY and A. MACLACHLAN. Trans. Faraday Soc. **57**, 1107 (1961).
26. R. F. BRIDGER and G. A. RUSSELL. J. Am. Chem. Soc. **85**, 3754 (1963).
27. G. A. RUSSELL. J. Am. Chem. Soc. **79**, 3871 (1957).
28. G. O. SCHENCK and K. H. SCHULTE-ELTE. Ann. Chem. **618**, 185 (1958).
29. C. S. FOOTE and S. WEXLER. J. Am. Chem. Soc. **86**, 3879 (1964).
30. W. F. BRILL. J. Am. Chem. Soc. **87**, 3286 (1965).
31. P. D. BARTLETT and T. G. TRAYLOR. J. Am. Chem. Soc. **85**, 2407 (1963).
32. T. G. TRAYLOR and C. A. RUSSELL. J. Am. Chem. Soc. **87**, 3698 (1965).
33. J. R. THOMAS. J. Am. Chem. Soc. **87**, 3935 (1965).
34. K. U. INGOLD. Intern. Symp. Free Radicals Soln., University of Michigan, Ann Arbor, Mich. 1966.
35. S. W. BENSON. J. Am. Chem. Soc. **87**, 972 (1965).
36. B. S. MIDDLETON and K. U. INGOLD. Can. J. Chem. **45**, 191 (1967).
37. R. E. PEARSON and J. C. MARTIN. J. Am. Chem. Soc. **85**, 354, 3142 (1963).
38. G. A. RUSSELL, C. DE BOER, and K. M. DESMOND. J. Am. Chem. Soc. **85**, 365, 3136, 3139 (1963).
39. C. WALLING, A. L. RIEGER, and D. D. TANNER. J. Am. Chem. Soc. **85**, 3129 (1963).
40. R. A. GREGG and F. R. MAYO. Discussions Faraday Soc. **2**, 328 (1947).

<sup>6</sup>For example, this technique has been used to show that bromine atoms are the abstracting species in N-bromosuccinimide brominations (37–39), and that trichloromethyl radicals are not the abstracting species in the aralkyl hydrocarbon retarded telomerization of 1-hexadecene and CCl<sub>4</sub> (38).

This article has been cited by:

1. Karen M. Schaich. Lipid Oxidation: New Perspectives on an Old Reaction 1-72. [[Crossref](#)]
2. Jan Honziček. 2019. Curing of Air-Drying Paints: A Critical Review. *Industrial & Engineering Chemistry Research* **58**:28, 12485-12505. [[Crossref](#)]
3. Andrea Baschieri, Romeo Pizzol, Yafang Guo, Riccardo Amorati, Luca Valgimigli. 2019. Calibration of Squalene, p -Cymene, and Sunflower Oil as Standard Oxidizable Substrates for Quantitative Antioxidant Testing. *Journal of Agricultural and Food Chemistry* **67**:24, 6902-6910. [[Crossref](#)]
4. Galina F. Fedorova, Viktoryia A. Lapina, Valery A. Menshov, Vladimir V. Naumov, Aleksei V. Trofimov, Yury B. Tsaplev, Rostislav F. Vasil'ev, Olga I. Yablonskaya. 2019. Updating the Chemiluminescence Oxygen-Aftereffect Method for Determining the Rate Constant of the Peroxy-Radical Self-Reaction: Oxidation of Cyclohexene. *Photochemistry and Photobiology* **95**:3, 780-786. [[Crossref](#)]
5. Shunji Kato, Naoki Shimizu, Yasuhiko Hanzawa, Yurika Otoki, Junya Ito, Fumiko Kimura, Susumu Takekoshi, Masayoshi Sakaino, Takashi Sano, Takahiro Eitsuka, Teruo Miyazawa, Kiyotaka Nakagawa. 2018. Determination of triacylglycerol oxidation mechanisms in canola oil using liquid chromatography–tandem mass spectrometry. *npj Science of Food* **2**:1. . [[Crossref](#)]
6. Radhika Mallepu, Laxmi Potlapally, Vijaya Lakshmi Gollapalli. 2018. Photo-oxidation of some flavonoids with photochemically generated t -BuO • radicals in a t -BuOH water system using a kinetic approach. *Journal of the Chinese Chemical Society* **65**:10, 1266-1273. [[Crossref](#)]
7. Chrysostomos Chatgililoglu, Vincent W. Bowry. 2018. Why Not Trans? Inhibited Radical Isomerization Cycles and Coupling Chains of Lipids and Alkenes with Alkane - thiols. *The Journal of Organic Chemistry* **83**:16, 9178-9189. [[Crossref](#)]
8. Etsuo Niki. 2018. Oxidant-specific biomarkers of oxidative stress. Association with atherosclerosis and implication for antioxidant effects. *Free Radical Biology and Medicine* **120**, 425-440. [[Crossref](#)]
9. Katarzyna Jodko-Piórecka, Jakub Cedrowski, Grzegorz Litwinienko. Physico-chemical principles of antioxidant action, including solvent and matrix dependence and interfacial phenomena 225-272. [[Crossref](#)]
10. L. Dubrulle, R. Lebeuf, M. Fressancourt-Collinet, V. Nardello-Rataj. 2017. Optimization of the vegetable oil composition in alkyd resins: A kinetic approach based on FAMEs autoxidation. *Progress in Organic Coatings* **112**, 288-294. [[Crossref](#)]
11. I. V. Tikhonov, I. V. Moskalenko, E. M. Pliss, M. A. Fomich, A. V. Bekish, V. V. Shmanai. 2017. Kinetic isotope H/D effect in the oxidation of ethers of linoleic acid in solutions. *Russian Journal of Physical Chemistry B* **11**:3, 395-399. [[Crossref](#)]
12. Nan Huang, Ting Wang, Wen-Long Wang, Qian-Yuan Wu, Ang Li, Hong-Ying Hu. 2017. UV/chlorine as an advanced oxidation process for the degradation of benzalkonium chloride: Synergistic effect, transformation products and toxicity evaluation. *Water Research* **114**, 246-253. [[Crossref](#)]
13. Karen M. Schaich. 17 Rethinking Lipid Oxidation 477-498. [[Crossref](#)]
14. Zosia A. M. Zielinski, Derek A. Pratt. 2017. Lipid Peroxidation: Kinetics, Mechanisms, and Products. *The Journal of Organic Chemistry* **82**:6, 2817-2825. [[Crossref](#)]
15. Riccardo Amorati, Andrea Baschieri, Luca Valgimigli. 2017. Measuring Antioxidant Activity in Bioorganic Samples by the Differential Oxygen Uptake Apparatus: Recent Advances. *Journal of Chemistry* **2017**, 1-12. [[Crossref](#)]
16. Chenyu Shen, Rongsong Zhu, Guangqun Zhai. 2016. Fluorescent oligo(N-phenylmaleimide)s via aerobic radical telomerization initiated by benzylic hydrocarbons: Catalytic effect of Co II /N-hydroxyphthalimide pair. *Journal of Polymer Science Part A: Polymer Chemistry* **54**:24, 3846-3857. [[Crossref](#)]
17. Ron Shah, Derek A. Pratt. 2016. Determination of Key Hydrocarbon Autoxidation Products by Fluorescence. *The Journal of Organic Chemistry* **81**:15, 6649-6656. [[Crossref](#)]
18. Evan A. Haidasz, Antonius T. M. Van Kessel, Derek A. Pratt. 2016. A Continuous Visible Light Spectrophotometric Approach To Accurately Determine the Reactivity of Radical-Trapping Antioxidants. *The Journal of Organic Chemistry* **81**:3, 737-744. [[Crossref](#)]
19. Jean-Philippe R. Chauvin, Evan A. Haidasz, Markus Griesser, Derek A. Pratt. 2016. Polysulfide-1-oxides react with peroxy radicals as quickly as hindered phenolic antioxidants and do so by a surprising concerted homolytic substitution. *Chemical Science* **7**:10, 6347-6356. [[Crossref](#)]
20. Adam H. Metherel, Ken D. Stark. 2016. The stability of blood fatty acids during storage and potential mechanisms of degradation: A review. *Prostaglandins, Leukotrienes and Essential Fatty Acids* **104**, 33-43. [[Crossref](#)]
21. Paul M. Rawson, Christy-Anne Stansfield, Renée L. Webster, David Evans, Ulas Yildirim. 2015. The oxidative stability of synthetic fuels and fuel blends with monoaromatic blending components. *Fuel* **161**, 97-104. [[Crossref](#)]

22. Verena Weidmann, Serge Kliewer, Marko Sick, Sergej Bycinskij, Margarethe Kleczka, Julia Rehbein, Axel G. Griesbeck, Holger Zorn, Wolfgang Maison. 2015. Studies towards the synthetic applicability of biocatalytic allylic oxidations with the lyophilisate of *Pleurotus sapidus*. *Journal of Molecular Catalysis B: Enzymatic* **121**, 15-21. [[Crossref](#)]
23. L. Xu, N. A. Porter. 2015. Free radical oxidation of cholesterol and its precursors: Implications in cholesterol biosynthesis disorders. *Free Radical Research* **49**:7, 835-849. [[Crossref](#)]
24. Chad R. Borges, Joshua W. Jeffs, Erandi P. Kapuruge. **1200**, 375. [[Crossref](#)]
25. R. A. Sakhautdinova, A. R. Gimadieva, L. R. Yakupova, I. M. Borisov, R. L. Safiullin. 2015. Kinetics of the radical-chain oxidation of methyl oleate inhibited by 5-amino-6-methyluracil and 5-amino-1,3,6-trimethyluracil. *Kinetics and Catalysis* **56**:2, 125. [[Crossref](#)]
26. K. U. Ingold. 60 Years of Research on Free Radical Physical Organic Chemistry 223-250. [[Crossref](#)]
27. Octavie Okamba-Diogo, Emmanuel Richaud, Jacques Verdu, François Fernagut, Jean Guilment, Bruno Fayolle. 2014. Molecular and macromolecular structure changes in polyamide 11 during thermal oxidation. *Polymer Degradation and Stability* **108**, 123-132. [[Crossref](#)]
28. Keith U. Ingold, Derek A. Pratt. 2014. Advances in Radical-Trapping Antioxidant Chemistry in the 21st Century: A Kinetics and Mechanisms Perspective. *Chemical Reviews* **114**:18, 9022-9046. [[Crossref](#)]
29. Libin Xu, Ned A. Porter. 2014. Reactivities and Products of Free Radical Oxidation of Cholestadienols. *Journal of the American Chemical Society* **136**:14, 5443-5450. [[Crossref](#)]
30. Jing Xiao, Luoming Wu, Ying Wu, Bing Liu, Lu Dai, Zhong Li, Qibin Xia, Hongxia Xi. 2014. Effect of gasoline composition on oxidative desulfurization using a phosphotungstic acid/activated carbon catalyst with hydrogen peroxide. *Applied Energy* **113**, 78-85. [[Crossref](#)]
31. Ned A. Porter. 2013. A Perspective on Free Radical Autoxidation: The Physical Organic Chemistry of Polyunsaturated Fatty Acid and Sterol Peroxidation. *The Journal of Organic Chemistry* 130409110713002. [[Crossref](#)]
32. Ken Akamatsu. 2013. Development of 'leaky' liposome triggered by radiation applicable to a drug reservoir and a simple radiation dosimeter. *Applied Radiation and Isotopes* **74**, 144-151. [[Crossref](#)]
33. Sharwatie Ramsaywack, Christopher M. Vogels, S. Laurie Ricker, Stephen A. Westcott, L. Ross C. Barclay. 2013. Pyridinones Are Not Antioxidants As Shown by Kinetics of Free Radical Autoxidation, but They Prevent Radical Oxidations Catalyzed by Toxic Heavy Metals. *Chemical Research in Toxicology* **26**:3, 399-409. [[Crossref](#)]
34. K.M. Schaich. Challenges in Elucidating Lipid Oxidation Mechanisms 1-52. [[Crossref](#)]
35. Karen M. Schaich, Fereidoon Shahidi, Ying Zhong, N. A. Michael Eskin. Lipid Oxidation 419-478. [[Crossref](#)]
36. Adam H Metherel, Ryan C Hogg, Lindy M Buzikievich, Ken D Stark. 2013. Butylated hydroxytoluene can protect polyunsaturated fatty acids in dried blood spots from degradation for up to 8 weeks at room temperature. *Lipids in Health and Disease* **12**:1, 22. [[Crossref](#)]
37. Verena Weidmann, Mathias Schaffrath, Holger Zorn, Julia Rehbein, Wolfgang Maison. 2013. Elucidation of the regio- and chemoselectivity of enzymatic allylic oxidations with *Pleurotus sapidus* – conversion of selected spirocyclic terpenoids and computational analysis. *Beilstein Journal of Organic Chemistry* **9**, 2233-2241. [[Crossref](#)]
38. Etsuo Niki. Lipid Peroxidation . [[Crossref](#)]
39. L. R. Yakupova, R. L. Safiullin. 2011. Kinetics of the initiated and inhibited oxidation of methyl oleate in homogeneous and aqueous emulsion media. *Kinetics and Catalysis* **52**:6, 785-792. [[Crossref](#)]
40. Huiyong Yin, Libin Xu, Ned A. Porter. 2011. Free Radical Lipid Peroxidation: Mechanisms and Analysis. *Chemical Reviews* **111**:10, 5944-5972. [[Crossref](#)]
41. Derek A. Pratt, Keri A. Tallman, Ned A. Porter. 2011. Free Radical Oxidation of Polyunsaturated Lipids: New Mechanistic Insights and the Development of Peroxyl Radical Clocks. *Accounts of Chemical Research* **44**:6, 458-467. [[Crossref](#)]
42. M. M. Silaev. 2011. Competition Kinetics of the Nonbranched-Chain Addition of Free Radicals to Olefins, Formaldehyde, and Oxygen. *International Journal of Chemical Engineering* **2011**, 1-19. [[Crossref](#)]
43. Riccardo Amorati, Orazio A. Attanasi, Gianfranco Favi, Stefano Menichetti, Gian Franco Pedulli, Caterina Viglianisi. 2011. Amphiphilic antioxidants from "cashew nut shell liquid" (CNSL) waste. *Organic & Biomolecular Chemistry* **9**:5, 1352. [[Crossref](#)]
44. Akio Fujisawa, Walter C. Dunlap, Yorihiro Yamamoto. 2010. Vitamin E protection in the biochemical adaptation of marine organisms to cold-water environments. *Comparative Biochemistry and Physiology Part B: Biochemistry and Molecular Biology* **157**:2, 145-158. [[Crossref](#)]

45. G. Vijayalakshmi, M. Adinarayana, P. Jayaprakash Rao, M. N. Kumara, D. G. Bhadregowda, K. S. Rangappa. 2010. Retracted: Kinetics of oxidation of adenosine by tert-butoxyl radical: protection and repair by chlorogenic acid. *Journal of Physical Organic Chemistry* **23**:10, 1017-1017. [[Crossref](#)]
46. Ivan Tichonov, Vitaly Roginsky, Evgeny Pliss. 2010. Natural polyphenols as chain-breaking antioxidants during methyl linoleate peroxidation. *European Journal of Lipid Science and Technology* **112**:8, 887-893. [[Crossref](#)]
47. Keith U. Ingold. The Only Stable Organic Sigma Radicals: Di-tert-Alkyliminoyls 231-244. [[Crossref](#)]
48. S. G. Yunusova, L. R. Yakupova, A. V. Ivanova, R. L. Safiullin, E. G. Galkin, M. S. Yunusov. 2010. Fatty acid composition of *Oenothera biennis* seed oil during storage. Antioxidant activity. *Chemistry of Natural Compounds* **46**:2, 278-282. [[Crossref](#)]
49. Stefano Menichetti, Riccardo Amorati, Maria Grazia Bartolozzi, Gian Franco Pedulli, Antonella Salvini, Caterina Vigliani. 2010. A Straightforward Hetero-Diels-Alder Approach to (2- ambo ,4' R ,8' R )- $\alpha/\beta/\gamma/\delta$ -4-Thiatocopherol. *European Journal of Organic Chemistry* **2010**:11, 2218-2225. [[Crossref](#)]
50. Xavier Colin, Emmanuel Richaud, Jacques Verdu, Carole Monchy-Leroy. 2010. Kinetic modelling of radiochemical ageing of ethylene-propylene copolymers. *Radiation Physics and Chemistry* **79**:3, 365-370. [[Crossref](#)]
51. Libin Xu, Zeljka Korade, Ned A. Porter. 2010. Oxysterols from Free Radical Chain Oxidation of 7-Dehydrocholesterol: Product and Mechanistic Studies. *Journal of the American Chemical Society* **132**:7, 2222-2232. [[Crossref](#)]
52. Di Hu, Derek A. Pratt. 2010. Secondary orbital interactions in the propagation steps of lipid peroxidation. *Chemical Communications* **46**:21, 3711. [[Crossref](#)]
53. S. Sarrabi, X. Colin, A. Tcharkhtchi. 2010. Kinetic modeling of polypropylene thermal oxidation during its processing by rotational molding. *Journal of Applied Polymer Science* **5**, n/a-n/a. [[Crossref](#)]
54. Ronny Neumann\*, Alexander M. Khenkin. Oxidation of Dienes and Polyenes . [[Crossref](#)]
55. Mario C. Foti, Riccardo Amorati. 2009. Non-phenolic radical-trapping antioxidants. *Journal of Pharmacy and Pharmacology* **61**:11, 1435-1448. [[Crossref](#)]
56. Jia Yao, Dushyanth K. Dokuru, Matthew Noestheden, SungAe S. Park, Bruce A. Kerwin, Janan Jona, Drazen Ostovic, Darren L. Reid. 2009. A Quantitative Kinetic Study of Polysorbate Autoxidation: The Role of Unsaturated Fatty Acid Ester Substituents. *Pharmaceutical Research* **26**:10, 2303-2313. [[Crossref](#)]
57. Libin Xu, Todd A. Davis, Ned A. Porter. 2009. Rate Constants for Peroxidation of Polyunsaturated Fatty Acids and Sterols in Solution and in Liposomes. *Journal of the American Chemical Society* **131**:36, 13037-13044. [[Crossref](#)]
58. S. Sarrabi, X. Colin, A. Tcharkhtchi. 2008. Isothermal oxidation kinetics of unstabilized polypropylene in the molten state. *Journal of Applied Polymer Science* **110**:4, 2030-2040. [[Crossref](#)]
59. Pascal Lignier, Stéphane Mangematin, Franck Morfin, Jean-Luc Rousset, Valérie Caps. 2008. Solvent and oxidant effects on the Au/TiO<sub>2</sub>-catalyzed aerobic epoxidation of stilbene. *Catalysis Today* **138**:1-2, 50-54. [[Crossref](#)]
60. Yuko Minami, Kanako Yokoyama, Noriko Bando, Yoshichika Kawai, Junji Terao. 2008. Occurrence of singlet oxygen oxygenation of oleic acid and linoleic acid in the skin of live mice. *Free Radical Research* **42**:3, 197-204. [[Crossref](#)]
61. S. Sarrabi, X. Colin, A. Tcharkhtchi. 2008. Dégradation thermique du polypropylène au cours du rotomoulage. *Matériaux & Techniques* **96**:6, 281-294. [[Crossref](#)]
62. Malgorzata Musialik, Marcin Kita, Grzegorz Litwinienko. 2008. Initiation of lipid autoxidation by ABAP at pH 4-10 in SDS micelles. *Organic & Biomolecular Chemistry* **6**:4, 677. [[Crossref](#)]
63. M. M. Silaev. 2007. Simulation of the initiated addition of hydrocarbon free radicals and hydrogen atoms to oxygen via a nonbranched chain mechanism. *Theoretical Foundations of Chemical Engineering* **41**:6, 831-838. [[Crossref](#)]
64. L. P. Stepovik, M. V. Gulenova, A. N. Tishkina, V. K. Cherkasov. 2007. System vanadium alkoxy compound-tert-butyl hydroperoxide-oxidant of hydrocarbon C-H bonds. *Russian Journal of General Chemistry* **77**:7, 1254-1262. [[Crossref](#)]
65. Levon Tavadyan, Avet Khachoyan, Gagik Martoyan, Afaf Kamal-Eldin. 2007. Numerical revelation of the kinetic significance of individual steps in the reaction mechanism of methyl linoleate peroxidation inhibited by  $\alpha$ -tocopherol. *Chemistry and Physics of Lipids* **147**:1, 30-45. [[Crossref](#)]
66. A. V. Chekanov, A. N. Osipov, Yu. A. Vladimirov, V. I. Sergienko, O. M. Panasencko. 2007. A comparative spin trapping study of the interaction of hypobromous and hypochlorous acids with tert-butyl hydroperoxide. *Biophysics* **52**:1, 1-7. [[Crossref](#)]
67. Yoshichika Kawai, Mariko Miyoshi, Jae-Hak Moon, Junji Terao. 2007. Detection of cholesteryl ester hydroperoxide isomers using gas chromatography-mass spectrometry combined with thin-layer chromatography blotting. *Analytical Biochemistry* **360**:1, 130-137. [[Crossref](#)]



68. John D. Hearn, Lindsay H. Renbaum, Xi Wang, Geoffrey D. Smith. 2007. Kinetics and products from reaction of Cl radicals with dioctyl sebacate (DOS) particles in O<sub>2</sub>: a model for radical-initiated oxidation of organic aerosols. *Physical Chemistry Chemical Physics* **9**:34, 4803. [[Crossref](#)]
69. X. Colin, B. Fayolle, L. Audouin, J. Verdu. 2006. The classical kinetic model for radical chain oxidation of hydrocarbon substrates initiated by bimolecular hydroperoxide decomposition. *International Journal of Chemical Kinetics* **38**:11, 666-676. [[Crossref](#)]
70. L. Charitha, M. Adinarayana. 2005. Kinetics of oxidation of adenosine byt-butoxyl radical: Protection and repair by caffeic acid. *International Journal of Chemical Kinetics* **37**:9, 515-521. [[Crossref](#)]
71. K. M. Schaich. Lipid Oxidation: Theoretical Aspects . [[Crossref](#)]
72. V. Bellenger, J. Decelle, N. Huet. 2005. Ageing of a carbon epoxy composite for aeronautic applications. *Composites Part B: Engineering* **36**:3, 189-194. [[Crossref](#)]
73. L. P. Stepovik, M. V. Gulenova, I. M. Martynova. 2005. Titanium(IV) tert-Butoxide-tert-Butyl Hydroperoxide System as Oxidant for C-H Bonds in Hydrocarbons and Oxygen-containing Compounds. *Russian Journal of General Chemistry* **75**:4, 507-513. [[Crossref](#)]
74. . Comments on Contemporary Methods of Oxygen and Nitrogen Free Radical Detection 1-284. [[Crossref](#)]
75. . Oxidation of Polymers 1-454. [[Crossref](#)]
76. Huiyong Yin, Ned A. Porter. 2005. New Insights Regarding the Autoxidation of Polyunsaturated Fatty Acids. *Antioxidants & Redox Signaling* **7**:1-2, 170-184. [[Crossref](#)]
77. Noriko Bando, Hiroki Hayashi, Saori Wakamatsu, Takahiro Inakuma, Mariko Miyoshi, Akihiko Nagao, Ryo Yamauchi, Junji Terao. 2004. Participation of singlet oxygen in ultraviolet-a-induced lipid peroxidation in mouse skin and its inhibition by dietary  $\beta$ -carotene: an ex vivo study. *Free Radical Biology and Medicine* **37**:11, 1854-1863. [[Crossref](#)]
78. Sergiu Coseri, Keith U. Ingold. 2004. Distinguishing between Abstraction and Addition as the First Step in the Reaction of a Nitroxyl Radical with Cyclohexene. *Organic Letters* **6**:10, 1641-1643. [[Crossref](#)]
79. Afaf Kamal-Eldin, Marjukka Mäkinen, Anna-Maija Lampi. The Challenging Contribution of Hydroperoxides to the Lipid Oxidation Mechanism . [[Crossref](#)]
80. Hayato SABA, Masuo NAKANO. 2003. Oxidation of Triacylglycerol by Laccase-Mediator Systems. *Journal of Oleo Science* **52**:1, 15-23. [[Crossref](#)]
81. Ryouta Maeba, Youltuz Yusufu, Hiroyuki Shimasaki, Nobuo Ueta. 2002. Comparison of the oxidizability of various glycerophospholipids in bilayers by the oxygen uptake method. *Lipids* **37**:9, 893-900. [[Crossref](#)]
82. Xing-Hua Wang, Toshiaki Ohshima, Hideki Ushio, Chiaki Koizumi. 1999. Proportion of geometrical hydroperoxide isomers generated by radical oxidation of methyl linoleate in homogeneous solution and in aqueous emulsion. *Lipids* **34**:7, 675-679. [[Crossref](#)]
83. Gian Franco Pedulli, Marco Lucarini, Emanuela Marchesi, Francesco Paolucci, Sergio Roffia, Diana Fiorentini, Laura Landi. 1999. Medium effects on the antioxidant activity of dipyridamole. *Free Radical Biology and Medicine* **26**:3-4, 295-302. [[Crossref](#)]
84. Gadi Rothenberg, Yoel Sasson. 1999. Application of perturbation theory to free-radical benzylic and allylic oxidation of unconjugated  $\pi$ -systems. *Tetrahedron* **55**:2, 561-568. [[Crossref](#)]
85. Catherine Belzung, Anne-Marie Leguisquet, Serge Barreau, Sylvie Delion-Vancassel, Sylvie Chalon, Georges Durand. 1998.  $\alpha$ -Linolenic Acid Deficiency Modifies Distractibility but Not Anxiety and Locomotion in Rats during Aging. *The Journal of Nutrition* **128**:9, 1537-1542. [[Crossref](#)]
86. Kazumasa Yazu, Yorihiro Yamamoto, Etsuo Niki, Keiji Miki, Koji Ukegawa. 1998. Mechanism of lower oxidizability of eicosapentaenoate than linoleate in aqueous micelles. II. Effect of antioxidants. *Lipids* **33**:6, 597-600. [[Crossref](#)]
87. Gadi Rothenberg, Yoel Sasson. 1998. Cyclic vs. acyclic allylic hydrogen abstraction: An entropy motivated process?. *Tetrahedron* **54**:20, 5417-5422. [[Crossref](#)]
88. B.Vernon Cheney. 1998. AM1 calculations on reactive oxygen species Part 3. Stereochemistry of hydroperoxide products obtained from oxidation of linoleic acid. *Journal of Molecular Structure: THEOCHEM* **424**:3, 207-223. [[Crossref](#)]
89. Božena Lánská. 1998. Termination of peroxy radicals of 6-hexanelactam. *Polymer Degradation and Stability* **61**:1, 27-31. [[Crossref](#)]
90. Kazuhiro KUBO, Morio SAITO, Tadahiro TADOKORO, Akio MAEKAWA. 1998. N-3 fatty acid intake and lipid peroxidation (4). Dietary Docosahexaenoic Acid Dose Not Promote Lipid Peroxidation in Rat Tissue to the Extent Expected from Peroxidizability Index of the Lipids. *Bioscience, Biotechnology, and Biochemistry* **62**:9, 1698-1706. [[Crossref](#)]
91. Gadi Rothenberg, Yael Yatziv, Yoel Sasson. 1998. Comparative autoxidation of 3-Carene and  $\alpha$ -Pinene: Factors governing regioselective hydrogen abstraction reactions. *Tetrahedron* **54**:3-4, 593-598. [[Crossref](#)]

92. Nagahiko Sakuma, Seiji Iwata, Takeshi Hibino, Nozomu Tamai, Kanna Sasai, Takayuki Yoshimata, Yoshinobu Kamiya, Masanobu Kawaguchi, Takao Fujinami. 1997. Effects of vitamin C and vitamin E on plasma levels of lipid hydroperoxides and thiobarbituric acid reactive substance in humans. *Current Therapeutic Research* **58**:5, 317-322. [[Crossref](#)]
93. Hiroyuki Haraguchi, Harumi Ishikawa, Yolanda Sanchez, Tetsuya Ogura, Yumi Kubo, Isao Kubo. 1997. Antioxidative constituents in *Heterotheca inuloides*. *Bioorganic & Medicinal Chemistry* **5**:5, 865-871. [[Crossref](#)]
94. H. Haraguchi, H. Ishikawa, S. Sakai, B. -P. Ying, I. Kubo. 1996. Inhibition of lipid peroxidation by diterpenoid from *Podocarpus nagi*. *Experientia* **52**:6, 573-576. [[Crossref](#)]
95. B. Vernon Cheney. 1996. AM1 calculations on reactive oxygen species. Part 2. Kinetics of lipid peroxidation based upon approximate rate constants from scaled energy barriers. *Journal of Molecular Structure: THEOCHEM* **364**:2-3, 239-270. [[Crossref](#)]
96. Kazumasa Yazu, Yorihiro Yamamoto, Koji Ukegawa, Etsuo Niki. 1996. Mechanism of lower oxidizability of eicosapentaenoate than linoleate in aqueous micelles. *Lipids* **31**:3, 337-340. [[Crossref](#)]
97. Fernando Antunes, Armindo Salvador, H. Susana Marinho, Rui Alves, Ruy E. Pinto. 1996. Lipid peroxidation in mitochondrial inner membranes. I. An integrative kinetic model. *Free Radical Biology and Medicine* **21**:7, 917-943. [[Crossref](#)]
98. Gian Franco Pedulli, Marco Lucarini, Pamela Pedrielli, Monica Sagrini, Marta Cipollone. 1996. The determination of the oxygen consumption in autoxidation studies by means of EPR spectroscopy. *Research on Chemical Intermediates* **22**:1, 1-14. [[Crossref](#)]
99. V. A. Pluzhnikov, G. A. Kovtun. 1996. Chain termination catalyzed by 1,4-benzoquinones in the oxidation of 9,10-dihydrophenanthrene. *Theoretical and Experimental Chemistry* **32**:1, 30-33. [[Crossref](#)]
100. Olga A. Ozhogina, Olga T. Kasaikina. 1995.  $\beta$ -carotene as an interceptor of free radicals. *Free Radical Biology and Medicine* **19**:5, 575-581. [[Crossref](#)]
101. Michael J. Thomas. 1995. The role of free radicals and antioxidants: How do we know that they are working?. *Critical Reviews in Food Science and Nutrition* **35**:1-2, 21-39. [[Crossref](#)]
102. Cheves Walling. Autoxidation 24-65. [[Crossref](#)]
103. Takafumi Nonogaki, Yoichi Noda, Yasuo Goto, Junji Kishi, Takahide Mori. 1994. Developmental blockage of mouse embryos caused by fatty acids. *Journal of Assisted Reproduction and Genetics* **11**:9, 482-488. [[Crossref](#)]
104. G. A. Kovtun, V. A. Pluzhnikov. 1994. The role of chelate substituent steric effects in reactions between hydroperoxyl radicals and copper bis(salicylalalkyliminates). *Theoretical and Experimental Chemistry* **29**:1, 51-53. [[Crossref](#)]
105. DANIEL C. LIEBLER. 1993. Antioxidant Reactions of Carotenoids. *Annals of the New York Academy of Sciences* **691**:1 Carotenoids i, 20-31. [[Crossref](#)]
106. Daniel C. Liebler. 1993. The Role of Metabolism in the Antioxidant Function of Vitamin E. *Critical Reviews in Toxicology* **23**:2, 147-169. [[Crossref](#)]
107. G. Vasv $\ddot{\text{u}}$ ri, D. G $\ddot{\text{u}}$ ll. 1993. Self-Reaction of Benzylperoxy Radicals in the Oxidation of Toluene. A Chemiluminescence and Product Study. *Berichte der Bunsengesellschaft f $\ddot{\text{u}}$ r physikalische Chemie* **97**:1, 22. [[Crossref](#)]
108. Bernard Le Tutour, Didier Guedon. 1992. Antioxidative activities of *Olea europaea* leaves and related phenolic compounds. *Phytochemistry* **31**:4, 1173-1178. [[Crossref](#)]
109. Ágnes Keszler, György Irinyi, Károly Héberger, Dezső Gál. 1992. Reactivity of HO $_2$  $\cdot$  Radicals in Alcohols. *Berichte der Bunsengesellschaft für physikalische Chemie* **96**:2, 175-179. [[Crossref](#)]
110. Rajkumar Kumarathasan, Amirithini B. Rajkumar, Norman R. Hunter, Hyman D. Gesser. 1992. Autoxidation and yellowing of methyl linolenate. *Progress in Lipid Research* **31**:2, 109-126. [[Crossref](#)]
111. Vladimir Andreevich Belyakov, Gerlinde Lauterbach, Wilhelm Pritzkow, Volkmar Voerckel. 1992. Kinetics and regioselectivity of the Autoxidation of Alkylaromatic Hydrocarbons. *Journal für Praktische Chemie/Chemiker-Zeitung* **334**:5, 373-382. [[Crossref](#)]
112. Kunihiro Tokumura, Hideyo Nosaka, Yoshihisa Fujiwara. 1991. Photodeoxygenation of transient p-methoxybenzyl-peroxy radical in solution at room temperature. *Chemical Physics Letters* **177**:6, 559-562. [[Crossref](#)]
113. Noboru Takekoshi, Eiji Murakami, Shinji Ohkubo. 1991. Effects of some radical scavengers on reperfusion-induced arrhythmias in the canine heart. *Cardiovascular Drugs and Therapy* **5**:S2, 289-295. [[Crossref](#)]
114. Wilhelm Pritzkow, Tarigopula Siva Sankara Rao. 1990. The Autoxidation of Acetylenic Hydrocarbons. *Zeitschrift für Chemie* **30**:12, 422-427. [[Crossref](#)]
115. Bernard Le Tutour. 1990. Antioxidative activities of algal extracts, synergistic effect with vitamin E. *Phytochemistry* **29**:12, 3759-3765. [[Crossref](#)]
116. S. L. Khursan, V. S. Martemyanov. 1989. Influence of substituents on rate constants for recombination of tertiary peroxy radicals. *Reaction Kinetics and Catalysis Letters* **40**:2, 253-258. [[Crossref](#)]

117. Matilde Maiorino, Mariagrazia Coassin, Antonella Roveri, Fulvio Ursini. 1989. Microsomal lipid peroxidation: Effect of vitamin E and its functional interaction with phospholipid hydroperoxide glutathione peroxidase. *Lipids* **24**:8, 721-726. [[Crossref](#)]
118. R.J. Hsieh, J.E. Kinsella. Oxidation of Polyunsaturated Fatty Acids: Mechanisms, Products, and Inhibition with Emphasis on Fish 233-341. [[Crossref](#)]
119. Ned A. Porter, Dennis G. Wujek. The Autoxidation of Polyunsaturated Lipids 55-79. [[Crossref](#)]
120. Wolf Bors, M. Erben-Russ, M. Saran. 1987. Fatty acid peroxy radicals: Their generation and reactivities. *Bioelectrochemistry and Bioenergetics* **18**:1-3, 37-49. [[Crossref](#)]
121. A. Rockenbauer, M. Györ. 1987. Electron spin resonance detection of oscillating peroxy radical concentration in the cumene—anthraquinone system. *Journal of Photochemistry* **38**, 109-116. [[Crossref](#)]
122. John P. Cosgrove, Daniel F. Church, William A. Pryor. 1987. The kinetics of the autoxidation of polyunsaturated fatty acids. *Lipids* **22**:5, 299-304. [[Crossref](#)]
123. Toru Takagi, Kazuo Miyashita. 1987. Autoxidative rates of nonmethylene-interrupted polyenoic fatty acids. *Journal of the American Oil Chemists' Society* **64**:3, 407-413. [[Crossref](#)]
124. Michael Erben-Russ, Wolf Bors, Manfred Saran. 1987. Reactions of Linoleic Acid Peroxyl Radicals with Phenolic Antioxidants: A Pulse Radiolysis Study. *International Journal of Radiation Biology and Related Studies in Physics, Chemistry and Medicine* **52**:3, 393-412. [[Crossref](#)]
125. Yorihiro Yamamoto, Michael H. Brodsky, Jeffrey C. Baker, Bruce N. Ames. 1987. Detection and characterization of lipid hydroperoxides at picomole levels by high-performance liquid chromatography. *Analytical Biochemistry* **160**:1, 7-13. [[Crossref](#)]
126. A. Fautitano, A. Buttafava, F. Martinotti, V. Comincioli, P. Bortolus. 1986. Decay constants of peroxy radicals from polypropylene and polypropylene model compounds: A kinetic ESR study. *Polymer Photochemistry* **7**:6, 491-502. [[Crossref](#)]
127. Wilhelm Pritzkow, Gerda Thomas, Lothar Willecke. 1985. Studies on the Regioselectivities in the Oxidation of Simple Alkyl Aromatic Hydrocarbons by Molecular Oxygen. *Journal für Praktische Chemie* **327**:5, 847-851. [[Crossref](#)]
128. Yamamoto Yorihiro, Nika Etsuo, Kamiya Yoshio, Shimasaki Hiroyuki. 1984. Oxidation of lipids. *Biochimica et Biophysica Acta (BBA) - Lipids and Lipid Metabolism* **795**:2, 332-340. [[Crossref](#)]
129. Michael P. Doyle, Jan W. Terpstra, Charles H. Winter, John H. Griffin. 1984. Rhodium(II) acetate catalyzed hydrocarbon oxidations by molecular oxygen. *Journal of Molecular Catalysis* **26**:2, 259-266. [[Crossref](#)]
130. Ned A. Porter. [32] Chemistry of lipid peroxidation 273-282. [[Crossref](#)]
131. Yamamoto Yorihiro, Nika Etsuo, Kamiya Yoshio, Shimasaki Hiroyuki, Paul K. Schick, Barbara P. Schick, Karen Foster, Andrew Block, Phillip R. Green, Robert M. Bell. 1984. Oxidation of lipids. *Biochimica et Biophysica Acta (BBA) - Lipids and Lipid Metabolism* **795**:2, 332. [[Crossref](#)]
132. F. Haslbeck, W. Grosch. 1984. Bildung von Hydroperoxy-epidioxiden bei der Autoxidation von Linolsäurephenylester. *Fette, Seifen, Anstrichmittel* **86**:10, 408-413. [[Crossref](#)]
133. Franz Haslbeck, Werner Grosch. 1983. Autoxidation of phenyl linoleate and phenyl oleate: HPLC analysis of the major and minor monohydroperoxides as phenyl hydroxystearates. *Lipids* **18**:10, 706-713. [[Crossref](#)]
134. J. Lukács, K. Héberger, D. Gál. 1983. Rate Coefficients of Elementary Steps in the Inhibited Oxidation of Ethylbenzene, I. Interaction between Peroxy Radicals and Inhibitor Molecules. *Berichte der Bunsengesellschaft für physikalische Chemie* **87**:7, 606-609. [[Crossref](#)]
135. M. Futu-Tangu, E. Lawson, W. Pritzkow, V. Voerckel. 1983. Kinetische Untersuchungen über die Autoxidation von Olefinen. *Journal für Praktische Chemie* **325**:4, 545-550. [[Crossref](#)]
136. Yorihiro Yamamoto, Etsuo Niki, Yoshio Kamiya. 1982. Oxidation of lipids: III. Oxidation of methyl linoleate in solution. *Lipids* **17**:12, 870-877. [[Crossref](#)]
137. Yorihiro Yamamoto, Etsuo Niki, Yoshio Kamiya. 1982. Oxidation of Lipids. I. Quantitative Determination of the Oxidation of Methyl Linoleate and Methyl Linolenate. *Bulletin of the Chemical Society of Japan* **55**:5, 1548-1550. [[Crossref](#)]
138. G. Moad, E. Rizzardo, D. H. Solomon. 1982. The Reaction of Benzoyloxy Radicals with Styrene—Implications Concerning the Structure of Polystyrene. *Journal of Macromolecular Science: Part A - Chemistry* **17**:1, 51-59. [[Crossref](#)]
139. K. Blau, W. Pritzkow. 1982. Über die Autoxidation einiger asymmetrischer p-Dialkylbenzene. *Journal für Praktische Chemie* **324**:1, 12-14. [[Crossref](#)]
140. William A. Pryor, Donald G. Prier, Daniel F. Church. 1981. Radical production from the interaction of ozone and PUFA as demonstrated by electron spin resonance spin-trapping techniques. *Environmental Research* **24**:1, 42-52. [[Crossref](#)]



141. G. A. Kovtun, D. L. Lysenko, A. S. Berenblyum, I. I. Moiseev. 1980. Mechanism of the inhibition of oxidation reactions by metal complexes. *Bulletin of the Academy of Sciences of the USSR Division of Chemical Science* **29**:4, 540-543. [[Crossref](#)]
142. Ned A. Porter, Richard A. Wolf, Hugo Weenen. 1980. The free radical oxidation of polyunsaturated lecithins. *Lipids* **15**:3, 163-167. [[Crossref](#)]
143. Ned A. Porter. Prostaglandin Endoperoxides 261-294. [[Crossref](#)]
144. William A. Pryor, Donald G. Prier, John W. Lightsey, Daniel F. Church. Initiation of the Autoxidation of Polyunsaturated Fatty Acids (PUFA) by Ozone and Nitrogen Dioxide 1-16. [[Crossref](#)]
145. John G. Huntington, Frank R. Mayo, Norman A. Kirshen. 1979. Mild oxidations of coal models. *Fuel* **58**:1, 31-36. [[Crossref](#)]
146. S. Barkin, M. Bixon, R. M. Noyes, K. Bar-eli. 1978. On the oxidation of malonic acid by ceric ions. *International Journal of Chemical Kinetics* **10**:6, 619-636. [[Crossref](#)]
147. Wing-sum Dominic Wong, E. G. Hammond. 1977. Analysis of oleate and linoleate hydroperoxides in oxidized ester mixtures. *Lipids* **12**:6, 475-479. [[Crossref](#)]
148. E. T. Denisov, N. I. Mitskevich, V. E. Agabekov. The Oxidation of Ethers 171-198. [[Crossref](#)]
149. Heinz Kropf, Dietrich Gürtler. 1976. Katalyse durch Phthalocyanine, XIX. Autoxidation von 1,1-Diphenyläthan in Gegenwart von Kupferphthalocyanin und Nickelphtalocyanin. *Justus Liebigs Annalen der Chemie* **1976**:3, 519-548. [[Crossref](#)]
150. Arne Holm, Niels Harrit, Nils Henrik Toubro. 1976. On the mechanism of photolysis of 4-phenyl-1,3,2-oxathiazolylio-5-oxide in ethanol. *Tetrahedron* **32**:21, 2559-2562. [[Crossref](#)]
151. Jan F. Rabek. Chapter 4 Oxidative Degradation of Polymers 425-538. [[Crossref](#)]
152. V. A. Belyakov, V. S. Rafikova. 1974. Determination by the method of chemiluminescence of the rate constant of quadratic recombination of peroxide radicals in reactions with a low luminescence yield. *Bulletin of the Academy of Sciences of the USSR Division of Chemical Science* **23**:11, 2359-2365. [[Crossref](#)]
153. N. Yanishlieva, A. Popov. 1974. Ermittlung der Initiierungsgeschwindigkeit in der Induktionsperiode bei der Autoxydation von ungesättigten Lipidsystemen. *Food / Nahrung* **18**:2, 195-205. [[Crossref](#)]
154. J. A. Maassen, Th. J. de Boer. 1973. C-Nitroso compounds. Part XXVI: Photochemistry of C-nitroso compounds. The photooxidation of ethylbenzene in the presence of 2-methyl-2-nitrosopropane. *Recueil des Travaux Chimiques des Pays-Bas* **92**:2, 185-194. [[Crossref](#)]
155. E. T. Denisov, R. L. Vardanyan. 1972. The role of the hydrogen bond in the reaction of peroxide radicals with inhibitors. *Bulletin of the Academy of Sciences of the USSR Division of Chemical Science* **21**:11, 2396-2399. [[Crossref](#)]
156. W.H. Richardson, H.E. O'Neal. Chapter 4 The Unimolecular Decomposition and Isomerization of Oxygenated Organic Compounds (other than Aldehydes and Ketones) 381-565. [[Crossref](#)]
157. N. Yanishlieva, A. Popov. 1971. Über einige Eigentümlichkeiten in der Kinetik zu Beginn der Autoxydation von Estern ungesättigter Fettsäuren. 3. Mitt. Inhibierte Autoxydation. *Food / Nahrung* **15**:6, 671-681. [[Crossref](#)]
158. J. A. Howard, J. C. Scaiano. 8.5.1.1.17 Polymeric peroxy radicals 241-244. [[Crossref](#)]
159. J. A. Howard, J. C. Scaiano. Part 2 283-292. [[Crossref](#)]
160. J. A. Howard, J. C. Scaiano. 8.5.1.1.1 Alkylperoxy radicals 206-214. [[Crossref](#)]
161. J. A. Howard, J. C. Scaiano. 8.5.3.2 Reactions of alkylperoxy radicals with alkenes 263-273. [[Crossref](#)]
162. J. A. Howard, J. C. Scaiano. References for 8.3 - 8.5 423-430. [[Crossref](#)]
163. J. A. Howard, J. C. Scaiano. Part 1 274-282. [[Crossref](#)]
164. J. A. Howard, J. C. Scaiano. 8.5.3.5 Reactions of alkylperoxy radicals with aralkenes 292-298. [[Crossref](#)]
165. J. A. Howard, J. C. Scaiano. 8.5.1.1.4 Alkylperoxy radicals 219-226. [[Crossref](#)]
166. J. A. Howard, J. C. Scaiano. 8.5.3.1 Reactions of alkylperoxy radicals with alkanes 257-263. [[Crossref](#)]
167. J. A. Howard, J. C. Scaiano. 8.5.1.1.3 Alkynylperoxy radicals 218-218. [[Crossref](#)]
168. J. A. Howard, J. C. Scaiano. 8.5.1.1.18 Group IVB centered peroxy radicals 244-245. [[Crossref](#)]
169. J. A. Howard, J. C. Scaiano. 8.5.1.1.2 Alkenylperoxy radicals 214-218. [[Crossref](#)]
170. Hiroshi Hotta, Nobutake Suzuki, Kazuo Isoda. 1969. The Effect of Organic Additives on the Autoxidation of Benzene and Toluene in the Aqueous Phase and the Kinetic Analysis. *Bulletin of the Chemical Society of Japan* **42**:11, 3056-3064. [[Crossref](#)]
171. Etsuo Niki, Yoshio Kamiya, Nobuto Ohta. 1969. Cooxidation of Hydrocarbons in Liquid Phase. VI. Reactivities of Hydrocarbons and Peroxy Radicals. *Bulletin of the Chemical Society of Japan* **42**:8, 2312-2318. [[Crossref](#)]

172. L. R. Mahoney. 1969. Antioxidantien. *Angewandte Chemie* **81**:15, 555-563. [[Crossref](#)]
173. D. J. Carlsson, K. U. Ingold, L. C. Bray. 1969. Kinetics and rate constants for the reaction of tri-n-butylgermanium hydride with methyl iodide and carbon tetrachloride. The combination of methyl and trichloromethyl radicals in solution. *International Journal of Chemical Kinetics* **1**:3, 315-323. [[Crossref](#)]
174. Keith U. Ingold. 1969. Peroxy radicals. *Accounts of Chemical Research* **2**:1, 1-9. [[Crossref](#)]
175. Kei Sin Wei, Albert H. Adelman. 1969. The photooxidation of toluene. The role of an excited charge transfer complex. *Tetrahedron Letters* **10**:38, 3297-3300. [[Crossref](#)]
176. Frank R. Mayo. 1968. Free radical autoxidations of hydrocarbons. *Accounts of Chemical Research* **1**:7, 193-201. [[Crossref](#)]