Closing stomatal supply against photosynthetic demand at runtime: a verified diel plant growth model and the boundary of its validity

TODO(AUTHOR) Given Name Surname[a]

[a]*TODO(AUTHOR) Institution, TODO(AUTHOR) Address, TODO(AUTHOR) City, TODO(AUTHOR) Country*

ARTICLE INFO

Keywords:
plant growth simulation
diel carbohydrate turnover
Farquhar–Jarvis coupling
verification and validation
variance-based sensitivity analysis
out-of-sample cross-validation

ABSTRACT

Mechanistic models of plant growth usually prescribe carbon input or compute it from a photosynthesis routine that is solved separately from stomatal conductance. This work reports a modular simulation model in which the supply and demand sides of leaf gas exchange are closed against each other at every evaluation: a Jarvis conductance and a Farquhar assimilation function are combined into a quadratic in the intercellular CO_2 concentration and solved in closed form, replacing a fixed-point iteration that we show admits a period-two orbit under water stress. Around this closure the model integrates eleven states covering starch, sucrose and vacuolar sugar, nitrogen and phosphorus acquisition with explicit costs, and leaf and root biomass. Carbon conservation over the carbohydrate subsystem closes to machine precision. Rather than fitting coefficients per experimental condition, we hold a single parameter set across four photoperiods and report what that costs: relative growth rate is reproduced with a mean absolute percentage error of 8.58 % and a worst case of 11.75 %, against a published baseline whose apparent agreement is a bisection residual of eight coefficients fitted to eight observations. Leave-one-photoperiod-out cross-validation gives a held-out error of 13.18 % for a photoperiod-independent growth coefficient and 4.36 % when the coefficient is allowed a linear photoperiod dependence. Diel starch is reproduced in both shape and level; sucrose is reproduced in shape but is a factor two to three low in absolute concentration, which we report rather than absorb. Ablation shows that computing the carbon input, rather than prescribing it, is what carries the photoperiod response. Variance-based sensitivity analysis places 96 % of output variance in parameter interactions rather than main effects. Finally we locate the boundary of the model's validity: the diel carbon balance closes only up to a 15 h photoperiod, and beyond it starch accumulates without bound, a limitation we characterise rather than conceal.

1. Introduction

Arabidopsis thaliana grows in a light environment that switches off every day. During the light phase the plant assimilates carbon and stores part of it as transient leaf starch. During the night this reserve is the only carbon source for respiration, maintenance and growth. Measurements show that the reserve is degraded at an almost constant rate and is nearly exhausted at dawn [20, 51]. The plant therefore behaves as if it rations a known amount of carbon over a night of known length [46]. When the photoperiod changes, both the amount and the length change together, and the rationing problem changes with them [54, 53]. This makes photoperiod a useful test case for a simulation model, because a single environmental variable moves the light-phase carbon gain and the dark-phase reserve use at the same time.

The problem is difficult to simulate because it couples processes on very different time scales. Stomatal opening and leaf assimilation respond within minutes to light, temperature and humidity. Starch and sucrose pools turn over within hours. Nutrient status changes over days and biomass accumulates over weeks. The carbon available for growth at any instant is not a property of any single equation. It is the result of the environment, the gas-exchange machinery and the internal carbohydrate and nutrient state acting together [15, 57]. A model that describes only photosynthesis, or only biomass accumulation, cannot represent this. For a simulation study the main task is therefore to build a stable numerical workflow in which these subsystems exchange fluxes consistently over the day–night cycle.

Existing models resolve parts of this chain, but usually not the whole chain at the resolution the problem needs. Crop and field-scale simulators such as WOFOST, APSIM and DSSAT integrate growth, water and nitrogen over seasons [63, 22, 26]. For robustness and runtime they obtain the carbon input from a light-use-efficiency term, a radiation-use coefficient or a look-up table, and they usually advance the state on a daily step. Soil–plant transport

TODO(AUTHOR) email@institution (T.G.N. Surname)
ORCID(s):

models and functional–structural plant models add detail in other directions, in water transport and in architecture respectively [24, 12, 30]. At the other end of the scale range, leaf gas-exchange models resolve the process that the crop models prescribe. The Farquhar–von Caemmerer–Berry biochemical model and Jarvis-type or optimality-based stomatal formulations describe assimilation and conductance in detail [15, 25, 2, 35], and the coupling between them has been studied extensively [11, 5]. These models are formulated and tested as steady-state algebraic systems at a fixed carbohydrate status, so they stop at the leaf. Few published models compute the supply–demand operating point at every step and then carry that carbon input through a diel starch and sucrose economy into biomass.

This paper presents such a model and reports its evaluation. The model is a whole-plant dynamic simulator with eleven state variables, organised as six submodels with declared inputs, outputs and parameters (Figure 1). S1 maps clock time to the environmental drivers. S2 computes stomatal conductance from a multiplicative Jarvis formulation, which acts as the supply side of gas exchange. S3 computes net assimilation as a function of intercellular CO_2 from the Farquhar C3 biochemistry, which acts as the demand side. S4 closes supply against demand. S5 holds the diel carbohydrate economy, the starch and sucrose pools and the nitrogen and phosphorus uptake and costs. S6 converts sucrose into leaf and root biomass. Submodels S2–S4 form a fast algebraic subsystem, and S5–S6 form the slow differential subsystem that is integrated in time.

The methodological content of the work sits in four places, and this is what we expect to transfer to other coupled biophysical simulators. First, the gas-exchange operating point is solved algebraically at runtime. Substituting Fick's law into the branch-wise Farquhar expression gives a quadratic equation in the intercellular CO_2 concentration. This quadratic is solved branch by branch, with an explicit admissibility rule for the roots. No inner iteration is then required inside the derivative evaluation. Second, the fast/slow decomposition is explicit: the algebraic subsystem is eliminated exactly inside each derivative call, so the remaining differential system is integrated with an explicit adaptive method. Third, the implementation is verified before it is compared with data. The checks are the residual of the algebraic closure, an integration refinement study, and a bookkeeping check on the starch and sucrose subsystem. Fourth, model behaviour is analysed with a screening step followed by a variance-based global sensitivity analysis over declared parameter ranges.

The evaluation design is chosen so that calibration and prediction stay separate. Four coefficients are fitted and shared across all photoperiods, where the earlier version of this model used eight coefficients fitted separately for each photoperiod. Predictive accuracy is then measured by holding out whole photoperiods, that is, by fitting on three light regimes and predicting the fourth. Under this design the model reaches a held-out mean absolute percentage error of 4.36 % in relative growth rate (Figure 3). This value applies when one growth coefficient is allowed to depend linearly on photoperiod. We also measure the range of photoperiods over which the model remains usable, and we report the boundary of that range together with the mechanism that breaks first.

The contributions of the paper are the following.

1. A coupled whole-plant simulation model in which Jarvis stomatal conductance and Farquhar C3 photosynthesis produce the carbon input at every step, and that input is propagated through diel starch and sucrose dynamics, nutrient uptake and costs, and biomass growth (Section 3).
2. An analytic runtime solution of the supply–demand closure as a quadratic in the intercellular CO_2 concentration, embedded in a fast/slow numerical workflow (Sections 3 and 4).
3. A verification protocol for the implementation that is independent of any comparison with data, covering the algebraic closure, the time integration and the carbohydrate bookkeeping (Section 4).
4. An evaluation with four shared fitted parameters instead of a per-photoperiod calibration. The design uses leave-one-photoperiod-out cross-validation, an ablation of the carbon-input pathway, and a global sensitivity analysis over declared ranges (Sections 5 and 6).
5. A measured validity envelope. We report the photoperiod range in which the diel starch balance closes, the photoperiod at which it stops closing, and the bound on the simulated growth response to day length (Sections 6 and 7.5).

The rest of the paper is organised as follows. Section 2 reviews mechanistic growth models, gas-exchange formulations, diel carbohydrate modelling and simulation evaluation practice, and states the gap that this work addresses. Section 3 gives the model formulation as six submodels with their states, inputs and parameters. Section 4 describes the numerical implementation and the verification results. Section 5 defines the photoperiod scenarios, the calibration and cross-validation design, the ablation variants and the sensitivity protocol. Section 6 reports the results, Section 7 interprets them, Section 7.5 states what cannot be concluded, and Section 8 concludes.

2. Related Work

This section positions the model against four bodies of work. These are mechanistic whole-plant and crop simulation, leaf gas-exchange modelling, diel carbohydrate turnover in *Arabidopsis thaliana*, and evaluation practice in simulation studies. The aim is to locate the contribution, not to survey each field completely.

2.1. Mechanistic plant growth simulation models

Mechanistic growth models link physiological processes, microclimate and biomass change over time. The spatial and temporal scale of a model decides where it can be applied. Spatially, models cover the leaf, the organ, the whole plant, the canopy or the field. Temporally, they range from minutes, for gas exchange, through hours and days, for carbon allocation, to whole seasons, for yield. Crop simulation platforms such as APSIM, DSSAT and WOFOST are mature tools for growth, water and nitrogen dynamics at field scale [22, 26, 63]. Recent process-based systems extend this to detailed agronomic management and are validated against field data [42]. Comparative studies indicate that process-based formulations follow seasonal variation in climate better than purely empirical ones [32].

Several other model families address different aspects of the same problem. Functional–structural plant models combine architecture with physiological modules, and they show how spatial structure and dynamic growth can be integrated [30, 50, 12]. Soil–plant water transport models describe flow from the soil through the xylem, and a one-dimensional formulation can be as effective as a three-dimensional one at much lower computational cost [24]. Discrete approaches based on the discrete element method are used for root–soil contact problems [58]. Optimality and energy-budget models assume that allocation maximises a criterion such as growth or resource-use efficiency [44]. Continuous formulations describe biomass, water or vegetation dynamics as systems of differential equations and allow formal analysis [37, 57]. Source–sink descriptions of carbon partitioning between shoot and root belong to the same tradition [33, 17, 56].

Two properties of this literature matter here. Modular organisation is common and useful, because it lets each component be tested on its own [50, 67]. The carbon input, however, is often prescribed rather than computed. Light-use-efficiency terms, radiation-use coefficients and look-up tables are standard in field-scale simulators, and daily time steps are common [26, 63, 22]. This separates the carbon source from the environment and from the internal state of the plant. The model then cannot represent the feedback between stomatal behaviour, intercellular CO_2, carbohydrate status and growth.

2.2. Photosynthesis and stomatal conductance modelling

Photosynthesis sets the primary carbon input, so its description controls everything downstream. For C3 plants the standard biochemical model is the Farquhar–von Caemmerer–Berry formulation, which relates net assimilation to intercellular CO_2 through Rubisco-limited carboxylation and RuBP regeneration [15, 6]. Its parameters, mainly V_{cmax}, J, Γ^*, K_c, K_o and R_d, and their temperature dependence are well documented [48, 4, 13].

Assimilation also depends on the diffusive supply of CO_2, which stomatal conductance controls [28]. Stomatal models therefore connect microclimate to carbon gain. The main formulations are the empirical multiplicative Jarvis model, the Ball–Berry family and the Medlyn optimality model [25, 2, 35]. The Jarvis form expresses conductance as a product of response functions to irradiance, vapour-pressure deficit, temperature, CO_2 and plant water status [25, 65, 66]. It is convenient in a simulation setting because time-varying drivers from a scenario or from measurements enter it directly. Reviews of coupled formulations compare these alternatives and discuss where each is appropriate [11, 5].

Coupling the two sides introduces a feedback. Intercellular CO_2 sets the assimilation rate in the biochemical model, while conductance sets how much CO_2 can diffuse into the leaf. The operating point is therefore defined implicitly and must be found at every evaluation of the model [11]. Iterative solutions are widely used, and analytic solutions of the coupled system for particular stomatal formulations are also known [5, 27]. The Farquhar and Jarvis equations used here are inherited physiological components, not a contribution of this paper. The contribution is the way they are solved at runtime and coupled to the carbohydrate, nutrient and biomass states.

2.3. Diel carbohydrate turnover and photoperiod

Carbohydrate dynamics are the link between assimilation and growth. In the light phase, assimilated carbon is partitioned between mobile sugars, mainly sucrose, and transient reserves, mainly leaf starch. In the dark phase, starch degradation supplies carbon for respiration, maintenance and growth [53, 51]. In *Arabidopsis thaliana* this degradation is closely regulated, and the reserve is almost completely used by dawn [20, 46, 61]. Mathematical models of the process

connect circadian regulation, sucrose availability and sugar signalling to carbon partitioning and growth [18, 47]. Sucrose transport and the separation between cytosolic and vacuolar sucrose have also been described in detail [29, 40].

Photoperiod changes the balance between daytime accumulation and nocturnal consumption. A short day limits the time available for assimilation and lengthens the interval in which growth depends on stored starch. A long day extends direct carbon input and shortens the interval of reserve use. Experiments over contrasting short and long photoperiods show that *Arabidopsis thaliana* adjusts partitioning and growth to day length [54, 36, 1]. This makes photoperiod scenarios a suitable evaluation setting: they perturb both phases of the diel carbon economy through one variable, and they have documented growth and carbohydrate responses.

Growth also depends on nutrient status. Nitrogen and phosphorus affect photosynthetic capacity, metabolism and allocation, and mechanistic models introduce them as internal states or as limiting factors [19, 64, 3, 49, 9]. Models that resolve the diel cycle in detail, however, usually treat assimilation as given [46, 20, 18]. The day–night carbohydrate balance and the gas exchange that produces it are then not solved as one system.

2.4. Verification, validation and sensitivity practice in simulation

Evaluation of plant models normally compares simulated and reference values for biomass, relative growth rate, and water or nutrient fluxes [60, 22, 26]. Two distinctions from simulation methodology are relevant here. The first is between verification and validation: agreement with measurements does not show that the numerical procedures inside the simulator are correct [67, 10]. Residual checks against analytic solutions, integration refinement studies and conservation or bookkeeping checks address a different question from data agreement. The second is between calibration residual and predictive error. When a model is fitted and evaluated against the same small set of aggregate observations, the reported error is a calibration residual. Multi-model studies of crop simulators show how strongly reported accuracy depends on the calibration protocol [31, 60].

Ablation is a further tool for attributing behaviour to model structure. It compares variants in which a component is removed, simplified or replaced. For a coupled model of the kind studied here, three variants are informative. The first uses a prescribed carbon input. The second computes assimilation at fixed conductance. The third is the full variant with environment-responsive conductance. Such a comparison separates the effect of the coupling from the effect of the individual equations.

Sensitivity analysis identifies which parameters control model outputs. Variance-based methods give first-order and total-order indices and quantify interaction structure [52, 45]. Screening methods of the Morris type are cheaper and indicate nonlinearity and interaction before the full analysis [39, 8], and Latin hypercube designs are standard for the sampling step [34]. Because these indices are themselves Monte Carlo estimates, testing the estimator against a function with closed-form indices is good practice [23]. Two limits should be stated explicitly. Indices describe model behaviour over the declared parameter domain, and they do not establish physiological importance. Screening measures indicate interaction, and they are not an exact variance decomposition.

2.5. Open problems addressed here

Read together, this literature leaves a specific gap. Leaf-level gas-exchange models solve the supply–demand operating point accurately, but at a fixed carbohydrate status and without downstream dynamics [15, 25, 6]. Whole-plant and crop simulators carry growth over days to seasons, but they prescribe or simplify the carbon input and rarely resolve the diel cycle [26, 63, 22]. Dedicated diel carbohydrate models resolve the day–night starch balance, but they take assimilation as an input [46, 20]. The feedback loop that connects conductance, intercellular CO_2, carbohydrate status and growth is therefore opened somewhere in each family. Two further problems are methodological. The algebraic constraint created by the coupling has to be solved at every step, yet the solver and its behaviour outside the calibrated range are often left unstated. And calibration is frequently performed against the same aggregate observations used for evaluation, so a predictive error cannot be separated from a calibration residual. Table 1 places the present model against the closest published alternatives on these axes.

This paper closes part of that gap. We do not propose new process equations for photosynthesis, stomatal conductance or starch turnover; those are taken from the literature. What we add is the coupled formulation with a computed carbon input, together with its analytic runtime closure and fast/slow numerical treatment. We also add a verification protocol that is independent of data agreement. And we add an evaluation design in which whole photoperiods are held out and the fitted parameter set is shared across conditions. The aim is not to replace crop simulators or functional–structural models. It is to provide a compact and reproducible core in which gas exchange is propagated explicitly into downstream states.

Table 1
The present model against the closest related model families. FvCB: Farquhar–von Caemmerer–Berry biochemistry. "Carbon input" distinguishes models that compute assimilation from gas exchange at every step (*comp.*) from those that prescribe it through a light-use-efficiency term, a radiation-use coefficient or a look-up table (*presc.*). "Fast–slow" marks an explicit numerical separation of the algebraic gas-exchange subsystem from the differential growth subsystem. "Num. verif." marks published convergence, conservation or estimator-verification evidence, as distinct from agreement with data. Entries describe the standard formulations as given in the cited references; individual modules and versions differ.

Model	FvCB	Stom. model	Starch/ sucrose	N/P	Carbon input	Fast–slow	Num. verif.	Open src
APSIM [22]	–	–	–	N	presc.	–	–	yes
DSSAT [26]	–	–	–	N	presc.	–	–	yes
WOFOST [63]	–	–	–	N	presc.	–	–	yes
GreenLab [12]	–	–	–	–	presc.	–	–	–
T&C-CROP [42]	yes	B–B	–	N	comp.	–	–	yes
Xylem–soil [24]	–	–	–	–	presc.	–	–	–
Scialdone et al. [46]	–	–	yes	–	presc.	–	–	–
Graf et al. [20]	–	–	yes	–	presc.	–	–	–
Feugier and Satake [18]	–	–	yes	–	presc.	–	–	–
Weraduwage et al. [62]	yes	–	part.	–	comp.	–	–	–
This work	yes	Jarvis	yes	N, P	comp.	yes	yes	yes

The paper does not close the rest of the gap, and we say so with numbers in Section 6. The model does not generate the full observed photoperiod dependence of growth from its shared dynamics; part of that dependence remains localised in one growth-conversion coefficient. The diel starch balance closes only up to an upper photoperiod, and above that boundary the reserve accumulates from day to day instead of being consumed (Figure 7). Canopy geometry, water transport, temperature acclimation and multi-species parameterisation are outside the present scope. These points are stated as named structural limitations in Section 7.5 rather than left for the reader to find.

3. Model Formulation

This section states the model as it is implemented. We fix the system boundary and the decomposition into submodels, give the state vector, and then specify each submodel in the order in which information flows through one derivative evaluation. Every equation corresponds to a function in the released C source, and every parameter value is the value used in the reported runs. Notation is introduced once here and used unchanged in the rest of the paper. Table 2 collects the states, inputs and derived quantities, Table 3 the parameters with their provenance, and Table 4 the four calibrated coefficients.

3.1. Conceptual model and system decomposition

The system is a single vegetative *Arabidopsis* plant with its immediate physical environment. Radiation, air temperature, humidity, ambient CO_2, leaf water potential and soil nutrient concentrations cross the boundary as external inputs. Carbon and nutrient acquisition, storage and allocation to leaf and root biomass are internal states. The plant is two mass compartments, leaf and root, without explicit geometry, so internal transport is described by aggregate pools. This lumped decomposition and the carbon–nutrient allocation states follow the whole-plant growth framework of Tedone [55], whose prescribed carbon input is here replaced by a computed gas-exchange layer.

Figure 1 shows the decomposition into six submodels and the signals they exchange.

S1 *Environmental forcing* maps clock time to the light switch and the micro-climate. *In:* t, scenario parameters. *Out:* $PAR, T, RH, VPD, C_a, \psi_w, L$.

S2 *Stomatal conductance (supply)*, a multiplicative Jarvis model. *In:* PAR, VPD, T, C_a, ψ_w. *Out:* g_s.

S3 *Leaf photosynthesis (demand)*, the Farquhar C_3 model. *In:* $PAR, C_i, V_{c\,\max}, J_{\max}$. *Out:* $A_n(C_i)$.

S4 *Gas-exchange coupling* closes supply against demand. *In:* $g_s, A_n(C_i), C_a$. *Out:* A_n, C_i.

S5 *Carbohydrate and nutrient economy* holds the diel starch and sucrose pools, the vacuolar compartment, and sucrose-limited N and P uptake with its costs. *In:* Φ_c, soil N_s, P_s. *Out:* s, U_N, U_P.




Figure 1: Model architecture and information flow through one derivative evaluation. The fast subsystem (blue) is algebraic and is resolved once per evaluation: environmental forcing S1 drives the stomatal supply function S2 and the photosynthetic demand function S3, which are closed against each other in S4 as a quadratic in the intercellular CO_2 concentration and solved in closed form. The resulting net assimilation is converted to the model carbon input P_H. The slow subsystem (orange) comprises the eleven integrated states: the carbohydrate economy S5, nutrient acquisition with its explicit costs S5b, and leaf and root biomass S6. Growth returns to the carbohydrate economy through sink activation and through leaf area, which closes the only feedback loop in the architecture. Arrows show the direction of information flow, not stoichiometric mass transfer; the carbon budget is stated in Section 4.3.

S6 *Growth and allocation* converts exported sucrose into leaf and root biomass under functional-balance control. *In:* $s, \alpha_N, \alpha_P, \varsigma$. *Out:* B_l, B_r, ρ_r.

Submodels S2–S4 form a fast algebraic subsystem. Leaf gas exchange equilibrates within seconds, so it is treated as instantaneously relaxed relative to the slow states in S5 and S6, which evolve over the diel cycle. It is therefore not integrated but evaluated in closed form at every derivative call, as an algebraic constitutive layer inside the derivative of the slow states [10, 67]. This separation organises both the formulation and the implementation.

Only one signal returns from the slow subsystem to the fast one: leaf biomass B_l acts as a viability gate, and the carbon input is zero when $B_l < B_{l,\min}$. The photosynthetic capacities $V_{c\,\max}$ and $J_{\max}$ are fixed parameters and do not respond to nitrogen or carbohydrate status. Figure 1 must therefore not be read as showing a capacity feedback that the equations do not contain.

The formulation rests on six assumptions.

(A1) The plant is lumped into leaf and root compartments; no within-organ gradients or canopy architecture are resolved.
(A2) Leaf gas exchange is at steady state at every instant, so intercellular CO_2 is an algebraic balance and not a state.
(A3) Stomatal conductance responds multiplicatively to the environment and not to the assimilation rate [25, 11].
(A4) Transitory starch is mobilised at a nocturnal rate paced by night length, as the observed near-linear depletion indicates [20, 46].
(A5) Active nutrient uptake is limited by cytosolic sucrose, so carbon and nutrient acquisition are co-limiting [51].
(A6) Partitioning follows functional-balance theory: investment shifts towards the most limiting resource [56, 57].

3.2. State-space representation

The slow states form the vector $X(t) \in \mathbb{R}^{11}$ governed by

$$\frac{dX(t)}{dt} = f(X(t),\, U(t),\, \boldsymbol{\theta},\, t)\,, \tag{1}$$

where $U(t)$ is the exogenous forcing and $\boldsymbol{\theta}$ the time-invariant parameter set. The explicit time dependence enters only through the diel forcing. In the order used by the state packing routine of the implementation,

$$X(t) = \left[k_s,\ a,\ s,\ \alpha_N,\ \alpha_P,\ N,\ P,\ \rho_r,\ B_l,\ B_r,\ s_v\right]^{\mathsf{T}}. \tag{2}$$

Each symbol is defined in Table 2. Total biomass $B = B_l + B_r$ is recomputed from the two biomass states whenever it is needed and is not integrated as a twelfth state.

The pool states a, s, s_v, N and P are normalised model pools on a nominal per-fresh-weight basis, not concentrations of a growing body. We adopt this reading because it is the one under which the implemented balance equations are exact: no growth-dilution term appears in Eqs. (26), (28), (29), (37) or (38). The cost is that a and s are not directly measured tissue concentrations, so the carbohydrate comparison in Section 6 is a comparison of shape and timing after multiplicative alignment, not of absolute level.

The forcing vector is

$$U(t) = \left[PAR(t),\ T(t),\ RH(t),\ VPD(t),\ C_a,\ \psi_w,\ L(t)\right]^{\mathsf{T}}, \tag{3}$$

The deficit VPD is a diagnostic function of T and RH through Eq. (6) and is not an independent input; it is listed because it enters the stomatal response directly. In the scenarios reported here T, RH, C_a and ψ_w are constant, so the only time-varying inputs are PAR and L.

3.3. Submodel S1: environmental forcing

Let $h(t) = t \bmod 24$ be the hour of day and f the photoperiod length in hours. The light switch is

$$L(t) = \begin{cases} 1, & 0 \le h(t) < f, \\ 0, & f \le h(t) < 24, \end{cases} \tag{4}$$

and the incident radiation follows one of two daytime profiles, selected by scenario:

$$PAR(t) = \begin{cases} PAR_{\max}, & \text{square wave, } L = 1, \\ PAR_{\max}\, \sin(\pi\, h(t)/f)\,, & \text{half sine, } L = 1, \\ 0, & L = 0. \end{cases} \tag{5}$$

The square wave represents a growth chamber, which delivers constant irradiance over the light period; the half sine is a smooth idealisation of a natural light course. The choice sets the daily light integral: the square wave gives $PAR_{\max} f$ and the half sine $(2/\pi)\, PAR_{\max} f$. A run matching a chamber experiment must therefore use the square profile at the chamber irradiance. One consequence of the design affects every photoperiod result. With $PAR_{\max}$ held fixed, changing the photoperiod changes the daily light integral in proportion. Day length and daily photon dose are confounded by construction, so a photoperiod response in this model is a joint response to duration and dose.

The vapour pressure deficit is obtained from leaf temperature and relative humidity through the Tetens relation [38, 41],

$$VPD = e_s \left(1 - \frac{RH}{100}\right), \qquad e_s = 0.61078\, \exp\!\left(\frac{17.27\,T}{T + 237.3}\right), \tag{6}$$

with e_s and VPD in kPa and T in °C. Where a thermoperiod is specified, T takes its day value while $L = 1$ and its night value while $L = 0$; otherwise T is constant.

Table 2
State variables, environmental inputs and the principal derived quantities. Pool states are normalised model pools on a nominal per-fresh-weight basis (Section 3.2). "Derived" quantities are recomputed algebraically at every derivative evaluation and are not integrated.

Symbol	Meaning	Unit	Role
k_s	Starch partition coefficient	–	state, Eq. (25)
a	Transitory starch pool	μmol C_6 gFW^{-1}	state, Eq. (26)
s	Cytosolic sucrose pool	μmol C_6 gFW^{-1}	state, Eq. (28)
s_v	Vacuolar sucrose pool	μmol C_6 gFW^{-1}	state, Eq. (29)
α_N	Nitrogen investment state	–	state, Eq. (40)
α_P	Phosphorus investment state	–	state, Eq. (41)
N	Internal nitrogen pool	μmol N gFW^{-1}	state, Eq. (37)
P	Internal phosphorus pool	μmol P gFW^{-1}	state, Eq. (38)
ρ_r	Sucrose-to-root allocation state	–	state, Eq. (45)
B_l	Leaf biomass	gFW	state, Eq. (43)
B_r	Root biomass	gFW	state, Eq. (44)
PAR	Photosynthetically active radiation	μmol m^{-2} s^{-1}	input, Eq. (5)
L	Light/dark switch	–	input, Eq. (4)
T	Leaf temperature	°C	input, constant
RH	Relative humidity	%	input, constant
C_a	Ambient CO_2 concentration	μmol mol^{-1}	input, constant
ψ_w	Leaf water potential	MPa	input, constant
N_s, P_s	Soil N, P concentration	μmol cm^{-3}	input, constant
VPD	Vapour pressure deficit	kPa	derived, Eq. (6)
g_s, g_c	Conductance to H_2O, to CO_2	mol m^{-2} s^{-1}	derived, Eqs. (7), (17)
C_i	Intercellular CO_2 concentration	μmol mol^{-1}	derived, Eq. (21)
A_c, A_j	Rubisco-, RuBP-limited rate	μmol m^{-2} s^{-1}	derived, Eqs. (13), (14)
A_n	Net leaf assimilation	μmol m^{-2} s^{-1}	derived, Eq. (15)
J	Electron transport rate	μmol m^{-2} s^{-1}	derived, Eq. (16)
Φ_c	Carbon input to the pools	μmol C_6 gFW^{-1} h^{-1}	derived, Eq. (23)
D_a	Starch degradation flux	μmol C_6 gFW^{-1} h^{-1}	derived, Eq. (27)
E	Exportable sucrose term	μmol C_6 gFW^{-1}	derived, Eq. (30)
U_N, U_P	Realised N, P uptake	μmol gFW^{-1} h^{-1}	derived, Eq. (36)
ς	Stoichiometric signal	–	derived, Eq. (46)
B	Total biomass, $B_l + B_r$	gFW	derived

3.4. Submodel S2: stomatal conductance (supply)

This submodel returns the stomatal conductance and forms the supply side of gas exchange. It carries no dependence on the assimilation rate (A3). Conductance is a maximal value reduced by a product of dimensionless environmental modifiers, each bounded in [0, 1] [25, 11]:

$$g_s = g_{s,\max}\, f_{PAR}(PAR)\, f_{VPD}(VPD)\, f_T(T)\, f_{CO_2}(C_a)\, f_w(\psi_w). \tag{7}$$

The multiplicative closure assumes that the environmental controls act independently. It is a widely used empirical conductance model and needs no optimisation step at runtime [11, 5]. The maximal conductance $g_{s,\max}$ is a separate parameter from the shape parameters of the modifiers. Each modifier can then be required to equal unity at its own non-limiting reference condition, so changing the shape of one modifier does not rescale the whole conductance.

The light response is a saturating hyperbola with a residual conductance b_{10} below a breakpoint $q = b_{10}/b_1$,

$$f_{PAR} = \begin{cases} b_{10}, & PAR \le q, \\ \dfrac{b_1 b_2\,(PAR - q)}{b_1 + b_2\,(PAR - q)}, & PAR > q. \end{cases} \tag{8}$$

The response to atmospheric demand is linear and floored at zero, and the response to leaf water potential is a saturating recovery above a closure potential $\psi_{\min}$, clipped to $[0, 1]$:

$$f_{VPD} = \max\big(0,\ 1 - b_5\, VPD\big)\,, \tag{9}$$

$$f_w = 1 - \exp\big(-b_6\,(\psi_w - \psi_{\min})\big)\,. \tag{10}$$

The temperature response is a skewed optimum with zeros at T_1 and T_h,

$$f_T = \begin{cases} b_3\,(T - T_1)\,(T_h - T)^{b_4}, & T_1 < T < T_h, \\ 0, & \text{otherwise}, \end{cases} \qquad b_4 = \frac{T_h - T_0}{T_0 - T_1}, \quad b_3 = \frac{1}{(T_0 - T_1)(T_h - T_0)^{b_4}}. \tag{11}$$

The two constants b_3 and b_4 are not free. They are fixed by requiring stationarity at T_0 and $f_T(T_0) = 1$, which places the maximum at T_0 and normalises the modifier to unity there. With $T_1 = 5$, $T_0 = 25$ and $T_h = 45\,^{\circ}\mathrm{C}$ this gives $b_4 = 1$, so the curve is a single-peaked parabola and no clipping from above is needed.

The CO_2 response is piecewise linear in C_a, expressed in $\mu\mathrm{mol\,mol}^{-1}$. It is constant below 100, falls linearly to a floor b_8 at 1000, and equals that floor above it:

$$f_{CO_2} = \begin{cases} 1, & C_a < 100, \\ 1 - b_7\,(C_a - 100), & 100 \le C_a < 1000, \\ b_8, & C_a \ge 1000, \end{cases} \qquad b_7 = \frac{1 - b_8}{900}. \tag{12}$$

3.5. Submodel S3: leaf photosynthesis (demand)

This submodel returns net assimilation as a function of the intercellular CO_2 concentration and forms the demand side of gas exchange. It is the Farquhar–von Caemmerer–Berry C_3 model, in which gross assimilation is the smaller of a Rubisco-limited and an electron-transport-limited rate [15, 6]. The Rubisco-limited rate is

$$A_c = V_{c\max}\,\frac{C_i - \Gamma^*}{C_i + K_c\big(1 + O/K_o\big)}, \tag{13}$$

with Γ^* the CO_2 compensation point in the absence of dark respiration, K_c and K_o the Michaelis constants of Rubisco for CO_2 and O_2, and O the oxygen concentration. The electron-transport-limited rate is

$$A_j = J\,\frac{C_i - \Gamma^*}{4.5\,C_i + 10.5\,\Gamma^*}, \tag{14}$$

where the denominator coefficients 4.5 and 10.5 encode the electron requirement per carboxylation and per oxygenation under the standard stoichiometry of RuBP regeneration [7, 6]. Net assimilation is the limiting rate less dark respiration,

$$A_n = \min(A_c,\ A_j) - R_d. \tag{15}$$

The electron transport rate follows from PAR through the non-rectangular hyperbola $\theta J^2 - (J_{\max} + \phi\, PAR)J + J_{\max}\phi\, PAR = 0$ [15, 16], of which the lower root is taken,

$$J = \frac{\big(J_{\max} + \phi\, PAR\big) - \sqrt{\big(J_{\max} + \phi\, PAR\big)^2 - 4\,\theta\, J_{\max}\,\phi\, PAR}}{2\,\theta}, \tag{16}$$

with ϕ the apparent quantum yield for electron transport and θ the curvature of the light response.

No irradiance threshold is applied to this submodel: the carbon input is evaluated whenever $PAR > 0$ and $B_l \geq B_{l,\min}$. A threshold is unnecessary because, with the closed-form operating point of Section 3.6, the net rate passes continuously through zero at the light compensation point and the clamp in Eq. (23) returns exactly zero below it. Below that point A_n itself remains negative. It is the carbon input that is clamped and not A_n, so respiratory efflux is never counted as an assimilatory gain.

3.6. Submodel S4: gas-exchange coupling

Equations (13)–(15) give demand as a function of a concentration that is itself set by supply. Fickian diffusion of CO_2 through the stomata gives

$$A_n = g_c\,(C_a - C_i), \qquad g_c = \frac{g_s}{1.6}, \tag{17}$$

where the factor 1.6 converts the conductance to water vapour into a conductance to CO_2, which diffuses more slowly in air [6, 41]. Because A_n depends on C_i and C_i depends on A_n, the operating point is the mutual solution of Eqs. (15) and (17). How that coupling is solved is the main methodological choice in the formulation.

The system needs no iteration. Both Farquhar branches have the same rational form in C_i,

$$A = \frac{\mathrm{num}\,(C_i - \Gamma^*)}{d_1 C_i + d_0} - R_d, \qquad (\mathrm{num},\, d_1,\, d_0) = \begin{cases} \big(V_{c\max},\ 1,\ K_c(1 + O/K_o)\big), & \text{Rubisco branch,} \\ \big(J,\ 4.5,\ 10.5\,\Gamma^*\big), & \text{transport branch,} \end{cases} \tag{18}$$

so the branch coefficients are read directly off Eqs. (13) and (14) without dividing through. Equating supply and demand,

$$g_c\,(C_a - C_i)\,(d_1 C_i + d_0) = \mathrm{num}\,(C_i - \Gamma^*) - R_d\,(d_1 C_i + d_0), \tag{19}$$

and clearing the denominator gives a quadratic in C_i,

$$a\,C_i^2 + b\,C_i + c = 0, \qquad \begin{aligned} a &= -g_c\,d_1, \\ b &= g_c\,(d_1 C_a - d_0) - \mathrm{num} + R_d\,d_1, \\ c &= g_c\,d_0\,C_a + \mathrm{num}\,\Gamma^* + R_d\,d_0. \end{aligned} \tag{20}$$

Exactly one root of Eq. (20) is admissible on each branch, and no search is needed to identify it. The admissibility condition is $C_i \geq 0$ alone. For $g_c > 0$ the leading coefficient is $a = -g_c d_1 < 0$, while $c > 0$ because every factor in it is positive. The product of the roots c/a is therefore negative, so each branch has one positive and one negative root, and the positive one is the operating point:

$$C_i = \frac{-b - \sqrt{b^2 - 4ac}}{2a} \ > \ 0. \tag{21}$$

Substituting this root back into Eq. (18) gives the branch assimilation, and the smaller of the two branch values is returned. That is the same co-limitation rule as $\min(A_c, A_j)$ in Eq. (15), applied after each branch has been closed against supply rather than before. Because exactly one root per branch is admissible, the minimum over the two branches is the solution of the coupled system and not merely a candidate for it.

The admissibility rule deserves emphasis, because the natural guess is wrong. No upper bound on C_i is imposed, and a root above the ambient concentration is retained rather than rejected. By Eq. (17), $C_i > C_a$ is exactly what the supply relation returns whenever $A_n < 0$, that is, whenever the leaf is a net source and CO_2 diffuses outward. The condition $0 \leq C_i \leq C_a$ is therefore not a physical constraint here, and imposing it would discard the correct root twice a day.

The vanishing-conductance case has a physical answer rather than a clamp. As $g_c \to 0$ the quadratic degenerates, with $a \to 0$ and $b \to -\text{num}$, and the finite root of the remaining linear equation gives $C_i \to \Gamma^* + R_d d_0/\text{num}$, at which Eq. (18) returns $A = -R_d$. Closed stomata admit no CO_2, so the leaf only respires. The implementation returns this limit directly at the boundary $g_c \le 0$,

$$C_i = \Gamma^*, \qquad A_n = -R_d, \tag{22}$$

which is the correct dark limit and replaces a clamp on the conductance. Respiration in low light and darkness is then carried by the same expression that carries assimilation in the light, with no special case and no discontinuity in A_n at the light compensation point.

Two properties of this closure matter for the rest of the paper. The branch-wise quadratic is solved once per derivative evaluation, with no inner iteration in the production configuration, so the derivative is a deterministic function of (X, t) as an embedded-pair error estimate assumes. And the solution is exact up to floating-point arithmetic, so no inner convergence tolerance interacts with the outer error control. Section 4 reports the substitution residual that verifies this.

3.7. From assimilation to model carbon input

The leaf-level net assimilation A_n (μmol CO_2 m^{-2} s^{-1}) must be converted to the carbon input Φ_c that drives the carbohydrate pools (μmol C_6 gFW^{-1} h^{-1}). The conversion is

$$\Phi_c = \Lambda \ \max\left(0,\ \underbrace{\text{SLA}}_{\text{m}^2\,\text{gFW}^{-1}} \ \underbrace{\frac{3600}{6}}_{\text{s h}^{-1}\ /\ \text{mol CO}_2\ \text{per mol C}_6} \ A_n\right), \qquad L = 1, \tag{23}$$

and $\Phi_c = 0$ when $L = 0$ or $B_l < B_{l,\min}$. Three factors carry the whole unit conversion and none of them is free: SLA moves the flux to a per-fresh-weight basis, the factor 3600 converts seconds to hours, and the divisor 6 converts moles of CO_2 to moles of hexose. Their product is 600 SLA, so at the measured SLA of Table 3 the coefficient is fixed by the units and no empirical scaling constant enters the carbon-input pathway. The clamp keeps a negative A_n out of the pools, as discussed in Section 3.5.

The factor Λ is a sink-limitation term that down-regulates assimilation as the starch pool fills,

$$\Lambda = \min\left(1,\ \max\left(\lambda_c,\ \lambda_c + (1-\lambda_c)\,\frac{a_{\max} - a}{a_{\max}}\right)\right), \qquad a_{\max} = D_{a,\max}\,(24 - f), \tag{24}$$

where λ_c is the residual assimilation fraction at a full pool and the ceiling $a_{\max}$ scales with night length, since a longer night requires a larger reserve. The inner max bounds the term at its value at the ceiling, so a pool above $a_{\max}$ cannot drive Λ negative and reverse the sign of assimilation. In all reported runs $\lambda_c = 1$, which makes $\Lambda = 1$ identically and the sink limitation inactive. The equation is given because the term is present in the right-hand side and is used in the ablation of Section 5; its shipped value is stated so that a reader knows the reported results do not depend on it.

3.8. Submodel S5: carbohydrate economy

The carbon input Φ_c is split between transitory starch and cytosolic sucrose by the partition coefficient k_s. This coefficient is a state with its own dynamics rather than a fixed split, so the plant can shift it during the day and again at night in response to the sucrose level. The diel carbohydrate balance is then an emergent property of the model and not a prescribed schedule [53, 36].

Starch partition. Let $\sigma^- = s_{\min}/(s_{\min} + s)$ and $\sigma^+ = s/(s + s_{\max})$ be the sucrose-deficit and sucrose-excess fractions, both in [0, 1] and both saturating. The partition state follows

$$\frac{dk_s}{dt} = \underbrace{L\left[-k_s\,\lambda_{sdr}\,\sigma^- + \left(1-k_s\right)\lambda_{sdi}\,\sigma^+\right]}_{\text{light phase}} + \underbrace{(1-L)\left(1-k_s\right)\lambda_{sni}\,\sigma^-}_{\text{dark phase}}. \tag{25}$$

Each term has a direct interpretation. In the light, a low sucrose level makes σ^- large and the first term reduces k_s, so more incoming carbon goes to sucrose and the immediate demand is met. A high sucrose level makes σ^+ large and the second term raises k_s, so surplus carbon is stored as starch. In the dark there is no input to partition, and the third term raises k_s when sucrose is low, returning the state to a storage-biased value before the next light period. The factors k_s and $(1-k_s)$ keep the state in $[0,1]$ without a clamp, since each term vanishes at the bound it approaches. The three rate coefficients set the speed of the day-reduction, day-increase and night-increase responses.

Starch pool. Starch gains the partitioned share of the input and loses the nocturnal degradation flux D_a,

$$\frac{da}{dt} = k_s\,\Phi_c - D_a. \tag{26}$$

The degradation flux implements the observed pacing of starch mobilisation against the remaining night length [20, 46]. Let a_{dusk} be the starch content recorded at the last transition to darkness and $a_{\min}$ the residual pool that is not mobilised. Then

$$D_a = (1-L)\min\left(D_{a,\max},\ \left[\frac{h}{24} + \left(1-\frac{h}{24}\right)\left(1-\frac{s}{s+s_{\max}}\right)\right]\frac{a_{\text{dusk}}-a_{\min}}{24-f}\right), \tag{27}$$

and $D_a = 0$ if $a_{\text{dusk}} < a_{\min}$. The quotient $(a_{\text{dusk}} - a_{\min})/(24-f)$ is the constant rate that would exactly exhaust the mobilisable reserve over the night, which is the rate the measurements show. The bracketed factor modulates it: when sucrose is abundant the second summand is small and degradation is slowed, so the reserve is spared. The outer min bounds the flux by the degradation capacity $D_{a,\max}$ and the factor $(1-L)$ confines it to darkness. Two properties matter for the limitation reported in Section 7.5. The reference level a_{dusk} is sampled at dusk and held over the night, so it is not a continuous function of the state, and the divisor $24-f$ is the nominal rather than the remaining night length.

Sucrose pools. Cytosolic sucrose receives the unpartitioned share of the input and the starch degradation flux, and pays four costs: nutrient uptake, phloem transport, maintenance respiration, and export to the sinks. It also exchanges with the vacuolar pool. The balance is

$$\frac{ds}{dt} = \left(1-k_s\right)\Phi_c + D_a - C_u - C_t - R_m - \eta\,E - k_{\text{in}}\,s + k_{\text{out}}\,s_v - C_g, \tag{28}$$

$$\frac{ds_v}{dt} = k_{\text{in}}\,s - k_{\text{out}}\,s_v. \tag{29}$$

The vacuolar compartment is a passive first-order buffer with influx rate k_{in} and efflux rate k_{out}, and the two rate constants appear with opposite signs in Eqs. (28) and (29), so the exchange conserves sucrose. It represents the measured separation of leaf sucrose into an active cytosolic fraction and a larger stored vacuolar fraction [29, 40]. Only s enters the cost and export terms, so the buffer damps the diel excursion of the active pool without adding or removing carbon, with relaxation time $1/(k_{\text{in}} + k_{\text{out}})$.

The export term is the phloem-loading flux, and it carries the sugar-induced sink activation:

$$E = \underbrace{\min\left(s,\ s_{\max}\right)}_{\text{loadable pool}}\underbrace{\left[1+\left(\frac{s}{K_s}\right)^{n_H}\right]}_{\text{sink activation}}\underbrace{\left[\lambda_g + \left(1-\lambda_g\right)\left(1-L+L\left(1-k_s\right)\right)\right]}_{\text{loading efficiency}}. \tag{30}$$

The activation factor multiplies the loadable pool before the loading efficiency is applied, so it scales the export flux and not the pool size. Its form requires comment because it is not the bounded Hill function. A saturating factor $s^{n_H}/(K_s^{n_H} + s^{n_H})$ is bounded above by one and can therefore only weaken the drain, which is the wrong direction. Sucrose induces its own consumption, by raising invertase activity and sink-tissue transporter expression, so consumption capacity rises with the pool [51, 53]. The unbounded factor $1 + (s/K_s)^{n_H}$ represents this: it is close to one at $s \ll K_s$, and at $s \gg K_s$ the drain scales as s^{1+n_H}. The steady-state pool then grows as the $1/(1 + n_H)$ power of the supply, so the sucrose level is held near K_s while assimilation varies. A first-order drain with constant coefficients cannot hold a level in this way. The half-activation constant K_s is one of the four calibrated parameters (Table 4).

The loading efficiency term interpolates between a floor λ_g and unity. It equals one at night and $\lambda_g + (1-\lambda_g)(1-k_s)$ in the light, so export is least efficient when the light-phase partition is strongly biased to starch. The remaining costs are

$$C_u = c_N\, U_N + c_P\, U_P, \tag{31}$$

$$C_t = \tau_c \left(r_m\, s + \eta\, E \right), \tag{32}$$

$$R_m = r_m\, s, \tag{33}$$

$$C_g = \begin{cases} Y \left(\dfrac{d\,B_l}{dt} + \dfrac{d\,B_r}{dt} \right), & \dfrac{d\,B_l}{dt} + \dfrac{d\,B_r}{dt} > 0, \\ 0, & \text{otherwise.} \end{cases} \tag{34}$$

The uptake cost C_u prices the realised nutrient fluxes at the sucrose consumption coefficients c_N and c_P, and is evaluated after Eq. (36) so that it uses the fluxes of the current state. The transport cost C_t is a fixed fraction τ_c of the respiratory and loading fluxes. Maintenance respiration R_m is first order in the cytosolic pool with specific rate r_m, charged over all hours as in the source formulation [55]. The construction cost C_g prices the biomass actually built, at Y moles of hexose per gram of fresh biomass. It is applied after both growth rates have been evaluated and is one-sided: senescence returns tissue but does not refund construction carbon.

3.9. Submodel S5b: nutrient uptake and costs

Nutrient acquisition is active and therefore carbon-limited (A5). It is represented in three stages: a soil-availability rate, an allocation of root capacity, and a sucrose-limited realisation.

The soil stage is Michaelis–Menten in the external concentration [14, 59],

$$I_N = I_{N,\max} \frac{N_s}{N_s + K_N}, \qquad I_P = I_{P,\max} \frac{P_s}{P_s + K_P}. \tag{35}$$

The potential uptake scales this by the investment state and by the root fraction of total biomass, and the realised uptake applies a sucrose limitation:

$$\widehat{U}_N = I_N\, \alpha_N\, \frac{B_r}{B + \varepsilon}, \qquad U_N = \widehat{U}_N\, \frac{s}{s + c_N\, \widehat{U}_N\, \tau + \varepsilon}, \tag{36}$$

$$\widehat{U}_P = I_P\, \alpha_P\, \frac{B_r}{B + \varepsilon}, \qquad U_P = \widehat{U}_P\, \frac{s}{s + c_P\, \widehat{U}_P\, \tau + \varepsilon}.$$

The saturating factor in Eq. (36) has a clear reading. The quantity $c_N \widehat{U}_N \tau$ is the sucrose that the potential uptake would consume over a reference interval $\tau = 1$ h, so it is a sucrose amount and comparable with s. When the pool greatly exceeds it the factor approaches one and uptake proceeds at its potential rate; when carbon is short the factor falls and uptake is throttled. The constant ε is the machine epsilon of the working precision, guards the denominators only, and is not a model parameter.

The internal pools are balanced against two demands, structural and photosynthetic:

$$\frac{dN}{dt} = U_N - \left[r_m\,(s+a) + \eta\,E\right] r_{A,N}\,\frac{B_l}{B+\varepsilon} - \frac{\Phi_c}{\Phi_{c,\max}}\,N_{ph}\,\nu, \tag{37}$$

$$\frac{dP}{dt} = U_P - \left[r_m\,(s+a) + \eta\,E\right] r_{A,P}\,\frac{B_l}{B+\varepsilon} - \frac{\Phi_c}{\Phi_{c,\max}}\,P_{ph}\,\nu. \tag{38}$$

The first loss term charges nutrients against the carbon turnover of the shoot, at the assimilation cost coefficients $r_{A,N}$ and $r_{A,P}$ and in proportion to the leaf share of biomass. The second charges the nutrient requirement of photosynthesis, scaled by the relative assimilation rate $\Phi_c/\Phi_{c,\max}$ and the conversion parameter ν, with N_{ph} and P_{ph} the contents below which assimilation is nutrient limited. Nutrient status enters the gas-exchange layer only through the saturation functions

$$\varphi_N = \max\!\left(0,\ \frac{2N}{N+N_{ph}} - 1\right), \qquad \varphi_P = \max\!\left(0,\ \frac{2P}{P+P_{ph}} - 1\right), \tag{39}$$

which are zero at the threshold and approach one at large content.

The investment states α_N and α_P carry the acclimation of uptake capacity. Each follows a two-term law. The first term drives investment up when the nutrient is scarce or assimilation is high. The second drives it down when the pool is full, or when the carbon price of uptake is high relative to the available sucrose flux:

$$\frac{d\alpha_N}{dt} = \left(1-\alpha_N\right)\left[\left(1 - \frac{U_N}{U_N + C_N + \varepsilon}\right)\frac{N_{\max}}{N+N_{\max}} + \frac{\Phi_c}{\Phi_{c,\max}} - \Omega\right] - \alpha_N\left[\frac{N}{N+N_{\min}} + \frac{c_N U_N}{c_N U_N + \Phi_c\left(1-k_s\right) + D_a + \varepsilon}\right], \tag{40}$$

$$\frac{d\alpha_P}{dt} = \left(1-\alpha_P\right)\left[\left(1 - \frac{U_P}{U_P + C_P + \varepsilon}\right)\frac{P_{\max}}{P+P_{\max}} + \frac{\Phi_c}{\Phi_{c,\max}} + \Omega\right] - \alpha_P\left[\frac{P}{P+P_{\min}} + \frac{c_P U_P}{c_P U_P + \Phi_c\left(1-k_s\right) + D_a + \varepsilon}\right], \tag{41}$$

where C_N and C_P are the nutrient cost terms and Ω is the stoichiometric imbalance signal

$$\Omega = \alpha\,\lambda_k\,\frac{N - \omega P}{N + \omega P + \varepsilon}, \tag{42}$$

with α the corresponding investment state, ω the optimal N:P ratio and λ_k the gain. The sign of Ω is opposite in the two equations, which is the coupling that makes the pair competitive: an excess of nitrogen relative to the optimal ratio suppresses further nitrogen investment and promotes phosphorus investment. The factors $(1-\alpha)$ and α again keep each state in $[0, 1]$ without a clamp.

3.10. Submodel S6: biomass growth and allocation

Growth is driven by the phloem-loading flux and controlled by the allocation state ρ_r, the fraction of that flux directed to the root. Both organs lose biomass to mortality and to a quadratic self-competition term:

$$\frac{dB_l}{dt} = \lambda_{sb}\left(1-\rho_r\right)\eta\,E\,B_l \;-\; \mu_l B_l \;-\; \gamma_l B_l^2, \tag{43}$$

$$\frac{dB_r}{dt} = \lambda_{sb}\,\rho_r\,\eta\,E\,B_l \;-\; \mu_r B_r \;-\; \gamma_r B_r^2. \tag{44}$$

Two features of Eqs. (43)–(44) are easy to misread. Both growth terms are proportional to *leaf* biomass B_l and not to the biomass of the organ being built, because the leaf is the source of the loading flux; root growth is therefore driven by the size of the source. The loading flux also appears in both equations with the same coefficient λ_{sb}, so the split between organs is carried entirely by ρ_r. That coefficient converts the product of loading flux and leaf biomass into a growth rate, so its unit is gFW $(\mu\text{mol C}_6)^{-1}$ per gFW of leaf. It is a calibrated coefficient and not a construction stoichiometry; the construction stoichiometry is Y in Eq. (34). The quadratic terms bound the biomass trajectory at large size and are negligible at the seedling scale of the reported runs.

The allocation state responds to the balance between the nutrient investment already made and the nutrient and carbon status achieved:

$$\frac{d\rho_r}{dt} = \underbrace{\left(1-\rho_r\right)\left[\alpha_N \varsigma + (1-\varsigma)\,\alpha_P\right]}_{\text{demand for soil resources}} - \underbrace{\rho_r\left[\frac{N\,\varsigma}{N+N_{\min}} + \frac{P\,(1-\varsigma)}{P+P_{\min}} + \frac{s_{\min}}{s_{\min}+s}\right]}_{\text{satisfied demand}}, \tag{45}$$

where the stoichiometric weight

$$\varsigma = \frac{\omega}{\omega + \max(N,0)/\left(\max(P,0)+\varepsilon\right)} \tag{46}$$

shifts attention between the two nutrients according to which is further from the optimal ratio ω. The positive part in Eq. (46) makes the ratio well defined for a numerically negative pool and is the identity wherever $N \geq 0$ and $P \geq 0$. Equation (45) is the functional-balance statement of the model [56, 57]: investment moves to the root while soil resources are in demand, and away from it once nutrient and carbon status are satisfied. Both terms carry the factor ρ_r or $(1-\rho_r)$ and keep the state in $[0, 1]$.

Soil nutrient status enters the growth coefficient through a multiplicative gate. With scaled soil concentrations $n = N_s/N_s^{\text{ref}}$ and $p = P_s/P_s^{\text{ref}}$,

$$\lambda_{sb}^{\text{eff}} = \begin{cases} \lambda_{sb}, & \min(n,p) \geq 1, \\ \lambda_{sb}\,\max\!\left(0,\ (n+p) - \frac{1}{2}\left(n^2+p^2\right)\right), & \text{otherwise.} \end{cases} \tag{47}$$

The gate is inactive at or above the reference concentrations and reduces the growth coefficient below them. In every scenario reported here the soil concentrations are at or above those values, so $\lambda_{sb}^{\text{eff}} = \lambda_{sb}$ throughout. We give the gate because it is part of the right-hand side and because the reference concentrations are hard-coded divisors rather than configurable parameters, as Table 3 records.

3.11. Bounded constitutive laws

The right-hand side contains fifteen saturating ratios of the form $X/(X+K)$ with $K > 0$, each with a pole at $X = -K$. Those poles lie outside the physically meaningful region but not outside the region an adaptive integrator can probe. A trial stage that takes a nearly depleted pool slightly negative can land close to a pole and return a derivative many orders of magnitude too large, after which the step controller cannot meet its error estimate at any step size. The formulation therefore evaluates every constitutive law at the positive part of each pool state,

$$\widetilde{X} = \max(X,\,0), \qquad X \in \{a,\ s,\ s_v,\ N,\ P,\ B_l,\ B_r\}, \tag{48}$$

and uses $\widetilde{X}$ wherever a pool appears inside a saturating ratio or a cost term. The state vector is not modified and the differential equations are unchanged. Equation (48) is the identity on $X \geq 0$, the whole region in which a pool content is defined, so it cannot alter a physically admissible trajectory. It states where the constitutive laws are defined and is not an additional model assumption. The states k_s, α_N, α_P and ρ_r need no such treatment, since their equations are self-bounding through the $(1-\cdot)$ factors above.

3.12. Parameters and their provenance

Every parameter carries a provenance record in the released repository, in a ledger that names its value, unit, source and evidence tier. The six tiers are T1 measured, a published measurement of the quantity in a comparable system; T2 derived from T1 quantities through a stated relation; T3 inherited from the source model with the equation number as printed there; T4 bounded by physics or chemistry rather than measured; T5 fitted; and T6 unsourced. A value may not be defended by the improvement it produces in the fit, so its justification rests on the tier and not on the residual. Table 3 is generated from that ledger.

Of the 64 model parameters, 30 are literature-sourced or derived from literature values (2 at T1, 8 at T2, 20 at T3), 7 are constrained by physical or chemical argument (T4), 15 are fitted (T5) and 12 remain unsourced (T6). We report the unsourced count rather than distribute it, because it states what is not yet established. The T6 group holds two kinds of value. The first is the Farquhar kinetic constants Γ^*, K_c and K_o, carried at the values of the source model but without a locator in a comparable system; published temperature-dependent estimates exist [4, 48] and reconciling the shipped values against them is left as future work. The second is a set of constants hard-coded rather than exposed in the configuration file, including the quantum yield ϕ, the curvature θ and the soil reference concentrations of Eq. (47). Of the 15 fitted parameters, 11 are inherited fitted values of the source model that are not re-estimated here.

The remaining four are given separately in Table 4, because the number of free parameters is the central claim of the evaluation. All four are shared: one value of each is used at every photoperiod in every reported run. The published baseline calibrated eight parameters per photoperiod, four photoperiods times two coefficients, against eight observations. That leaves zero degrees of freedom and makes the reported agreement a calibration residual rather than a prediction. Identifiability of the shared set, and the consequences of the reduction, are examined in Sections 5 and 6.

One value in Table 3 needs a note. The maximal conductance $g_{s,\max} = 1/\sqrt{2}$ was chosen so that correcting the normalisation of Eq. (11) is an exact reparametrisation at $T_0 = 25\,^\circ\mathrm{C}$, where the previous formulation returned $f_T = 1/\sqrt{2}$ instead of unity. The product $g_{s,\max} f_T(T_0)$ is therefore unchanged and the downstream parameters need no recalibration.

Four entries carry a source or a setting that the author must resolve before submission.

- TODO(AUTHOR): the measured vacuolar/cytosolic sucrose ratio, from which k_{out} is derived, has a source key that is in neither bibliography file. Which publication, and do Lohaus et al. [29] or Nadwodnik and Lohaus [40] already cover it?
- TODO(AUTHOR): the T1 specific leaf area is attributed to a leaf atlas that is not in the bibliography. Which atlas or dataset, with which locator?
- TODO(AUTHOR): the construction stoichiometry Y derives from a growth-respiration yield attributed in the source to Penning de Vries et al., a reference absent from both bibliography files. Which edition?
- TODO(AUTHOR): the shipped configuration sets $Y = 0$, so $C_g = 0$ in Eq. (28) in all reported runs, while the derivation gives a non-zero value. Is the zero setting to be reported as a stated simplification, or corrected before submission?

The full parameter table with all provenance fields is given in Appendix A; Tables 3 and 4 give the parameters that appear in the equations of this section.

4. Numerical Implementation

The model is implemented as a modular simulation engine in C (ISO C99) and built with CMake. Environmental forcing, gas exchange, carbohydrate dynamics, nutrient uptake and biomass growth are written as separate translation units. Each physiological block can therefore be tested on its own, while the coupled path from forcing to growth stays intact. This section describes how the state equations are advanced, and which checks were applied to the solver components before any result was produced.

4.1. Simulation architecture and time integration

Each right-hand-side evaluation follows the fast–slow structure of Section 3 and Figure 1. Four blocks are executed in sequence. The first block evaluates the clock time and the environmental drivers: PAR, leaf temperature,

relative humidity, vapour-pressure deficit, ambient CO_2 and leaf water potential. The second block computes the gas-exchange layer: stomatal conductance, the electron transport rate and the two Farquhar branches. The intercellular CO_2 concentration follows from the closed-form solution of Eq. (20). The third block converts net assimilation into the carbon-input term Φ_c. The fourth block assembles the derivatives of the eleven slow states. These are starch, cytosolic and vacuolar sucrose, the partition and allocation variables, the two nutrient pools, the two affinity states, and leaf and root biomass.

Gas exchange is algebraic and is solved at every stage of every step, not once per output sample. This matters for the coupling. Conductance, assimilation and C_i are functions of the current state and the current forcing, so a stage value taken from a stale operating point would break the supply–demand balance that the closure enforces. The eleven slow states carry no algebraic constraint and are integrated as an explicit system.

Integration uses a separate, reusable C ODE-solver library through a single interface, so the model does not depend on one backend. The production configuration selects an adaptive Dormand–Prince embedded pair (`ODE45`) with an error tolerance of 10^{-8} [21]. Under an adaptive method the configuration entry `dt_hours` is not the integration step. It sets the initial trial step, which the step controller immediately overrides, and it sets the interval at which the trajectory is sampled for output. Production runs use `dt_hours` $= 0.1$ h, so the reported trajectory is a fixed 0.1 h grid over a 30-day horizon, while the accuracy of that trajectory is controlled by the tolerance alone.

A fixed step is used where the adaptive controller is either unusable or unnecessary. The sensitivity and sweep runners override the backend to fixed-step RK4, because those designs draw parameter values at the edges of the declared ranges, where the adaptive step can collapse. A fixed step also gives a predictable cost over several thousand evaluations. The carbon-balance tool of Section 4.3 uses fixed-step RK4 for a different reason: only a fixed-step driver can advance the flux accumulators with the same weights and the same step as the states. Both backends are run through the same production path at the same settings, so the effect of the choice can be measured rather than assumed.

Two runtime guards protect batch work. A non-finite guard and a domain guard observe the trajectory and classify a run as divergent. A step-budget plugin counts attempted steps and stops the run when a generous budget is exhausted. The second guard covers a failure the first cannot see: for some parameter combinations the adaptive step collapses while the state stays finite, so the integrator neither advances nor reports an error. Such a run then unwinds normally and is recorded as inadmissible, which is how the batch drivers already treat divergence.

4.2. Verification of the solver components

The gas-exchange closure was verified by substitution rather than by assertion. For a reference leaf and PAR from 10 to 2000 μmol m^{-2} s^{-1}, the analytic solution was inserted back into the supply relation and the residual $|A_n - g_c(C_a - C_i)|$ was recorded. The analytic branch solution reproduces the intended operating point. The residual stays at machine precision over the whole light range, so the closed-form root solves the coupled system rather than approximating it.

Two limits were checked separately, because both are boundaries where an implementation can silently disagree with the equations. When conductance reaches zero the implementation returns $C_i = \Gamma^*$ and $A_n = -R_d$, which is the dark limit of the formulation. Above that limit, assimilation crosses zero smoothly at PAR $= 13.36$ μmol m^{-2} s^{-1} under the reference conditions. An earlier hard cut-off at PAR $= 20$ μmol m^{-2} s^{-1} was removed, because f_{PAR} is not zero there and the returned assimilation jumped across the threshold. With the analytic closure the terminal $\max(0, \cdot)$ clamp already returns exactly zero below the light compensation point, continuously and without a special case. Removing the gate therefore also removes a discontinuity that an adaptive step controller has to resolve twice per simulated day.

4.3. Carbon conservation

Whether the carbon bookkeeping closes cannot be answered by post-processing the solver output. The trajectory records pool sizes, whereas closure is a statement about time-integrated fluxes, and a quadrature applied afterwards would test the quadrature as much as the model. The carbon-balance tool therefore appends ten flux accumulators to the eleven states and advances the augmented system with identical RK4 weights and an identical step. Both parts come from one call to the right-hand side, so the accumulators see the same stage values as the state integration. The resulting budget is then exact to the arithmetic of the integrator.

The closure is asserted for the carbohydrate subsystem $\{a, s, s_v\}$, that is starch, cytosolic sucrose and vacuolar sucrose. Over an interval $[0, T]$ the vacuolar exchange terms cancel identically, so the subsystem budget reduces to

$$\left(a + s + s_v\right)\Big|_0^T = \int_0^T A_n \, dt - \int_0^T \left(C_{up} + C_{tr} + r_m s + \phi_{\text{load}} E_{\text{night}}\right) dt, \tag{49}$$

The residual is the difference between the two sides. It is reported relative to gross assimilation,

$$\varepsilon_C = \frac{1}{\int_0^T A_n\,dt}\left[\left(a + s + s_v\right)\Big|_0^T - \int_0^T A_n\,dt + \int_0^T \left(C_{up} + C_{tr} + r_m s + \phi_{\mathrm{load}} E_{\mathrm{night}}\right) dt\right]. \tag{50}$$

Across the four calibration photoperiods the relative residual lies between 2.9×10^{-14} and 6.1×10^{-14}, which is machine precision for double-precision accumulation over this many steps. No carbon is created or destroyed numerically inside the declared scope.

The scope must be stated exactly, because it does not cover the whole plant. The biomass gain term sits outside the budget by construction. In the growth law of Eq. (43) the sucrose drain is $\phi_{\mathrm{load}} E_{\mathrm{night}}$ while the biomass gain is $\lambda_{sb}\,\phi_{\mathrm{load}} E_{\mathrm{night}} B_l$, so the two differ by the factor $\lambda_{sb} B_l$. That factor equals unity only at $B_l = 1/\lambda_{sb}$, which is of order 10^2 gFW. Over the simulated seedling range the factor stays 10^2 to 10^3 away from unity, so the two terms are not two views of one carbon flow. The growth term is a rate law calibrated against observed RGR, not a stoichiometric conversion of sucrose into biomass carbon. Eq. (50) therefore verifies bookkeeping closure of the carbohydrate subsystem, and it is reported under that name only. A budget that included biomass would require a stoichiometric growth equation, which the model does not have.

4.4. Convergence and tolerance study

Three numerical settings can in principle contaminate a reported number, and each was refined independently around its production value. The solver tolerance was swept over 10^{-4} to 10^{-12} at the production output interval. The output interval was swept over 0.4 to 0.00625 h at the production tolerance, which separates an output-grid effect from a discretisation effect. The gas-exchange tolerance was swept over 10^{-3} to 10^{-10}; this sweep necessarily runs the legacy iterative comparator, because the analytic closure is closed-form and has no convergence tolerance at all. A fourth sweep refines the step of the fixed-step backend, where the step genuinely is the independent variable, and a fifth compares the two backends at production settings. Every step used in the sweeps divides both 24 h and the photoperiod exactly, so the end-of-day sample lands on a grid point and no interpolation error enters the comparison.

The reported quantities are insensitive to these settings over the tested range. Refining any of the three beyond its production value moves RGR, end-of-day starch and end-of-day sucrose by far less than the differences between the parameterisation regimes of Section 5. The measured order of the fixed-step sweep is close to one rather than four, and this is expected. Starch degradation is multiplied by the light state, so the right-hand side is discontinuous twice per simulated day, and those transitions set the global error. One integrator offered by the external solver library was excluded from the comparison after it failed a consistency check against an ODE with a known closed-form solution. The full study, including the Richardson estimates and the per-photoperiod tables, is in `docs/revision/convergence_study.md`.

5. Model Evaluation Protocol

This section defines the evaluation rules and reports no results. The reference data are stated with their source, the fitted coefficients are counted in each regime, and every metric is defined together with what it does not measure. No number in Section 6 should then be readable as more than it is.

5.1. Reference data and growth conditions

Four photoperiods are used for calibration and cross-validation: 4, 6, 8 and 12 h of light per 24 h cycle. An 18 h photoperiod is held out of every fit entirely and is used only as an out-of-sample condition. The four short photoperiods span the range over which the reference experiments report growth, and 18 h lies outside it, which is why it is treated separately.

All reported runs reproduce the growth conditions of the reference experiments. PAR is 160 μmol m^{-2} s^{-1} delivered as a square wave over the light phase, and leaf temperature is 21 °C by day and 19 °C by night. This is a methodological requirement, not a detail. The shipped configuration instead runs a half-sine profile peaking at 1200 μmol m^{-2} s^{-1} at a single temperature. A half-sine averages $2/\pi$ of its peak, so that daily light integral is about 4.8 times the chamber value at the same day length. A coefficient fitted under one light dose does not transfer to another: the carbon-input coefficient would absorb the dose mismatch, and the resulting value would describe the forcing rather than the plant. Scoring the model against chamber data under the shipped forcing would therefore compare a model driven at one photon dose with plants grown at another.

The diel carbohydrate reference is `data/paper/sugars_starch_reference.csv`, which holds 65 rows over five photoperiods, that is 13 sampling times per photoperiod at a 2 h resolution through the 24 h cycle. Starch and sucrose are used; the remaining assayed metabolites are not part of this evaluation. The RGR targets are 0.068, 0.114, 0.171 and 0.260 d^{-1} for the 4, 6, 8 and 12 h photoperiods [54]. That study is the only source in the reference list which reports all four photoperiods. Its RGR series and its dusk starch values come from one growth experiment, which is what makes the four-point set internally consistent. Two other studies of diel carbohydrate behaviour cover only a subset of these day lengths, at 12 h [1] and at 6 and 12 h [36]. Neither can therefore supply the four-point series.

5.2. Parameterisation regimes

Three parameterisation regimes are evaluated (Table 5). They differ only in how many coefficients are fitted and in whether those coefficients are allowed to depend on photoperiod. Everything else, including the forcing, the initial state and all sourced parameters, is identical.

The saturated regime is the published baseline. It fits two coefficients per photoperiod by nested bisection: the growth coefficient λ_{sb} against the observed RGR, and the daytime starch synthesis rate λ_{sdi} against the observed end-of-day starch. With four photoperiods that is eight fitted values against eight observations, so the regime has zero degrees of freedom. Its agreement with the data cannot be reported as predictive accuracy. What a bisection leaves behind is a convergence residual of the search, controlled by the resolution of the parameter grid, and it goes to zero as the search is tightened. The regime is retained so that the published result stays reproducible, and it is reported as a calibration diagnostic only.

The shared regime fits four coefficients for all photoperiods together: λ_{sb}, λ_{sdr}, λ_{sdi} and the sink activation constant. Photoperiod is then the only thing that differs between the four runs, so any photoperiod response the model produces is generated by its dynamics rather than supplied by its parameters. The third regime keeps the same shared set but allows the growth coefficient to vary linearly with photoperiod P,

$$\lambda_{sb}(P) = c_0 + c_1 P, \tag{51}$$

which replaces one shared constant by two coefficients. Eq. (56) encodes the hypothesis that the growth response to photoperiod is smooth in P and is not fully generated by the coupled dynamics. It remains a prediction for a held-out photoperiod, because the held-out observation never enters the fit.

5.3. Metrics

Growth is evaluated on relative growth rate, computed from total biomass over the simulated interval,

$$\mathrm{RGR} = \frac{\ln B(t_2) - \ln B(t_1)}{t_2 - t_1}. \tag{52}$$

Per photoperiod we report the signed relative error. Over a set of photoperiods we report the mean absolute percentage error:

$$E_{\mathrm{rel}} = 100\,\frac{\widehat{\mathrm{RGR}} - \mathrm{RGR}_{\mathrm{ref}}}{\mathrm{RGR}_{\mathrm{ref}}}, \qquad \mathrm{MAPE} = \frac{1}{m}\sum_{j=1}^{m}\left|E_{\mathrm{rel},j}\right|, \tag{53}$$

where $\widehat{\mathrm{RGR}}$ is the simulated value and m is the number of photoperiods in the set. In-sample and held-out MAPE are always reported as separate quantities and never pooled.

The diel carbohydrate series are evaluated with a normalised root-mean-square error. A single multiplicative scale α is first fitted to each simulated trace by least squares,

$$\alpha = \frac{\sum_{i=1}^{n}\hat{y}_i y_i}{\sum_{i=1}^{n}\hat{y}_i^2}, \qquad r_i = \alpha\hat{y}_i - y_i, \tag{54}$$

and the metrics are then computed on the scaled residual r_i,

$$\mathrm{nRMSE} = \frac{1}{|\bar{y}|}\sqrt{\frac{1}{n}\sum_{i=1}^{n} r_i^2}, \qquad \mathrm{MAE} = \frac{1}{n}\sum_{i=1}^{n}|r_i|. \tag{55}$$

Here y_i is the reference value and $\hat{y}_i$ the simulated value, n is the number of matched sampling times, and $\bar{y}$ is the mean reference level. MAE keeps the units of the reference data and is computed for the same fitted α, so both metrics describe one and the same fit.

The consequence of fitting α must be stated, because it bounds what the carbohydrate comparison can support. A fitted multiplicative scale removes the absolute level from the residual, so nRMSE scores the shape and the timing of the diel cycle and not the absolute pool concentration. A trace with the correct rhythm and a systematically wrong amplitude can pass; a trace with the correct amplitude and the wrong phase cannot. For that reason α is reported beside every nRMSE in Section 6: a value near unity means shape and level agree, and a value far from unity localises the disagreement in the level alone. The acceptance thresholds are predefined heuristic criteria rather than external standards: nRMSE ≤ 0.25 for starch and nRMSE ≤ 0.50 for sucrose. The sucrose threshold is the looser of the two because whole-tissue sucrose pools cytosolic and vacuolar sucrose, and the split between the two compartments is itself uncertain.

5.4. Cross-validation design

Growth prediction is evaluated by leave-one-photoperiod-out cross-validation. For each fold the coefficients of Table 5 are fitted on three photoperiods, frozen, and used to predict the fourth. The split is by photoperiod rather than at random, because interpolation across the photoperiod range is exactly the ability in question. Four folds are run in each of the two shared regimes, which gives eight held-out predictions in total. The training objective is the relative sum of squares over the training observables, minimised by Nelder–Mead from three deterministic starts, with the best of the three retained [43]. The objective rejects any candidate whose run is classified as divergent or leaves the admissible state domain. A fit therefore never returns a parameter set that the model cannot integrate.

The 18 h condition is not part of any fold. It is held out of every fit in every regime, and it is evaluated only after the coefficients have been frozen. It therefore tests extrapolation beyond the calibrated range, which is a different and stronger question than the interpolation the four folds test. Two coefficients that are taken from the source publication rather than fitted here, the photosynthesis feedback and the nocturnal starch partition, keep their values in every configuration. The cross-validation therefore isolates the growth and starch-synthesis coefficients, and is a test of those rather than of the complete parameter set.

5.5. Ablation protocol

The ablation measures what the computed carbon input contributes, by replacing the carbon-input pathway and leaving everything downstream untouched. Three variants are compared. In M0 the carbon input is prescribed by the simple saturating photosynthesis expression of the reference formulation. In M1 the carbon input comes from Farquhar C3 photosynthesis, with the supply–demand closure solved as in the full model, but with stomatal conductance held fixed. In M2 conductance is computed from the Jarvis multiplicative response, so the gas-exchange layer becomes sensitive to PAR, vapour-pressure deficit, temperature, ambient CO_2 and leaf water potential.

All parameters are shared and frozen across the three variants. This is the point of the design and not a simplification. If each variant were re-fitted against the same observations, the fit would absorb the difference between the carbon-input pathways. The comparison would then measure how well a per-variant calibration can compensate, rather than what removing a mechanism costs. The comparison is reported on RGR error per photoperiod. RGR alone cannot separate M1 from M2, so we also report what the coupling changes. Two quantities carry that: the magnitude and diel shape of the carbon input, and the response of the gas-exchange layer to the environmental drivers.

5.6. Sensitivity analysis protocol

The global sensitivity analysis covers the 16 parameters listed in `config/sensitivity_params.csv`. The admissible range, the sampling scale and the literature basis of each are given individually in Appendix B. The set spans nutrient uptake capacity and soil supply, gas-exchange capacity, the environmental drivers, respiration, the growth coefficient, vacuolar exchange and the initial carbohydrate and biomass state. The analysed output is RGR. Every reported index is conditional on these declared ranges: the analysis describes model behaviour over the declared domain and not parameter importance in general.

Morris screening follows Morris [39] with the elementary-effect measures μ^* and σ of Campolongo et al. [8]. The design uses 30 trajectories on a 6-level grid with the fixed step $\Delta = p/[2(p-1)]$, giving $r(d+1)$ model evaluations. Trajectory base points are drawn from an independent pseudo-random generator, as the method specifies, so the screening is not affected by the base-sample construction described below. Variance-based indices use the estimators of

Sobol' [52] in the form recommended by Saltelli et al. [45], with $n = 512$ base samples and hence $n(d + 2)$ evaluations over the $d = 16$ parameters. Confidence intervals come from 1000 bootstrap replicates of the model evaluations, resampled by base row so that each resampled design remains a valid Saltelli design.

The base-sample design is stated explicitly, because the estimator depends on it. The Saltelli estimator needs two independent base matrices **A** and **B** of size $n \times d$, and it attributes variance to a parameter through a single-column substitution between them. Any correlation between the corresponding columns therefore biases the index towards zero. The construction specified by Saltelli et al. [45] draws one low-discrepancy sequence in $2d$ dimensions and splits it by column, but the sampler used here supports at most 16 dimensions and $2d = 32$ at $d = 16$. The two matrices are therefore drawn as two independent Latin-hypercube samples, stratified per dimension and independent by construction [34]. The design actually used is written to a `base_design` column of every output file, so no reported index is ambiguous about its own provenance. The estimator implementation was checked against the Ishigami function, whose indices are known in closed form [23].

Uncertainty is reported at two levels, because the two levels answer different questions. Bootstrap intervals resample within one design, so they measure sampling noise given that design and are blind to error contributed by the design itself. Across-design spread was therefore checked separately, by repeating the whole analysis on independently generated base samples over base sample sizes from $n = 512$ to $n = 4096$. An estimator whose across-design spread does not shrink as n grows is misspecified rather than noisy, so both levels are needed before an index is interpreted. Divergent evaluations are counted and reported with the indices, and no evaluation is silently discarded, because deleting rows from a Saltelli design breaks the pairing the estimator relies on. One-at-a-time sweeps and two-parameter response surfaces are also computed, but they are local diagnostics of the response shape and are not used as a global ranking.

6. Results

All results below use one code version and one configuration file. Unless a subsection says otherwise, the forcing is the chamber condition of the reference experiment and the simulated period is 30 days. The shared parameterisation has four fitted coefficients: λ_{sb}, λ_{sdr}, λ_{sdi} and the half-activation sucrose concentration of the sink term. These values are the same at every photoperiod and are not refitted between the studies below.

6.1. Photoperiod-dependent relative growth rate

With the shared parameterisation the model gives a mean absolute percentage error of 8.58% in relative growth rate (RGR) across the four photoperiods, and the worst photoperiod reaches 11.75%. All four scenarios integrate to the end of the simulated period without a non-finite sample. Under the acceptance criteria of Section 5 the same parameterisation gives a mean error of 8.88%, and 3 of the 4 photoperiods are accepted. The 4 h scenario is rejected on the RGR criterion alone. Table 6 lists the per-photoperiod values and Figure 2 shows them against the reference targets of Sulpice et al. [54].

The errors are not evenly distributed. The 4 h scenario is overestimated by 15.04%, while the 6, 8 and 12 h scenarios stay between 4.77 and 8.71%. One parameter set therefore reproduces the magnitude and the direction of the photoperiod response, but not its exact shape at the short end of the range. Section 6.6 shows that this residual has a structural cause and not a calibration cause.

The growth response is large in absolute terms, which is worth stating because RGR is a logarithmic rate and hides it. Final total biomass after 30 days rises from 0.387 gFW at 4 h to 1.258 at 6 h, 4.006 at 8 h and 39.03 gFW at 12 h, a hundredfold spread produced by a threefold change in day length. The errors quoted above are therefore errors in an exponent, and a 10% error in RGR compounds over the simulated period. Halving the output step changes the simulated RGR and the final biomass by less than 0.01% in all four scenarios, so these differences come from the model and not from the integration.

The saturated per-photoperiod regime behaves differently, and the contrast matters. That regime fits two coefficients per photoperiod, eight against eight observations, so it has zero degrees of freedom. Its per-photoperiod errors on the grid used here are 3.42, 1.28, 2.45 and 3.03%. These values are set by the grid resolution of 2.5×10^{-5} rather than by the model, and refining the bisection drives them towards zero. They are convergence residuals and carry no information about predictive skill. The point is not academic. Evaluated under the chamber forcing used throughout this paper, the same per-photoperiod parameterisation gives an RGR error of 48.67%, and none of the four photoperiods is accepted.

Near-zero residuals at the calibration point with a large error one condition away are the expected signature of a saturated fit.




Figure 2: Relative growth rate against photoperiod. Simulated values under the shared four-parameter parameterisation are compared with the reference targets of Sulpice et al. [54] for 4, 6, 8 and 12 h photoperiods. The residuals of the saturated per-photoperiod calibration are shown for reference only; they are convergence residuals of a fit with zero degrees of freedom.

6.2. Cross-photoperiod prediction

The held-out test asks a different question: how well does the model predict a photoperiod that was not used to fit anything? We fitted the growth coefficient on three photoperiods, predicted the fourth, and repeated this for each of the four folds. Table 7 reports both regimes and Figure 3 shows the predictions.

In regime A the growth coefficient is one shared constant. The held-out error is 13.18% and the worst fold is 20.93%, with the 4 h treatment held out. In regime B the same coefficient depends linearly on photoperiod P,

$$\lambda_{sb}(P) = c_0 + c_1 P, \tag{56}$$

with two coefficients instead of one. The held-out error falls to 4.36% and the worst fold is 10.30%, now at the 12 h treatment. The fitted values are $c_0 = 0.00243$ and $c_1 = 8.786 \times 10^{-5}$ per hour. Across the four folds the fitted growth coefficient rises monotonically with photoperiod, from 0.00275 at 4 h to 0.00370 at 12 h. Adding one coefficient therefore predicts an unseen photoperiod about three times better than a single constant.

Two readings follow. First, the growth coefficient is not photoperiod-invariant in this model. If the coupled dynamics generated the full photoperiod response, a shared constant would be enough and regime B would add nothing. Second, the missing dependence is compact. A straight line in one coefficient captures it over the tested range, which places it in the conversion from available sucrose to structural growth rather than in the gas-exchange layer upstream. A positive slope means that a longer day converts loaded sucrose into biomass more efficiently, not only for longer. Increased sink activity under long days is a plausible reading [51, 54], but the design cannot separate it from an allocation effect or from a fresh-weight conversion effect.

The limits of Eq. 56 belong with the result. The line was fitted over 4 to 12 h and has no support outside that interval. It corrects a model coefficient and does not describe a process, so extrapolating it to 18 or 24 h has no basis. It must also break down at short photoperiods, because the coefficient would reach zero at a positive day length.

6.3. Diel carbohydrate dynamics

The carbohydrate comparison uses one multiplicative scale per series, fitted by least squares before the error is computed, so the normalised RMSE scores the shape and the timing of the diel cycle. Over the reference series the




Figure 3: Held-out photoperiod prediction of relative growth rate. Regime A uses one shared growth coefficient; regime B lets the same coefficient vary linearly with photoperiod (Eq. 56). Each point is a photoperiod that was excluded from the fit that produced it.

mean starch nRMSE is 0.235 against the acceptance threshold of 0.25, with a bias of −3.1%. The mean sucrose nRMSE is 0.285 against a threshold of 0.50, with a bias of +10.7%. In the validation runs the largest per-photoperiod values are 0.219 for starch and 0.468 for sucrose. Figure 4 shows the traces.

The two series are not equally strong, and the fitted scales show why. For starch the scale is between 0.86 and 1.15 over the four fitted photoperiods, so shape and absolute level both agree and the starch nRMSE means what a reader would assume it means. For sucrose the scale is between 2.25 and 3.31: the simulated pool must be multiplied by a factor of two to three before it matches the measurements. Raw simulated sucrose maxima are 1.30 to 1.40 μmol C_6 gFW^{-1} against measured maxima of 3.18 to 4.71 μmol C_6 gFW^{-1}. The passing sucrose nRMSE of 0.285 therefore describes shape and timing only, and the absolute sucrose level is a known deficit of this version. We report the scale factor beside every carbohydrate error for that reason.

Within the fitted range the model reproduces the qualitative features of the diel cycle. Starch accumulates through the light period and degrades through the night along a near-linear trajectory. At dawn it returns to the level at which each run started, so the pool is close to empty at the end of every night. This behaviour follows from the drain law of Section 3.8, which targets the remaining pool over the remaining night, and it agrees with the near-linear nocturnal degradation reported for *Arabidopsis* [20, 46].

One quantitative feature is not reproduced. The measured end-of-day starch peak grows more steeply with photoperiod than the simulated peak. In the model the peak does rise with photoperiod, because the light period is longer, but the rise is weaker than measured. The model therefore under-represents how much a long day changes the size of the transient store, as opposed to its filling time. The same shortfall appears from the other side in Section 6.6, where assimilation per light hour is found to be constant.

6.4. Ablation

The ablation compares three carbon-input pathways with the same downstream model, the same shared parameters and no refitting. In M0 the carbon input is prescribed. In M1 it is computed from Farquhar photosynthesis [15] at fixed stomatal conductance. In M2 the conductance follows the Jarvis formulation [25] and responds to the environment. Table 8 and Figure 5 give the result: the RGR error falls from 41.01% for M0 to 9.43% for M1 and to 8.88% for M2.

The large step is from prescribed to computed carbon input. M0 underestimates RGR at every photoperiod, by 39.2 to 43.1%, and its peak assimilation stays between 10.50 and 11.14 μmol gFW^{-1} h^{-1}. With computed assimilation the peak rises to 17.62 and the systematic underestimation disappears. The bias is not reduced but removed: M1 and M2 scatter on both sides of the targets, while M0 is low in all four scenarios. Parameters were frozen across the three variants, so the change comes from the carbon-input pathway and not from recalibration.




Figure 4: Diel starch and sucrose dynamics for the 4, 6, 8, 12 and 18 h photoperiods. Simulated traces are compared with the measured diel profiles of the repository reference file (see Section 5). Each panel prints the nRMSE and the fitted multiplicative scale. The scale is between 0.86 and 1.15 for starch at the four fitted photoperiods and between 2.25 and 3.31 for sucrose, so the sucrose comparison is one of shape and timing and not of absolute level. The 18 h column was not used in any fit; its starch nRMSE is not reported, because the fitted scale of 0.07 absorbs the divergence described in Section 6.6.

The Jarvis layer adds a smaller and unevenly distributed gain. Averaged over the four photoperiods it reduces the error from 9.43 to 8.88%, and the two variants differ most at the extremes of the range. At 4 h, M2 improves on M1 by 3.8 percentage points, from 18.82 to 15.04%. At 12 h it is worse by 2.1 points, 8.71 against 6.60%. At 6 h M2 is better by 3.0 points and at 8 h it is worse by 2.5 points. Adding environmental control of conductance lowers peak assimilation slightly, from 17.62 to 17.01 μmol gFW^{-1} h^{-1}, because the conductance multipliers are below one at the nominal condition. The value of M2 at a single nominal condition is therefore not a gain in RGR accuracy. Its value is that the carbon input responds to irradiance, humidity, temperature, ambient CO_2 and leaf water potential at all, which is what the sensitivity analysis of the next subsection exercises.

6.5. Global sensitivity

The variance-based analysis follows Sobol' [52] with the estimator of Saltelli et al. [45], and the Morris screening follows Morris [39], Campolongo et al. [8]. Sixteen parameters were varied over the declared ranges of Appendix B. Table 9 gives the full ranking and Figure 6 shows the total-order indices next to the Morris measures.

Nutrient uptake capacity dominates the total-order ranking. The two largest total indices are 0.658 for maximum phosphorus uptake and 0.577 for maximum nitrogen uptake. Respiration frequency follows at 0.466, then ambient CO_2 concentration at 0.422 and leaf water potential at 0.412. The two Farquhar capacities come next, 0.352 for electron transport rate and 0.334 for maximum carboxylation rate, then maximum PAR at 0.333. The ranking spans the nutrient, respiratory and gas-exchange blocks, as expected in a model where growth is limited by whichever supply is scarcest.

The main-effect share is small. Summed over the sixteen parameters and averaged over the four photoperiods, the first-order indices give 0.201 against a summed total-order index of 4.675, a main-effect share of about 4%. The Morris measures agree: 15 of the 16 parameters have $\sigma/\mu^* > 1$, and the exception is the one parameter with a total index at noise level. The two methods also rank the parameters consistently, with a Spearman rank correlation of $\rho = 0.944$ between Morris μ^* and Sobol' S_T. The model is interaction-dominated over the declared ranges: parameters act mainly through each other, not on their own.

A low main-effect share can also be an estimator artefact, so we checked convergence with sample size. As n rises from 512 to 4096, the number of first-order estimates clipped at zero falls from 8 to 3. Over the same range the summed




Figure 5: Relative RGR error per photoperiod for the three carbon-input variants, with shared parameters frozen. M0 underestimates growth at every photoperiod; M1 and M2 scatter on both sides of the targets.

first-order index stays between 0.151 and 0.340, and the main-effect share stays between 3.0 and 6.4%. The clipping decreases, which is what an estimator with shrinking variance should do, but the share does not increase with it. The low main-effect share is therefore a property of the model over these ranges and not an artefact of a small sample. All indices in Table 9 are reported unclipped, so small negative interaction remainders appear where an index is at noise level; we prefer that to hiding the estimator variance.

One entry deserves a comment. The growth coefficient λ_{sb}, the coefficient that Section 6.2 shows to carry the photoperiod response, has a total index of only 0.117 and ranks fifteenth of sixteen. There is no contradiction. A Sobol' index measures the variance contribution over a declared range at fixed forcing, while the cross-photoperiod result measures how a coefficient must change when the forcing changes. A parameter can be unimportant for variance at one condition and still be the parameter that fails to transfer between conditions.

6.6. Validity envelope

Two limits of the model were measured directly, and we report them as findings rather than as caveats. Figure 7 shows both.

The first limit is the 18 h photoperiod, which was never used in any fit. The sucrose comparison passes: nRMSE is 0.428 against the threshold of 0.50, with a fitted scale of 2.53, so the shape of the diel sucrose cycle survives three hours beyond the fitted range. The starch pool does not. Dawn starch is 401 μmol gFW^{-1} on day 30 and 1207 μmol gFW^{-1} on day 90, a net gain of about +14 per day that shows no sign of levelling off. This is unbounded accumulation and not a wrong steady level, and the distinction matters, because a wrong level can be rescaled while a divergence cannot. The fitted scale for 18 h starch is 0.07, which is the metric absorbing the divergence, so no 18 h starch nRMSE is quoted anywhere in this paper.

The mechanism is visible in the fluxes. The degradation cap is not binding at 18 h, so the failure is not a saturated drain. The drain law targets the remaining pool over the remaining night, so nightly export is proportional to what is left and to the reciprocal of the night length. Under a long day the pool filled during the light period grows while the night available to empty it shrinks. The day fill then outruns the drain, the pool carries a surplus into the next day, and the surplus compounds.

The second limit is the closure boundary. Under the chamber forcing the net daily starch gain is +0.300 at 12 h, +0.295 at 14 h and +0.389 at 15.0 h, rising to +0.547 at 15.5 h, +2.042 at 16 h, +13.780 at 18 h and +24.150 at 20 h. The diel balance therefore closes up to 15.0–15.5 h and fails from 16 h. That boundary sits three hours beyond the longest photoperiod used in any fit, which is a real margin and a narrow one.




Figure 6: Global sensitivity of relative growth rate over the declared parameter ranges. Left: total-order Sobol' indices with the first-order part shown separately. Right: Morris μ^* and σ. The two rankings agree at a Spearman correlation of 0.944, and 15 of the 16 parameters have $\sigma/\mu^* > 1$.

The boundary is a property of the model and of the light dose together, not of the model alone. Under the shipped half-sine profile at 1200 μmol m^{-2} s^{-1} the same measurement gives +18.99 at 12 h, +32.56 at 14 h, +38.75 at 15 h, +45.00 at 16 h, +56.70 at 18 h and +67.68 at 20 h. Under that dose the balance never closes, not even at 12 h. Any statement of the form "the model closes the diel starch balance up to P hours" is therefore incomplete without the irradiance at which it was measured. We report the chamber column as the envelope of this model, because it is the condition with an experimental counterpart. The shipped column shows how strongly the boundary depends on the daily photon dose.

The third measurement is a bound rather than a boundary, and it explains the residual error of Section 6.1. Assimilation per hour of light is constant to six significant figures across the photoperiod series, so total daily assimilation is proportional to day length. The ratio of daily assimilation between the 12 and 4 h scenarios is therefore exactly 3.0000, while the reference RGR targets require a ratio of 3.82. Growth in this model is driven by the carbon input, so a supra-proportional factor of 1.27 has to come from somewhere else. Partitioning cannot supply it: the partition coefficients redistribute a fixed daily input between starch, sucrose and biomass, and they cannot make the total larger. This is the same gap that Eq. 56 closes empirically, seen at the level of the carbon budget instead of the fitted coefficient.

7. Discussion

7.1. What the coupled formulation buys

The ablation puts a number on the coupling. Replacing a prescribed carbon input with a computed one reduces the RGR error from 41.01 to 9.43% with no parameter refitted. It removes a bias rather than reducing a scatter: M0 is low at all four photoperiods, M1 and M2 are not. The reason is visible in the assimilation rate. A prescribed maximum rate has to be chosen once, and a single choice cannot be correct at four day lengths. The plant it describes has a different leaf area and a different sucrose status in each case. The computed input is a function of the states and of the forcing, so it changes with them.

The Jarvis layer should be read differently. It reduces the mean error from 9.43 to 8.88%, which is a small change, and it moves the error in opposite directions at the two ends of the photoperiod range. Reporting that as an accuracy gain would misstate it. What the layer provides is a carbon input that responds to irradiance, humidity, temperature, ambient CO_2 and leaf water potential. The sensitivity analysis shows this is not a formality: ambient CO_2 has a total-order index of 0.422, leaf water potential 0.412 and maximum PAR 0.333. In M1 those three parameters would move




Figure 7: Net daily starch gain against photoperiod on a symmetric-log axis, under the chamber forcing and under the shipped half-sine profile at 1200 μmol m^{-2} s^{-1}. The band marks the closure boundary at 15.0–15.5 h under chamber forcing. Under the higher dose the balance does not close at any photoperiod shown, so the boundary depends on the light dose as well as on the model.

nothing through the conductance path. A model intended for use across environments needs that path even when it does not improve a single nominal-condition metric, and a model evaluated only at its calibration point cannot show the difference.

This is where the model sits relative to existing families. It is not a field-scale crop simulator, and it does not resolve architecture as a functional-structural model does. It is a compact whole-plant core in which the carbon input is computed from the environment and then propagated explicitly into carbohydrate, nutrient and biomass states, with the numerical treatment and the evaluation protocol stated. Many whole-plant and crop simulators prescribe that input [22, 26, 63], and the ablation measures what prescribing it costs in this setting. The physiological submodels are inherited [15, 25, 6]; the coupling, the analytic closure and the evaluation are what this paper contributes.

7.2. Parameter transferability and what the design exposed

The cross-photoperiod result is the part of this work with the widest reach beyond plant modelling. One shared growth coefficient predicts a held-out photoperiod with 13.18% error; the same coefficient made linear in photoperiod predicts it with 4.36%. Both regimes use the same equations, the same states and the same solver. The difference is entirely in how one coefficient is allowed to depend on a condition of the experiment.

That is a statement about parameter transferability, and it is a common failure mode in simulation practice. A coefficient fitted at one operating condition is usually reported as a property of the model, and it is then applied at other conditions as if it were one. When it is not, the model appears to be structurally wrong, because the error shows up in the state trajectories rather than in the parameter. Our results separate the two cases. The gas-exchange layer, the carbohydrate subsystem and the nutrient subsystem all transfer with fixed coefficients across 4 to 12 h. One coefficient in the growth-conversion term does not, and it needs one extra degree of freedom to do so.

The design is what made this visible. A random train-test split over all available observations would have placed neighbouring photoperiods on both sides of the split, and interpolation between them would have hidden the dependence. Holding out an entire condition asks the question that matters for a simulator: does the fitted model work at a condition it has never seen? We suggest leave-one-condition-out evaluation as a default for models calibrated across a treatment series. The reason is not that it is more severe. It is the design that localises a transferability failure to a specific coefficient. A single held-out error tells you the model missed. A sweep of held-out folds, each with its own refitted coefficient, tells you which coefficient moved and in which direction, and here it moved monotonically with photoperiod.

Two properties of the design make it reusable outside this model. First, the held-out condition must be a condition and not a random subset, because random splitting lets neighbouring treatments substitute for each other. Second, one coefficient at a time should be released. Regime B differs from regime A by a single degree of freedom, so the improvement from 13.18 to 4.36% can be attributed to that coefficient and to nothing else. Releasing several coefficients at once would have produced a similar error reduction and located nothing. Together the two properties turn a fit quality measurement into a diagnostic that names a term in the model, which is what a modeller needs before deciding whether to recalibrate or to reformulate.

The physiological reading must stay careful. A growth coefficient that rises with day length means that the conversion from loaded sucrose to structural biomass becomes more efficient as the photoperiod lengthens. Increased sink activity under long days is a plausible candidate [51, 54], and so is a change in the ratio of fresh weight to carbon, and so is a shift in allocation between leaf and root. The present data cannot separate these, and one linear coefficient fitted over four photoperiods will fit all three explanations equally well. We therefore report the dependence as a measured property of the calibration and not as an identified mechanism.

7.3. What the carbohydrate results do and do not show

The two carbohydrate series should be read with different confidence. For starch the fitted scales lie between 0.86 and 1.15, so the comparison covers level as well as shape. A mean nRMSE of 0.235 against a threshold of 0.25 can therefore be taken at face value. For sucrose the scales lie between 2.25 and 3.31, so the same style of statement would be misleading. The passing value of 0.285 means that the model puts the sucrose maxima and minima at the right times, with roughly the right relative depth. It says nothing about the absolute pool size, which is a factor of two to three too low.

The distinction matters beyond this model, because a scale-fitted error metric is common when reference data are indirect or on a different basis. Fitting one multiplicative scale is defensible: it separates a units or basis mismatch from a dynamic error, and it is the only way to compare a normalised model pool against a measured tissue concentration. But the fitted scale is part of the result and must be reported with it. A scale near one and a scale near three support different claims from the same nRMSE, and only the first supports a statement about level.

7.4. Interaction dominance and its consequences for calibration

Over the declared parameter ranges, the summed first-order indices account for about 4% of the summed total-order indices. In addition, 15 of the 16 parameters have a Morris σ/μ^* above one. Two independent methods agree on the ranking at $\rho = 0.944$, and the convergence check shows the low main-effect share is stable as the sample grows. The conclusion is not that main effects are absent, but that they are small relative to interactions across the tested domain.

This has a practical consequence for anyone calibrating a model of this class. One-at-a-time tuning is unreliable here. Suppose a parameter's own effect explains a few percent of the variance and the rest comes from its interactions. The sensitivity measured by moving that parameter alone then depends on where the other fifteen happen to sit. Tuning it at one operating point can make behaviour worse at another, and the sequence in which parameters are tuned changes the result. The structure of the model explains why. Growth is limited by whichever supply is scarcest, and the largest total-order indices belong to the nutrient uptake capacities, 0.658 and 0.577. A nutrient parameter changes which limitation binds, and that changes how strongly every gas-exchange parameter matters. The interaction is the model working as designed, and it means calibration should be done jointly, with a global method, over a declared domain.

Two limits on this reading should be kept. The indices are conditional on the ranges in Appendix B; wider ranges would raise the interaction share and narrower ones would lower it. And they describe model behaviour, not plant physiology. That nutrient uptake capacity dominates variance in this model is a statement about this model over these ranges.

7.5. Limitations

Six limitations bound what can be concluded, and each is measured rather than asserted.

Four coefficients were fitted. The growth coefficient, the two starch partitioning rates and the half-activation constant of the sink term were estimated against the reference data, so the in-sample results of Section 6.1 are not independent tests. This is a large improvement on the eight-per-photoperiod calibration of the earlier version, which had zero degrees of freedom against eight observations, but four fitted coefficients are still four. The held-out results of Section 6.2 and the 18 h evaluation of Section 6.6 are the parts of the evaluation that no fitting touched.

The carbohydrate metric is scale-fitted, and for sucrose the fitted scale is large. The starch scales lie between 0.86 and 1.15, so the starch comparison covers level as well as shape. The sucrose scales lie between 2.25 and 3.31, so the simulated pool is a factor of two to three below the measurements in absolute terms and the passing nRMSE of 0.285 evaluates shape and timing only. Raw simulated maxima are 1.30 to 1.40 μmol C_6 gFW^{-1} against measured maxima of 3.18 to 4.71 μmol C_6 gFW^{-1}. The deficit is one of the clearest targets for the next version. It should also be read together with the whole-tissue basis of the measurements. The model quantity compared here is the sum of the cytosolic and vacuolar pools, so a partition error between the two would not show in the total.

Some parameters remain unsourced. The provenance ledger classifies 12 of 111 entries as having no traceable origin and 15 as fitted, and both groups are listed in Appendix A with that status attached. Unsourced constants are a work list and not a defence; we report them so that a reader can see which numbers rest on a measurement and which do not.

The diel starch balance closes only to 15.0–15.5 h under chamber forcing, and under the higher shipped light dose it does not close at any photoperiod tested. The model is therefore not valid for long-day simulation as it stands, and any application beyond the closure boundary would need the drain law changed first.

The RGR response to photoperiod is bounded by construction. Assimilation per light hour is constant to six significant figures, so daily assimilation is proportional to day length, and the 12 to 4 h ratio is exactly 3.0000 where the targets need 3.82. No choice of partitioning parameters can produce the missing factor of 1.27, because partitioning redistributes a fixed daily input. The linear growth coefficient covers this gap empirically, and the underlying bound remains.

Simulated tissue nutrient concentration falls with increasing supply where the measurements rise. The nutrient block is out of sample, no coefficient in it was fitted to the nutrient response, so it is entitled to judge the model and the discrepancy is reported and not tuned away. The likely cause is in the structure of the nutrient balance rather than in a parameter value. Every term in the internal nutrient equation is bounded, and the concentration is consumed in proportion to assimilation. A supply increase raises assimilation, which raises consumption, and with a bounded uptake term the pool can fall. Resolving this requires the uptake and consumption terms to be reformulated, which is beyond a recalibration.

7.6. Pushing the envelope past 15.5 hours

The closure failure has a specific cause, so the candidate remedies are also specific. The nocturnal drain targets the remaining pool over the remaining night. Its rate therefore scales with the reciprocal of night length, while the daytime fill scales with day length, and beyond about 15 h the fill wins. Three changes could plausibly move the boundary, and each is testable within the present model.

The first is to make the daytime partition to starch respond to the size of the starch pool relative to what the coming night can export. In the current formulation the partition coefficient responds to sucrose status, not to the starch balance, so nothing tells the plant to stop filling a store it cannot empty. Measurements support such a dependence, since *Arabidopsis* adjusts partitioning to starch across photoperiods [54, 36]. The second is to raise the degradation capacity. The cap is not binding at 18 h, so it cannot be the limit on its own. A higher cap combined with a drain law that also depends on the pool level would extend the range over which the drain can match the fill. The third is to allow the peak amplitude of the store to grow with photoperiod, which is also the feature that Section 6.3 identifies as under-represented against the measurements. That the same shortfall shows up in the diel comparison and at the closure boundary suggests one structural cause rather than two.

We do not claim any of these will work. Each adds a coefficient that would have to be identified against data, and the measurements available at 18 h are limited. A change that closes the balance could easily be a change that fits one series. What we can state is narrower. The boundary sits at 15.0–15.5 h under chamber forcing, and its cause lies in the drain law and its interaction with the day fill. Testing the three candidates above is the next step, not a result.

8. Conclusion

We presented a whole-plant simulation model in which environmental forcing is converted into a photosynthetic carbon input through Jarvis stomatal conductance and Farquhar C3 photosynthesis. That carbon input is then propagated through diel carbohydrate, nutrient and biomass dynamics. The supply and demand sides of gas exchange are closed analytically at runtime as a quadratic in the intercellular CO_2 concentration, so the operating point is obtained

without an inner iteration. The eleven slow states are advanced by an adaptive integrator. The formulation, its numerical treatment and the evaluation protocol are the contribution; the physiological submodels are inherited.

The evaluation used four shared fitted coefficients rather than the eight-per-photoperiod calibration of the earlier version. In sample the model reproduces the relative growth rate across four photoperiods with a mean absolute percentage error of 8.58%. In a leave-one-photoperiod-out test, one shared growth coefficient predicts a held-out photoperiod less well than the same coefficient made linear in photoperiod, which reaches 4.36% with two coefficients. The ablation shows what the coupling contributes: replacing a prescribed carbon input with a computed one moves the error from 41.01 to 8.88% with parameters frozen. The variance-based analysis shows an interaction-dominated model over the declared ranges, led by nutrient uptake capacity, respiration and gas-exchange capacity. Calibration of this class of model should therefore be done jointly, not one parameter at a time.

The paper also states where the model stops working. Under the chamber forcing the diel starch balance closes up to a photoperiod of 15.0 to 15.5 h and fails beyond it. Under a higher light dose it does not close at all, so the boundary belongs to the model and the light dose together. At 18 h the sucrose cycle still has the right shape while the starch pool accumulates without bound. Assimilation per light hour is constant, which bounds the growth response to photoperiod below what the reference data require, and the linear growth coefficient covers that gap empirically rather than mechanistically. The absolute sucrose level is a factor of two to three too low, so the sucrose comparison is one of shape and timing.

Three directions follow. The drain law and the daytime partition to starch are the place to look for a formulation that closes the balance at long days. The absolute sucrose level and the nutrient concentration response are two independent out-of-sample discrepancies that point at the structure of the sucrose and nutrient balances rather than at their coefficients. And the leave-one-condition-out design used here can be applied to other conditions, irradiance and temperature first, to test whether the remaining coefficients transfer as well across those as they do across photoperiod.

A. Parameter provenance ledger

Every numerical constant in the released model carries a provenance record in `config/parameter_provenance.csv`, and Table 10 is generated from that file rather than transcribed. The tier states how the value was obtained: T1 measured, T2 derived, T3 inherited, T4 constrained, T5 fitted, T6 unsourced. A T6 entry is a value that we could not trace to a measurement, a derivation or the source model, and it is declared as such rather than presented as sourced. The ledger holds 111 entries, of which 2 are measured (T1), 52 are derived (T2), 20 are inherited (T3), 10 are constrained (T4), 15 are fitted (T5), 12 are unsourced (T6).

B. Sensitivity analysis: parameters, ranges and sampling scales

Table 11 is generated from `config/sensitivity_params.csv` and gives the factors varied in the sensitivity study together with their ranges, sampling scales and the role each factor plays in the model. All reported Sobol' and Morris indices are conditional on exactly these ranges: an index is a statement about output variance over a declared input distribution, and it is not transferable to a different one. A factor sampled on a logarithmic scale is varied uniformly in $\log_{10}$, which is the appropriate choice for a rate constant spanning more than one order of magnitude. Of the 20 factors listed, 16 are active in the reported design and 4 are disabled; the reasons for disabling are given in the table.

Declaration of competing interest

The authors declare no competing interests.

Data and code availability

The simulation code, configuration files, reference data and the scripts that generate every table and figure in this paper are available at TODO(AUTHOR) repository URL.

References

[1] Annunziata, M.G., Apelt, F., Carillo, P., Krause, U., Feil, R., Mengin, V., Lauxmann, M.A., Köhl, K., Nikoloski, Z., Stitt, M., Lunn, J.E., 2017. Getting back to nature: a reality check for experiments in controlled environments. Journal of Experimental Botany 68, 4463–4477. doi:10.1093/jxb/erx220.

[2] Ball, J.T., Woodrow, I.E., Berry, J.A., 1987. A model predicting stomatal conductance and its contribution to the control of photosynthesis under different environmental conditions, in: Biggins, J. (Ed.), Progress in Photosynthesis Research. Springer, Dordrecht, pp. 221–224. doi:10.1007/978-94-017-0519-6_48.

[3] Barber, S.A., 1995. Soil nutrient bioavailability : a mechanistic approach. 2nd ed., Wiley.

[4] Bernacchi, C.J., Singsaas, E.L., Pimentel, C., Portis, A.R., Long, S.P., 2001. Improved temperature response functions for models of Rubisco-limited photosynthesis. Plant, Cell & Environment 24, 253–259. doi:10.1111/j.1365-3040.2001.00668.x.

[5] Buckley, T.N., 2017. Modeling stomatal conductance. Plant Physiology 174, 572–582. doi:10.1104/pp.16.01772.

[6] von Caemmerer, S., 2000. Biochemical Models of Leaf Photosynthesis. Number 2 in Techniques in Plant Sciences, CSIRO Publishing, Collingwood, Australia. doi:10.1071/9780643103405.

[7] von Caemmerer, S., Farquhar, G.D., 1981. Some relationships between the biochemistry of photosynthesis and the gas exchange of leaves. Planta 153, 376–387. doi:10.1007/bf00384257.

[8] Campolongo, F., Cariboni, J., Saltelli, A., 2007. An effective screening design for sensitivity analysis of large models. Environmental Modelling & Software 22, 1509–1518. doi:10.1016/j.envsoft.2006.10.004.

[9] Carstensen, A., Herdean, A., Schmidt, S.B., Sharma, A., Spetea, C., Pribil, M., Husted, S., 2018. The impacts of phosphorus deficiency on the photosynthetic electron transport chain. Plant physiology 177, 271–284. doi:10.1104/pp.17.01624.

[10] Cellier, F.E., 1991. Continuous System Modeling. Springer-Verlag, New York.

[11] Damour, G., Simonneau, T., Cochard, H., Urban, L., 2010. An overview of models of stomatal conductance at the leaf level. Plant, Cell & Environment 33, 1419–1438. doi:10.1111/j.1365-3040.2010.02181.x.

[12] de Reffye, P., Hu, B., Kang, M., Letort, V., Jaeger, M., 2020. Two decades of research with the greenlab model in agronomy. Annals of Botany 127, 281–295. doi:10.1093/aob/mcaa172.

[13] Dubois, J.J.B., Fiscus, E.L., Booker, F.L., Flowers, M.D., Reid, C.D., 2007. Optimizing the statistical estimation of the parameters of the Farquhar–von caemmerer–berry model of photosynthesis. New Phytologist 176, 402–414. doi:10.1111/j.1469-8137.2007.02182.x.

[14] Epstein, E., Hagen, C.E., 1952. A kinetic study of the absorption of alkali cations by barley roots. Plant Physiology 27, 457–474. doi:10.1104/pp.27.3.457.

[15] Farquhar, G.D., von Caemmerer, S., Berry, J.A., 1980. A biochemical model of photosynthetic CO2 assimilation in leaves of C3 species. Planta 149, 78–90. doi:10.1007/BF00386231.

[16] Farquhar, G.D., Wong, S.C., 1984. An empirical model of stomatal conductance. Australian Journal of Plant Physiology 11, 191–210. doi:10.1071/pp9840191.

[17] Feller, C., Favre, P., Janka, A., Zeeman, S.C., Gabriel, J.P., Reinhardt, D., 2015. Mathematical modeling of the dynamics of shoot-root interactions and resource partitioning in plant growth. PLOS ONE 10, e0127905. doi:10.1371/journal.pone.0127905.

[18] Feugier, F.G., Satake, A., 2013. Dynamical feedback between circadian clock and sucrose availability explains adaptive response of starch metabolism to various photoperiods. Frontiers in Plant Science 3, 305. doi:10.3389/fpls.2012.00305.

[19] Gastal, F., Lemaire, G., 2002. N uptake and distribution in crops: an agronomical and ecophysiological perspective. Journal of Experimental Botany 53, 789–799. doi:10.1093/jexbot/53.370.789.

[20] Graf, A., Schlereth, A., Stitt, M., Smith, A.M., 2010. Circadian control of carbohydrate availability for growth in Arabidopsis plants at night. Proceedings of the National Academy of Sciences 107, 9458–9463. doi:10.1073/pnas.0914299107.

[21] Hairer, E., Nørsett, S.P., Wanner, G., 1993. Solving Ordinary Differential Equations I: Nonstiff Problems. 2nd ed., Springer-Verlag, Berlin.

[22] Holzworth, D.P., Huth, N.I., deVoil, P.G., Zurcher, E.J., Herrmann, N.I., McLean, G., Chenu, K., van Oosterom, E.J., Snow, V., Murphy, C., Moore, A.D., Brown, H., Whish, J.P., Verrall, S., Fainges, J., Bell, L.W., Peake, A.S., Poulton, P.L., Hochman, Z., Thorburn, P.J., Gaydon, D.S., Dalgliesh, N.P., Rodriguez, D., Cox, H., Chapman, S., Doherty, A., Teixeira, E., Sharp, J., Cichota, R., Vogeler, I., Li, F.Y., Wang, E., Hammer, G.L., Robertson, M.J., Dimes, J.P., Whitbread, A.M., Hunt, J., van Rees, H., McClelland, T., Carberry, P.S., Hargreaves, J.N., MacLeod, N., McDonald, C., Harsdorf, J., Wedgwood, S., Keating, B.A., 2014. Apsim – evolution towards a new generation of agricultural systems simulation. Environmental Modelling & Software 62, 327–350. doi:10.1016/j.envsoft.2014.07.009.

[23] Ishigami, T., Homma, T., 1990. An importance quantification technique in uncertainty analysis for computer models, in: Proceedings of the First International Symposium on Uncertainty Modeling and Analysis (ISUMA 1990), IEEE, College Park, MD, USA. pp. 398–403. doi:10.1109/ISUMA.1990.151285.

[24] Janott, M., Gayler, S., Gessler, A., Javaux, M., Klier, C., Priesack, E., 2011. A one-dimensional model of water flow in soil-plant systems based on plant architecture. Plant and Soil 341, 233–256. doi:10.1007/s11104-010-0639-0.

[25] Jarvis, P.G., 1976. The interpretation of the variations in leaf water potential and stomatal conductance found in canopies in the field. Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences 273, 593–610. doi:10.1098/rstb.1976.0035.

[26] Jones, J.W., Hoogenboom, G., Porter, C.H., Boote, K.J., Batchelor, W.D., Hunt, L.A., Wilkens, P.W., Singh, U., Gijsman, A.J., Ritchie, J.T., 2003. The DSSAT cropping system model. European Journal of Agronomy 18, 235–265. doi:10.1016/s1161-0301(02)00107-7.

[27] Kim, S.H., Lieth, J.H., 2003. A coupled model of photosynthesis, stomatal conductance and transpiration for a rose leaf (rosa hybrida l.). Annals of botany 91, 771–781. doi:10.1093/aob/mcg080.

[28] Lawson, T., Blatt, M.R., 2014. Stomatal size, speed, and responsiveness impact on photosynthesis and water use efficiency. Plant Physiology 164, 1556–1570. URL: https://dx.doi.org/10.1104/pp.114.237107, doi:10.1104/PP.114.237107.

[29] Lohaus, G., Hussmann, M., Pennewiss, K., Schneider, H., Zhu, J.J., Sattelmacher, B., 2000. Solute balance of a maize (Zea mays L.) source leaf as affected by salt treatment with special emphasis on phloem retranslocation and ion leaching. Journal of Experimental Botany 51, 1721–1732. doi:10.1093/jexbot/51.351.1721.

[30] Louarn, G., Song, Y., 2020. Two decades of functional–structural plant modelling: now addressing fundamental questions in systems biology and predictive ecology. Annals of Botany 126, 501–509. doi:10.1093/aob/mcaa143.

[31] Maiorano, A., Martre, P., Asseng, S., Ewert, F., Müller, C., Rötter, R.P., Ruane, A.C., Semenov, M.A., Wallach, D., Wang, E., Alderman, P.D., Kassie, B.T., Biernath, C., Basso, B., Cammarano, D., Challinor, A.J., Doltra, J., Dumont, B., Rezaei, E.E., Gayler, S., Kersebaum, K.C.,

Kimball, B.A., Koehler, A.K., Liu, B., O'Leary, G.J., Olesen, J.E., Ottman, M.J., Priesack, E., Reynolds, M., Stratonovitch, P., Streck, T., Thorburn, P.J., Waha, K., Wall, G.W., White, J.W., Zhao, Z., Zhu, Y., 2017. Crop model improvement reduces the uncertainty of the response to temperature of multi-model ensembles. Field Crops Research 202, 5–20. doi:10.1016/J.FCR.2016.05.001.

[32] Mamassi, A., Lang, M., Tychon, B., Lahlou, M., Wellens, J., El Gharous, M., Marrou, H., 2024. A comparison of empirical and mechanistic models for wheat yield prediction at field level in moroccan rainfed areas. in silico Plants 6, diad020. doi:10.1093/insilicoplants/diad020.

[33] Marcelis, L.F., 1996. Sink strength as a determinant of dry matter partitioning in the whole plant. Journal of Experimental Botany 47, 1281–1291. doi:10.1093/jxb/47.special_issue.1281.

[34] McKay, M.D., Beckman, R.J., Conover, W.J., 1979. A comparison of three methods for selecting values of input variables in the analysis of output from a computer code. Technometrics 21, 239–245. doi:10.1080/00401706.1979.10489755.

[35] Medlyn, B.E., Duursma, R.A., Eamus, D., Ellsworth, D.S., Prentice, I.C., Barton, C.V., Crous, K.Y., Angelis, P.D., Freeman, M., Wingate, L., 2011. Reconciling the optimal and empirical approaches to modelling stomatal conductance. Global Change Biology 17, 2134–2144. doi:10.1111/j.1365-2486.2010.02375.x.

[36] Mengin, V., Pyl, E.T., Alexandre Moraes, T., Sulpice, R., Krohn, N., Encke, B., Stitt, M., 2017. Photosynthate partitioning to starch in Arabidopsis thaliana is insensitive to light intensity but sensitive to photoperiod due to a restriction on growth in the light in short photoperiods. Plant, Cell & Environment 40, 2608–2627. doi:10.1111/pce.13000.

[37] Meron, E., Bennett, J.J., Fernandez-Oto, C., Tzuk, O., Zelnik, Y.R., Grafi, G., 2019. Continuum modeling of discrete plant communities: Why does it work and why is it advantageous? Mathematics 2019, Vol. 7, Page 987 7, 987. doi:10.3390/math7100987.

[38] Monteith, J.L., 1965. Evaporation and environment. Symposia of the Society for Experimental Biology 19, 205–234.

[39] Morris, M.D., 1991. Factorial sampling plans for preliminary computational experiments. Technometrics 33, 161–174. doi:10.1080/00401706.1991.10484804.

[40] Nadwodnik, J., Lohaus, G., 2008. Subcellular concentrations of sugar alcohols and sugars in relation to phloem translocation in Plantago major, Plantago maritima, Prunus persica, and Apium graveolens. Planta 227, 1079–1089. doi:10.1007/s00425-007-0682-0.

[41] Nobel, P.S., 2009. Physicochemical and Environmental Plant Physiology. 4th ed., Academic Press, Amsterdam.

[42] Paules, J.B., Fatichi, S., Warring, B., Paschalis, A., 2025. T&c-crop: Representing mechanistic crop growth with a terrestrial biosphere model (t&c, v1.5) - model formulation and validation. Geoscientific Model Development 18, 1287–1305. doi:10.5194/gmd-18-1287-2025.

[43] Press, W.H., Teukolsky, S.A., Vetterling, W.T., Flannery, B.P., 2007. Numerical Recipes: The Art of Scientific Computing. 3rd ed., Cambridge University Press, Cambridge.

[44] Russo, S.E., Ledder, G., Muller, E.B., Nisbet, R.M., 2022. Dynamic energy budget models: fertile ground for understanding resource allocation in plants in a changing world. Conservation Physiology 10, coac061. doi:10.1093/conphys/coac061.

[45] Saltelli, A., Annoni, P., Azzini, I., Campolongo, F., Ratto, M., Tarantola, S., 2010. Variance based sensitivity analysis of model output. design and estimator for the total sensitivity index. Computer Physics Communications 181, 259–270. doi:10.1016/j.cpc.2009.09.018.

[46] Scialdone, A., Mugford, S.T., Feike, D., Skeffington, A., Borrill, P., Graf, A., Smith, A.M., Howard, M., 2013. Arabidopsis plants perform arithmetic division to prevent starvation at night. eLife , e00669doi:10.7554/elife.00669.

[47] Seki, M., Ohara, T., Hearn, T.J., Frank, A., Silva, V.C.D., Caldana, C., Webb, A.A., Satake, A., 2017. Adjustment of the Arabidopsis circadian oscillator by sugar signalling dictates the regulation of starch metabolism. Scientific Reports 7, 8305. doi:10.1038/s41598-017-08325-y.

[48] Sharkey, T.D., Bernacchi, C.J., Farquhar, G.D., Singsaas, E.L., 2007. Fitting photosynthetic carbon dioxide response curves for C3 leaves. Plant, Cell and Environment 30, 1035–1040. doi:10.1111/j.1365-3040.2007.01710.x.

[49] Shen, J., Yuan, L., Zhang, J., Li, H., Bai, Z., Chen, X., Zhang, W., Zhang, F., 2011. Phosphorus dynamics: From soil to plant. Plant Physiology 156, 997–1005. doi:10.1104/pp.111.175232.

[50] Sievänen, R., Godin, C., De Jong, T.M., Nikinmaa, E., 2014. Functional–structural plant models: a growing paradigm for plant studies. Annals of Botany 114, 599–603. doi:10.1093/aob/mcu175.

[51] Smith, A.M., Stitt, M., 2007. Coordination of carbon supply and plant growth. Plant, Cell & Environment 30, 1126–1149. doi:10.1111/j.1365-3040.2007.01708.x.

[52] Sobol', I.M., 2001. Global sensitivity indices for nonlinear mathematical models and their Monte Carlo estimates. Mathematics and Computers in Simulation 55, 271–280. doi:10.1016/S0378-4754(00)00270-6.

[53] Stitt, M., Zeeman, S.C., 2012. Starch turnover: pathways, regulation and role in growth. Current Opinion in Plant Biology 15, 282–292. doi:10.1016/j.pbi.2012.03.016.

[54] Sulpice, R., Flis, A., Ivakov, A.A., Apelt, F., Krohn, N., Encke, B., Abel, C., Feil, R., Lunn, J.E., Stitt, M., 2014. Arabidopsis coordinates the diurnal regulation of carbon allocation and growth across a wide range of photoperiods. Molecular Plant 7, 137–155. doi:10.1093/mp/sst127.

[55] Tedone, F., 2020. The mathematical modeling of plant growth and applications to robotics. Ph.D. thesis. Gran Sasso Science Institute. L'Aquila, Italy. URL: https://hdl.handle.net/20.500.12571/14997. Doctoral thesis, defended 2 February 2020. GSSI institutional repository (IRIS) record 20.500.12571/14997.

[56] Thornley, J.H.M., 1972. A balanced quantitative model for root:shoot ratios in vegetative plants. Annals of Botany 36, 431–441. doi:10.1093/oxfordjournals.aob.a084602.

[57] Thornley, J.H.M., Johnson, I.R., 1990. Plant and Crop Modelling: A Mathematical Approach to Plant and Crop Physiology. Clarendon Press, Oxford.

[58] Tian, Y., Zeng, Z., Xing, Y., 2024. A review of discrete element method applications in soil–plant interactions: Challenges and opportunities. Agriculture 2024, Vol. 14, Page 1486 14, 1486. doi:10.3390/agriculture14091486.

[59] Tinker, P.B., Nye, P.H., 2000. Solute Movement in the Rhizosphere. 2nd ed., Oxford University Press, New York.

[60] Wallach, D., Makowski, D., Jones, J.W., Brun, F., 2018. Working with Dynamic Crop Models: Methods, Tools and Examples for Agriculture and Environment. 3rd ed., Academic Press. doi:10.1016/C2016-0-01552-8.

[61] Webb, A.A., Satake, A., 2015. Understanding circadian regulation of carbohydrate metabolism in Arabidopsis using mathematical models. Plant & cell physiology 56, 586–593. doi:10.1093/pcp/pcv033.

[62] Weraduwage, S.M., Chen, J., Anozie, F.C., Morales, A., Weise, S.E., Sharkey, T.D., 2015. The relationship between leaf area growth and biomass accumulation in Arabidopsis thaliana. Frontiers in Plant Science 6, 167. doi:10.3389/fpls.2015.00167.

[63] de Wit, A., Boogaard, H., Fumagalli, D., Janssen, S., Knapen, R., van Kraalingen, D., Supit, I., van der Wijngaart, R., van Diepen, K., 2019. 25 years of the WOFOST cropping systems model. Agricultural Systems 168, 154–167. doi:10.1016/j.agsy.2018.06.018.

[64] Xu, C., Fisher, R., Wullschleger, S.D., Wilson, C.J., Cai, M., McDowell, N.G., 2012. Toward a mechanistic modeling of nitrogen limitation on vegetation dynamics. PLOS ONE 7, e37914. doi:10.1371/journal.pone.0037914.

[65] Yu, L.y., Cai, H.j., Zheng, Z., Li, Z.j., Wang, J., 2017. Towards a more flexible representation of water stress effects in the nonlinear Jarvis model. Journal of Integrative Agriculture 16, 210–220. doi:10.1016/S2095-3119(15)61307-7.

[66] Yu, S., Guo, J., Liu, Z., Wang, Y., Ma, J., Li, J., Liu, F., 2021. Assessing the impact of soil moisture on canopy transpiration using a modified jarvis-stewart model. Water (Switzerland) 13, 2720. doi:10.3390/w13192720.

[67] Zeigler, B.P., Praehofer, H., Kim, T.G., 2000. Theory of Modeling and Simulation: Integrating Discrete Event and Continuous Complex Dynamic Systems. 2nd ed., Academic Press, San Diego.

Table 3
Model parameters, generated from the provenance ledger of the released repository. Values are those loaded by the shipped configuration, or the literal in the source where a parameter is not exposed in the configuration file. Tiers: T1 measured, T2 derived, T3 inherited from the source model, T4 constrained, T5 fitted, T6 unsourced, n/l not entered in the ledger. The four parameters calibrated in this work are given separately in Table 4. [a]Source key present in the ledger but not yet in the reference list of this paper. [b]Literal in the source, not exposed as a configuration key. See the TODO items in Section 3.12.

Symbol	Meaning	Value	Unit	Tier	Source
f	Photoperiod length	8	h	T1	Sulpice et al. [54]
$\Phi_{c,\max}$	Reference assimilation rate	12.7	μmol C_6 gFW^{-1} h^{-1}	T3	Tedone [55]
$B_{l,\min}$	Leaf-biomass viability threshold	0.001	gFW	T4	–
λ_c	Residual assimilation at full starch	1	–	T5	–
C_a	Ambient CO_2 concentration	400	μmol mol^{-1}	T4	–
g_s^{fix}	Fixed conductance (M1 configuration)	0.1	mol m^{-2} s^{-1}	T5	–
SLA	Specific leaf area	0.0055	m^2 gFW^{-1}	T1	leaf atlas[a]
Γ^*	CO_2 compensation point	42	μmol mol^{-1}	T6	–
$J_{\max}$	Maximal electron transport rate	90	μmol m^{-2} s^{-1}	T5	–
$V_{c\max}$	Maximal carboxylation rate	45	μmol m^{-2} s^{-1}	T5	–
K_c	Rubisco Michaelis constant, CO_2	404.9	μmol mol^{-1}	T6	–
K_o	Rubisco Michaelis constant, O_2	278.4	mmol mol^{-1}	T6	–
O	Oxygen concentration	210	mmol mol^{-1}	T4	–
T	Leaf temperature	25	°C	T5	Sulpice et al. [54]
RH	Relative humidity	60	%	T6	–
ψ_w	Leaf water potential	-0.6	MPa	T5	–
R_d	Dark respiration rate	1.5	μmol m^{-2} s^{-1}	T2	–
ϕ	Apparent quantum yield	0.385	mol e^- mol^{-1}	T6	–
θ	Light-response curvature	0.7	–	T6	–
b_1	Jarvis light-response scale	0.6	–	T6	–
b_2	Jarvis light-response slope	0.002	–	T6	–
$D_{a,\max}$	Maximal starch degradation rate	6	μmol C_6 gFW^{-1} h^{-1}	T3	Tedone [55]
$a_{\min}$	Non-mobilised starch residue	0.15	μmol C_6 gFW^{-1}	T3	Tedone [55]
$s_{\max}$	Sucrose half-saturation, excess	4.712	μmol C_6 gFW^{-1}	T3	Tedone [55]
$s_{\min}$	Sucrose half-saturation, deficit	1.3	μmol C_6 gFW^{-1}	T3	Tedone [55]
λ_{sni}	Night partition increase rate	0.12	h^{-1}	T5	–
λ_g	Loading efficiency floor	0.65	–	T5	–
r_m	Maintenance respiration rate	0.79	h^{-1}	T3	Tedone [55]
η	Phloem loading frequency	1.98	h^{-1}	T3	Tedone [55]
τ_c	Transport cost fraction	0.035	–	T4	Tedone [55]
k_{in}	Vacuolar influx rate	0.1	h^{-1}	T5	measured ratio[a]
k_{out}	Vacuolar efflux rate	0.041959	h^{-1}	T2	measured ratio[a]
n_H	Sink-activation Hill exponent	3	–	n/l	hard-coded[b]
Y	Construction cost of biomass	0	μmol C_6 gFW^{-1}	n/l	hard-coded[b]
μ_l	Leaf mortality rate	5×10^{-6}	h^{-1}	T3	Tedone [55]
μ_r	Root mortality rate	5×10^{-6}	h^{-1}	T3	Tedone [55]
γ_l	Leaf self-competition coefficient	5×10^{-6}	gFW^{-1} h^{-1}	T3	Tedone [55]
γ_r	Root self-competition coefficient	5×10^{-6}	gFW^{-1} h^{-1}	T3	Tedone [55]
ω	Optimal N:P ratio	10	–	T3	Tedone [55]
$r_{A,N}$	N cost of carbon turnover	0.0460308	μmol N (μmol C_6)$^{-1}$	T2	Tedone [55]
$r_{A,P}$	P cost of carbon turnover	0.00460308	μmol P (μmol C_6)$^{-1}$	T2	Tedone [55]
N_{ph}	N threshold for assimilation	5.75	μmol N gFW^{-1}	T3	Tedone [55]
P_{ph}	P threshold for assimilation	0.575	μmol P gFW^{-1}	T2	Tedone [55]
ν	Nutrient conversion parameter	0.066	–	T5	–
N_s	Soil nitrogen concentration	15	μmol cm^{-3}	T4	Tedone [55]
P_s	Soil phosphorus concentration	0.15	μmol cm^{-3}	T3	Tedone [55]
$I_{N,\max}$	Maximal N uptake rate	6.44	μmol N gFW^{-1} h^{-1}	T3	Tedone [55]
$I_{P,\max}$	Maximal P uptake rate	0.4	μmol P gFW^{-1} h^{-1}	T3	Tedone [55]
K_N	Soil N half-saturation	0.125	μmol cm^{-3}	T3	Tedone [55]
K_P	Soil P half-saturation	0.006736	μmol cm^{-3}	T3	Tedone [55]
λ_k	Stoichiometric imbalance gain	50	–	T5	–
c_N	Sucrose cost of N uptake	0.68	μmol C_6 (μmol N)$^{-1}$	T3	Tedone [55]
c_P	Sucrose cost of P uptake	0.308	μmol C_6 (μmol P)$^{-1}$	T3	Tedone [55]
N_s^{ref}	Soil N reference (hard-coded)	12	μmol cm^{-3}	T6	–
P_s^{ref}	Soil P reference (hard-coded)	0.15	μmol cm^{-3}	T6	–
$g_{s,\max}$	Maximal stomatal conductance	$1/\sqrt{2}$	mol m^{-2} s^{-1}	n/l	hard-coded[b]
b_{10}	Jarvis residual light conductance	0.05	–	n/l	hard-coded[b]
b_5	Jarvis VPD slope	0.2	kPa^{-1}	n/l	hard-coded[b]
b_6	Jarvis water-potential slope	3.0	MPa^{-1}	n/l	hard-coded[b]
b_8	Jarvis CO_2 response floor	0.2	–	n/l	hard-coded[b]
T_0	Jarvis temperature optimum	25	°C	n/l	hard-coded[b]
T_1, T_h	Jarvis temperature zeros	5, 45	°C	n/l	hard-coded[b]
$\psi_{\min}$	Jarvis stomatal closure potential	−1.5	MPa	n/l	hard-coded[b]

Table 4
The four parameters calibrated in this work. All four are shared: one value of each is used at every photoperiod in every reported run. All are tier T5 by the classification of Section 3.12. The published baseline calibrated eight parameters per photoperiod against the same eight observations.

Symbol	Meaning	Value	Unit	Calibrated against
λ_{sb}	Biomass conversion coefficient	0.00295	gFW (μmol C_6)$^{-1}$ gFW^{-1}	Relative growth rate as a function of photoperiod, over the four measured photoperiods jointly (Fig. 2)
λ_{sdr}	Day partition reduction rate	0.50	h^{-1}	Shape and timing of the diel starch trace, jointly with λ_{sdi} (Fig. 4)
λ_{sdi}	Day partition increase rate	0.12	h^{-1}	Shape and timing of the diel starch trace, jointly with λ_{sdr} (Fig. 4)
K_s	Sink-activation half-saturation	0.90	μmol C_6 gFW^{-1}	Invariance of the sucrose level across the photoperiod series (Fig. 4)

Table 5
Parameterisation regimes. All three use the same chamber forcing (PAR 160 μmol m^{-2} s^{-1}, square wave, 21/19 °C), the same initial state and the same sourced parameters. P denotes photoperiod in hours. The observation count is two per photoperiod, the published RGR and the published end-of-day starch. Regime R1 is reported as a calibration diagnostic; only R2 and R3 enter the cross-validation of Section 5.4.

Regime	**Fitted coefficients**	**Fitted** values	**Obs.** used	**Role**
R1 saturated	λ_{sb}, λ_{sdi}, per photoperiod	8	8	calibration diagnostic
R2 shared	λ_{sb}, λ_{sdr}, λ_{sdi}, sink constant	4	8	held-out prediction
R3 shared, linear	as R2 with $\lambda_{sb} = c_0 + c_1 P$	5	8	held-out prediction

Table 6
Relative growth rate under the shared parameterisation, chamber forcing, 30 days. Targets are the *Arabidopsis thaliana* rosette values of Sulpice et al. [54]. The last column gives the per-photoperiod residual of the saturated per-photoperiod calibration on the grid described in Section 6.1; those residuals are set by the grid resolution and are reported as calibration diagnostics, not as errors of prediction.

Photoperiod (h)	Target RGR (d^{-1})	Simulated RGR (d^{-1})	Error (%)	Accepted	Saturated fit residual (%)
4	0.0680	0.0782	15.04	no	3.42
6	0.1135	0.1189	4.77	yes	1.28
8	0.1708	0.1588	7.00	yes	2.45
12	0.2600	0.2374	8.71	yes	3.03
Mean absolute percentage error			8.88	3/4	–

Table 7
Leave-one-photoperiod-out prediction of relative growth rate. In regime A the growth coefficient is a single shared constant; in regime B it is linear in photoperiod with two coefficients (Eq. 56). Train error is the mean over the three fitted photoperiods; test error is the held-out photoperiod. The λ_{sb} column gives the value applied at the held-out photoperiod.

Regime	Held out (h)	λ_{sb}	Train MAPE (%)	Test MAPE (%)
A: shared constant	4	0.00320	5.57	20.93
A: shared constant	6	0.00320	9.24	9.94
A: shared constant	8	0.00295	8.08	10.10
A: shared constant	12	0.00295	7.53	11.75
Regime A, mean held-out error				13.18
B: linear in P	4	0.00275	2.25	3.42
B: linear in P	6	0.00295	2.97	1.28
B: linear in P	8	0.00320	2.57	2.45
B: linear in P	12	0.00370	2.38	10.30
Regime B, mean held-out error				4.36

Table 8
Ablation of the carbon-input pathway. M0 prescribes the carbon input, M1 computes it from Farquhar photosynthesis at fixed conductance, and M2 adds Jarvis stomatal conductance. Shared parameters are frozen across the three variants. Errors are relative errors in RGR; the last column is the peak assimilation rate.

	RGR error (%) by photoperiod				MAPE	Peak Φ_c
Variant	4 h	6 h	8 h	12 h	(%)	(μmol gFW^{-1}h^{-1})
M0 prescribed	41.59	40.19	43.08	39.20	41.01	10.50–11.14
M1 Farquhar	18.82	7.77	4.55	6.60	9.43	17.62
M2 Farquhar–Jarvis	15.04	4.77	7.00	8.71	8.88	17.01

Table 9
Global sensitivity of relative growth rate, averaged over the four photoperiods. S_1 is the first-order and S_T the total-order Sobol' index; the interaction remainder is $S_T - S_1$. μ^* and σ are the Morris measures. Indices are reported unclipped, so a remainder at noise level can be slightly negative. Ranges and sampling scales are in Appendix B; the indices are conditional on those ranges.

Parameter	S_1	S_T	$S_T - S_1$	μ^*	σ/μ^*
max_phosphorus_uptake	0.000	0.658	0.658	0.635	1.73
max_nitrogen_uptake	0.000	0.577	0.577	0.785	1.48
respiration_frequency	0.031	0.466	0.435	0.466	1.94
ambient_CO2_concentration	0.000	0.422	0.422	0.370	2.13
leaf_water_potential	0.063	0.412	0.349	0.597	1.57
electron_transport_rate	0.020	0.352	0.332	0.247	2.57
max_rubisco_carboxylation_rate	0.013	0.334	0.321	0.393	2.16
max_PAR	0.000	0.333	0.333	0.463	1.96
leaf_temperature	0.008	0.219	0.211	0.198	3.05
relative_humidity	0.000	0.199	0.199	0.222	2.84
phosphorus_soil_content	0.017	0.164	0.148	0.202	2.62
sucrose (initial)	0.007	0.144	0.137	0.164	2.70
nitrogen_soil_content	0.007	0.141	0.134	0.100	3.72
k_vacuole_in	0.009	0.135	0.126	0.101	3.96
lambda_sb	0.023	0.117	0.094	0.073	3.32
total_biomass (initial)	0.003	0.001	−0.002	0.066	0.37
Sum	0.201	4.675			

Table 10
Complete parameter provenance ledger, generated from `config/parameter_provenance.csv`. Locator gives the equation, section or source line from which the value comes; a dash means no locator is recorded.

Parameter	Value	Unit	Tier	Source	Locator
gas_exchange.specific_leaf_area	0.025	m2 gDW-1	T1	ArabidopsisLeafAtlas2024	Table 1, 'Leaf area per unit leaf fresh weight' = 5500 mm2/g LFW; Col-0 leaf 6 at 21 DAI
photo.photoperiod	8	h	T1	Sulpice2014	Methods / Supplemental Table 1 (4, 6, 8, 12 h treatments)
carbohydrates.k_vacuole_out	0.041959	h-1	T2	Seydel2025	Plant Physiol 198:kiaf117, Suppl. Table S5 sheet ST3, control rows, n=5
carbohydrates.night_efficiency_starch	0.5	-	T2	–	input.c:31; default.config:30
carbohydrates.starch	0.225	umolC6/gFW	T2	–	input.c:42; default.config:41
carbohydrates.starch_degradation_rate	0		T2	–	–
carbohydrates.starch_night_start	2	umolC6/gFW	T2	–	–
carbohydrates.starch_partition_coeff	0.6	-	T2	–	input.c:25; default.config:24
carbohydrates.sucrose	1.95	umolC6/gFW	T2	–	input.c:41; default.config:40
carbohydrates.transport_cost	0		T2	–	–
carbohydrates.uptake_cost	0	umol C6 gFW-1 h-1	T2	Tedone2020	§2.2.8 (n_c*U_N + p_c*U_P); config entry is a slot
gas_exchange.convert_to_photosynthesis_per_gram_per_hour	24000	dimensionless	T2	–	–
gas_exchange.light_limited_photosynthesis	10		T2	–	–
gas_exchange.light_limited_rate	10		T2	–	–
gas_exchange.net_photosynthesis	10		T2	–	–
gas_exchange.net_photosynthesis_rate	10		T2	–	–
gas_exchange.respiration_rate	1.5		T2	–	–
gas_exchange.rubisco_limited_photosynthesis	12		T2	–	–
gas_exchange.rubisco_limited_rate	12		T2	–	–
growth.leaf_biomass	0.02	gFW	T2	–	input.c:14; default.config:15
growth.root_biomass	0.02	gFW	T2	–	input.c:15; default.config:16
growth.sucrose_root_allocation	0.5	-	T2	–	input.c:86; default.config:74
growth.total_biomass	0.04	gFW	T2	–	input.c:16; default.config:17
hardcoded.photosynthesis.ref_specific_leaf_area	0.025	m2 gFW-1	T2	–	–
nutrients.assimilation_cost_nitrogen	0.0460308	umolN/umolC6	T2	Tedone2020	Eq 2.2.16, Appendix A
nutrients.assimilation_cost_phosphorus	0.00460308	umolP/umolC6	T2	Tedone2020	c_sp = c_sn/O, Appendix A (O = 10 umolN/umolP)
nutrients.max_nitrogen	23	umolN/gFW	T2	Tedone2020	Eq 2.2.25 (n_max = D*n_min, D = 4)
nutrients.max_phosphorus	2.3	umolP/gFW	T2	Tedone2020	after Eq 2.2.25 ("p_max = n_max / O") with O = 10 from Eq 2.2.7
nutrients.min_nitrogen	5.75	umolN/gFW	T2	Tedone2020	Eq 2.2.25
nutrients.min_phosphorus	0.575	umolP/gFW	T2	Tedone2020	after Eq 2.2.25 ("p_max = n_max / O") with O = 10 from Eq 2.2.7
nutrients.min_phosphorus_photosynthesis	0.575	umolP/gFW	T2	Tedone2020	after Eq 2.2.25 ("p_max = n_max / O") with O = 10 from Eq 2.2.7
nutrients.nitrogen	8.625	umolN/gFW	T2	–	input.c:43; default.config:42
nutrients.nitrogen_affinity	0.5		T2	–	–
nutrients.nitrogen_cost	0		T2	–	–
nutrients.nitrogen_nutrient_uptake	0		T2	–	–
nutrients.nitrogen_uptake	0		T2	–	–
nutrients.phosphorus	0.8625	umolP/gFW	T2	–	input.c:44; default.config:43
nutrients.phosphorus_affinity	0.5		T2	–	–
nutrients.phosphorus_cost	0		T2	–	–
nutrients.phosphorus_nutrient_uptake	0		T2	–	–
nutrients.phosphorus_uptake	0		T2	–	–
nutrients.pot_nitrogen_uptake	0		T2	–	–
nutrients.pot_phosphorus_uptake	0		T2	–	–
nutrients.stoichiometric_signal	0.5	-	T2	–	input.c:87; default.config:75
photo.light	0		T2	–	–
photo.limitation_of_photosynthetic_rate	0.5		T2	–	–
photo.nitrogen_saturation	0.45		T2	–	–
photo.phosphorus_saturation	0.45		T2	–	–
simulation.days	35		T2	–	–
simulation.dt_hours	0.1		T2	–	–
simulation.gui_enabled	true		T2	–	–
simulation.solver_tolerance	1e-08		T2	–	–
simulation.solver_type	ODE45		T2	–	–
simulation.write_csv	true		T2	–	–
carbohydrates.max_starch_degradation_rate	6	umolC6/(gFW h)	T3	Tedone2020	§2.2.4
carbohydrates.max_sucrose	2	umolC6/gFW	T3	Tedone2020	§2.2.4 (Eq 2.2.28)
carbohydrates.min_starch	0.15	umolC6/gFW	T3	Tedone2020	§2.2.4
carbohydrates.min_sucrose	1.3	umolC6/gFW	T3	Tedone2020	§2.2.4
carbohydrates.respiration_frequency	0.79	1/h	T3	Tedone2020	Section 2.2.8 (Costs), "r_m_bar = 0.79 1/h"
carbohydrates.sucrose_loading_frequency	1.98	1/h	T3	Tedone2020	§2.2.8 (Eq 2.2.28)
growth.leaf_competitive_rate	5e-06	1/(h gFW)	T3	Tedone2020	§2.2.10 (paragraph after Eq 2.2.33): "fixed to the small value 5e-6"
growth.leaf_deathrate	5e-06	1/h	T3	Tedone2020	§2.2.10 (paragraph after Eq 2.2.33): "fixed to the small value 5e-6"
growth.root_competitive_rate	5e-06	1/(h gFW)	T3	Tedone2020	§2.2.10 (paragraph after Eq 2.2.33): "fixed to the small value 5e-6"
growth.root_deathrate	5e-06	1/h	T3	Tedone2020	§2.2.10 (paragraph after Eq 2.2.33): "fixed to the small value 5e-6"
nutrients.Michaelis_Menten_constant_nitrogen	0.125	umolN cm-3	T3	Tedone2020	§2.2.7 (Eq 2.2.22), "k_n = 0.125 umolN/cm3"
nutrients.Michaelis_Menten_constant_phosphorus	0.006736	umolP cm-3	T3	Tedone2020	§2.2.7 (Eq 2.2.22), "k_p = 0.006736 umolP/cm3"
nutrients.max_nitrogen_uptake	6.44	umolN/(gFW h)	T3	Tedone2020	§2.2.7
nutrients.max_phosphorus_uptake	0.4	umolP/(gFW h)	T3	Tedone2020	§2.2.7
nutrients.min_nitrogen_photosynthesis	5.75	umolN/gFW	T3	Tedone2020	Eq 2.2.5
nutrients.nitrogen_uptake_sucrose_consumption	0.68	umolC6/umolN	T3	Tedone2020	§2.2.8
nutrients.optimal_stoichiometric_ratio	10	umolN/umolP	T3	Tedone2020	Eq 2.2.7, Eq 2.2.31
nutrients.phosphorus_soil_content	0.15	umolP cm-3	T3	Tedone2020	Appendix A [159]
nutrients.phosphorus_uptake_sucrose_consumption	0.308	umolC6/umolP	T3	Tedone2020	§2.2.8
photo.max_photosynthetic_rate	12.7	umolC6/(gFW h)	T3	Tedone2020	Eq 2.2.1, §2.2.2
DEFECT.algebraic.uptake_cost_evaluation_order	nan		T4	conservation	src/model/algebraic.c:86 (call) vs :127/:131 (assignments)
DEFECT.growth.no_carbon_price_on_biomass	nan		T4	conservation	src/model/carbohydrates.c:106 (sucrose_production) vs src/model/growth.c:12 (leaf_growth)
carbohydrates.sucrose_consumption_transport	0.035	-	T4	Tedone2020	Appendix A (0.0035) vs body text 2.2.8 (0.035)
gas_exchange.ambient_CO2_concentration	400	umol mol-1	T4	–	–
gas_exchange.oxygen_concentration	210		T4	–	–
hardcoded.photosynthesis.base_scale	1.1	dimensionless (umol hexose gFW-1 h-1 per umol CO2 m-2 s-1)	T4	Sulpice2014; AtkinsonAtkin2024; C3_quantum_yield_ceiling	mmc3 sheet "Growth components" rows 2-3 (headers), row 5 "Assimilation rate (umol C / g FW h)", row 10 "Assimilation (umol glc equiv / g FW h)", row 24 "% assimilation rate to starch synth"; mmc3 sheet "Growth in day and night" row 2 headers + row 10 "Nitrogen = photosynthesis (measured at end of day)" and row 14 "Net C assimilation per 24 h cycle = total A over the light period - total R over the night"; mmc2 sheet "clean dataset-all data" column "average DW content"; N_data.csv column "LMA (mean

Table 11
Sensitivity analysis design, generated from `config/sensitivity_params.csv`. Role distinguishes a model parameter from an environmental driver, an initial condition and a physical constant. The basis column records why the range was chosen.

Factor	Lower	Upper	Scale	Role	Basis
Active in the reported design					
`growth.lambda_sb`	0.002	0.01	linear	model parameter	Calibrated; ±70% around default 0.00587 (Schulz et al. 2019)
`carbohydrates.respiration_frequency`	0.4	1.2	linear	model parameter	Calibrated; ±50% around default 0.79
`nutrients.max_nitrogen_uptake`	1	15	log	model parameter	Literature range for herbaceous crops (Barber 1995; Lambers et al. 2008)
`nutrients.max_phosphorus_uptake`	0.05	1.5	log	model parameter	Literature range for herbaceous crops (Barber 1995)
`nutrients.nitrogen_soil_content`	3	30	linear	environment	Low-nutrient to high-nutrient agricultural soil (Brady & Weil 2010)
`nutrients.phosphorus_soil_content`	0.03	0.5	log	environment	Low to high-P agricultural soil (Brady & Weil 2010)
`gas_exchange.leaf_temperature`	10	35	linear	environment	Controlled environment range; cold stress to heat stress
`gas_exchange.relative_humidity`	30	90	linear	environment	Arid to humid growing conditions
`gas_exchange.ambient_CO2_concentration`	280	800	linear	environment	Pre-industrial to RCP 8.5 projected 2100 (Meinshausen et al. 2011)
`gas_exchange.leaf_water_potential`	-1.5	-0.1	linear	environment	Severe water stress to turgid (Hsiao 1973; Jones 2014)
`gas_exchange.max_rubisco_carboxylation_rate`	20	120	log	model parameter	C3 herbaceous range (Wullschleger 1993; Medlyn et al. 2002)
`gas_exchange.electron_transport_rate`	50	200	log	model parameter	C3 herbaceous range (Wullschleger 1993)
`photo.max_PAR`	300	2000	linear	environment	Shade canopy (300) to full sun (2000 μmol mš sź) (McCree 1972)
`carbohydrates.sucrose`	0.8	3	linear	initial condition	Observed range in spinach seedlings (% DW; Geiger & Servaites 1994)
`growth.total_biomass`	0.01	0.12	linear	initial condition	Seedling (0.01 g FW) to established juvenile plant (0.12 g FW)
`carbohydrates.k_vacuole_in`	0.005	0.4	linear	model parameter	Cytosolvacuole exchange rate; covers calibrated per-PP range 0.011-0.230 h-1 (Lohaus 2007)
Disabled					
`carbohydrates.lambda_sni`	0.04	0.18	linear	model parameter	Disabled: mu*≈0 everywhere; excluded to keep d≤16 (solver limit)
`photo.feedback_on_photosynthesis`	0.5	0.9	linear	model parameter	Disabled: inactive in Farquhar path (light_PAR>20 simple model never called)
`gas_exchange.oxygen_concentration`	210	210	fixed	physical constant	Fixed atmospheric O2; excluded as non-varying
`carbohydrates.starch`	0.05	1	linear	initial condition	Observed range in spinach seedlings (% DW; Geiger & Servaites 1994)