

QuTiP lecture: Correlation functions

Author: J. R. Johansson, robert@riken.jp

<http://dml.riken.jp/~rob/>

Latest version of this ipython notebook lecture is available at: <http://github.com/jrjohansson/qutip-lectures>

In [47]: `%pylab inline`

```
Welcome to pylab, a matplotlib-based Python environment [backend:
module://IPython.zmq.pylab.backend_inline].
For more information, type 'help(pylab)'.
```

In [48]: `from qutip import *`

Introduction

First-order coherence function

This example demonstrates how to calculate a correlation function on the form $\langle A(\tau)B(0) \rangle$ for a non-steady initial state. Consider an oscillator that is interacting with a thermal environment. If the oscillator initially is in a coherent state, it will gradually decay to a thermal (incoherent) state. The amount of coherence can be quantified using the first-order optical coherence function

$$g^{(1)}(\tau) = \frac{\langle a^\dagger(\tau)a(0) \rangle}{\sqrt{\langle a^\dagger(\tau)a(\tau) \rangle \langle a^\dagger(0)a(0) \rangle}}$$

For a coherent state $|g^{(1)}(\tau)| = 1$, and for a completely incoherent (thermal) state $g^{(1)}(\tau) = 0$. The following code calculates and plots $g^{(1)}(\tau)$ as a function of τ .

Example: Decay of a coherent state to an incoherent (thermal) state

```
In [49]: N = 15
taulist = linspace(0,10.0,200)
a = destroy(N)
H = 2*pi*a.dag()*a

# collapse operator
G1 = 0.75
n_th = 2.00 # bath temperature in terms of excitation number
c_ops = [sqrt(G1*(1+n_th)) * a, sqrt(G1*n_th) * a.dag()]

# start with a coherent state
rho0 = coherent_dm(N, 2.0)

# first calculate the occupation number as a function of time
n = mesolve(H, rho0, taulist, c_ops, [a.dag() * a]).expect[0]


# calculate the correlation function G1 and normalize with n to obtain g1
G1 = correlation(H, rho0, None, taulist, c_ops, a.dag(), a)
g1 = G1 / sqrt(n[0] * n)
```

```
In [50]: fig, axes = subplots(2, 1, sharex=True, figsize=(12,6))

axes[0].plot(taulist, real(g1), 'b', label=r'First-order coherence function $g^{\{1\}}(\tau)$')
axes[1].plot(taulist, real(n), 'r', label=r'occupation number $n(\tau)$')
axes[0].legend()
```

```
axes[1].legend()
axes[1].set_xlabel(r'$\tau$')
```

Out [50]: <matplotlib.text.Text at 0x3aec6c10>



Leggett-Garg inequality

Definition: Given an observable $Q(t)$ that is bound below and above by $|Q(t)| \leq 1$, the assumptions of

- macroscopic realism
- noninvasive measurements

implies that

$$L(t_1, t_2) = \langle Q(t_1)Q(0) \rangle + \langle Q(t_1 + t_2)Q(t_1) \rangle - \langle Q(t_1 + t_2)Q(0) \rangle \leq 1$$

If Q is at a steady state at the initial time of measurement, we can set $\tau = t_1 = t_2$ and the Leggett-Garg inequality then reads

$$L(\tau) = 2\langle Q(\tau)Q(0) \rangle - \langle Q(2\tau)Q(0) \rangle \leq 1$$

References

- [A. J. Leggett and A. Garg, Phys. Rev. Lett. 54, 857 \(1985\)](#)
- [A. J. Leggett, J. Phys. Condens. Matter 14, R415 \(2002\)](#)

```
In [51]: def leggett_garg(c_mat):
    """
    For a given correlation matrix c_mat = <Q(t1+t2)Q(t1)>, calculate the Leggett-Garg
    inequality.
    """

    N, M = shape(c_mat)

    lg_mat = zeros([N/2, M/2], dtype=complex)
    lg_vec = zeros(N/2, dtype=complex)

    for i in range(N/2):
        # i == j
        lg_vec[i] = 2*c_mat[0, i] - c_mat[0, 2*i];
```

```

    for j in range(M/2):

        # c_mat(i, j) = <Q(dt i+dt j)Q(dt i)>
        #
        # LG = <Q(t_1)Q(0)> + <Q(t_1+t_2)Q(t_1)> - <Q(t_1+t_2)Q(0)>

        lg_mat[i, j] = c_mat[0, i] + c_mat[i, j] - c_mat[0, i+j];

    return lg_mat, lg_vec

```

Example: Leggett-Garg inequality for two coupled resonators (optomechanical system)

References:

- [N. Lambert, J.R. Johansson, F. Nori, Phys. Rev. B 82, 245421 \(2011\).](#)

```

In [97]: wc = 1.0 * 2 * pi # cavity frequency
        wa = 1.0 * 2 * pi # resonator frequency
        g = 0.3 * 2 * pi # coupling strength
        kappa = 0.075 # cavity dissipation rate
        gamma = 0.005 # resonator dissipation rate
        Na = Nc = 3 # number of cavity fock states
        n_th = 0.0 # avg number of thermal bath excitation

        tlist = linspace(0, 7.5, 251)
        tlist_sub = tlist[0:(len(tlist)/2)]

```

```

In [98]: # start with an excited resonator
        rho0 = tensor(fock_dm(Na,0), fock_dm(Nc,1))

        a = tensor(qeye(Nc), destroy(Na))
        c = tensor(destroy(Nc), qeye(Na))

        na = a.dag() * a
        nc = c.dag() * c

        H = wa * na + wc * nc - g * (a + a.dag()) * (c + c.dag())

```

```

In [102]: # measurement operator on resonator
        Q = na # photon number resolving detector
        #Q = tensor(qeye(Nc), 2 * fock_dm(Na, 1) - qeye(Na)) # fock-state |1> detector
        #Q = tensor(qeye(Nc), qeye(Na) - 2 * fock_dm(Na, 0)) # click or no-click detector

```

```

In [103]: c_op_list = []

        rate = kappa * (1 + n_th)
        if rate > 0.0:
            c_op_list.append(sqrt(rate) * c)

        rate = kappa * n_th
        if rate > 0.0:
            c_op_list.append(sqrt(rate) * c.dag())

        rate = gamma * (1 + n_th)
        if rate > 0.0:
            c_op_list.append(sqrt(rate) * a)

        rate = gamma * n_th
        if rate > 0.0:
            c_op_list.append(sqrt(rate) * a.dag())

```

```

In [104]: c_mat = correlation(tlist, rho0, tlist, tlist, c_op_list, 0, 0)

```

```
In [104]: corr_mat = correlation(H, rnow, tlist, tlist, c_op_list, Q, Q)
```

```
In [105]: LG_tt, LG_t = leggett_garg(corr_mat)
```

Plot results

```
In [106]: fig, axes = subplots(1, 2, figsize=(12,4))

axes[0].pcolor(tlist,tlist,abs(corr_mat),edgecolors='none')
axes[0].set_xlabel(r'$t_1 + t_2$')
axes[0].set_ylabel(r'$t_1$')
axes[0].autoscale(tight=True)

axes[1].pcolor(tlist_sub,tlist_sub,abs(LG_tt),edgecolors='none')
axes[1].set_xlabel(r'$t_1$')
axes[1].set_ylabel(r'$t_2$')
axes[1].autoscale(tight=True)

fig, axes = subplots(1, 1, figsize=(12,4))
axes.plot(tlist_sub, diag(LG_tt), label=r'$\tau = t_1 = t_2$')
axes.plot(tlist_sub, ones(shape(tlist_sub)), 'k', label=r'quantum boundary')
axes.fill_between(tlist_sub, diag(LG_tt), 1, where=(diag(LG_tt)>1), color="blue",
alpha=0.5)
axes.set_xlim([0, max(tlist_sub)])
axes.legend(loc=0)
axes.set_xlabel(r'$\tau$')
axes.set_ylabel(r'$LG(\tau)$')
```

Out [106]: <matplotlib.text.Text at 0x48571190>

