Mathematics

Oliver Brady

April 2, 2023

Contents

1	Cal 1.1	culus Differe	entiation	3											
2	Seri	ies													
3	Mu	ltivaria	able Calculus	4											
4	Vector Calculus 5														
	4.1	Operat	tors	5											
		4.1.1	Grad	5											
		4.1.2	Div	5											
		4.1.3	Curl	5											
		4.1.4	Laplacian	5											
	4.2	Integra	al Theorems	5											
		4.2.1	Divergence Theorem	5											
		4.2.2	Stokes's Theorem	5											
5	Fluid Mechanics 6														
	5.1	Kinem	atics	6											
		5.1.1	Coordinates	6											
		5.1.2	Velocity	6											
		5.1.3	Stagnation Points	6											
		5.1.4	Streamlines	6											
		5.1.5	Particle Paths	7											
		5.1.6	Steady Flow	7											
		5.1.7	Convective Derivative	7											
		5.1.8	Vorticity	8											
		5.1.9	Velocity Potential	8											
		5.1.10	Incompressible Flow	8											
	5.2	Pressu	re in a Fluid	8											
	5.3		Oynamics	8											
	5.4		imensional Flow	8											
	5.5		ity Dynamics	8											

5.6	Free Surface	Waves														8
IJ.U	Tice pullace	vvaves														$^{\circ}$

- 1 Calculus
- 1.1 Differentiation

- 2 Series
- 3 Multivariable Calculus

4 Vector Calculus

- 4.1 Operators
- 4.1.1 Grad
- 4.1.2 Div
- 4.1.3 Curl
- 4.1.4 Laplacian
- 4.2 Integral Theorems
- 4.2.1 Divergence Theorem
- 4.2.2 Stokes's Theorem

5 Fluid Mechanics

5.1 Kinematics

5.1.1 Coordinates

Lagrangian $\underline{x}(\underline{a}, t)$: The motion of individual particles is studied; the position \underline{x} of a particle at time t is related to its position at a reference point in time \underline{a} (typically at t = 0).

Eulerian (\underline{x}, t) : The 'flow field' is considered as a whole and the state of a fluid is described in terms of the values at a fixed location \underline{x} and at a fixed time

5.1.2 Velocity

In Cartesian coordinates the velocity of a fluid particle at position $\underline{x}(x,y,z)$ is given by:

$$\underline{u}(x,y,z) = u(x,y,z)\underline{\hat{i}} + v(x,y,z)\underline{\hat{j}} + w(x,y,z)\underline{\hat{k}}$$

5.1.3 Stagnation Points

Stagnation points occur when the velocity vector \underline{u} is equal to $\underline{0}$


$$u = 0$$

$$v = 0$$

$$w = 0$$

5.1.4 Streamlines

A streamline is a curve C drawn at one point in time such that the fluid velocity vector \underline{u} is tangent to C at every point along C.



$$\frac{d\underline{x}}{ds} = \underline{u}$$

$$\frac{dx}{ds} = u, \frac{dy}{ds} = v, \frac{dz}{ds} = w$$

$$\frac{dx}{u} = \frac{dy}{v} = \frac{dz}{w} (= ds)$$

5.1.5 Particle Paths

Particle path is obtained by solving the initial value problem:

$$\frac{d\underline{x}}{dt} = \underline{u}(\underline{x}, t) , \underline{x} = x_0 \text{ at } t = 0$$

$$\frac{dx}{dt} = u , x(0) = x_0$$

$$\frac{dy}{dt} = v, y(0) = y_0$$

$$\frac{dz}{dt} = w, z(0) = z_0$$

5.1.6 Steady Flow

Steady Flow: The flow velocity vector \underline{u} is independent of time t **Unsteady Flow**: \underline{u} depends on t; the pattern of streamlines changes with t

5.1.7 Convective Derivative

The convective derivative tells us how a property changes as it moves with a flow.

General

$$\boxed{\frac{D*}{Dt} = \frac{\partial *}{\partial t} + (\underline{u} \cdot \nabla)* = \frac{\partial *}{\partial t} + u\frac{\partial *}{\partial x} + v\frac{\partial *}{\partial y} + w\frac{\partial *}{\partial z}}$$

Scalar

$$\frac{D\rho}{Dt} = \frac{\partial\rho}{\partial t} + (\underline{u}\cdot\nabla)\rho = \frac{\partial\rho}{\partial t} + u\frac{\partial\rho}{\partial x} + v\frac{\partial\rho}{\partial y} + w\frac{\partial\rho}{\partial z}$$

Vector

$$\frac{D\underline{\mathbf{u}}}{Dt} = \frac{\partial\underline{\mathbf{u}}}{\partial t} + (\underline{u} \cdot \nabla)\underline{\mathbf{u}} = \frac{\partial\underline{\mathbf{u}}}{\partial t} + u\frac{\partial\underline{\mathbf{u}}}{\partial x} + v\frac{\partial\underline{\mathbf{u}}}{\partial y} + w\frac{\partial\underline{\mathbf{u}}}{\partial z}$$

5.1.8 Vorticity

Vorticity $\underline{\omega}$ is a measure of the local rotation of fluid particles in flow.

$$\boxed{\underline{\omega} = \nabla \times \underline{u}}$$

$$\nabla \times \underline{u} = \begin{vmatrix} \underline{i} & \underline{j} & \underline{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ u & v & w \end{vmatrix}$$

Irrotational Flow:

$$\underline{\omega} = \underline{0}$$

5.1.9 Velocity Potential

For an irrotational flow the velocity can be described as the gradient of a scalar field known as the *Velocity Potential*.

$$\nabla \times \underline{u} = 0$$

The curl of the gradient of a scalar field is zero:

$$\nabla \times (\nabla \phi) = 0$$
$$\underline{u} = \nabla \phi$$

5.1.10 Incompressible Flow

$$\nabla \cdot \underline{u} = 0$$

When this is true the convective derivative of the fluid density is zero:

$$\frac{D\rho}{Dt} = 0$$

- 5.2 Pressure in a Fluid
- 5.3 Flow Dynamics
- 5.4 Tow-dimensional Flow
- 5.5 Vorticity Dynamics
- 5.6 Free Surface Waves