

Part III Project Notebook

Thomas Holland (th675)

October 2025 - 2026

Contents

1	Introduction	1
2	Plan for the project in outline	1
3	Weekly progress/goals	2
3.1	Meeting 1: 13/Oct	2
3.2	Meeting 2: 17/Oct	2
3.3	Meeting 3: 21/Oct	2
3.4	Meeting 4: 24/ Oct	2
3.5	Meeting 5: 31/Oct	2
3.6	Meeting 6: 3/Nov	3
4	Literature log	3
5	Thoughts log	3
5.1	2025-10-10 22:01	3
5.2	2025-10-18	3
6	Code notes	3
6.1	2025-10-17	3
6.2	2025-10-26	3
6.3	2025-11-01 14:48	3
7	Maths	3

1 Introduction

This is my project log, which is updated regularly to document my progress, thoughts, and reflections throughout the duration of my project. It serves as a personal journal where I can track my journey, challenges faced, solutions implemented, and insights gained.

It also pulls in my progress and goals from the shared Google Doc that I have with my supervisor, acting as a bridge between my personal reflections and the collaborative aspects of my project.

2 Plan for the project in outline

- Implementing the traditional method(s) for estimating sea level change from satellite altimetry observations, and investigating its accuracy. For global mean sea level, this method is just to spatially average sea surface height changes over the oceans.
 - Roughly first 2-4 weeks
- Application of Bayesian inversion methods that incorporate sea level physics. First done for estimates at a single time. Comparing new methods against old.
 - Roughly second half of Michaelmas term
- Extension to consider time-dependent estimates.

- Running over Christmas break
- Consider building in feed forwards mechanisms (Kalman filter-like)
- Possible extension to consider:
 - Adding ice altimetry data and/or other data types
 - Effect of knocking out %'s of the networks (e.g. esp tidal guages → look at their average downtime)
 - Comparison of the data types (more data vs more types of data)
 - Consider if different data types have different abilities to resolve features (e.g. tidal gauges trade off within East and West Antarctic Ice ... does this exist within altimetry data?)
 - Early Lent

3 Weekly progress/goals

3.1 Meeting 1: 13/Oct

- Start reading through the literature esp. focusing on:
 - Horton 2018 for general overview
 - Lickley 2018 for sea level to surface
 - The Reciprocity Paper (TRP) §4.6 (and then the references within this)
- Look at implementing equivalent of eq. 81 from TRP to convert the SL change from fingerprint to SSH to then average this over the ocean for GMSL change

3.2 Meeting 2: 17/Oct

- Discussion of scientific method
- Explore the error space for conventional methods
 - Error for each ice sheet over satellite availability bands
 - Ternary space for major ice sheet contributions at satellite availability bands
- Future explore adding:
 - Noise based on satellite sampling
 - Ocean dynamic signals
- Writing useful scripts/functions that can be reused in the future

3.3 Meeting 3: 21/Oct

- Went through error exploration code
- Look into how changing lengths scales can change end distributions
- Explore forcibly pushing forward distribution

3.4 Meeting 4: 24/ Oct

- Work through PyGeoInf tutorials 1→5
- Discussed distribution mapping and pushing forward through function theory
- Parallelise the samples function
- Look at non-zero means

3.5 Meeting 5: 31/Oct

- Spoke through random field operators and pushing forward distributions
- Spoke through workflow of error quantification
- Speak to Dan wrt. Ocean dynamics and ice sheet change sensible distributions
 - Ratios of contribution to error of melt vs dynamics vs noise (... another ternary?)
- Goal to work on implementing a workflow for estimating errors

3.6 Meeting 6: 3/Nov

4 Literature log

5 Thoughts log

5.1 2025-10-10 22:01

To what extent can the noise/artifacts that were filtered out by @coulson2022 be build into the output to see what the “real world” observable would be?

5.2 2025-10-18

Build a load across multiple latitudes and then plot this against error at different satellite segments to see how error varies with load and latitude.

6 Code notes

6.1 2025-10-17

For parallel processing, using joblib’s Parallel and delayed is much simpler than multiprocessing.Pool. Just wrap the function call in delayed() and pass to Parallel.

Tried using ternary-plot but found mpternary worked much better.

6.2 2025-10-26

Using ternary plots works well but it is important to provide contextual side plots that show a selection from the ternary location in actual space.

For example for melt contribution plots show a side plot of what each ternary space plot looks like on a map.

Similarly having bands of satalite latitude ranges showing what is being sampled by different plots can help contextualise the differences seen in distribution ternary plots.

6.3 2025-11-01 14:48

7 Maths




Figure 1: alt text