

Solution 1 (a)

$$\boldsymbol{\mu} = [\mu_x, \mu_y]^T$$

$$\boldsymbol{\Sigma} = \begin{bmatrix} \sigma_x^2 & \rho\sigma_x\sigma_y \\ \rho\sigma_x\sigma_y & \sigma_y^2 \end{bmatrix}$$

- Mean vector: $\boldsymbol{\mu}$
- Covariance matrix: $\boldsymbol{\Sigma}$

Let $\mu_x = 2$, $\mu_y = 2$, $\sigma_x = 1$, $\sigma_y = 0.5$, and $\rho = -0.5$

Solve for mean vector $\boldsymbol{\mu}$:

$$\boldsymbol{\mu} = [\mu_x \quad \mu_y]^T = \begin{bmatrix} \mu_x \\ \mu_y \end{bmatrix} = \begin{bmatrix} 2 \\ 2 \end{bmatrix}$$

Solve for covariance matrix $\boldsymbol{\Sigma}$:

$$\boldsymbol{\Sigma} = \begin{bmatrix} \sigma_x^2 & \rho\sigma_x\sigma_y \\ \rho\sigma_x\sigma_y & \sigma_y^2 \end{bmatrix} = \begin{bmatrix} 1^2 & (-0.5)(1)(0.5) \\ (-0.5)(1)(0.5) & 0.5^2 \end{bmatrix} = \begin{bmatrix} 1 & -0.25 \\ -0.25 & 0.25 \end{bmatrix}$$

\therefore the bivariate Gaussian has parameters:

$$\mathcal{N}(\boldsymbol{\mu}, \boldsymbol{\Sigma}) = \mathcal{N}\left(\begin{bmatrix} 2 \\ 2 \end{bmatrix}, \begin{bmatrix} 1 & -0.25 \\ -0.25 & 0.25 \end{bmatrix}\right)$$

Solution 1 (b)

For a bivariate Gaussian distribution, we need to specify the following parameters:

$$\boldsymbol{\mu} = [\mu_x, \mu_y]^T$$

$$\boldsymbol{\Sigma} = \begin{bmatrix} \sigma_x^2 & \rho\sigma_x\sigma_y \\ \rho\sigma_x\sigma_y & \sigma_y^2 \end{bmatrix}$$

- Mean vector: $\boldsymbol{\mu}$
- Covariance matrix: $\boldsymbol{\Sigma}$

Let $\mu_x = 1$, $\mu_y = 2$, $\sigma_x = 1$, $\sigma_y = 0$

Solve for mean vector $\boldsymbol{\mu}$:

$$\boldsymbol{\mu} = [\mu_x \quad \mu_y]^T = \begin{bmatrix} \mu_x \\ \mu_y \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \end{bmatrix}$$

Solve for covariance matrix $\boldsymbol{\Sigma}$:

$$\boldsymbol{\Sigma} = \begin{bmatrix} \sigma_x^2 & \rho\sigma_x\sigma_y \\ \rho\sigma_x\sigma_y & \sigma_y^2 \end{bmatrix} = \begin{bmatrix} 1^2 & 0 \\ 0 & 0^2 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$$

\therefore the bivariate Gaussian has parameters:

$$\mathcal{N}(\boldsymbol{\mu}, \boldsymbol{\Sigma}) = \mathcal{N}\left(\begin{bmatrix} 1 \\ 2 \end{bmatrix}, \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}\right)$$

Solution 2

Step 1

The Bayesian decision rule for generative classifiers is defined as:

$$\hat{y} = \arg \max_{y \in \{+, -\}} p(y|x) = \arg \max_{y \in \{+, -\}} \frac{p(x|y)p(y)}{p(x)}$$

- Since $p(x)$ is constant for both classes, the decision rule simplifies to:

$$\hat{y} = \arg \max_{y \in \{+, -\}} p(x|y)p(y)$$

Step 2

The (+) class is predicted when the following holds:

$$p(x|+)p(+) > p(x|-)p(-)$$

Step 3

Identify possible reasons for always predicting the positive class.

1. **Highly imbalanced prior probabilities:** If $p(+) \gg p(-)$, the classifier might always predict the positive class because the prior term dominates the decision, regardless of the likelihood term. This occurs when the training data contains many more + examples than - ones.
2. **Poor estimation of class-conditional densities:** If $p(x|+)$ is consistently overestimated or $p(x|-)$ is consistently underestimated across the input space, the classifier will favor the + class.

\therefore the classifier that predicts + for all points x in the input space is likely due to a combination of highly imbalanced prior probabilities and poor estimation of class-conditional densities.

Solution 3 (a)

Step 1

From Bayes Theorem we have:

$$p(y|x) = \frac{p(x|y)p(y)}{p(x)}$$

or

$$p(y|x) \propto p(x|y)p(y)$$

Step 2

The class probabilities are given as:

$$\pi_1 = 0.33 \quad \pi_2 = 0.39 \quad \pi_3 = 0.28$$

From the plot the density for each class is:

$$P(x = 12.0|Class_1) = 0.4 \quad P(x = 12.0|Class_2) = 0.05 \quad P(x = 12.0|Class_3) = 0.0025$$

Step 3

Use class probabilities and densities from *Step 2*, the apply Bayes Theorem

$$P(Class_1|x = 12.0) \propto P(x = 12.0|Class_1)P(Class_1) = 0.4 \times 0.33 = 0.132$$

$$P(Class_2|x = 12.0) \propto P(x = 12.0|Class_2)P(Class_2) = 0.05 \times 0.39 = 0.0195$$

$$P(Class_3|x = 12.0) \propto P(x = 12.0|Class_3)P(Class_3) = 0.0025 \times 0.28 = 0.0007$$

\therefore the label $Class_1$ would be assigned at $x=12.0$

Solution 3 (b)

Step 1

From Bayes Theorem we have:

$$p(y|x) = \frac{p(x|y)p(y)}{p(x)}$$

or

$$p(y|x) \propto p(x|y)p(y)$$

Step 2

The class probabilities are given as:

$$\pi_1 = 0.33 \quad \pi_2 = 0.39 \quad \pi_3 = 0.28$$

From *Figure 1*, the density for each class is:

$$P(x = 12.5|Class_1) = 0.6 \quad P(x = 12.5|Class_2) = 0.3 \quad P(x = 12.5|Class_3) = 0.05$$

Step 3

Use class probabilities and densities from *Step 2*, then apply Bayes Theorem

$$P(Class_1|x = 12.5) \propto P(x = 12.5|Class_1)P(Class_1) = 0.6 \times 0.33 = 0.198$$

$$P(Class_2|x = 12.5) \propto P(x = 12.5|Class_2)P(Class_2) = 0.3 \times 0.39 = 0.117$$

$$P(Class_3|x = 12.5) \propto P(x = 12.5|Class_3)P(Class_3) = 0.05 \times 0.28 = 0.014$$

\therefore the label $Class_1$ would be assigned at $x=12.5$

Solution 3 (c)

Step 1

From Bayes Theorem we have:

$$p(y|x) = \frac{p(x|y)p(y)}{p(x)}$$

or

$$p(y|x) \propto p(x|y)p(y)$$

Step 2

The class probabilities are given as:

$$\pi_1 = 0.33 \quad \pi_2 = 0.39 \quad \pi_3 = 0.28$$

From *Figure 1*, the density for each class is:

$$P(x = 13.0|Class_1) = 0.3 \quad P(x = 13.0|Class_2) = 0.6 \quad P(x = 13.0|Class_3) = 0.2$$

Step 3

Use class probabilities and densities from *Step 2*, then apply Bayes Theorem

$$P(Class_1|x = 13.0) \propto P(x = 13.0|Class_1)P(Class_1) = 0.3 \times 0.33 = 0.099$$

$$P(Class_2|x = 13.0) \propto P(x = 13.0|Class_2)P(Class_2) = 0.6 \times 0.39 = 0.234$$

$$P(Class_3|x = 13.0) \propto P(x = 13.0|Class_3)P(Class_3) = 0.2 \times 0.28 = 0.056$$

\therefore the label $Class_2$ would be assigned at $x=13.0$

Solution 3 (d)

Step 1

From Bayes Theorem we have:

$$p(y|x) = \frac{p(x|y)p(y)}{p(x)}$$

or

$$p(y|x) \propto p(x|y)p(y)$$

Step 2

The class probabilities are given as:

$$\pi_1 = 0.33 \quad \pi_2 = 0.39 \quad \pi_3 = 0.28$$

From *Figure 1*, the density for each class is:

$$P(x = 13.5|Class_1) = 0.1 \quad P(x = 13.5|Class_2) = 0.7 \quad P(x = 13.5|Class_3) = 0.4$$

Step 3

Use class probabilities and densities from *Step 2*, then apply Bayes Theorem

$$P(Class_1|x = 13.5) \propto P(x = 13.5|Class_1)P(Class_1) = 0.1 \times 0.33 = 0.033$$

$$P(Class_2|x = 13.5) \propto P(x = 13.5|Class_2)P(Class_2) = 0.7 \times 0.39 = 0.273$$

$$P(Class_3|x = 13.5) \propto P(x = 13.5|Class_3)P(Class_3) = 0.4 \times 0.28 = 0.112$$

\therefore the label $Class_2$ would be assigned at $x=13.5$

Solution 3 (e)

Step 1

From Bayes Theorem we have:

$$p(y|x) = \frac{p(x|y)p(y)}{p(x)}$$

or

$$p(y|x) \propto p(x|y)p(y)$$

Step 2

The class probabilities are given as:

$$\pi_1 = 0.33 \quad \pi_2 = 0.39 \quad \pi_3 = 0.28$$

From *Figure 1*, the density for each class is:

$$P(x = 14.0|Class_1) = 0.05 \quad P(x = 14.0|Class_2) = 0.2 \quad P(x = 14.0|Class_3) = 0.8$$

Step 3

Use class probabilities and densities from *Step 2*, then apply Bayes Theorem

$$P(Class_1|x = 14.0) \propto P(x = 14.0|Class_1)P(Class_1) = 0.05 \times 0.33 = 0.0165$$

$$P(Class_2|x = 14.0) \propto P(x = 14.0|Class_2)P(Class_2) = 0.2 \times 0.39 = 0.078$$

$$P(Class_3|x = 14.0) \propto P(x = 14.0|Class_3)P(Class_3) = 0.8 \times 0.28 = 0.224$$

\therefore the label $Class_3$ would be assigned at $x=14.0$

Solution 4 (a)

Step 1Analyze μ

- Since $\mu = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$, the center of the Gaussian is at the origin.

Step 2Analyze Σ


- The covariance matrix $\Sigma = \begin{bmatrix} 9 & 0 \\ 0 & 1 \end{bmatrix}$ indicates that the variance in the x-direction is 9 and in the y-direction is 1. This means that the Gaussian will be elongated along the x-axis.

Step 3Look at the standard deviation to determine the shape, since Σ is a diagonal matrix

- The standard deviation in the x-direction is $\sigma_x = \sqrt{9} = 3$
- The standard deviation in the y-direction is $\sigma_y = \sqrt{1} = 1$

Step 4

Sketch the Gaussian



Solution 4 (b)

Step 1Analyze μ

- Since $\mu = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$, the center of the Gaussian is at the origin.

Step 2Analyze Σ

- The covariance matrix $\Sigma = \begin{bmatrix} 1 & -0.75 \\ -0.75 & 1 \end{bmatrix}$ has non-zero off-diagonal elements.
- This indicates a correlation between variables.
- The negative correlation (-0.75) means that as one variable increases, the other tends to decrease.
- Look at the eigenvalues and eigenvectors to determine the shape since we don't have a diagonal matrix in this case.


Step 3

Find eigenvalues and eigenvectors

- The eigenvalues of Σ are $\lambda_1 = 1 + 0.75 = 1.75$ and $\lambda_2 = 1 - 0.75 = 0.25$.
- The corresponding eigenvectors are $v_1 = \begin{bmatrix} 1 \\ -1 \end{bmatrix}$ and $v_2 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$.
- The standard deviations along the principal axes are $\sigma_1 = \sqrt{1.75} \approx 1.32$ and $\sigma_2 = \sqrt{0.25} = 0.5$.

Step 4

Sketch the Gaussian



Solution 5

A unit vector in \mathbb{R}^2 is a vector of the form:

$$\begin{bmatrix} x \\ y \end{bmatrix}$$

To be orthogonal to $\begin{bmatrix} 1 \\ 1 \end{bmatrix}$, the dot product must equal zero:

$$\begin{bmatrix} 1 \\ 1 \end{bmatrix} \cdot \begin{bmatrix} x \\ y \end{bmatrix} = 0$$

This gives the equation:

$$x + y = 0$$

This means that $y = -x$.

Now, we need to find the unit vectors. A unit vector has a magnitude of 1:

$$\sqrt{x^2 + y^2} = 1$$

Substituting $y = -x$ into the equation:

$$\sqrt{x^2 + (-x)^2} = 1$$

$$\sqrt{2x^2} = 1$$

$$\sqrt{2}|x| = 1$$

$$|x| = \frac{1}{\sqrt{2}}$$

This gives two solutions for x :

$$x = \frac{1}{\sqrt{2}} \quad \text{or} \quad x = -\frac{1}{\sqrt{2}}$$

Substituting back to find y :

$$y = -\frac{1}{\sqrt{2}} \quad \text{or} \quad y = \frac{1}{\sqrt{2}}$$

\therefore the unit vectors orthogonal to $\begin{bmatrix} 1 \\ 1 \end{bmatrix}$ are:

$$\begin{bmatrix} \frac{1}{\sqrt{2}} \\ -\frac{1}{\sqrt{2}} \end{bmatrix} \quad \text{and} \quad \begin{bmatrix} -\frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} \end{bmatrix}$$

Solution 6

Let $x \cdot x = 25$, $\forall x \in \mathbb{R}^d$

$$x \cdot x = \|x\|^2 = 25 \rightarrow \sqrt{\|x\|^2} = \sqrt{25} \rightarrow \|x\| = 5$$

Hence, the vector x has a magnitude of 5.

\therefore the set of all points $x \in \mathbb{R}^d$ with $x \cdot x = 25$ is a sphere of radius 5 centered at the origin in d -dimensional space.

Solution 7

The function $f(x) = 2x_1 - x_2 + 6x_3$ can be expressed in the form of a dot product:

$$f(x) = w \cdot x$$

Where w is a vector in \mathbb{R}^3 and x is a vector in \mathbb{R}^3 .

We can then express $f(x)$ as the dot product of two matrices:

$$f(x) = \begin{bmatrix} w_1 \\ w_2 \\ w_3 \end{bmatrix} \cdot \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = w_1 \cdot x_1 + w_2 \cdot x_2 + w_3 \cdot x_3$$

It follows that: $w_1 = 2$ $w_2 = -1$ $w_3 = 6$

\therefore the vector w is:

$$w = \begin{bmatrix} 2 \\ -1 \\ 6 \end{bmatrix}$$

Solution 8

Let the dimensions of A be $m \times n$ and the dimensions of B be $n \times p$.

The product AB will have dimensions $m \times p$.

Given that AB has dimensions 10×20 , and $A = m \times 30$.

$$m = 10 \qquad p = 20 \qquad n = 30$$

We know that the product AB can be expressed as:

$$AB = (m \times n) \times (n \times p)$$

\therefore the dimensions of A and B are:

$$A : 10 \times 30$$

$$B : 30 \times 20$$

Solution 9 (a)

The matrix X has n rows and d columns, so the dimension of X is:

$$X \in \mathbb{R}^{n \times d}$$

This means that X has n data points, each with d features.

\therefore the dimension of X is $n \times d$.

Solution 9 (b)

The matrix XX^T is the product of an $n \times d$ matrix and a $d \times n$ matrix.

The resulting matrix will have dimensions $n \times n$.

\therefore the dimension of XX^T is $n \times n$.

Solution 9 (c)

The (i, j) entry of $X^T X$ is the dot product of the i -th row of X and the j -th column of X^T

This is simply the sum of the products of the corresponding elements:

$$(X^T X)_{ij} = \sum_{k=1}^d x_k^{(i)} x_k^{(j)}$$

This is the inner product of the i -th and j -th data points.

\therefore the (i, j) entry of $X^T X$ is the inner product of the i -th and j -th data points or $(x^{(i)}, x^{(j)})$.

Solution 10

Step 1Compute x^T

$$x = \begin{bmatrix} 1 \\ 3 \\ 5 \end{bmatrix} \rightarrow x^T = [1 \quad 3 \quad 5]$$

Step 2Compute $x^T x$

$$x^T x = [1 \quad 3 \quad 5] \begin{bmatrix} 1 \\ 3 \\ 5 \end{bmatrix} = 1^2 + 3^2 + 5^2 = 1 + 9 + 25 = 35$$

Step 3Compute xx^T

$$xx^T = \begin{bmatrix} 1 \\ 3 \\ 5 \end{bmatrix} [1 \quad 3 \quad 5] = \begin{bmatrix} 1^2 & 1 \cdot 3 & 1 \cdot 5 \\ 3 \cdot 1 & 3^2 & 3 \cdot 5 \\ 5 \cdot 1 & 5 \cdot 3 & 5^2 \end{bmatrix} = \begin{bmatrix} 1 & 3 & 5 \\ 3 & 9 & 15 \\ 5 & 15 & 25 \end{bmatrix}$$

\therefore the result of $x^T x$ is a scalar 35 and the result of xx^T is a matrix:

$$\begin{bmatrix} 1 & 3 & 5 \\ 3 & 9 & 15 \\ 5 & 15 & 25 \end{bmatrix}$$

Solution 11

$$\text{Let } f(x) = 3x_1^2 + 2x_1x_2 - 4x_1x_3 + 6x_3^2$$

Step 1

Define x , x^T , and M

$$x = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix}$$

$$x^T = [x_1 \quad x_2 \quad x_3]$$

$$M = \begin{bmatrix} m_{11} & m_{12} & m_{13} \\ m_{21} & m_{22} & m_{23} \\ m_{31} & m_{32} & m_{33} \end{bmatrix}$$

Step 2

Expand $x^T M x$ so it is in the same form as $f(x)$

$$x^T M x = m_{11}x_1^2 + m_{12}x_1x_2 + m_{13}x_1x_3 + m_{21}x_2x_1 + m_{22}x_2^2 + m_{23}x_2x_3 + m_{31}x_3x_1 + m_{32}x_3x_2 + m_{33}x_3^2.$$

Step 3

Match coefficients of $x^T M x$ with $f(x)$

$$m_{11} = 3 \quad m_{12} = 2 \quad m_{13} = -4$$

$$m_{21} = 2 \quad m_{22} = 0 \quad m_{23} = 0$$

$$m_{31} = -4 \quad m_{32} = 0 \quad m_{33} = 6$$

Step 4

Plug in values and express the matrix M

$$M = \begin{bmatrix} m_{11} & m_{12} & m_{13} \\ m_{21} & m_{22} & m_{23} \\ m_{31} & m_{32} & m_{33} \end{bmatrix} = \begin{bmatrix} 3 & 2 & -4 \\ 2 & 0 & 0 \\ -4 & 0 & 6 \end{bmatrix}$$

\therefore the matrix M is:

$$M = \begin{bmatrix} 3 & 2 & -4 \\ 2 & 0 & 0 \\ -4 & 0 & 6 \end{bmatrix}$$

Solution 12 (a)

The determinant of a diagonal matrix is the product of its diagonal elements:

$$|A| = 1 \cdot 2 \cdot 3 \cdot 4 \cdot 5 \cdot 6 \cdot 7 \cdot 8 = 8! = 40320$$

\therefore the determinant of $A = 40320$

Solution 12 (b)

The inverse of a diagonal matrix is obtained by taking the reciprocal of each diagonal element:

$$A^{-1} = \text{diag}\left(\frac{1}{1}, \frac{1}{2}, \frac{1}{3}, \frac{1}{4}, \frac{1}{5}, \frac{1}{6}, \frac{1}{7}, \frac{1}{8}\right)$$

\therefore the inverse of A is:

$$A^{-1} = \begin{bmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & \frac{1}{2} & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & \frac{1}{3} & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & \frac{1}{4} & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & \frac{1}{5} & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & \frac{1}{6} & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & \frac{1}{7} & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & \frac{1}{8} \end{bmatrix}$$

Solution 13 (a)

Pseudocode for training procedure:

1. Input

- x : training data
- y : training labels
- c : smoothing constant for covariance matrices

2. Initialize

- x_{train} : 80% of x
- x_{val} : 20% of x
- y_{train} : 80% of y
- y_{val} : 20% of y
- π_k : class k frequencies
- μ_k : class k mean vector
- Σ_k : class k covariance matrices
- qda: Gaussian generative model

3. Iterate

- For each class k in 0, 1, ..., 9:
 - Compute π_k as the fraction of training points in class k
 - Compute μ_k as the mean of training points in class k
 - Compute Σ_k as the covariance of training points in class k
 - Smooth Σ_k by adding cI to it
 - Store each π_k , μ_k , and Σ_k

4. Evaluate

- For each point in the validation set(x_{val}) and each c value tested:
 - Compute average accuracy of the model over all classes
 - Store the accuracy for each c value
 - Choose the c value that gives the best accuracy
 - Store the best c value
- Compute the final model using the best c value
- Store each π_k , μ_k , and Σ_k for the final model

Solution 13 (b)

- A single value of $c = 0.95$ for all ten classes.
- The value of c was chosen based on the validation set accuracy.
- The accuracy for $c = 0.95$ was 0.8742 or 87.42%.





Figure 1: Results from c value testing


Solution 13 (c)

- The model with $c = 0.95$ predicted 1178 out of 10,000 incorrectly.
- Therefore, the error rate on the MNIST test set was 0.1178 or 11.78%.
- Note: The accuracy of the model on the MNIST test set (88.22%) was very close to accuracy on the training set (87.42)%.


Solution 13 (d)




(a) 5 classified as 3




(b) 3 classified as 2



(c) 5 classified as 8



(d) 1 classified as 8



(e) 5 classified as 8

Figure 2: Examples of misclassified digits from the test set