

**Solution 1 (a)**

---

$$\boldsymbol{\mu} = [\mu_x, \mu_y]^T$$

$$\boldsymbol{\Sigma} = \begin{bmatrix} \sigma_x^2 & \rho\sigma_x\sigma_y \\ \rho\sigma_x\sigma_y & \sigma_y^2 \end{bmatrix}$$

- Mean vector:  $\boldsymbol{\mu}$
- Covariance matrix:  $\boldsymbol{\Sigma}$

Let  $\mu_x = 2$ ,  $\mu_y = 2$ ,  $\sigma_x = 1$ ,  $\sigma_y = 0.5$ , and  $\rho = -0.5$

Solve for mean vector  $\boldsymbol{\mu}$ :

$$\boldsymbol{\mu} = [\mu_x \quad \mu_y]^T = \begin{bmatrix} \mu_x \\ \mu_y \end{bmatrix} = \begin{bmatrix} 2 \\ 2 \end{bmatrix}$$

Solve for covariance matrix  $\boldsymbol{\Sigma}$ :

$$\boldsymbol{\Sigma} = \begin{bmatrix} \sigma_x^2 & \rho\sigma_x\sigma_y \\ \rho\sigma_x\sigma_y & \sigma_y^2 \end{bmatrix} = \begin{bmatrix} 1^2 & (-0.5)(1)(0.5) \\ (-0.5)(1)(0.5) & 0.5^2 \end{bmatrix} = \begin{bmatrix} 1 & -0.25 \\ -0.25 & 0.25 \end{bmatrix}$$

$\therefore$  the bivariate Gaussian has parameters:

$$\mathcal{N}(\boldsymbol{\mu}, \boldsymbol{\Sigma}) = \mathcal{N}\left(\begin{bmatrix} 2 \\ 2 \end{bmatrix}, \begin{bmatrix} 1 & -0.25 \\ -0.25 & 0.25 \end{bmatrix}\right)$$

---

**Solution 1 (b)**

---

For a bivariate Gaussian distribution, we need to specify the following parameters:

$$\boldsymbol{\mu} = [\mu_x, \mu_y]^T$$

$$\boldsymbol{\Sigma} = \begin{bmatrix} \sigma_x^2 & \rho\sigma_x\sigma_y \\ \rho\sigma_x\sigma_y & \sigma_y^2 \end{bmatrix}$$

- Mean vector:  $\boldsymbol{\mu}$
- Covariance matrix:  $\boldsymbol{\Sigma}$

Let  $\mu_x = 1$ ,  $\mu_y = 2$ ,  $\sigma_x = 1$ ,  $\sigma_y = 0$

Solve for mean vector  $\boldsymbol{\mu}$ :

$$\boldsymbol{\mu} = [\mu_x \quad \mu_y]^T = \begin{bmatrix} \mu_x \\ \mu_y \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \end{bmatrix}$$

Solve for covariance matrix  $\boldsymbol{\Sigma}$ :

$$\boldsymbol{\Sigma} = \begin{bmatrix} \sigma_x^2 & \rho\sigma_x\sigma_y \\ \rho\sigma_x\sigma_y & \sigma_y^2 \end{bmatrix} = \begin{bmatrix} 1^2 & 0 \\ 0 & 0^2 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$$

$\therefore$  the bivariate Gaussian has parameters:

$$\mathcal{N}(\boldsymbol{\mu}, \boldsymbol{\Sigma}) = \mathcal{N}\left(\begin{bmatrix} 1 \\ 2 \end{bmatrix}, \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}\right)$$

---

## Solution 2

---

### Step 1

The Bayesian decision rule for generative classifiers is defined as:

$$\hat{y} = \arg \max_{y \in \{+, -\}} p(y|x) = \arg \max_{y \in \{+, -\}} \frac{p(x|y)p(y)}{p(x)}$$

- Since  $p(x)$  is constant for both classes, the decision rule simplifies to:

$$\hat{y} = \arg \max_{y \in \{+, -\}} p(x|y)p(y)$$

### Step 2

The (+) class is predicted when the following holds:

$$p(x|+)p(+) > p(x|-)p(-)$$

### Step 3

Identify possible reasons for always predicting the positive class.

1. **Highly imbalanced prior probabilities:** If  $p(+) \gg p(-)$ , the classifier might always predict the positive class because the prior term dominates the decision, regardless of the likelihood term. This occurs when the training data contains many more + examples than - ones.
2. **Poor estimation of class-conditional densities:** If  $p(x|+)$  is consistently overestimated or  $p(x|-)$  is consistently underestimated across the input space, the classifier will favor the + class.

$\therefore$  the classifier that predicts + for all points  $x$  in the input space is likely due to a combination of highly imbalanced prior probabilities and poor estimation of class-conditional densities.

---

**Solution 3 (a)**

---

**Step 1**

From Bayes Theorem we have:

$$p(y|x) = \frac{p(x|y)p(y)}{p(x)}$$

or

$$p(y|x) \propto p(x|y)p(y)$$

**Step 2**

The class probabilities are given as:

$$\pi_1 = 0.33 \quad \pi_2 = 0.39 \quad \pi_3 = 0.28$$

From the plot the density for each class is:

$$P(x = 12.0|Class_1) = 0.4 \quad P(x = 12.0|Class_2) = 0.05 \quad P(x = 12.0|Class_3) = 0.0025$$

**Step 3**

Use class probabilities and densities from *Step 2*, the apply Bayes Theorem

$$P(Class_1|x = 12.0) \propto P(x = 12.0|Class_1)P(Class_1) = 0.4 \times 0.33 = 0.132$$

$$P(Class_2|x = 12.0) \propto P(x = 12.0|Class_2)P(Class_2) = 0.05 \times 0.39 = 0.0195$$

$$P(Class_3|x = 12.0) \propto P(x = 12.0|Class_3)P(Class_3) = 0.0025 \times 0.28 = 0.0007$$

$\therefore$  the label  $Class_1$  would be assigned at  $x=12.0$

---

**Solution 3 (b)**

---

**Step 1**

From Bayes Theorem we have:

$$p(y|x) = \frac{p(x|y)p(y)}{p(x)}$$

or

$$p(y|x) \propto p(x|y)p(y)$$

**Step 2**

The class probabilities are given as:

$$\pi_1 = 0.33 \quad \pi_2 = 0.39 \quad \pi_3 = 0.28$$

From *Figure 1*, the density for each class is:

$$P(x = 12.5|Class_1) = 0.6 \quad P(x = 12.5|Class_2) = 0.3 \quad P(x = 12.5|Class_3) = 0.05$$

**Step 3**

Use class probabilities and densities from *Step 2*, then apply Bayes Theorem

$$P(Class_1|x = 12.5) \propto P(x = 12.5|Class_1)P(Class_1) = 0.6 \times 0.33 = 0.198$$

$$P(Class_2|x = 12.5) \propto P(x = 12.5|Class_2)P(Class_2) = 0.3 \times 0.39 = 0.117$$

$$P(Class_3|x = 12.5) \propto P(x = 12.5|Class_3)P(Class_3) = 0.05 \times 0.28 = 0.014$$

$\therefore$  the label  $Class_1$  would be assigned at  $x=12.5$

---

**Solution 3 (c)**

---

**Step 1**

From Bayes Theorem we have:

$$p(y|x) = \frac{p(x|y)p(y)}{p(x)}$$

or

$$p(y|x) \propto p(x|y)p(y)$$

**Step 2**

The class probabilities are given as:

$$\pi_1 = 0.33 \quad \pi_2 = 0.39 \quad \pi_3 = 0.28$$

From *Figure 1*, the density for each class is:

$$P(x = 13.0|Class_1) = 0.3 \quad P(x = 13.0|Class_2) = 0.6 \quad P(x = 13.0|Class_3) = 0.2$$

**Step 3**

Use class probabilities and densities from *Step 2*, then apply Bayes Theorem

$$P(Class_1|x = 13.0) \propto P(x = 13.0|Class_1)P(Class_1) = 0.3 \times 0.33 = 0.099$$

$$P(Class_2|x = 13.0) \propto P(x = 13.0|Class_2)P(Class_2) = 0.6 \times 0.39 = 0.234$$

$$P(Class_3|x = 13.0) \propto P(x = 13.0|Class_3)P(Class_3) = 0.2 \times 0.28 = 0.056$$

$\therefore$  the label  $Class_2$  would be assigned at  $x=13.0$

---

**Solution 3 (d)**

---

**Step 1**

From Bayes Theorem we have:

$$p(y|x) = \frac{p(x|y)p(y)}{p(x)}$$

or

$$p(y|x) \propto p(x|y)p(y)$$

**Step 2**

The class probabilities are given as:

$$\pi_1 = 0.33 \quad \pi_2 = 0.39 \quad \pi_3 = 0.28$$

From *Figure 1*, the density for each class is:

$$P(x = 13.5|Class_1) = 0.1 \quad P(x = 13.5|Class_2) = 0.7 \quad P(x = 13.5|Class_3) = 0.4$$

**Step 3**

Use class probabilities and densities from *Step 2*, then apply Bayes Theorem

$$P(Class_1|x = 13.5) \propto P(x = 13.5|Class_1)P(Class_1) = 0.1 \times 0.33 = 0.033$$

$$P(Class_2|x = 13.5) \propto P(x = 13.5|Class_2)P(Class_2) = 0.7 \times 0.39 = 0.273$$

$$P(Class_3|x = 13.5) \propto P(x = 13.5|Class_3)P(Class_3) = 0.4 \times 0.28 = 0.112$$

$\therefore$  the label  $Class_2$  would be assigned at  $x=13.5$

---

**Solution 3 (e)**

---

**Step 1**

From Bayes Theorem we have:

$$p(y|x) = \frac{p(x|y)p(y)}{p(x)}$$

or

$$p(y|x) \propto p(x|y)p(y)$$

**Step 2**

The class probabilities are given as:

$$\pi_1 = 0.33 \quad \pi_2 = 0.39 \quad \pi_3 = 0.28$$

From *Figure 1*, the density for each class is:

$$P(x = 14.0|Class_1) = 0.05 \quad P(x = 14.0|Class_2) = 0.2 \quad P(x = 14.0|Class_3) = 0.8$$

**Step 3**

Use class probabilities and densities from *Step 2*, then apply Bayes Theorem

$$P(Class_1|x = 14.0) \propto P(x = 14.0|Class_1)P(Class_1) = 0.05 \times 0.33 = 0.0165$$

$$P(Class_2|x = 14.0) \propto P(x = 14.0|Class_2)P(Class_2) = 0.2 \times 0.39 = 0.078$$

$$P(Class_3|x = 14.0) \propto P(x = 14.0|Class_3)P(Class_3) = 0.8 \times 0.28 = 0.224$$

$\therefore$  the label  $Class_3$  would be assigned at  $x=14.0$

---



**Solution 4 (a)**

---

**Step 1**Analyze  $\mu$ 

- Since  $\mu = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$ , the center of the Gaussian is at the origin.

**Step 2**Analyze  $\Sigma$ 


- The covariance matrix  $\Sigma = \begin{bmatrix} 9 & 0 \\ 0 & 1 \end{bmatrix}$  indicates that the variance in the x-direction is 9 and in the y-direction is 1. This means that the Gaussian will be elongated along the x-axis.

**Step 3**Look at the standard deviation to determine the shape, since  $\Sigma$  is a diagonal matrix

- The standard deviation in the x-direction is  $\sigma_x = \sqrt{9} = 3$
- The standard deviation in the y-direction is  $\sigma_y = \sqrt{1} = 1$

**Step 4**

Sketch the Gaussian



**Solution 4 (b)**

---

**Step 1**Analyze  $\mu$ 

- Since  $\mu = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$ , the center of the Gaussian is at the origin.

**Step 2**Analyze  $\Sigma$ 

- The covariance matrix  $\Sigma = \begin{bmatrix} 1 & -0.75 \\ -0.75 & 1 \end{bmatrix}$  has non-zero off-diagonal elements.
- This indicates a correlation between variables.
- The negative correlation ( $-0.75$ ) means that as one variable increases, the other tends to decrease.
- Look at the eigenvalues and eigenvectors to determine the shape since we don't have a diagonal matrix in this case.


**Step 3**

Find eigenvalues and eigenvectors

- The eigenvalues of  $\Sigma$  are  $\lambda_1 = 1 + 0.75 = 1.75$  and  $\lambda_2 = 1 - 0.75 = 0.25$ .
- The corresponding eigenvectors are  $v_1 = \begin{bmatrix} 1 \\ -1 \end{bmatrix}$  and  $v_2 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$ .
- The standard deviations along the principal axes are  $\sigma_1 = \sqrt{1.75} \approx 1.32$  and  $\sigma_2 = \sqrt{0.25} = 0.5$ .

**Step 4**

Sketch the Gaussian



**Solution 5**

---

A unit vector in  $\mathbb{R}^2$  is a vector of the form:

$$\begin{bmatrix} x \\ y \end{bmatrix}$$

To be orthogonal to  $\begin{bmatrix} 1 \\ 1 \end{bmatrix}$ , the dot product must equal zero:

$$\begin{bmatrix} 1 \\ 1 \end{bmatrix} \cdot \begin{bmatrix} x \\ y \end{bmatrix} = 0$$

This gives the equation:

$$x + y = 0$$

This means that  $y = -x$ .

Now, we need to find the unit vectors. A unit vector has a magnitude of 1:

$$\sqrt{x^2 + y^2} = 1$$

Substituting  $y = -x$  into the equation:

$$\sqrt{x^2 + (-x)^2} = 1$$

$$\sqrt{2x^2} = 1$$

$$\sqrt{2}|x| = 1$$

$$|x| = \frac{1}{\sqrt{2}}$$

This gives two solutions for  $x$ :

$$x = \frac{1}{\sqrt{2}} \quad \text{or} \quad x = -\frac{1}{\sqrt{2}}$$

Substituting back to find  $y$ :

$$y = -\frac{1}{\sqrt{2}} \quad \text{or} \quad y = \frac{1}{\sqrt{2}}$$

$\therefore$  the unit vectors orthogonal to  $\begin{bmatrix} 1 \\ 1 \end{bmatrix}$  are:

$$\begin{bmatrix} \frac{1}{\sqrt{2}} \\ -\frac{1}{\sqrt{2}} \end{bmatrix} \quad \text{and} \quad \begin{bmatrix} -\frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} \end{bmatrix}$$

---

**Solution 6**

---

Let  $x \cdot x = 25$ ,  $\forall x \in \mathbb{R}^d$

$$x \cdot x = \|x\|^2 = 25 \rightarrow \sqrt{\|x\|^2} = \sqrt{25} \rightarrow \|x\| = 5$$

Hence, the vector  $x$  has a magnitude of 5.

$\therefore$  the set of all points  $x \in \mathbb{R}^d$  with  $x \cdot x = 25$  is a sphere of radius 5 centered at the origin in  $d$ -dimensional space.

---

**Solution 7**

---

The function  $f(x) = 2x_1 - x_2 + 6x_3$  can be expressed in the form of a dot product:

$$f(x) = w \cdot x$$

Where  $w$  is a vector in  $\mathbb{R}^3$  and  $x$  is a vector in  $\mathbb{R}^3$ .

We can then express  $f(x)$  as the dot product of two matrices:

$$f(x) = \begin{bmatrix} w_1 \\ w_2 \\ w_3 \end{bmatrix} \cdot \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = w_1 \cdot x_1 + w_2 \cdot x_2 + w_3 \cdot x_3$$

It follows that:  $w_1 = 2$   $w_2 = -1$   $w_3 = 6$

$\therefore$  the vector  $w$  is:

$$w = \begin{bmatrix} 2 \\ -1 \\ 6 \end{bmatrix}$$

---

**Solution 8**

---

Let the dimensions of  $A$  be  $m \times n$  and the dimensions of  $B$  be  $n \times p$ .

The product  $AB$  will have dimensions  $m \times p$ .

Given that  $AB$  has dimensions  $10 \times 20$ , and  $A = m \times 30$ .

$$m = 10 \qquad p = 20 \qquad n = 30$$

We know that the product  $AB$  can be expressed as:

$$AB = (m \times n) \times (n \times p)$$

$\therefore$  the dimensions of  $A$  and  $B$  are:

$$A : 10 \times 30$$

$$B : 30 \times 20$$

---

**Solution 9 (a)**

---

The matrix  $X$  has  $n$  rows and  $d$  columns, so the dimension of  $X$  is:

$$X \in \mathbb{R}^{n \times d}$$

This means that  $X$  has  $n$  data points, each with  $d$  features.

$\therefore$  the dimension of  $X$  is  $n \times d$ .

---

**Solution 9 (b)**

---

The matrix  $XX^T$  is the product of an  $n \times d$  matrix and a  $d \times n$  matrix.

The resulting matrix will have dimensions  $n \times n$ .

$\therefore$  the dimension of  $XX^T$  is  $n \times n$ .

---



**Solution 9 (c)**

---

The  $(i, j)$  entry of  $X^T X$  is the dot product of the  $i$ -th row of  $X$  and the  $j$ -th column of  $X^T$

This is simply the sum of the products of the corresponding elements:

$$(X^T X)_{ij} = \sum_{k=1}^d x_k^{(i)} x_k^{(j)}$$

This is the inner product of the  $i$ -th and  $j$ -th data points.

$\therefore$  the  $(i, j)$  entry of  $X^T X$  is the inner product of the  $i$ -th and  $j$ -th data points or  $(x^{(i)}, x^{(j)})$ .

---

**Solution 10**

---

**Step 1**Compute  $x^T$ 

$$x = \begin{bmatrix} 1 \\ 3 \\ 5 \end{bmatrix} \rightarrow x^T = [1 \quad 3 \quad 5]$$

**Step 2**Compute  $x^T x$ 

$$x^T x = [1 \quad 3 \quad 5] \begin{bmatrix} 1 \\ 3 \\ 5 \end{bmatrix} = 1^2 + 3^2 + 5^2 = 1 + 9 + 25 = 35$$

**Step 3**Compute  $xx^T$ 

$$xx^T = \begin{bmatrix} 1 \\ 3 \\ 5 \end{bmatrix} [1 \quad 3 \quad 5] = \begin{bmatrix} 1^2 & 1 \cdot 3 & 1 \cdot 5 \\ 3 \cdot 1 & 3^2 & 3 \cdot 5 \\ 5 \cdot 1 & 5 \cdot 3 & 5^2 \end{bmatrix} = \begin{bmatrix} 1 & 3 & 5 \\ 3 & 9 & 15 \\ 5 & 15 & 25 \end{bmatrix}$$

$\therefore$  the result of  $x^T x$  is a scalar 35 and the result of  $xx^T$  is a matrix:

$$\begin{bmatrix} 1 & 3 & 5 \\ 3 & 9 & 15 \\ 5 & 15 & 25 \end{bmatrix}$$

---

**Solution 11**

---

$$\text{Let } f(x) = 3x_1^2 + 2x_1x_2 - 4x_1x_3 + 6x_3^2$$

**Step 1**

Define  $x$ ,  $x^T$ , and  $M$

$$x = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix}$$

$$x^T = [x_1 \quad x_2 \quad x_3]$$

$$M = \begin{bmatrix} m_{11} & m_{12} & m_{13} \\ m_{21} & m_{22} & m_{23} \\ m_{31} & m_{32} & m_{33} \end{bmatrix}$$

**Step 2**

Expand  $x^T M x$  so it is in the same form as  $f(x)$

$$x^T M x = m_{11}x_1^2 + m_{12}x_1x_2 + m_{13}x_1x_3 + m_{21}x_2x_1 + m_{22}x_2^2 + m_{23}x_2x_3 + m_{31}x_3x_1 + m_{32}x_3x_2 + m_{33}x_3^2.$$

**Step 3**

Match coefficients of  $x^T M x$  with  $f(x)$

$$m_{11} = 3 \quad m_{12} = 2 \quad m_{13} = -4$$

$$m_{21} = 2 \quad m_{22} = 0 \quad m_{23} = 0$$

$$m_{31} = -4 \quad m_{32} = 0 \quad m_{33} = 6$$

**Step 4**

Plug in values and express the matrix  $M$

$$M = \begin{bmatrix} m_{11} & m_{12} & m_{13} \\ m_{21} & m_{22} & m_{23} \\ m_{31} & m_{32} & m_{33} \end{bmatrix} = \begin{bmatrix} 3 & 2 & -4 \\ 2 & 0 & 0 \\ -4 & 0 & 6 \end{bmatrix}$$

$\therefore$  the matrix  $M$  is:

$$M = \begin{bmatrix} 3 & 2 & -4 \\ 2 & 0 & 0 \\ -4 & 0 & 6 \end{bmatrix}$$

---

**Solution 12 (a)**

---

The determinant of a diagonal matrix is the product of its diagonal elements:

$$|A| = 1 \cdot 2 \cdot 3 \cdot 4 \cdot 5 \cdot 6 \cdot 7 \cdot 8 = 8! = 40320$$

$\therefore$  the determinant of  $A = 40320$

---

## Solution 12 (b)

---

Since all values above and below the diagonal are zero the inverse of  $A$  is obtained by taking the reciprocal of each diagonal element:

$$A^{-1} = \text{diag}\left(\frac{1}{1}, \frac{1}{2}, \frac{1}{3}, \frac{1}{4}, \frac{1}{5}, \frac{1}{6}, \frac{1}{7}, \frac{1}{8}\right)$$

Checking the definition of the inverse  $AA^{-1} = I$ :

$$AA^{-1} = \begin{bmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 2 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 3 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 4 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 5 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 6 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 7 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 8 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & \frac{1}{2} & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & \frac{1}{3} & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & \frac{1}{4} & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & \frac{1}{5} & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & \frac{1}{6} & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & \frac{1}{7} & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & \frac{1}{8} \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$

$\therefore A^{-1}$  is:

$$A^{-1} = \begin{bmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & \frac{1}{2} & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & \frac{1}{3} & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & \frac{1}{4} & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & \frac{1}{5} & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & \frac{1}{6} & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & \frac{1}{7} & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & \frac{1}{8} \end{bmatrix}$$


---

## Solution 13 (a)

Pseudocode for training procedure:

```

1  FUNCTION fit_generative_model(x, y):
2      SET k = 10 (number of classes)
3      SET d = number of features in x
4      INITIALIZE mu AS kxd matrix of zeros
5      INITIALIZE sigma AS kxdxd matrix of zeros
6      INITIALIZE pi AS vector of k zeros
7
8      Normalize x to range [0,1]
9
10     Split data into training (80%) and validation (20%) sets
11
12     FUNCTION calc_priors_means_covariances(x, y, c):
13         FOR each class j from 0 to k-1:
14             Find all samples where y equals j
15             CALCULATE prior probability pi[j] AS fraction of samples in class j
16             CALCULATE mean vector mu[j] from samples in class j
17             IF c > 0:
18                 CALCULATE covariance matrix sigma[j] with regularization c
19             ELSE:
20                 CALCULATE covariance matrix sigma[j] without regularization
21         RETURN mu, sigma, pi
22
23     FUNCTION fit(x_train, y_train, x_val, y_val, c):
24         CALCULATE mu, sigma, pi using training data with regularization c
25         INITIALIZE score matrix for validation samples
26
27         FOR each class label from 0 to k-1:
28             Create multivariate normal distribution with mean mu[label] and covariance
                sigma[label]
29             CALCULATE log probability for all validation samples (vectorized)
30
31         Make predictions by finding class with highest score for each sample
32         Count number of errors (predictions != y_val)
33         RETURN number of errors
34
35     FUNCTION find_best_c():
36         INITIALIZE list of c values to test
37         INITIALIZE empty list for errors
38         INITIALIZE dictionary to store c values and errors
39
40         FOR each c value:
41             CALCULATE error using fit function
42             Store c and error in lists and dictionary
43
44         Find c with minimum error
45         RETURN best c, c_error_dictionary
46
47     CALCULATE best c using find_best_c function
48
49     CALCULATE final mu, sigma, pi using entire dataset with best c
50
51     RETURN mu, sigma, pi
52 END FUNCTION

```

Python code for training procedure:

```

1  def fit_generative_model(x,y):
2      k = 10 # labels 0,1,...,k-1
3      d = (x.shape)[1] # number of features
4      mu = np.zeros((k,d))
5      sigma = np.zeros((k,d,d))
6      pi = np.zeros(k)
7
8      ## normalize data [0,1]
9      x = x.astype(np.float32)/255.0

```

```

10
11     ## initialize training set and validation set to find best regularization parameter
12     c
13     ## note: we will do a 80/20 split of x,y
14     x_train, x_val, y_train, y_val = train_test_split(x, y, test_size=0.2,
15         random_state=17)
16
17     def calc_priors_means_covariances(x,y,c):
18         ## calc priors, means, and covariances
19         if c > 0:
20             for j in range(k):
21                 indices = (y==j).flatten()
22                 x_j = x[indices]
23                 pi[j] = x_j.shape[0]/x.shape[0]
24                 mu[j] = np.mean(x_j, axis=0)
25                 sigma[j] = np.cov(x_j.T) + c*np.eye(d)
26         else:
27             for j in range(k):
28                 indices = (y==j).flatten()
29                 x_j = x[indices]
30                 pi[j] = x_j.shape[0]/x.shape[0]
31                 mu[j] = np.mean(x_j, axis=0)
32                 sigma[j] = np.cov(x_j.T)
33         return mu, sigma, pi
34
35     ## evaluate c using multivariate normal distribution to the training set and
36     validation set
37     def fit(x_train,y_train,x_val,y_val,c):
38         print(f'calculating mu, sigma, pi for c: {c}')
39         mu, sigma, pi = calc_priors_means_covariances(x_train,y_train,c)
40         score = np.zeros((len(x_val),k))
41         print(f'calc score for c: {c}')
42         for label in range(0,k):
43             rv = multivariate_normal(mean=mu[label],
44                 cov=sigma[label],allow_singular=True)
45             score[:,label] = np.log(pi[label]) + rv.logpdf(x_val[:,:])
46             #for i in range(0,len(x_val)):
47                 #score[i,label] = np.log(pi[label]) + rv.logpdf(x_val[i,:])
48         predictions = np.argmax(score, axis=1)
49         errors = np.sum(predictions != y_val)
50         print(f'c: {c}, errors: {errors}')
51         return errors
52
53     ## find best c
54     def find_best_c():
55         ## initialize c values to test and an empty list to store errors
56         c_values = [0,0.0001,0.001, 0.01, 0.1, 1, 10, 100, 1000, 10000]
57         errors = []
58         ## loop through c values and calculate error
59         for c in c_values:
60             print("Calculating error for c: ", c)
61             error = fit(x_train,y_train,x_val,y_val,c)
62             errors.append(error)
63             print("c: ", c, " error: ", error)
64             c_error_dict['c_value'].append(c)
65             c_error_dict['error'].append(error)
66         ## find best c where error is minimum
67         best_c = c_values[np.argmin(errors)]
68
69         return best_c
70
71     ## calculate best c
72     best_c, c_error_dict = find_best_c()
73
74     ## calc mu, sigma, pi using best c
75     mu, sigma, pi = calc_priors_means_covariances(x,y,best_c)
76
77     # Halt and return parameters
78     return mu, sigma, pi

```

### Solution 13 (b)

---

- A single value of  $c = 0.1$  for all ten classes.
- The value of  $c$  was chosen based on the validation set error rate.
  - The model returned 608 errors for the 12000 validation labels from the training labels.




Figure 1: Results from  $c$  value testing

---



**Solution 13 (c)**

---


- The accuracy for  $c = 0.1$  on the test data was 0.9423 or 94.23%.
- The model with  $c = 0.1$  predicted 577 out of 10,000 incorrectly.

$\therefore$  The error rate on the MNIST test set was 0.0577 or 5.77%.


---

**Solution 13 (d)**


---




(a) 1 classified as 8




(b) 9 classified as 7



(c) 9 classified as 8



(d) 9 classified as 8



(e) 2 classified as 7

Figure 2: Examples of misclassified digits from the test set

---