

Fighting the Enemy Within

Basic Life Science and Issues : Presentation

4th Team

November, 2019

Chungnam National University

- **Chae Eun Kim**
College of Medicine, 19'
- **Jongkwan Bae**
Dept of EE Comm. Engineering Education, 17'
- **Seungmin Lee**
College of Medicine, 19'
- **Kangjun Heo**
Dept of Computer Science and Engineering, 17'



Fighting the Enemy Within

11th chapter of *The Epigenetics Revolution*

"Epigenetic perspective of Cancer and its treatment"



Healthy cells, have two types of genes:

- proto-oncogenes for cell proliferation
- tumor suppressor genes for regulation



Introduction: Cancer

Healthy cells, have two types of genes:

- proto-oncogenes for cell proliferation
- tumor suppressor genes for regulation

However, cancer cells lost balance of these, For example,

- proto-oncogenes is over-activated
- tumor suppressor genes is inactivated



Introduction: Cancer

Healthy cells, have two types of genes:

- proto-oncogenes for cell proliferation
- tumor suppressor genes for regulation

However, cancer cells lost balance of these, For example,

- proto-oncogenes is over-activated
- tumor suppressor genes is inactivated



Epigenetic Approach for Oncogenesis

- DNA Methylation

Hypermethylation of CpG island


- Repressive Histone Modification

Histone deacetylation



CpG Hypermethylation

CpG dinucleotide cluster (CpG island, CGI) are usually located in the promoter regions of genes in a DNA sequence.



Dense methylation for CpG island disables specific gene expression.



Histone deacetylation



Approach for Treatment



No easy wins



Alternative Approach



Conclusion



References

Carey, N. (2012). *The Epigenetics Revolution*. Columbia University Press

Kakumani, R., et al. (2012). *Identification of CpG islands in DNA sequences using statistically optimal null filters*, EURASIP Journal on Bioinformatics and Systems Biology



Q & A

Thank you!