Fighting the Enemy Within

Basic Life Science and Issues: Presentation

Group 4

November, 2019

Chungnam National University

Group Members

- Chaeeun Kim
 College of Medicine, 19'
- Jongkwan Bae
 Dept of EE Comm. Engineering Education, 17'
- Seungmin Lee
 College of Medicine, 19'
- Kangjun Heo
 Dept of Computer Science and Engineering, 17'



Chapter Abstraction

Fighting the Enemy Within

11th chapter of The Epigenetics Revolution

"Epigenetic perspective of Cancer and its treatment"



Introduction: Cancer

Healthy cells, have two types of genes:

- proto-oncogenes for cell proliferation
- tumor suppressor genes for regulation



Introduction: Cancer

Healthy cells, have two types of genes:

- proto-oncogenes for cell proliferation
- tumor suppressor genes for regulation

However, cancer cells lost balance of these, For example,

- · proto-oncogenes is over-activated
- tumor suppressor genes is inactivated



Introduction: Cancer

Healthy cells, have two types of genes:

- proto-oncogenes for cell proliferation
- tumor suppressor genes for regulation

However, cancer cells lost balance of these, For example,

- · proto-oncogenes is over-activated
- tumor suppressor genes is inactivated



Epigenetic Approach for Oncogenesis

DNA Methylation
 Hypermethylation of CpG island

Repressive Histone Modification
 Histone deacetylation



DNA Methylation


Cytosine before Guanine can be methylated

Methyl group is bond on 5' carbon atom



DNA Methylation

CpG dinuclotide cluster (CpG island, CGI) are usually located in the promoter regions of genes in a DNA sequence.



Hypermethylated CGI disables specific gene expression.



Histone deacetylation



Characteristics of Oncogenesis

- Multi-step process
- Defections must be accumulated
 Inherited oncogenes are slowly expressed
 e.g.) BRCA1 mutation
- Tumour suppressor gene Switched off
- Alteration with epigenetic access



Approach for Treatment

DNMT enzyme inhibitors
 5-azacytidine, 2-aza-5'-deoxycytidine

methylation inhibited by 5-azacytidine



No easy wins

HDAC inhibitor
 SAHA, Romidepsin



Alternative Approach




Conclusion



References

- [1] Carey, N. (2012). The Epigenetics Revolution. Columbia University Press
- [2] Kakumani, R.; et al. (2012). *Identification of CpG islands in DNA sequences* using statistically optimal null filters, EURASIP Journal on Bioinformatics and Systems Biology
- [3] Kazantsev, Aleksey G; et al. (2008). Therapeutic application of histone deacetylase inhibitors for central nervous system disorders, Nature Reviews. Drug Discovery London Vol. 7 Iss. 10 854-68.





Thank you!