Please check the examination de	etails below before entering	your candidate information	
Candidate surname	Ot	her names	
Pearson Edexcel Level 3 GCE	Centre Number	Candidate Number	
Wednesday 7 October 2020			
Afternoon (Time: 2 hours)	Paper Refe	rence 9MA0/01	
Mathematics Advanced Paper 1: Pure Mathen	natics 1		
You must have: Mathematical Formulae and St	atistical Tables (Greer	Total Marks	

Candidates may use any calculator allowed by Pearson regulations.

Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B).
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided
 - there may be more space than you need.
- You should show sufficient working to make your methods clear.
 Answers without working may not gain full credit.
- Inexact answers should be given to three significant figures unless otherwise stated.

Information

- A booklet 'Mathematical Formulae and Statistical Tables' is provided.
- There are 16 questions in this question paper. The total mark for this paper is 100.
- The marks for each question are shown in brackets
 - use this as a guide as to how much time to spend on each question.

Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.

Turn over ▶







1. (a) Find the first four terms, in ascending powers of x, of the binomial expansion of

$$(1+8x)^{\frac{1}{2}}$$

giving each term in simplest form.

(3)

(2)

(b) Explain how you could use $x = \frac{1}{32}$ in the expansion to find an approximation for $\sqrt{5}$ There is no need to carry out the calculation.

Question 1 continued	
	(Total for Question 1 is 5 marks)
	(



2.	By taking logarithms of both sides, solve the equation					
	$4^{3p-1} = 5^{210}$					
	giving the value of p to one decimal place.	(3)				

Question 2 continued	
	(Total for Question 2 is 3 marks)
	(Total for Question 2 is 3 marks)



- 3. Relative to a fixed origin O
 - point A has position vector $2\mathbf{i} + 5\mathbf{j} 6\mathbf{k}$
 - point B has position vector $3\mathbf{i} 3\mathbf{j} 4\mathbf{k}$
 - point C has position vector $2\mathbf{i} 16\mathbf{j} + 4\mathbf{k}$
 - (a) Find \overrightarrow{AB}

(2)

(b) Show that quadrilateral OABC is a trapezium, giving reasons for your answer.

(2)

Question 3 continued
(Total for Question 3 is 4 marks)



4. The function f is defined by

$$f(x) = \frac{3x - 7}{x - 2} \qquad x \in \mathbb{R}, x \neq 2$$

(a) Find $f^{-1}(7)$

(2)

(b) Show that $ff(x) = \frac{ax + b}{x - 3}$ where a and b are integers to be found.

(3)

1						
1						
1						
1	_		_	_		
1						
1						
1						
1						
-	_	_	_	_	_	_

Question 4 continued
(Total for Question 4 is 5 marks)
(======================================



5.	A car has six forward gears.	
	The fastest speed of the car	
	• in 1 st gear is 28 km h ⁻¹	
	• in 6 th gear is 115 km h ⁻¹	
	Given that the fastest speed of the car in successive gears is modelled by an arithmetic sequence ,	
	(a) find the fastest speed of the car in 3 rd gear.	
		(3)
	Given that the fastest speed of the car in successive gears is modelled by a geometric sequence ,	
	(b) find the fastest speed of the car in 5 th gear.	
		(3)



Question 5 continued
(Total for Question 5 is 6 marks)
(20m 101 Question e is o mains)



6. (a) Express $\sin x + 2\cos x$ in the form $R\sin(x + \alpha)$ where R and α are constants, R > 0 and $0 < \alpha < \frac{\pi}{2}$

Give the exact value of R and give the value of α in radians to 3 decimal places. (3)

The temperature, θ °C, inside a room on a given day is modelled by the equation

$$\theta = 5 + \sin\left(\frac{\pi t}{12} - 3\right) + 2\cos\left(\frac{\pi t}{12} - 3\right) \qquad 0 \leqslant t < 24$$

where t is the number of hours after midnight.

Using the equation of the model and your answer to part (a),

(b) deduce the maximum temperature of the room during this day,

(1)

(c) find the time of day when the maximum temperature occurs, giving your answer to the nearest minute.

(3)

Question 6 continued



Question 6 continued

Question 6 continued
(Total for Organian 6 is 7 marks)
(Total for Question 6 is 7 marks)



7.




Figure 1

Figure 1 shows a sketch of a curve C with equation y = f(x) and a straight line l.

The curve C meets l at the points (-2,13) and (0,25) as shown.

The shaded region R is bounded by C and l as shown in Figure 1.

Given that

- f(x) is a quadratic function in x
- (-2, 13) is the minimum turning point of y = f(x)

use inequalities to define R.

•	(5)

$\times\!\!\times\!\!\times\!\!\times$
$\times\!\!\times\!\!\times\!\!\times$
$\times\!\!\times\!\!\times\!\!\times$
$\times\!\!\times\!\!\times\!\!\times$
$\times\!\!\times\!\!\times\!\!\times$
$\times\!\!\times\!\!\times\!\!\times$
\times
AREA
REA
\times MM \times
\times
THIS A
\times
\times
\times
Z
\times
\times
XX XX
WRITE
\times
\times
DO NO
\times
XXXX
$\times\!\!\times\!\!\times\!\!\times$
\times
$\times\!\!\times\!\!\times\!\!\times$
$\times\!\!\times\!\!\times\!\!\times$
\times
$\times\!\!\times\!\!\times\!\!\times$
\times
$\times\!\!\times\!\!\times\!\!\times$
$\times \times \times \times$
XXXXX
XX75X
養
夏
E Z
NN TH
E IN TH
TE IN TH
NTE IN TH
RITE IN TH
WRITE IN TH
WRITEINTH
F WRITE IN TH
DT WRITE IN TH
OT WRITE IN TH
NOT WRITE IN TH
NOT WRITE IN THIS
O NOT WRITE IN TH
SO NOT WRITE IN TH
DO
A DO
EA DO
EA DO
REA DO
AREA DO
AREA DO
IS AREA DO
IIS AREA DO
IIS AREA DO
IIS AREA DO
IS AREA DO
N THIS AREA DO
IN THIS AREA DO
E IN THIS AREA DO
TE IN THIS AREA DO
TE IN THIS AREA DO
TE IN THIS AREA DO
TE IN THIS AREA DO
TE IN THIS AREA DO
WRITE IN THIS AREA DO
TE IN THIS AREA DO
OT WRITE IN THIS AREA DO
WRITE IN THIS AREA DO

Question / continued	



Question 7 continued	

Question 7 continued	
(Tota	l for Question 7 is 5 marks)



8.	A new smartphone was released by a company.	
	The company monitored the total number of phones sold, n , at time t days after the phone was released.	
	The company observed that, during this time,	
	the rate of increase of n was proportional to n	
	Use this information to write down a suitable equation for n in terms of t .	
	(You do not need to evaluate any unknown constants in your equation.)	(2)



Question 8 continued
(Total for Question 8 is 2 marks)



9.




Figure 2

Figure 2 shows a sketch of the curve C with equation y = f(x) where

$$f(x) = 4(x^2 - 2)e^{-2x}$$
 $x \in \mathbb{R}$

(a) Show that $f'(x) = 8(2 + x - x^2)e^{-2x}$

(3)

(b) Hence find, in simplest form, the exact coordinates of the stationary points of C.

(3)

The function g and the function h are defined by

$$g(x) = 2f(x)$$
 $x \in \mathbb{R}$

$$h(x) = 2f(x) - 3 \qquad x \geqslant 0$$

(c) Find (i) the range of g

(ii) the range of h

(3)

Question 9 continued



Question 9 continued

Question 9 continued
(Total for Question 9 is 9 marks)
(20m 101 Yucomon > 10 > mar ha)



10. (a) Use the substitution $x = u^2 + 1$ to show that

$$\int_{5}^{10} \frac{3 \, dx}{(x-1)(3+2\sqrt{x-1})} = \int_{p}^{q} \frac{6 \, du}{u(3+2u)}$$

where p and q are positive constants to be found.

(4)

(b) Hence, using algebraic integration, show that

$$\int_{5}^{10} \frac{3 \, dx}{(x-1)(3+2\sqrt{x-1})} = \ln a$$

where a is a rational constant to be found.

(6)

Question 10 continued



Question 10 continued	

Question 10 continued	
(Tota	al for Question 10 is 10 marks)
(100	20 20 20 20 20 20 20 20 20 20 20 20 20 2



11.




Figure 3

Circle C_1 has equation $x^2 + y^2 = 100$

Circle C_2 has equation $(x-15)^2 + y^2 = 40$

The circles meet at points A and B as shown in Figure 3.

(a) Show that angle AOB = 0.635 radians to 3 significant figures, where O is the origin.

(4)

The region shown shaded in Figure 3 is bounded by C_1 and C_2

(b) Find the perimeter of the shaded region, giving your answer to one decimal place.



CXXXXX	
****	1
	ı
*****	ı
	ı
$\sim\sim\sim\sim$	ı
******* I	
*****	ı
XXXXXXX	ı
NAME OF THE PERSON OF THE PERS	ı
M.	ı
	ı
XXXXXX	ı
*	ı
	ı
	ı
	ı
	ı
XX 44	ı
	ı
	ı
S	ı
	ı
	ı
0	Г
ă	Г
	Г
	Г
	Г
	ı
\times	ı
(XXXXXX)	ı
******* I	ı
	ı
	ı
	ı
	ı
	ı
XXXXXX	ı
SS	ı
	ı
	ı
~~ ? ~~~ I	ı
2	ı
	ı
	ı
	ı
	ı
	ı
5	ı
	ı
	ı
X2X	ı
	ı
******	ı
XXXX	ı
	ı
****** I	ĺ
XXXXX	ı
	ĺ
****** I	ı
****** I	ı
	ı
XXXX	ı
*****	ı
	ı
<u></u>	ı
	ı
XXXXXX	ĺ
∞	
	ĺ
₩	ĺ
XXX	ĺ
	ĺ
	ĺ
	ĺ
	ĺ
	ĺ
×9×	ĺ
	l
×9×	ĺ
	ı
~~~~~ I	1



Question 11 continued

Question 11 continued	
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
(Total for Question 11 is 8 marks)	_



12	. In this question you must show all stages of your working.	
	Solutions relying entirely on calculator technology are not acceptable.	
	(a) Show that	
	$\csc \theta - \sin \theta \equiv \cos \theta \cot \theta$ $\theta \neq (180n)^{\circ}$ $n \in \mathbb{Z}$	(3)
	(b) Hence, or otherwise, solve for $0 < x < 180^{\circ}$	,
	$\csc x - \sin x = \cos x \cot (3x - 50^{\circ})$	(5)



		1			
				ĺ	
				į	
	i				
					1
				ŀ	
				ì	
				ú	
					ij
				5	7
	d			5	3
	í	ì		5	
	ĺ	ĺ	2	2	3
⟨		ŀ	>	5	3
>		ŀ	2	2	
>		ŀ		>	
>		ŀ		>	
?		ŀ	3		
>					
>					
>					
>					
>					
>					
>					
>					
>>>>>					
>					
>>>>>>>>>>>>>>>>>>>>>>>>>>>>>>>>>>>>>>>					
\ \ \ \					
> > > >					
> > >					
> > >					
> > > >					
> > > >					

Question 12 continued

Question 12 continued	

Question 12 continued	
	_
	-
	-
	-
	-
	-
	-
	_
	-
	-
	_
	-
	-
	-
(Total for Question 12 is 8 marks)	
(2000 TOT QUODUON 12 NO MARKED)	



13. A sequence of numbers  $a_1, a_2, a_3, \dots$  is defined by

$$a_{n+1} = \frac{k(a_n + 2)}{a_n} \qquad n \in \mathbb{N}$$

where k is a constant.

Given that

- the sequence is a periodic sequence of order 3
- $a_1 = 2$
- (a) show that

$$k^2 + k - 2 = 0$$

**(3)** 

(b) For this sequence explain why  $k \neq 1$ 

**(1)** 

(c) Find the value of

$$\sum_{r=1}^{80} a_r$$

**(3)** 

					ğ	
		ġ				
		į				
		ď				
					2	
					2	
					Ì	
	K		Ì		1	
					ģ	
	А					
′	SI					
Ś	Į	9	7	7	7	
5		ì	S	Z	S	
<		į	Ē	3	S	
>	2			Ž	Ş	
{	K			2	5	
5						
>						
< > >						
< >		í				
		í				
< > >		í				
< > >		í				
< >>		í				
<>>>		í				
< > >		í				
<>>>		ŀ	>			
<		ŀ	>			
<		ŀ	>			
		ŀ	>			
		ŀ	>			
		ŀ	>			
		ŀ	>			
		í				
		ŀ				
		ŀ				
		ŀ				
		ŀ				
		ŀ				
		ŀ				

<b>Question 13 continued</b>		

Question 13 continued

Question 13 continued	
	(Total for Question 13 is 7 marks)



14. A large spherical balloon is deflating.

At time t seconds the balloon has radius rcm and volume Vcm³

The volume of the balloon is modelled as decreasing at a constant rate.

(a) Using this model, show that

$$\frac{\mathrm{d}r}{\mathrm{d}t} = -\frac{k}{r^2}$$

where k is a positive constant.

(3)

Given that

- the initial radius of the balloon is 40 cm
- after 5 seconds the radius of the balloon is 20 cm
- the volume of the balloon continues to decrease at a constant rate until the balloon is empty
- (b) solve the differential equation to find a complete equation linking r and t.

**(5)** 

(c) Find the limitation on the values of t for which the equation in part (b) is valid. (2)

Question 14 continued



Question 14 continued	

Question 14 continued	
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
(Total for Question 14 is 10 marks)	_



**15.** The curve C has equation

$$x^2 \tan y = 9 \qquad 0 < y < \frac{\pi}{2}$$

(a) Show that

$$\frac{\mathrm{d}y}{\mathrm{d}x} = \frac{-18x}{x^4 + 81}$$

**(4)** 

(b) Prove that C has a point of inflection at  $x = \sqrt[4]{27}$ 

12	١
w	J





$\times\!\!\times\!\!\times$
$\times\!\!\times\!\!\times\!\!\times$
$\times\!\!\times\!\!\times\!\!\times$
$\times\!\!\times\!\!\times\!\!\times$
*****
XXXX
>>>>>
$\times\!\!\times\!\!\times$
$\times\!\!\times\!\!\times\!\!\times$
X III
AREA
$\times$
INTHIS
$\times \times \times \times$
***
$\times \mapsto$
WRITE
$\times$
$\times$
$\times \bigcirc \times$
×2×
$\times\!\!\times\!\!\times\!\!\times$
DO NOT
$\times \triangle$
$\times\!\!\times\!\!\times$
$\times\!\!\times\!\!\times\!\!\times$
****
>>>>
****
****
>>>>>
$\times\!\!\times\!\!\times$
$\times\!\!\times\!\!\times$
$\times\!\!\times\!\!\times\!\!\times$
$\times\!\!\times\!\!\times$
$\times\!\!\times\!\!\times\!\!\times$
$\times\!\!\times\!\!\times\!\!\times$
◁
Ä
REA
AREA
S AREA
IIS AREA
HIS AREA
THIS AREA
THIS AREA
IN THIS AREA
IN THIS AREA
E IN THIS AREA
TE IN THIS AREA
RITE IN THIS AREA
VRITE IN THIS AREA
WRITE IN THIS
NOT WRITE IN THIS
NOT WRITE IN THIS
WRITE IN THIS
NOT WRITE IN THIS
NOT WRITE IN THIS
NOT WRITE IN THIS
NOT WRITE IN THIS
NOT WRITE IN THIS
NOT WRITE IN THIS
NOT WRITE IN THIS
NOT WRITE IN THIS
NOT WRITE IN THIS
NOT WRITE IN THIS
NOT WRITE IN THIS
NOT WRITE IN THIS
NOT WRITE IN THIS
NOT WRITE IN THIS
NOT WRITE IN THIS
DO NOT WRITE IN THIS
A DO NOT WRITE IN THIS
A DO NOT WRITE IN THIS
A DO NOT WRITE IN THIS
A DO NOT WRITE IN THIS
A DO NOT WRITE IN THIS
DO NOT WRITE IN THIS

		Š		
	į			
N		Я	۲	
	È			
N	ø	Ę	ú	h
2	5	ž	Š	ž
3	S	ξ	Š	2
3	5	7	3	2
3				
3				
3				
3	Ė	ì	2	ì
3	Ė	ì	2	ì
3	Ė	ì	2	ì
3	Ė	ì	2	ì
3	Ė	ì	2	ì
3	Ė	ì	2	ì
3	Ė	ì	2	ì
3	Ė	ì	2	ì
	Ė	ì	2	ì
	Ė	ì	2	ì
	Ė	ì	2	ì
	Ė	ì	2	ì
	Ė	ì	2	ì
	Ė	ì	2	ì
	Ė	ì		ì
	Ė	ì	2	ì
				ì
				ì
				ì
				ì
		ì		ì
				ì
				ì
				ì
				ì
				ì
				ì
				ì
				ì
				ì
				ì
				ì
				ì
				ì
				ì
				ì
				ì

Question 15 continued

Question 15 continued

Question 15 continued	
	(Total for Question 15 is 7 marks)



<b>16.</b> Prove by contradiction that there are no positive integers $p$ and $q$ such that				
		$4p^2 - q^2 = 25$		
		ip q 25		(4)

Question 16 continued



(Total for Question 16 is 4 marks)  TOTAL FOR PAPER IS 100 MARKS	estion 16 continued	
		(Total for Question 16 is 4 marks)
TOTAL FOR PAPER IS 100 MARKS		
		TOTAL FOR PAPER IS 100 MARKS