

Forward Finite Difference Series

Definition

Corban Amouzou

Abstract. We prove the equivalence of a series definition of the finite difference operation to its standard. The implication of approximation to the derivative is more explicit, and as a result, direct claims made of approximation-related properties of the finite difference operation can be proven with greater ease.

1 Theorem and Proof

Theorem 1:

Given some $x \in \mathbb{R}$ and $h \in \mathbb{R}$, let $f : \mathbb{R} \rightarrow \mathbb{R}$ be a continuous polynomial function on the interval $[x, x + h]$, $h \in \mathbb{R}$. The following relationship then holds:

$$\Delta_h[f](x) = \sum_{k=1}^{\deg f} \frac{h^k}{k!} f^{(k)}(x), \forall x \in \mathbb{R} \quad (1.1)$$

Proof. By induction on the degree of the input function f , we'll show [Theorem 1.1](#) to be true.

The base case is when f is a constant function (ie: has a degree of 0). [Theorem 1.1](#) is trivial for this case as the difference between terms of a constant is 0, and likewise is the derivative of a constant. To support the induction, we'll first assume that [Theorem 1.1](#) is true for some $n \in \mathbb{N}$. Let's first redefine $f(x)$ to have a degree of $m = n + 1$. Using the Taylor series expansion of a general polynomial function and the definition of the finite Forward Difference Operation (f.d.o), we can express the f.d.o of $f(x)$ as the following:

$$\Delta_h[f](x) = \sum_{k=0}^m \left(C_k (x + h)^k \right) - \sum_{k=0}^m (C_k x^k) \quad (1.2)$$

Using the binomial theorem, we have this expansion of $f(x+h)$:

$$f(x+h) = \sum_{k=0}^m \left(C_k \sum_{j=0}^k \binom{k}{j} x^{k-j} h^j \right) \quad (1.3)$$

To simplify the **f.d.o** expression, we'll need to extract $f(x)$ from the summation. This is done simply by incrementing j by 1 in $f(x+h)$ and writing in the first term explicitly:

$$f(x+h) = \sum_{k=0}^m \left(C_k \left[x^k + \sum_{j=1}^k \binom{k}{j} x^{k-j} h^j \right] \right) \quad (1.4)$$

From here we split the summation into two parts:

$$f(x+h) = \sum_{k=0}^m (C_k x^k) + \sum_{k=0}^m \left(C_k \sum_{j=1}^k \binom{k}{j} x^{k-j} h^j \right) \quad (1.5)$$

This allows us to cancel out $f(x)$ in the **f.d.o** giving us the following equality:

$$\Delta_h[f](x) = \sum_{k=0}^m \left(C_k \sum_{j=1}^k \binom{k}{j} x^{k-j} h^j \right) \quad (1.6)$$

Noticing now that the inner summation is empty for $k=0$, we can increment k by 1:

$$\Delta_h[f](x) = \sum_{k=1}^m \left(C_k \sum_{j=1}^k \binom{k}{j} x^{k-j} h^j \right) \quad (1.7)$$

Once again using the binomial theorem we expand the inner summation and move the coefficient into the summation. Further, we can move $\frac{1}{j!}$ into the h^j term:

$$\Delta_h[f](x) = \sum_{k=1}^m \left(\sum_{j=1}^k \left(\frac{h^j}{j!} \right) C_k \left(\frac{k!}{(k-j)!} \right) x^{k-j} \right) \quad (1.8)$$

We'll now need to use the concept of expressing a polynomial function as a sum of its ordered parts to reconcile the inner summation with a derivative over the entire function. To do this, we'll first need to define the ordered parts of a polynomial function. Then we'll need to define the n 'th derivative of a polynomial function. We'll use these definitions to simplify the inner summation.

Definition 1. Let f_n be the notation for the n 'th degree part of a polynomial function $f : \mathbb{R} \rightarrow \mathbb{R}$ of degree m . Simply put:

$$f_n(x) = C_n x^n \quad (1.9)$$

The sum of these parts is equivalent to the original function:

$$f(x) = \sum_{k=0}^m f_k(x) \quad (1.10)$$

Therefore the derivative of these parts is equivalent to the derivative of the original function:

$$f^{(n)}(x) = \sum_{k=0}^m f_k^{(n)}(x) \quad (1.11)$$

Definition 2. The n th derivative of some function $f : \mathbb{R} \rightarrow \mathbb{R}$ of degree m can be expressed as the following:

$$f^{(n)}(x) = \sum_{k=n}^m \frac{k!}{(k-n)!} C_k x^{k-n} \quad (1.12)$$

Returning to the proof, we notice that the inner summation is equivalent to the j 'th derivative of the k 'th degree part of f , so we can use [Definition 1](#) to simplify the expression:

$$\Delta_h[f](x) = \sum_{k=1}^m \left(\sum_{j=1}^k \left(\frac{h^j}{j!} \right) f_k^{(j)}(x) \right) \quad (1.13)$$

Finally, we can use [Definition 1 \(1.11\)](#) to conclude that the inner summation is equivalent to the k 'th derivative of f multiplied by $\frac{h^k}{k!}$:

$$\Delta_h[f](x) = \sum_{k=1}^m \left(\frac{h^k}{k!} f^{(k)}(x) \right) \quad (1.14)$$

[Equation 1.14](#) is equivalent to our inductive hypothesis, so we can conclude that [Theorem 1.1](#) is true for all $n \in \mathbb{N}$. \square