




## Relations & Functions

Part 4




## Binary Relations

How Stuff Compares to Stuff

### Relations

- A *binary relation* is a stated fact between on two objects
- "fact" is called a *predicate*
- Evaluates to true or false
- These are the foundation of most programming tasks



4/6/2017

Sacramento State - Cook - CSc 28 - Spring 2017

3

### Example Relations

- "x is taller than y"
- "x lives less than 50 miles from y"
- " $x \leq y$ "
- "x and y are siblings"
- "x has a y"

4/6/2017

Sacramento State - Cook - CSc 28 - Spring 2017

4

### Relations


- A *binary relation* from  $A$  to  $B$  is a subset of  $A \times B$
- So, a relation from  $A$  to  $B$  is a set of ordered pairs  $(a, b)$  where  $a \in A$  and  $b \in B$
- We can use the shorthand notation of  $a R b$  to denote that  $(a, b) \in R$

4/6/2017

Sacramento State - Cook - CSc 28 - Spring 2017

5

### Relationship Chart




|   | a | b |
|---|---|---|
| 1 | ✓ | ✓ |
| 2 | ✓ |   |
| 3 |   | ✓ |

4/6/2017

Sacramento State - Cook - CSc 28 - Spring 2017

6

## Relationship Chart



4/6/2017

Sacramento State - Cook - CSc 28 - Spring 2017

7

## Example: Capitols

- $A$  is a set of all cities in the World
- $B$  is a set of all states in the World
- The relation  $a R b$  specifies that  $a$  is the capitol of  $b$

4/6/2017

Sacramento State - Cook - CSc 28 - Spring 2017

8

## Example: Capitol Members

- (London, Britain)
- (Sacramento, California)
- (Madrid, Spain)
- (Tokyo, Japan)
- (New Delhi, India)
- (Albany, New York)


4/6/2017

Sacramento State - Cook - CSc 28 - Spring 2017

9

## Let's Draw One!

- Let's draw a relationship graph
- It will be between students and stuff they have



4/6/2017

Sacramento State - Cook - CSc 28 - Spring 2017

10

## Relation Domain

- The *domain* of a relation is a set of all the first elements of each tuple
- So, it is the elements that the make up the left-hand side of  $a R b$

$\{ a \mid (a, b) \in R \text{ for some } b \}$

4/6/2017

Sacramento State - Cook - CSc 28 - Spring 2017

11

## Relation Range

- The *range* of a relation is a set of all second elements from each tuple
- So, it is the elements that the make up the right-hand side of  $a R b$

$\{ b \mid (a, b) \in R \text{ for some } a \}$

4/6/2017

Sacramento State - Cook - CSc 28 - Spring 2017

12

## Example

$$R = \{ (1,1), (1,2), (1,3), (1,4), (2,4), (4,4) \}$$

Domain of  $R = \{ 1, 2, 4 \}$

Range of  $R = \{ 1, 2, 3, 4 \}$

4/6/2017

Sacramento State - Cook - CSc 28 - Spring 2017

13

## Inverse Relation

- The *inverse* of a relation swaps first and last element for each tuple
- The number of elements are the same, but the range and domain are reversed

$$R^{-1} = \{ (b,a) \mid (a,b) \in R \}$$

4/6/2017

Sacramento State - Cook - CSc 28 - Spring 2017

14

## Inverse Example


$$R = \{ (1,1), (1,2), (1,3), (1,4), (2,4), (4,4) \}$$

$$R^{-1} = \{ (1,1), (2,1), (3,1), (4,1), (4,2), (4,4) \}$$

4/6/2017

Sacramento State - Cook - CSc 28 - Spring 2017


15



How a Set Sees Itself

## "Relation On"

- Some relations of a set  $A$  are on itself
- In other words, each object in the related to the same "type" of object
- This is called a *relation on  $A$*
- ...and it is important to examine its properties




4/6/2017

Sacramento State - Cook - CSc 28 - Spring 2017

17

## Example Relationship Chart



4/6/2017

Sacramento State - Cook - CSc 28 - Spring 2017

18

## Example Relation Chart


- The previous chart represents when  $a$  divides  $b$
- In other words,  $a$  times some integer equals the value  $b$
- So,  $R = \{ (1, 1), (1, 2), (1, 3), (1, 4), (2, 2), (2, 4), (3, 3), (4, 4) \}$

4/6/2017

Sacramento State - Cook - CSc 28 - Spring 2017

19

## Reflexive Relations



- A *reflexive* relationship means that there is  $aRa$  for every  $a$
- Basically, everything has to be related to itself

4/6/2017

Sacramento State - Cook - CSc 28 - Spring 2017

20

## To Determine Reflexive...

- Look for some  $a \in A$  where there isn't a  $aRa$
- If found, not reflexive
- Otherwise reflexive



4/6/2017

Sacramento State - Cook - CSc 28 - Spring 2017

21

## Reflexive Example

Relation on set  $\{1, 2, 3, 4\}$

$R = \{ (1,1), (1,4), (2,2), (2,3), (3,3), (4,4) \}$

4/6/2017

Sacramento State - Cook - CSc 28 - Spring 2017

22

## Reflexive Example

Relation on set  $\{1, 2, 3, 4\}$


$R = \{ (1,1), (1,4), (2,2), (2,3), (3,3), (4,4) \}$

4/6/2017

Sacramento State - Cook - CSc 28 - Spring 2017

23

## Reflexive Example



|   | 1 | 2 | 3 | 4 |
|---|---|---|---|---|
| 1 | ✓ |   |   | ✓ |
| 2 |   | ✓ | ✓ |   |
| 3 |   |   | ✓ |   |
| 4 |   |   |   | ✓ |

4/6/2017

Sacramento State - Cook - CSc 28 - Spring 2017

24

## Nonreflexive Example

Relation on set  $\{1, 2, 3, 4\}$


$R = \{ (1,1), (1,4), (2,2), (2,3), (3,1), (4,4) \}$

4/6/2017

Sacramento State - Cook - CSc 28 - Spring 2017

25

## Nonreflexive Example




|   | 1 | 2 | 3 | 4 |
|---|---|---|---|---|
| 1 | ✓ | ✓ |   |   |
| 2 |   | ✓ | ✓ |   |
| 3 |   |   |   | ✓ |
| 4 |   |   |   | ✓ |

4/6/2017

Sacramento State - Cook - CSc 28 - Spring 2017

26

## Symmetric Relations




- A *symmetric* relationship means that for every  $aRb$  there is a  $bRa$
- So, if  $(a, b)$  exists in the relation, so must  $(b, a)$

4/6/2017

Sacramento State - Cook - CSc 28 - Spring 2017

27

## Symmetric Relations



- Note: Unlike the definition for reflexive, not every element in the domain needs to exist
- If a relation is not symmetric, it is *antisymmetric*

4/6/2017

Sacramento State - Cook - CSc 28 - Spring 2017

28

## To Determine Symmetric

- Look for an  $a$  and  $b$  where there is a  $aRb$  but no  $bRa$
- If found, nonsymmetric
- Otherwise symmetric



4/6/2017

Sacramento State - Cook - CSc 28 - Spring 2017

29

## Symmetric Example

Relation on set  $\{1, 2, 3, 4\}$

$R = \{ (1,1), (1,2), (2,1), (2,4), (3,3), (4,2) \}$

4/6/2017

Sacramento State - Cook - CSc 28 - Spring 2017

30

## Symmetric Example

Relation on set  $\{1, 2, 3, 4\}$


$R = \{ (1,1), (1,2), (2,1), (2,4), (3,3), (4,2) \}$

4/6/2017

Sacramento State - Cook - CSc 28 - Spring 2017

31

## Symmetric Example



|   | 1 | 2 | 3 | 4 |
|---|---|---|---|---|
| 1 | ✓ | ✓ |   |   |
| 2 | ✓ |   |   | ✓ |
| 3 |   |   | ✓ |   |
| 4 |   | ✓ |   |   |

4/6/2017

Sacramento State - Cook - CSc 28 - Spring 2017

32

## Nonsymmetric Example

Relation on set  $\{1, 2, 3, 4\}$


$R = \{ (1,1), (1,4), (2,2), (2,3), (3,3), (4,4) \}$

4/6/2017

Sacramento State - Cook - CSc 28 - Spring 2017

33

## Nonsymmetric Example



|   | 1 | 2 | 3 | 4 |
|---|---|---|---|---|
| 1 | ✓ |   |   | ✓ |
| 2 |   | ✓ | ✓ |   |
| 3 |   |   | ✓ |   |
| 4 |   |   |   | ✓ |

4/6/2017

Sacramento State - Cook - CSc 28 - Spring 2017

34

## To Determine Transitive

- Look for an  $a, b, c$  where there is a  $aRb$  and  $bRc$  but no  $aRc$
- If found, non transitive
- Otherwise transitive



4/6/2017

Sacramento State - Cook - CSc 28 - Spring 2017

35

## Transitive Example

Relation on set  $\{1, 2, 3, 4\}$

$R = \{ (2,1), (3,1), (3,2), (4,1), (4,2), (4,3) \}$

4/6/2017

Sacramento State - Cook - CSc 28 - Spring 2017

36

## Transitive Example

Relation on set  $\{1, 2, 3, 4\}$

$R = \{ (2,1), (3,1), (3,2), (4,1), (4,2), (4,3) \}$

Starting with (4,3)

4/6/2017

Sacramento State - Cook - CSc 28 - Spring 2017

37

## Transitive Example

Relation on set  $\{1, 2, 3, 4\}$

Starting with (3,2)

$R = \{ (2,1), (3,1), (3,2), (4,1), (4,2), (4,3) \}$

4/6/2017

Sacramento State - Cook - CSc 28 - Spring 2017

38

## Transitive Example

Relation on set  $\{1, 2, 3, 4\}$

$R = \{ (2,1), (3,1), (3,2), (4,1), (4,2), (4,3) \}$


Starting (again) with (4,3)

4/6/2017

Sacramento State - Cook - CSc 28 - Spring 2017

39

## Transitive Example



|   | 1 | 2 | 3 | 4 |
|---|---|---|---|---|
| 1 |   |   |   |   |
| 2 | ✓ |   |   |   |
| 3 | ✓ | ✓ |   |   |
| 4 | ✓ | ✓ | ✓ |   |

4/6/2017

Sacramento State - Cook - CSc 28 - Spring 2017

40

## Equivalence Relations

- If a relation is:
  - reflexive
  - symmetric
  - transitive
- It is an equivalence



4/6/2017

Sacramento State - Cook - CSc 28 - Spring 2017

41

## Example Table

| Relation            | Reflexive | Symmetric | Transitive | Equivalent |
|---------------------|-----------|-----------|------------|------------|
| $\leq$              |           |           |            |            |
| $\subset$           |           |           |            |            |
| Perpendicular lines |           |           |            |            |
| Parallel lines      |           |           |            |            |

4/6/2017

Sacramento State - Cook - CSc 28 - Spring 2017

42

## Example Table

| Relation            | Reflexive | Symmetric | Transitive | Equivalent |
|---------------------|-----------|-----------|------------|------------|
| $\leq$              | ✓         | ✗         | ✓          | ✗          |
| $\subset$           | ✗         | ✗         | ✓          | ✗          |
| Perpendicular lines | ✗         | ✓         | ✗          | ✗          |
| Parallel lines      | ✓         | ✓         | ✓          | ✓          |

4/6/2017

Sacramento State - Cook - CSc 28 - Spring 2017

43



## Manipulating Relations

How a Set Sees Itself

## Manipulating Relations

- Because relations are representable as sets, we can use set notation to define them
- We can also use set notation to manipulate them



4/6/2017

Sacramento State - Cook - CSc 28 - Spring 2017

45

## Example

$$\mathbf{A} = \{ (1,1), (2,2), (3,3) \}$$

$$\mathbf{B} = \{ (1,1), (2,4), (3,9) \}$$

4/6/2017

Sacramento State - Cook - CSc 28 - Spring 2017

46

## Example

$$\mathbf{A} \cup \mathbf{B} = \{ (1,1), (2,2), (2,4), (3,3), (3,9) \}$$

$$\mathbf{A} \cap \mathbf{B} = \{ (1,1) \}$$

$$\mathbf{A} - \mathbf{B} = \{ (2,2), (3,3) \}$$

$$\mathbf{B} - \mathbf{A} = \{ (2,4), (3,9) \}$$

4/6/2017

Sacramento State - Cook - CSc 28 - Spring 2017

47

## Finite Sets

- On a finite set, relations are quite simple...
- For a set with  $n$  elements, the maximum number of relations is simply  $n \times n = n^2$
- However, many relations are defined over an infinite set – e.g. all integers, reals, etc...

4/6/2017

Sacramento State - Cook - CSc 28 - Spring 2017

48



## Representing Relations

- We can represent a relation using set notation
- However, some are required to be denoted using set builder notation

```
R1 = { (a, b) | a is taller than b }
R2 = { (a, b) | a ≤ b }
```


4/6/2017

Sacramento State - Cook - CSc 28 - Spring 2017

49

## Let's Examine Some...

- Let's use students to create two relations
- Classes you plan to take
- Classes you enjoy



4/6/2017

Sacramento State - Cook - CSc 28 - Spring 2017

50

## Let's Examine Some...

- Let's examine two relations over a simple set  $A$  of  $\{1, 2, 3\}$  using set operators
- We can check if it is:
  - reflexive
  - symmetric
  - transitive



4/6/2017

Sacramento State - Cook - CSc 28 - Spring 2017

51

## Let's Examine Some...

$A = \{1, 2, 3\}$ :  $R, S$  relations.

$R = \{ (1, 1), (1, 2), (2, 2), (2, 3), (3, 1), (3, 3) \}$

$S = \{ (1, 1), (1, 2), (1, 3), (2, 1), (2, 3), (3, 2) \}$

4/6/2017

Sacramento State - Cook - CSc 28 - Spring 2017

52



## Closures

Making a relation "complete"

## Closure

- **Closure** of relation  $R$  is the **smallest** set (when unioned) gives  $R$  the desired property
- So, the closure of  $R$  is  $R \cup C$ , where  $C$  is the smallest set giving  $R \cup C$  the desired property



4/6/2017

Sacramento State - Cook - CSc 28 - Spring 2017

54

## Some Examples

- For the following examples, the relation is over the set  $\{1, 2, 3, 4\}$
- The slides will show how to make the closures for reflexive, symmetric, and (the hard one) transitive

4/6/2017

Sacramento State - Cook - CSc 28 - Spring 2017

55

## Example Reflexive Closure

$R = \{ (1,2), (2,3), (3,4) \}$

$C = \{ (1,1), (2,2), (3,3), (4,4) \}$

Missing (1,1) (2,2), (3,3) and (4,4)

4/6/2017

Sacramento State - Cook - CSc 28 - Spring 2017

56

## Example Reflexive Closure

$R \cup C = \{ (1,2), (2,3), (3,4), (1,1), (2,2), (3,3), (4,4) \}$

4/6/2017

Sacramento State - Cook - CSc 28 - Spring 2017

57

## Example Symmetric Closure

$R = \{ (1,2), (2,3), (3,4) \}$

$C = \{ (2,1), (3,2), (4,3) \}$

4/6/2017

Sacramento State - Cook - CSc 28 - Spring 2017

58

## Example Symmetric Closure

$R \cup C = \{ (1,2), (2,3), (3,4), (2,1), (3,2), (4,3) \}$

4/6/2017

Sacramento State - Cook - CSc 28 - Spring 2017

59

## Example Transitive Closure (1 of 3)

$R = \{ (1,2), (2,3), (3,4) \}$

$C = \{ (1,3) \}$

Added due to (1,2) and (2,3)

4/6/2017

Sacramento State - Cook - CSc 28 - Spring 2017

60

## Example Transitive Closure (2 of 3)

$$R = \{ (1, 2), (2, 3), (3, 4) \}$$

$$C = \{ (1, 3), (2, 4) \}$$

Added due to (2,3) and (3,4)

4/6/2017

Sacramento State - Cook - CSc 28 - Spring 2017

61

## Example Transitive Closure (3 of 3)

$$R = \{ (1, 2), (2, 3), (3, 4) \}$$

$$C = \{ (1, 3), (2, 4), (1, 4) \}$$

Had to add after we added (2,4) since R contains (1,2)

4/6/2017

Sacramento State - Cook - CSc 28 - Spring 2017

62

## Example Transitive Closure

$$R \cup C = \{ (1, 2), (2, 3), (3, 4), (1, 3), (2, 4), (1, 4) \}$$

4/6/2017

Sacramento State - Cook - CSc 28 - Spring 2017

63

## Let's Try Some


- Set is  $\{1, 2, 3, 4, 5\}$
- $R = \{ (1, 2), (2, 3), (3, 4) \}$
- What is the closures for this relations?



4/6/2017

Sacramento State - Cook - CSc 28 - Spring 2017

64




## Functions

Math Friendly Relations

## Functions

- We have all seen functions – which take inputs and produce output
- Example:  $f(x) = x^2$ 
  - $f(1) = 1$
  - $f(2) = 4$
  - $f(3) = 9 \dots$




4/6/2017

Sacramento State - Cook - CSc 28 - Spring 2017

66

## Functions

- Sets give a way to document "types" in mathematical functions
- A *function* from set  $X$  to set  $Y$  is a mapping from each element in  $X$  to elements in  $Y$



4/6/2017

Sacramento State - Cook - CSc 28 - Spring 2017

67

## Functions

- We will restrict a functions inputs and outputs by giving a "signature" for it
- $f$  is the function name

$f: \mathbb{N} \rightarrow \mathbb{N}$

4/6/2017

Sacramento State - Cook - CSc 28 - Spring 2017

68

## Functions

- The first  $\mathbb{N}$  is the function domain
- The second  $\mathbb{N}$  is the function range (codomain)

$f: \mathbb{N} \rightarrow \mathbb{N}$

4/6/2017

Sacramento State - Cook - CSc 28 - Spring 2017

69

## Domain and Range Definitions

- The domain and range of  $f$  is defined exactly as we saw for relations
- Which is not surprising given what a function really is

$\text{domain}(f) = \{ x \mid (x, y) \in f \text{ for some } y \}$   
 $\text{range}(f) = \{ y \mid (x, y) \in f \text{ for some } x \}$

4/6/2017

Sacramento State - Cook - CSc 28 - Spring 2017

70

## Function Attributes

- Function Rules:
  - must be defined for *every element in domain*
  - each value in domain *maps to one element*
- Notice that a function defines a set of ordered pairs: e.g. (1,1) (2,4) (3,9) ...
- We can therefore think of a function as a special kind of relation.

4/6/2017

Sacramento State - Cook - CSc 28 - Spring 2017

71

## Definition of a Function

Let  $f$  be a relation from  $A \rightarrow B$


Where, each  $a \in A$  appears exactly once in an ordered pair  $(a, b) \in f$

4/6/2017

Sacramento State - Cook - CSc 28 - Spring 2017

72

## Relationship Review



4/6/2017

Sacramento State - Cook - CSc 28 - Spring 2017

73

## Relations vs. Functions

- Each domain element, in a relation, can specify *many* relationships
- While, each element in a function domain only specifies *one* relationship
- So....
  - every* function is a relation
  - but not every relation is a function

4/6/2017

Sacramento State - Cook - CSc 28 - Spring 2017

74

## Relations vs. Functions


- Not that in the example (with 1,2,3 and a, b) that some elements in *A* had multiple values in *B*
- In a function, each member in *A* maps to exactly one value in *B*
- So, that relation was not a function!

4/6/2017

Sacramento State - Cook - CSc 28 - Spring 2017

75

## Function Example



4/6/2017

Sacramento State - Cook - CSc 28 - Spring 2017

76

## Examples

- For the following examples, let each example be defined as a relation from *A* to *B*
- Domain and range (codomain) are defined as:

$A = \{1, 2, 3\}$   
 $B = \{x, y, z\}$

4/6/2017

Sacramento State - Cook - CSc 28 - Spring 2017

77

## Is This a Function?

Let  $f = \{ (1, x), (2, y) \}$

**No**, the domain value 3 is missing as a first ordered-pair element

4/6/2017

Sacramento State - Cook - CSc 28 - Spring 2017

78

## Is This a Function?

Let  $g = \{ (1,x), (2,y), (3,z), (1,y) \}$

**No**, the domain element 1 is listed twice.

4/6/2017

Sacramento State - Cook - CSc 28 - Spring 2017

79

## Is This a Function?

Let  $h = \{ (1,x), (2,y), (3,x) \}$

**Yes**, each domain element of A is the first element once.


(There is no restriction on the second element)

4/6/2017

Sacramento State - Cook - CSc 28 - Spring 2017

80

## Is This a Function?



**No!**


(1,a) and (1,b)  
both exist

4/6/2017

Sacramento State - Cook - CSc 28 - Spring 2017

81

## Is This a Function?



**No!**


The domain element  
2 is not mapped to  
a or b

4/6/2017

Sacramento State - Cook - CSc 28 - Spring 2017

82

## Is This a Function?



**Yes**

4/6/2017

Sacramento State - Cook - CSc 28 - Spring 2017

83



## Function Definitions

The Mapping of Sets

## Function Definitions

- Functions are usually defined using a formula
- You should be able to tell that these match a Java method definition – header and body

```
f: Z → Z
f(x) = x * x
```

4/6/2017

Sacramento State - Cook - CSc 28 - Spring 2017

85

## Function Definitions

- First part tells us that f maps every integer to an integer
- Second part tells us f(x) and x<sup>2</sup> are the same thing

```
f: Z → Z
f(x) = x * x
```

4/6/2017

Sacramento State - Cook - CSc 28 - Spring 2017

86

## Example

- In the following, is *g* a function?
- R is a set of reals
- sqrt() is the square root function

```
g: R → R
g(x) = sqrt(x)
```

4/6/2017

Sacramento State - Cook - CSc 28 - Spring 2017

87

## Example

- **No.**
- Not every element of R maps to something in R
- For example, g(-1)  $\notin$  R

```
g: R → R
g(x) = sqrt(x)
```

4/6/2017

Sacramento State - Cook - CSc 28 - Spring 2017

88



## Composition

The Inception of Functions

## Composition

- *Composition* of two functions means the output of one function is used as the input as another
- This is very common in programming – you use the result of one expression as input to another



4/6/2017

Sacramento State - Cook - CSc 28 - Spring 2017

90

## Notation

- Notation for composition is straight forward – it simply consists of a empty circle operator
- Sometimes the (x) is put in front of the first function, but this is not always the case

$$f \circ g(x) \equiv f(g(x))$$

4/6/2017

Sacramento State - Cook - CSc 28 - Spring 2017

91

## Composition Example

$$f(x) = x + 4$$

$$g(x) = x^2$$

$$\begin{aligned} f \circ g(z) &= f(g(z)) \\ &= f(z^2) \\ &= z^2 + 4 \end{aligned}$$

4/6/2017

Sacramento State - Cook - CSc 28 - Spring 2017

92

## Composition Example 2

$$f(x) = x + 4$$

$$g(x) = x^2$$

$$\begin{aligned} g \circ f(z) &= g(f(z)) \\ &= g(z + 4) \\ &= z^2 + 8z + 16 \end{aligned}$$

4/6/2017

Sacramento State - Cook - CSc 28 - Spring 2017

93

## Composite Example

$$R = \{ (1,2), (3,1), (5,3) \}$$

$$S = \{ (2,3), (2,6), (3,9) \}$$

$$R \circ S = \{ (1,3), (1,6), (5,9) \}$$

4/6/2017

Sacramento State - Cook - CSc 28 - Spring 2017

94