

## MOLECULAR AND DEVELOPMENTAL NEUROSCIENCE

# $\beta$ -Amyloid activates presynaptic $\alpha 7$ nicotinic acetylcholine receptors reconstituted into a model nerve cell system: involvement of lipid rafts

Ghous M. Khan,<sup>1,\*</sup> Mei Tong,<sup>1,2</sup> Michelle Jhun,<sup>2,3</sup> Komal Arora<sup>2</sup> and Robert A. Nichols<sup>2</sup><sup>1</sup>Department of Pharmacology & Physiology, Drexel University College of Medicine, Philadelphia, PA 19102, USA<sup>2</sup>Department of Cell and Molecular Biology, University of Hawai'i at Manoa, Honolulu, HI 96813, USA<sup>3</sup>Molecular Bioscience and Bioengineering Program, University of Hawai'i at Manoa, Honolulu, HI 96813, USA**Keywords:** Alzheimer's disease, beta amyloid, calcium imaging, nicotinic acetylcholine receptor, presynaptic receptor

## Abstract

Beta amyloid ( $A\beta$ ) plays a central role in the pathogenesis of Alzheimer's disease.  $A\beta$  is the major constituent of senile plaques, but there is a significant presence of  $A\beta$  in the brain in soluble forms. The results of functional studies indicate that soluble  $A\beta$  interacts with the  $\alpha 7$  nicotinic acetylcholine receptor (nAChR) complex with apparent high affinity. However, conflicting data exist as to the nature of the  $A\beta$ – $\alpha 7$  nAChR interaction, and whether it is the result of specific binding. Moreover, both agonist-like and antagonist-like effects have been reported. In particular, agonist-like effects have been observed for presynaptic nAChRs. Here, we demonstrate  $A\beta_{1-42}$ -evoked stimulatory changes in presynaptic  $Ca^{2+}$  level via exogenous  $\alpha 7$  nAChRs expressed in the axonal varicosities of differentiated hybrid neuroblastoma NG108-15 cells as a model, presynaptic system. The  $A\beta_{1-42}$ -evoked responses were concentration-dependent and were sensitive to the highly selective  $\alpha 7$  nAChR antagonist  $\alpha$ -bungarotoxin. Voltage-gated  $Ca^{2+}$  channels and internal  $Ca^{2+}$  stores were both involved in  $A\beta_{1-42}$ -evoked increases in presynaptic  $Ca^{2+}$  following activation of  $\alpha 7$  nAChRs. In addition, disruption of lipid rafts by cholesterol depletion led to substantially attenuated responses to  $A\beta_{1-42}$ , whereas responses to nicotine were largely intact. These results directly implicate the nicotinic receptor complex as a target for the agonist-like action of pico- to nanomolar concentrations of soluble  $A\beta_{1-42}$  on the presynaptic nerve terminal, including the possible involvement of receptor-associated lipid rafts. This interaction probably plays an important neuromodulatory role in synaptic dynamics.

## Introduction

Beta amyloid ( $A\beta$ ) has been proposed to be a key player in the development of Alzheimer's disease (Selkoe, 2002; Selkoe & Schenk, 2003).  $A\beta$  is visible as the primary component of senile plaques in the brains of Alzheimer's patients.  $A\beta$  is also significantly present in the brain in soluble form, and, in this state, was found to interact with  $\alpha 7$  nicotinic acetylcholine receptors ( $\alpha 7$  nAChRs) with apparent high affinity (Wang *et al.*, 2000a,b), resulting in functional regulation (Dineley *et al.*, 2001; Liu *et al.*, 2001; Pettit *et al.*, 2001).

There is, however, conflicting evidence as to the nature of the  $A\beta$ – $\alpha 7$  nAChR interaction. Agonist-like effects have been observed for presynaptic nAChRs. Picomolar concentrations of  $A\beta_{1-42}$  were found to directly evoke sustained increases in presynaptic  $Ca^{2+}$  level in isolated presynaptic nerve endings (Dougherty *et al.*, 2003) in a manner largely dependent upon the presence of nAChRs (Mehta *et al.*, 2009). Relatively low concentrations of  $A\beta_{1-42}$  elicited were also found to

increase the overflow of dopamine in prefrontal cortex in the presence of tetrodotoxin (TTX), and this stimulatory effect was sensitive to antagonists of  $\alpha 7$  nAChRs and was lost in  $\alpha 7$  null mutant mice (Wu *et al.*, 2007). Interestingly,  $A\beta$  was also found to stimulate tau protein phosphorylation via  $\alpha 7$  nAChRs (Wang *et al.*, 2003). On the other hand,  $A\beta$  peptides were shown to inhibit  $\alpha 7$  nAChRs on rat hippocampal neurons in culture (Liu *et al.*, 2001) and slices (Pettit *et al.*, 2001).  $A\beta$  was also found to inhibit nAChR-mediated  $Ca^{2+}$  influx in synaptosomes (Lee & Wang, 2003; Wang *et al.*, 2009). Interaction of  $A\beta_{1-42}$  with  $\alpha 7$  nAChRs was found to facilitate  $A\beta$  internalization (Nagele *et al.*, 2002). Finally, a recent study reports a stimulatory effect of picomolar  $A\beta$  via presynaptic  $\alpha 7$  nAChRs on long-term potentiation in the Schaffer collateral pathway of hippocampal slices, whereas an inhibitory effect was observed with high nanomolar  $A\beta$  in a manner independent of nAChRs (Puzzo *et al.*, 2008). These findings, derived from the same system, demonstrate that the stimulatory and inhibitory effects of  $A\beta$  are concentration-dependent.

By contrast, a rigorous examination of  $A\beta$  interaction with  $\alpha 7$  nAChRs failed to reveal specific binding of the peptide to the receptor, nor did  $A\beta$  affect ligand interactions with  $\alpha 7$  nAChRs (Small *et al.*, 2007). The major target for  $A\beta$  was found to be membrane lipids, with the main binding energy probably the result of the interaction of

Correspondence: Dr R. A. Nichols, as above.

E-mail: robert.nichols@hawaii.edu

\*Present address: College of Pharmacy, University of Appalachia, Oakwood, VA 24631, USA.

Received 24 April 2009, revised 9 December 2009, accepted 21 December 2009

hydrophobic domains on A $\beta$  for the membrane. Thus, the definitive nature of the interaction between  $\alpha 7$  nAChRs and A $\beta$  remain to be defined. Furthermore, the role of an interaction of A $\beta$  with the membrane, and its resultant impact on nAChR regulation, also needs to be clarified. This study was therefore undertaken to study functional changes in presynaptic Ca<sup>2+</sup> occurring in response to A $\beta$  activation in a model neuronal system expressing exogenous presynaptic  $\alpha 7$  nAChRs. In addition, a potential role for lipid rafts in A $\beta$ -evoked presynaptic Ca<sup>2+</sup> responses was assessed.

## Materials and methods

### Cell culture and expression of exogenous $\alpha 7$ nAChRs

NG108-15 cells (from Dr. William Atchison, Michigan State University) were cultured in high-glucose Dulbecco's modified Eagle's medium (DMEM) containing 15% fetal bovine serum (FBS), 0.1 mM hypoxanthine, 1  $\mu$ M aminopterin and 16  $\mu$ M thymidine in the presence of 5% CO<sub>2</sub> at 37°C (Nelson *et al.*, 1976). Cells were plated at low density onto Cell-Tak-coated coverslips in 35-mm dishes and then differentiated with dibutyl-AMP (1 mM) (Sigma-Aldrich) in DMEM with 1% FBS. After 2–3 days, a pcDNA3.1 construct containing the mouse  $\alpha 7$  nAChR sequence (courtesy of Dr Jerry Stitzel, University of Colorado) was transfected into differentiated cells using the transfectant reagent FuGENE 6 (Roche Diagnostics), obtaining typically > 80% transfection efficiency as previously described (Nichols *et al.*, 2007). As primary control, empty pcDNA3.1 constructs were transfected as well. Mock-transfected control cells were incubated only with the transfection reagent.

### Confocal imaging

After 1–2 days, the transfected, differentiated cells were loaded with fluorescent Ca<sup>2+</sup> indicator dye Fluo-4/AM (Invitrogen) at 5  $\mu$ M in HEPES-buffered saline (HBS) containing (in mM) 142 NaCl, 2.4 KCl, 1.2 K<sub>2</sub>PO<sub>4</sub>, 1 MgCl<sub>2</sub>, 1 CaCl<sub>2</sub>, 5 D-glucose, pH 7.4, and saturated with O<sub>2</sub> at 37°C for 1–1.5 h in preparation for confocal imaging, as described (Nichols *et al.*, 2007). Cells were mounted in a rapid-exchange Warner perfusion system (36  $\mu$ L) for confocal imaging using either a Nikon Diaphot microscope attached to a Nikon PCM 2000 or a Zeiss Axiovert 200M attached to a Zeiss LSM 5 Pascal imaging system. Perfusion with HBS containing 100 nM TTX at 3–5 mL/min was started. Atropine (1  $\mu$ M) was added to the HBS as required. Imaging was commenced (excitation: 488nm; emission: 515–565 nm band-pass; 40 $\times$ /1.3 epifluorescence, oil-immersion Plan-Neofluar objective) and after obtaining a baseline series of four images, stimulatory and/or inhibitory agents were applied by rapid switching between manifolds on the perfusion system. Images were typically collected at 2- or 5-s intervals unless otherwise noted.

### Cholesterol depletion

Depletion of membrane cholesterol to disrupt lipid rafts in the differentiated NG108-15 cells was performed using methyl- $\beta$ -cyclodextrin (Zidovetzki & Levitan, 2007). In brief, cultures on Cell-Tak-coated coverslips in 35-mm dishes were washed with HBS and then incubated with 0.5 or 15 mM methyl- $\beta$ -cyclodextrin (CTD, Inc.) in HBS for 15 min at room temperature (Barrantes, 2007). For 15 mM methyl- $\beta$ -cyclodextrin, ~ 50% of cell cholesterol will be extracted (Zidovetzki & Levitan, 2007), preferentially from the plasma membrane, which is estimated to have roughly half of the total cellular

cholesterol (Warnock *et al.*, 1993), without substantial effect on surface nAChR expression (Barrantes, 2007). Cultures were rapidly washed three times with HBS to remove the methyl- $\beta$ -cyclodextrin. To assess disruption of lipid rafts, cultures were fixed with 4% paraformaldehyde in HBS at room temperature for 30 min, rinsed with phosphate-buffered saline (PBS), and then stained for 15 min with Alexa<sup>488</sup>-conjugated recombinant Cholera Toxin subunit B using the Vybrant staining kit (Invitrogen). The labeled Cholera Toxin subunit B selectively stains GM1 ganglioside, a key lipid raft component. To assess the effect of disruption of lipid rafts on A $\beta$ -evoked Ca<sup>2+</sup> responses, the cultures were preloaded with Fluo-4, as described previously, prior to treatment with methyl- $\beta$ -cyclodextrin, briefly washed with HBS and then imaged using confocal microscopy as described.

### Immunostaining

Cell cultures were fixed with 4% paraformaldehyde in HBS at room temperature for 30 min, rinsed with PBS and then permeabilized with 0.1% Triton X-100 in PBS for 15 min. The cultures were then incubated in PBS containing 5% bovine serum albumin, 10% normal goat serum and 0.1% TritonX-100 for 30 min to block non-specific binding. Affinity-purified anti- $\alpha 7$  nAChR antibody (1 : 100; Chemicon) was then added to the cultures and incubated for 60 min at room temperature. The cultures were then washed with 10% goat serum plus 0.1% TritonX-100 in PBS, and incubated with the FITC-conjugated goat anti-rabbit IgG (1 : 200; Jackson ImmunoResearch) as secondary antibody, for 60 min at room temperature. Background staining was assessed using the secondary antibody alone (Nayak *et al.*, 2000). The cultures were finally washed with 10% normal goat serum and PBS, plated onto glass microscope slides and sealed for imaging via confocal microscopy.

### Chemicals

Cell Tak was from BD Biosciences (Bedford, MA, USA). Nicotine tartrate citrate, acetylcholine chloride, choline chloride, caffeine, nitrendipine and atropine chloride were all purchased from Sigma-Aldrich. Nicotine, acetylcholine, choline and caffeine solutions were freshly prepared. A $\beta$ <sub>1–42</sub> was purchased from BACHEM (King of Prussia, PA, USA), while  $\omega$ -conotoxin GVIA, TTX and  $\alpha$ -BgTx were from Calbiochem (La Jolla, CA, USA). Methyl- $\beta$ -cyclodextrin was from Cyclodextrin Technologies (CTD, Inc., Gainesville, FL, USA). The Vybrant Lipid Raft staining kit was from Invitrogen (Carlsbad, CA, USA). All drugs, except A $\beta$ <sub>1–42</sub> and nitrendipine, were made into solutions in HBS. Stock nitrendipine (20 mM) was dissolved in dimethyl sulfoxide. Stock solutions of A $\beta$ <sub>1–42</sub> (1 mM) were made in water with vigorous vortexing (Dougherty *et al.*, 2003). Native and Western immunoblot analysis of nanomolar concentration of A $\beta$ <sub>1–42</sub> in HBS demonstrated the presence of several oligomeric forms, with little evidence of monomer under native conditions (see Fig. S1), consistent with prior analyses (Bell *et al.*, 2004).

### Data analysis and statistics

The fluorescent intensities associated with individual varicosities were determined from digitized images by a blind observer using MetaMorph image analysis software and were expressed as normalized values ( $F/F_0$ , where  $F_0$  is the fluorescence intensity at time zero). All time series were corrected for photobleaching (typically < 3%). Averaged data are presented as means  $\pm$  SEM, where  $n$  refers to the

number of (pooled) varicosities examined. Peak amplitudes (as  $F/F_0$ ) were determined from averaged maximum plateau values. Peak amplitudes were compared using one-way ANOVA with Bonferroni multiple comparison *post hoc* test or two-tailed Student's *t*-test for two groups, using GRAPHPAD INSTAT 3. Significance was indicated when  $P$  was minimally  $< 0.05$ .

## Results

### *Nicotinic agonists elicit dose-dependent $Ca^{2+}$ responses in NG108-15 neuritic presynaptic-like varicosities expressing $\alpha 7$ nAChRs*

Differentiated NG108-15 cells elaborate long, branched axonal processes containing large varicosities ( $\sim 5$ – $15 \mu\text{m}$ ) that are fully capable of forming synapses with an appropriate target (Nelson *et al.*, 1976), but do not synapse on each other. In the absence of synapse formation, the axonal varicosities otherwise display features of a presynaptic element (Han *et al.*, 1991), including synaptic vesicles capable of supporting neurotransmitter release (McGee *et al.*, 1978) and voltage-gated  $Ca^{2+}$  channels (VGCCs) typical of presynaptic terminals (Lukyanetz *et al.*, 1998), and thus permit assessment of presynaptic regulation in discretely identifiable structures. There are equivalent structures existing in the central nervous system, as axonal varicosities proposed to release neurotransmitter at a distance (so-called volume transmission) from postsynaptic targets have been noted in the brain for several neurotransmitter systems (e.g. Umbriaco *et al.*, 1994). A propos of the present study, these cells (and their varicosities) do not express functional nicotinic receptors, but do express 5-HT<sub>3</sub> receptors on their cell bodies (Yakel *et al.*, 1991) and axonal varicosities (Ronde & Nichols, 2001). These receptors are both members of the ligand-gated ion channel superfamily. Hence, the cells were predicted to have all of the necessary components for their expression and could thus serve as a neuronal system for 'reconstituting' presynaptic nicotinic receptors via exogenous expression.

The presence of  $\alpha 7$  nAChRs in axonal varicosities (arrows) of differentiated NG108-15 cells transfected with pcDNA3.1 vector containing the mouse  $\alpha 7$  nAChR sequence (Fig. 1B and C) was detected by immunocytochemistry using a recently developed anti-mouse  $\alpha 7$  nAChR antibody (Chemicon) and visualized via confocal

imaging. The background fluorescence resulting from immunostaining of differentiated NG108-15 cells transfected with empty vector using the same anti-mouse  $\alpha 7$  nAChR antibody is shown for comparison in Fig. 1A.

The effects of nAChR agonists such as nicotine, acetylcholine and choline were tested on the axonal varicosities of NG108-15 cells transfected with murine  $\alpha 7$  cDNA, measuring  $Ca^{2+}$  responses using confocal imaging typically 24–48 h after transfection. Increasing concentrations of the nAChR agonist nicotine (500 nM, 1  $\mu\text{M}$  and 10  $\mu\text{M}$ ) were applied in the presence of TTX (100 nM to block regenerative responses), and induced rapid-onset, sustained increases in  $[Ca^{2+}]$  in the neuritic varicosities, as detected by relative increases in fluorescence signal (Fig. 2A). To varying degrees, nicotine also induced  $Ca^{2+}$  responses in cell bodies. Both 1 and 10  $\mu\text{M}$  nicotine (Fig. 2B) produced maximum responses in the varicosities ( $2.86 \pm 0.33 F/F_0$ ,  $F_{4,38} = 38$ ,  $P < 0.001$  and  $2.96 \pm 0.39 F/F_0$ ,  $F_{4,38} = 37$ ,  $P < 0.001$ , respectively, compared with control), which were significantly more pronounced than that of 500 nM nicotine ( $1.58 \pm 0.01 F/F_0$ ,  $P > 0.05$  vs. control). Responses to all the concentrations of nicotine used in the experiment were completely blocked by perfusion with  $\alpha$ -bungarotoxin ( $\alpha$ -BgTx; 30 nM) (Fig. 2B,  $1.24 \pm 0.02 F/F_0$  for 1  $\mu\text{M}$  nicotine,  $P > 0.05$  vs. control), a highly selective  $\alpha 7$  nAChRs receptor antagonist. Extending the length of perfusion of nicotine (10  $\mu\text{M}$ ) for 7 min evoked sustained increases in  $Ca^{2+}$  response that peaked around 3 min and then gradually returned to baseline (Fig. S2). No response was observed to nicotine in mock-transfected preparations (Figs 2B and S2).

The specific  $\alpha 7$  nAChR agonist choline also evoked dose-dependent  $Ca^{2+}$  responses in axonal varicosities expressing  $\alpha 7$  nAChRs (Fig. 2C,  $1.82 \pm 0.08 F/F_0$ ,  $F_{3,36} = 86$ ,  $P < 0.001$  and  $1.42 \pm 0.06 F/F_0$ ,  $F_{3,36} = 27$ ,  $P < 0.001$  for 500 and 50  $\mu\text{M}$  choline, respectively, compared with control). Similarly, the AChR agonist acetylcholine, applied in the presence of atropine to isolate its nicotinic action, evoked concentration-dependent  $Ca^{2+}$  responses (Fig. 2D,  $1.36 \pm 0.05 F/F_0$ ,  $F_{3,24} = 17$ ,  $P < 0.01$  and  $1.50 \pm 0.50 F/F_0$ ,  $F_{3,24} = 34.7$ ,  $P < 0.001$  for 1 and 5 mM acetylcholine, respectively, compared with control), although they were not as consistently robust as those seen with nicotine or choline. Responses to both choline and acetylcholine were blocked by  $\alpha$ -BgTx (50 nM;  $1.066 \pm 0.02 F/F_0$ ,  $P > 0.05$  vs. control and  $1.12 \pm 0.02 F/F_0$ ,  $P > 0.05$  vs. control, respectively).




FIG. 1. Representative images demonstrating the presence of  $\alpha 7$  nAChR in axonal varicosities (e.g. arrow) of differentiated NG108-15 cells transfected with pcDNA3.1 vector containing the mouse  $\alpha 7$  nAChR sequence (B), as detected by immunocytochemistry using a recently developed anti-mouse  $\alpha 7$  nAChR antibody (Chemicon) and visualized with confocal imaging. Panel A shows the background fluorescence resulting from immunocytochemistry with the same anti-mouse  $\alpha 7$  nAChR antibody applied to differentiated NG108-15 cells transfected with empty pcDNA3.1 vector (A). Panel C shows a magnified view of an immunopositive varicosity (B, upper right). Scale bar = 10  $\mu\text{m}$ .






FIG. 2. Nicotinic agonist-evoked increases in  $[Ca^{2+}]_i$  in axonal varicosities of NG108-15 cells transfected with mouse  $\alpha 7$  nAChR. (A) Representative confocal image sequence (left) and individual  $Ca^{2+}$  responses (right) to  $1 \mu M$  nicotine (bar) in NG108-15 varicosities expressing  $\alpha 7$  nAChRs. Scale bar =  $10 \mu m$ . Averaged time series of changes in  $Ca^{2+}$  in varicosities in response to increasing concentrations of (B) nicotine ( $500 \text{ nM}$ ,  $n = 10$ ;  $1 \mu M$ ,  $n = 8$ ;  $10 \mu M$ ,  $n = 6$ ), as compared with  $1 \mu M$  nicotine applied to mock-transfected cultures (control,  $n = 6$ ); (C) choline ( $500 \mu M$ ,  $n = 7$ ;  $50 \mu M$ ,  $n = 14$ ), a specific  $\alpha 7$  nAChR agonist, as compared with control cultures ( $n = 6$ ); (D) acetylcholine (ACh:  $5 \text{ mM}$ ,  $n = 8$ ;  $1 \text{ mM}$ ,  $n = 9$ ), as compared with control ( $n = 6$ ). For all agonists, pretreatment with  $\alpha$ -BgTx (30 or  $50 \text{ nM}$ ) for 30 min completely blocked the agonist-evoked  $Ca^{2+}$  responses ( $1 \mu M$  nicotine,  $n = 13$ ;  $50 \mu M$  choline,  $n = 13$ ; ACh,  $n = 5$ ). For acetylcholine, atropine ( $1 \mu M$ ) was added to block muscarinic AChRs.  $Ca^{2+}$  responses were assessed via confocal imaging following loading with the  $Ca^{2+}$ -sensitive fluorescent dye Fluo-4.

#### *A $\beta_{1-42}$ induces $Ca^{2+}$ responses in NG108-15 neuritic varicosities expressing $\alpha 7$ nAChRs*

Previous work employing functional approaches suggested that the beta amyloid peptide regulates  $\alpha 7$  nAChRs (Liu *et al.*, 2001; Pettit *et al.*, 2001; Dougherty *et al.*, 2003; Wu *et al.*, 2007; Puzzo *et al.*, 2008; Mehta *et al.*, 2009), but none of the studies demonstrated definitively a direct interaction, and pharmacological inhibition of A $\beta$ -regulated responses using nicotinic antagonists was often partial. Moreover, specific binding of A $\beta$  to nAChRs has been questioned (Small *et al.*, 2007). Using NG108-15 cells expressing exogenous  $\alpha 7$  nAChR, we tested the acute effect of soluble A $\beta_{1-42}$ . Soluble A $\beta_{1-42}$  ( $100 \text{ nM}$ ) evoked sharp and sustained responses (Fig. 3,  $2.89 \pm 0.35 F/F_0$ ,  $F_{2,25} = 18$ ,  $P < 0.001$  vs. control reverse peptide), similar to those obtained using nicotine. The responses were nearly completely blocked by  $\alpha$ -BgTx (Fig. 3B,  $P > 0.05$  vs. control peptide). Soluble A $\beta_{1-42}$  evoked no response in mock-transfected varicosities and/or cell bodies. The small, slow increase in  $Ca^{2+}$  observed with the reverse peptide A $\beta_{42-1}$  in NG108-15 neuritic varicosities expressing  $\alpha 7$  nAChRs (Fig. 3B) may indicate a direct membrane effect of the hydrophobic elements in A $\beta$  (Small *et al.*, 2007). However, the block by  $\alpha$ -BgTx would indicate an absence of a non-specific membrane effect by A $\beta$  when the receptor is blocked. Extending the length of perfusion of A $\beta_{1-42}$  ( $100 \text{ nM}$ ) for 7 min evoked sustained increases in  $Ca^{2+}$  response that peaked around 3 min and then gradually returned to baseline (Fig. S3), similar to that

observed for nicotine. The A $\beta_{1-42}$ -evoked  $Ca^{2+}$  responses were concentration-dependent (Fig. 3C), with an  $EC_{50}$  between 1 and  $100 \text{ nM}$ . Addition of nicotine ( $500 \text{ nM}$ ) at the peak of the A $\beta_{1-42}$ -evoked  $Ca^{2+}$  response had no impact (Fig. 3D), consistent with prior evidence (Dougherty *et al.*, 2003; Mehta *et al.*, 2009) showing that activation by A $\beta$  occludes the response to subsequent co-application of nicotine.

#### *Nature of $Ca^{2+}$ responses to A $\beta_{1-42}$*

In NG108-15 axonal varicosities, 5-HT<sub>3</sub> receptor agonist-evoked  $Ca^{2+}$  responses were previously shown to depend on VGCCs and  $Ca^{2+}$  stores (Rondé & Nichols, 2001), indicating that the receptors trigger depolarization via  $Na^+$  influx, opening VGCCs, and  $Ca^{2+}$ -induced  $Ca^{2+}$  release (CICR) subsequent to the resultant  $Ca^{2+}$  influx. Because  $\alpha 7$  nAChR activation will also result in substantial  $Ca^{2+}$  influx and depolarization (Role & Berg, 1996), it was important to examine the roles of VGCCs and internal  $Ca^{2+}$  stores in A $\beta_{1-42}$ -evoked  $Ca^{2+}$  responses via nAChRs in NG108-15 varicosities. To test the role of VGCCs in A $\beta_{1-42}$ -evoked  $Ca^{2+}$  responses, we co-applied two  $Ca^{2+}$  channel antagonists, nitrendipine ( $20 \mu M$ ), an L-type VGCC blocker, and  $\omega$ -conotoxin GVIA ( $500 \text{ nM}$ ), an N-type VGCC blocker, which block depolarization-induced  $Ca^{2+}$  responses (see Rondé & Nichols, 2001). The  $Ca^{2+}$  channel blockers significantly inhibited A $\beta$ -evoked responses in axonal varicosities of NG108-15 cells expressing  $\alpha 7$




FIG. 3. Aβ-evoked increases in  $[Ca^{2+}]_i$  in axonal varicosities of NG108-15 cells transfected with mouse  $\alpha 7$  nAChR. (A) Representative confocal image sequence (left) and individual  $Ca^{2+}$  responses (right) to 100 nM Aβ<sub>1-42</sub> (bar) in NG108-15 varicosities expressing  $\alpha 7$  nAChRs. Scale bar = 10  $\mu m$ . (B) Averaged time series of changes in  $Ca^{2+}$  in varicosities in response to 100 nM Aβ<sub>1-42</sub> in the absence ( $n = 13$ ) or presence of 50 nM  $\alpha$ -BgTx (30-min pretreatment;  $n = 9$ ). For comparison, the effect of reverse sequence peptide Aβ<sub>42-1</sub> was assessed (control;  $n = 6$ ). There was no effect of Aβ<sub>1-42</sub> on mock-transfected or empty vector-transfected cultures (not shown). (C) Histogram of average peak amplitudes of  $Ca^{2+}$  responses to increasing concentrations of Aβ<sub>1-42</sub> (Control,  $n = 6$ ; 100 pM,  $n = 7$ ; 1 nM,  $n = 6$ ; 100 nM,  $n = 9$ ,  $F_{3,24} = 23.8$ ,  $P < 0.001$ ).  $Ca^{2+}$  responses were assessed via confocal imaging following loading with the  $Ca^{2+}$ -sensitive fluorescent dye Fluo-4. (D) Averaged time series of changes in  $Ca^{2+}$  in varicosities in response to 100 nM Aβ<sub>1-42</sub> followed by co-administration of 500 nM nicotine, where indicated ( $n = 9$ ). Inset shows effect of stimulation with nicotine alone in a sister culture ( $n = 9$ ).

nAChRs (Fig. 4A,  $1.2 \pm 0.05 F/F_0$ ,  $F_{2,25} = 24$ ,  $P < 0.001$  vs. control response to Aβ). The small increase in the presence of the VGCC blockers is probably the result of  $Ca^{2+}$  influx directly via the receptor channel (Rondé & Nichols, 2001).

$Ca^{2+}$  influx through presynaptic nAChRs was found to lead to mobilization of  $Ca^{2+}$  from internal  $Ca^{2+}$  stores via CICR in hippocampal terminals (Sharma & Vijayaraghavan, 2003;

La Magueresse & Cherubini, 2007; Sharma *et al.*, 2008) and glutamatergic prefrontal cortical terminals (Dickinson *et al.*, 2008), contributing to the modulation of neurotransmitter release. To test this pathway, intracellular  $Ca^{2+}$  stores were first depleted using millimolar caffeine in the presence of EGTA and then perfused with Aβ<sub>1-42</sub>. No significant responses were elicited by Aβ<sub>1-42</sub> in  $Ca^{2+}$  store-depleted varicosities (Fig. 4A,  $F_{2,25} = 37$ ,  $P < 0.001$  vs. control




FIG. 4.  $A\beta$ -evoked increases in  $[Ca^{2+}]_i$  in axonal varicosities of NG108-15 cells transfected with mouse  $\alpha 7$  nAChR depend on VGCCs and CICR. (A) Averaged time series of changes in  $Ca^{2+}$  in varicosities in response to 100 nM  $A\beta_{1-42}$  in the presence of 20  $\mu M$  nitrendipine and 500 nM  $\omega$ -conotoxin GVIA (10-min pretreatment;  $n = 16$ ) compared with prior depletion of  $Ca^{2+}$  stores using 10 mM caffeine in the absence of external  $Ca^{2+}$  (blue bar)  $n = 6$ . (B) Effect of pretreatment with 20  $\mu M$  nitrendipine and 500 nM  $\omega$ -conotoxin GVIA ( $n = 5$ ) compared with prior depletion of  $Ca^{2+}$  stores using 10 mM caffeine ( $n = 6$ ) on nicotine-induced  $Ca^{2+}$  responses (peak amplitudes normalized to control nicotine response,  $n = 8$ ; see Fig. 2).

response to  $A\beta$ ). Similar blockade by the  $Ca^{2+}$  channel antagonists ( $F_{2,17} = 15.2$ ,  $P < 0.01$ ) or caffeine-induced store depletion ( $F_{2,17} = 14.6$ ,  $P < 0.01$ ) was found for nicotine-induced  $Ca^{2+}$  responses (Fig. 4B). In summary, these results indicate that the  $\alpha 7$  nAChR activation upon  $A\beta$  application to the NG108-15 varicosities results in  $Ca^{2+}$  influx via VGCCs, which, together with  $Ca^{2+}$  influx via the receptor channel, triggers CICR.

#### Lipid raft disruption abolished $A\beta$ -evoked responses

In view of the noted interaction of  $A\beta$  with membrane lipids (Small *et al.*, 2007), particularly GM1 ganglioside (Kakio *et al.*, 2002) and the noted small but detectable effect of  $A\beta_{42-1}$  (see Fig. 3B), we tested whether disruption of lipid rafts, proposed to associate with  $\alpha 7$  nAChRs (Brusés *et al.*, 2001), had any impact on  $A\beta_{1-42}$ -evoked  $Ca^{2+}$  responses via nAChRs in NG108-15 varicosities. Prior application of 0.5 or 15 mM methyl- $\beta$ -cyclodextrin for 15 min to deplete membrane cholesterol and hence disrupt associated lipid rafts (Fig. 5A, right) attenuated  $A\beta_{1-42}$ -evoked responses in  $\alpha 7$  nAChR-transfected varicosities (Fig. 5A, left,  $F_{2,16} = 41$ ,  $P < 0.001$  for 15 mM cyclodextrin vs. control response to  $A\beta$ ). The higher concentration (15 mM) of methyl- $\beta$ -cyclodextrin has been shown to substantially deplete the membrane of cholesterol, whereas the lower concentration (0.5 mM) will only partially deplete the membrane (Zidovetzki & Levitan, 2007), and the results are generally consistent with this difference. Cholesterol depletion with 15 mM methyl- $\beta$ -cyclodextrin only partially attenuated nicotine-induced responses (Fig. 5B, left; nicotine: 68%, as compared with initial stimulation,  $P = 0.002$ , two-tailed unpaired *t*-test), indicating that nicotine's interaction with the receptor is largely intact. Interestingly, ACh-induced  $Ca^{2+}$  responses were substantially reduced, but not eliminated, following cholesterol depletion (Fig. 5B, right; ACh: 39%, as compared with initial stimulation,  $P = 0.002$ , two-tailed unpaired *t*-test). That robust responses to  $K^+$ -depolarization remained after treatment of the cells with methyl- $\beta$ -cyclodextrin verified the viability of the preparations (Fig. 5B). Disruption of lipid rafts by methyl- $\beta$ -cyclodextrin did not appear to alter the overall expression of  $\alpha 7$  nAChRs (Fig. 5C). These

results suggest an important role for associated lipids in the agonist-like effect of  $A\beta$  on presynaptic  $\alpha 7$  nAChRs.

#### Discussion

Soluble  $A\beta$  can aggregate into soluble prefibrillar intermediates, largely as low-molecular weight oligomers (Walsh & Selkoe, 2007). Soluble oligomers are proposed to play significant roles in the pathogenesis of Alzheimer's disease (Selkoe, 2008) and their levels have been found to strongly correlate with the neurodegenerative process in patients with Alzheimer's disease (McLean *et al.*, 1999). We report here the acute stimulatory effects of soluble  $A\beta$  in the presynaptic-like axonal varicosities of NG108-15 cells expressing  $\alpha 7$  nAChRs, comparable with  $Ca^{2+}$  responses evoked by nicotinic agonists. These effects were assessed as changes in  $Ca^{2+}$  levels in the axonal varicosities, as the regulation of presynaptic  $Ca^{2+}$  is integral to synaptic function. The  $Ca^{2+}$  responses to soluble  $A\beta_{1-42}$  applied to varicosities expressing  $\alpha 7$  nAChRs were blocked by  $\alpha$ -BgTx, a potent and highly selective antagonist of  $\alpha 7$  nAChRs. In addition, the  $A\beta_{1-42}$ -elicited responses were not observed in axonal varicosities of mock-transfected cells. Together, these results confirm a direct interaction of  $A\beta$  with presynaptic  $\alpha 7$  nAChRs. To our knowledge, this is the first instance of a putative receptor target for pico- to nanomolar  $A\beta$ , identified in *ex vivo* and *in vivo* studies of presynaptic terminals (Dougherty *et al.*, 2003; Wu *et al.*, 2007; Puzzo *et al.*, 2008; Abbott *et al.*, 2008; Mehta *et al.*, 2009; Wang *et al.*, 2009), being 'reconstituted' in a neuronal culture system. Other studies describe effects of  $A\beta$  on exogenous nAChRs expressed *in vitro*, but the systems used did not reconstitute the equivalent environment, here the presynaptic nerve terminal.

The pronounced reduction in  $A\beta$ -evoked  $Ca^{2+}$  responses following the application of  $Ca^{2+}$  channel blockers to the differentiated cultures suggest that presynaptic VGCCs are a major component in the overall  $Ca^{2+}$  entry in cases where nAChRs evoke significant changes in membrane potential. In addition, the loss of  $A\beta$ -evoked responses in varicosities in which internal  $Ca^{2+}$  stores were depleted suggests that CICR may play a dominant role in  $A\beta$ -triggered presynaptic






FIG. 5. Cholesterol depletion via methyl- $\beta$ -cyclodextrin to disrupt lipid rafts abolishes  $\text{A}\beta_{1-42}$ -evoked responses in  $\alpha 7$  nAChR-transfected varicosities in NG108-15 cells. (A) Left: averaged time series of  $\text{Ca}^{2+}$  responses to 100 nM  $\text{A}\beta_{1-42}$  in varicosities transfected with  $\alpha 7$  nAChR pretreated (closed circles) or not (control: triangles) with methyl- $\beta$ -cyclodextrin at 0.5 mM ( $n = 6$ ) or 15 mM ( $n = 7$ ) for 15 min (Barrantes, 2007). To maintain identical experimental conditions, the untreated (control) cultures ( $n = 6$ ) were pre-incubated for the same length of time in HBS before perfusing with  $\text{A}\beta_{1-42}$ . Right: confocal micrographs showing lipid raft disruption after treatment with methyl- $\beta$ -cyclodextrin schematic to deplete membrane cholesterol, using Alexa<sup>488</sup>-conjugated Cholera Toxin B staining (Vybrant Lipid Raft kit) for GM1 ganglioside, a key lipid raft component. Lipid rafts appeared as patchy staining on control cells, shifting to uniform, diffuse staining after cholesterol depletion. Upper sequence: scale bar = 30  $\mu\text{m}$ ; lower micrograph: scale bar = 10  $\mu\text{m}$ . (B) Averaged time series of  $\text{Ca}^{2+}$  responses to 1  $\mu\text{M}$  nicotine (left sequence;  $n = 18$ ) or 0.5  $\mu\text{M}$  ACh in the presence of 1  $\mu\text{M}$  atropine (right sequence;  $n = 14$ ) in varicosities transfected with  $\alpha 7$  nAChR following sequential treatment with HBS, methyl- $\beta$ -cyclodextrin at 15 mM for 15 min, and an additional 10-min wash with HBS followed by stimulation with nicotine or ACh (blue curves). Each sequence was followed by stimulation with 30 mM KCl (red curves). Preparations were washed for 5–10 min between each stimulation. (C) Immunostaining NG108-15 cells and varicosities (arrows) for  $\alpha 7$  nAChRs before (control) and after ( $\text{M}\beta\text{C}$ ) treatment (two fields). Scale bar = 30  $\mu\text{m}$ .

responses. Together, these data show that the activation of presynaptic  $\alpha 7$  nAChRs by pico- to nanomolar  $\text{A}\beta$  has a two-component  $\text{Ca}^{2+}$  entry into the varicosities, similar to that observed for nicotinic

agonist activation of the receptor. The first is from a direct permeation of  $\text{Ca}^{2+}$  via the nAChR channel, while the second results from entry via VGCCs opened in response to simultaneous nAChR-evoked

membrane depolarization. In response to this  $\text{Ca}^{2+}$  entry, CICR is evoked, resulting in a substantial amplification and prolongation of the signal.

Ionic currents mediated by  $\alpha 7$  nAChRs expressed in cell bodies typically display rapid desensitization (Role & Berg, 1996). In contrast, presynaptic nicotinic receptor activation has been found to result in a prolonged stimulatory effect (McGehee *et al.*, 1995; Gray *et al.*, 1996; Nayak *et al.*, 2001; Sharma & Vijayaraghavan, 2003) depending on the preparation (Mansvelder *et al.*, 2002; Mehta *et al.*, 2009). As noted, CICR, where it occurs, will further prolong the signal. The apparent kinetics of an agonist-like effect of  $\text{A}\beta$  on presynaptic signals also indicate a prolonged action. This effect is in sharp contrast to the effect of  $\text{A}\beta$  on nicotinic receptors expressed in cell bodies or dendrites (e.g. Liu *et al.*, 2001, 2009; Pettit *et al.*, 2001; Wu *et al.*, 2004). One possible explanation for the difference may be that  $\text{A}\beta$  rapidly induces, or stabilizes, the inactivated state of the nAChRs when expressed at postsynaptic or somatic sites, resulting in an antagonist-like action, whereas a slower rate of desensitization of presynaptic nAChRs may permit the agonist-like effect of  $\text{A}\beta$ . This possibility might be addressed, ideally, by examining ionic nicotinic currents at presynaptic and postsynaptic sites in the same system.

Regardless, the basis for the distinct behavior of presynaptic nAChRs remains to be elucidated. We have speculated that the microenvironment of the nAChR complex in the presynaptic terminal may be unique (Nayak *et al.*, 2001). Axons and terminals of neurons have been deemed to be large extensions of classical apical membranes, particularly in view of the abundant presence of lipid rafts (Tsui-Pierchala *et al.*, 2002). We thus examined whether cholesterol depletion, to disrupt lipid rafts, had an impact on  $\text{A}\beta$ -evoked presynaptic  $\text{Ca}^{2+}$  responses. Remarkably, the  $\text{A}\beta$ -evoked responses in the varicosities were dramatically attenuated following cholesterol depletion by methyl- $\beta$ -cyclodextrin, whereas nicotine-evoked responses were largely intact. Although cholesterol depletion will also occur in non-raft membrane (Zidovetzki & Levitan, 2007), clear disruption of lipid rafts, as detected via staining for GM1 ganglioside, which is a prominent component of rafts, was observed. In view of the hydrophobic domain of  $\text{A}\beta$  (Walsh & Selkoe, 2007), derived from part of the predicted transmembrane domain of the amyloid precursor protein, we postulate that an interaction of  $\text{A}\beta$  directly with the membrane (Small *et al.*, 2007) may influence its targeting and regulation of presynaptic nAChRs. There is evidence that GM1 ganglioside interacts with  $\text{A}\beta$  (Kakio *et al.*, 2002) and thus the lipid raft microenvironment may serve to guide  $\text{A}\beta$  in its interaction with the receptor, positioning it to interact with the ligand-binding (activation) domain. In addition, a dual lipid raft–receptor interaction may also explain the apparent picomolar potency of  $\text{A}\beta$  for the nAChRs in the varicosities, as the apparent affinity of  $\text{A}\beta$  will be a power-function composite of its individual affinities for the raft/membrane and the receptor.

The precise target(s) for  $\text{A}\beta$  in the membrane, the precise form (oligomeric?) of  $\text{A}\beta$  interacting with the membrane targets, and how the membrane target(s) contributes to the targeting and/or regulation of presynaptic nAChRs remain to be determined, particularly as lipid rafts are also present in the somata. Lipid–nAChR interactions have been noted (Barrantes, 2004), as an annular shell around the transmembrane core of the receptor, and, in addition, such lipid shells may participate in mediating receptor-raft targeting (Anderson & Jacobson, 2002). The interaction of rafts with the receptor complex and the interaction of  $\text{A}\beta$  with unique components in the rafts may provide a link among the disparate receptor systems found to be regulated by soluble  $\text{A}\beta$ .

## Supporting Information

Additional supporting information may be found in the online version of this article:

Fig. S1. SDS–PAGE and native gel analysis of the  $\text{A}\beta$  preparation solubilized in aqueous buffered solution.

Fig. S2. Averaged time series of  $\text{Ca}^{2+}$  responses in the varicosities of differentiated NG108-15 cells expressing exogenous  $\alpha 7$  nAChRs or not (mock-transfected) to an extended length of perfusion ( $> 5$  min) with nicotine ( $10 \mu\text{M}$ ).

Fig. S3. Averaged time series of  $\text{Ca}^{2+}$  responses in the varicosities of differentiated NG108-15 cells expressing exogenous  $\alpha 7$  nAChRs to an extended length of perfusion ( $> 5$  min) with  $\text{A}\beta_{1-42}$  ( $100 \text{ nM}$ ).

Please note: As a service to our authors and readers, this journal provides supporting information supplied by the authors. Such materials are peer-reviewed and may be re-organized for online delivery, but are not copy-edited or typeset by Wiley-Blackwell. Technical support issues arising from supporting information (other than missing files) should be addressed to the authors.

## Acknowledgements

We thank Dr Jerry Stitzel, University of Colorado, for providing the mouse  $\alpha 7$  nAChR sequence in the pcDNA3.1zeo expression vector. We thank Michelle Guerin for excellent technical assistance. We also thank Dr Frederick Bellinger for critical reading of the manuscript. This work was supported by grants from the American Health Assistance Foundation and the National Institute on Aging (AG21586).

## Abbreviations

$\text{A}\beta$ , beta amyloid; CICR,  $\text{Ca}^{2+}$ -induced  $\text{Ca}^{2+}$  release; HBS, HEPES-buffered saline; nAChR, nicotinic acetylcholine receptors; TTX, tetrodotoxin; VGCC, voltage-gated calcium channel;  $\alpha$ -BgTx,  $\alpha$ -bungarotoxin.

## References

- Abbott, J.J., Howlett, D.R., Francis, P.T. & Williams, R.J. (2008)  $\text{A}\beta_{1-42}$  modulation of Akt phosphorylation via  $\alpha 7$  nAChR and NMDA receptors. *Neurobiol. Aging*, **29**, 992–1001.
- Anderson, R.G. & Jacobson, K. (2002) A role for lipid shells in targeting proteins to caveolae, rafts, and other lipid domains. *Science*, **296**, 1821–1825.
- Barrantes, F.J. (2004) Structural basis for lipid modulation of nicotinic acetylcholine receptor function. *Brain Res. Rev.*, **47**, 71–95.
- Barrantes, F.J. (2007) Cholesterol effects on nicotinic acetylcholine receptor. *J. Neurochem.*, **103**, 72–80.
- Bell, K.A., O’Riordan, K.J., Sweatt, J.D. & Dineley, K.T. (2004) MAPK recruitment by  $\beta$ -amyloid in organotypic hippocampal slice cultures depends on physical state and exposure time. *J. Neurochem.*, **91**, 349–361.
- Brusés, J.L., Chauvet, N. & Rutishauser, U. (2001) Membrane lipid rafts are necessary for the maintenance of the  $\alpha 7$  nicotinic acetylcholine receptor in somatic spines of ciliary neurons. *J. Neurosci.*, **21**, 504–512.
- Dickinson, J.A., Kew, J.N.C. & Wonnacott, S. (2008) Presynaptic  $\alpha 7$  and  $\beta 2$ -containing nicotinic acetylcholine receptors modulate excitatory amino acid release from rat prefrontal cortex nerve terminals via distinct cellular mechanisms. *Mol. Pharmacol.*, **74**, 348–359.
- Dineley, K.T., Westerman, M., Bui, D., Bell, K., Ashe, K.H. & Sweatt, J.D. (2001)  $\beta$ -amyloid activates the mitogen-activated protein kinase cascade via hippocampal  $\alpha 7$  nicotinic acetylcholine receptors: in vitro and in vivo mechanisms related to Alzheimer’s disease. *J. Neurosci.*, **21**, 4125–4133.
- Dougherty, J.J., Wu, J. & Nichols, R.A. (2003)  $\beta$ -amyloid regulation of presynaptic nicotinic receptors in rat hippocampus and neocortex. *J. Neurosci.*, **23**, 6740–6747.
- Gray, R., Rajan, A.S., Radcliffe, K.A., Yakehiro, M. & Dani, J.A. (1996) Hippocampal synaptic transmission enhanced by low concentrations of nicotine. *Nature*, **383**, 713–716.
- Han, H.Q., Nichols, R.A., Rubin, M., Böhler, M. & Greengard, P. (1991) Induction of formation of presynaptic terminals in neuroblastoma cells by synapsin IIB. *Nature*, **349**, 697–700.



- Kakio, A., Nishimoto, S.I., Yanagisawa, K., Kozutsumi, Y. & Matsuzaki, K. (2002) Interactions of amyloid  $\beta$ -protein with various ganglioside-bound form as an endogenous seed for Alzheimer amyloid. *Biochemistry*, **41**, 7385–7390.
- La Magueresse, C. & Cherubini, E. (2007) Presynaptic calcium stores contribute to nicotine-elicited potentiation of evoked synaptic transmission at CA3-CA1 connections in the neonatal rat hippocampus. *Hippocampus*, **17**, 316–325.
- Lee, D.H.S. & Wang, H.Y. (2003) Differential physiologic responses of  $\alpha 7$  nicotinic acetylcholine receptors to  $\beta$ -amyloid<sub>1–40</sub> and  $\beta$ -amyloid<sub>1–42</sub>. *J. Neurobiol.*, **55**, 25–30.
- Liu, Q., Kawai, H. & Berg, D.K. (2001)  $\beta$ -Amyloid peptide blocks the response of  $\alpha 7$ -containing nicotinic receptors on hippocampal neurons. *Proc. Natl Acad. Sci. USA*, **98**, 4734–4739.
- Liu, Q., Huang, Y., Xue, F., Simard, A., DeChon, J., Li, G., Zhang, J., Lucier, L., Wang, M., Sierks, M., Hu, G., Chang, Y., Lukas, R.J. & Wu, J. (2009) A novel nicotinic acetylcholine receptor subtype in basal forebrain cholinergic neurons with high sensitivity to amyloid peptides. *J. Neurosci.*, **29**, 918–929.
- Lukyanetz, E.A., Piper, T.P. & Sihra, T.S. (1998) Calcineurin involvement in the regulation of high-threshold  $\text{Ca}^{2+}$  channels in NG108-15 (rodent neuroblastoma X glioma hybrid) cells. *J. Physiol.*, **510**, 371–385.
- Mansvelder, H.D., Keath, J.R. & McGehee, D.S. (2002) Synaptic mechanisms underlie nicotine-induced excitability of brain reward areas. *Neuron*, **33**, 905–919.
- McGee, R., Simpson, P., Christian, C., Mata, M., Nelson, P. & Nirenberg, M. (1978) Regulation of acetylcholine release from neuroblastoma X glioma hybrid cells. *Proc. Natl Acad. Sci. USA*, **75**, 1314–1318.
- McGehee, D.S., Heath, M.J., Gelber, S., Devay, P. & Role, L.W. (1995) Nicotine enhancement of fast excitatory synaptic transmission in CNS by presynaptic receptors. *Science*, **269**, 1692–1696.
- McLean, C.A., Cherny, R.A., Fraser, F.W., Fuller, S.J., Smith, M.J., Beyreuther, K., Bush, A.I. & Masters, C.L. (1999) Soluble pool of A $\beta$  amyloid as a determinant of severity of neurodegeneration in Alzheimer's disease. *Ann. Neurol.*, **46**, 860–866.
- Mehta, T.K., Dougherty, J.J., Wu, J., Choi, C.H., Khan, G.K. & Nichols, R.A. (2009) Defining presynaptic nicotinic receptors regulated by beta amyloid in mouse cortex and hippocampus with receptor null mutants. *J. Neurochem.*, **109**, 1452–1458.
- Nagele, R.G., D'Andrea, M.R., Anderson, W.J. & Wang, H.Y. (2002) Intracellular accumulation of  $\beta$ -amyloid<sub>1–42</sub> in neurons is facilitated by the  $\alpha 7$  nicotinic acetylcholine receptor in Alzheimer's disease. *Neuroscience*, **110**, 199–211.
- Nayak, S.V., Rondé, P., Spier, A.D., Lummis, S.C.R. & Nichols, R.A. (2000) Nicotinic receptors co-localize with 5-HT<sub>3</sub> serotonin receptors on striatal nerve terminals. *Neuropharmacology*, **39**, 2681–2690.
- Nayak, S.V., Dougherty, J.J., McIntosh, J.M. & Nichols, R.A. (2001)  $\text{Ca}^{2+}$  changes induced by different presynaptic nicotinic receptors in separate populations of individual striatal nerve terminals. *J. Neurochem.*, **76**, 1860–1870.
- Nelson, P., Christian, C. & Nirenberg, M. (1976) Synapse formation between clonal neuroblastoma X glioma hybrid cells and striated muscle cells. *Proc. Natl Acad. Sci. USA*, **73**, 123–127.
- Nichols, R.A., Dengler, A.F., Nakagawa, E.M., Bashkin, M., Paul, B.T., Wu, J. & Khan, G.M. (2007) A constitutive, transient receptor potential-like  $\text{Ca}^{2+}$  influx pathway in presynaptic nerve endings independent of voltage-gated  $\text{Ca}^{2+}$  channels and  $\text{Na}^{+}/\text{Ca}^{2+}$  exchange. *J. Biol. Chem.*, **282**, 36102–36111.
- Pettit, D.L., Shao, Z. & Yakel, J.L. (2001)  $\beta$ -Amyloid<sub>1–42</sub> peptide directly modulates nicotinic receptors in the rat hippocampal slice. *J. Neurosci.*, **21**, RC120.
- Puzzo, D., Privitera, L., Leznik, E., Fà, M., Staniszewski, A., Palmeri, A. & Arancio, O. (2008) Picomolar amyloid- $\beta$  positively modulates synaptic plasticity and memory in hippocampus. *J. Neurosci.*, **28**, 14537–14545.
- Role, L. & Berg, D.K. (1996) Nicotinic receptors in the development and modulation of CNS synapses. *Neuron*, **16**, 1077–1085.
- Rondé, P. & Nichols, R.A. (2001) Postsynaptic target regulates functional responses induced by 5-HT<sub>3</sub> serotonin receptors on axonal varicosities of NG108-15 hybrid neuroblastoma cells. *Neuroscience*, **102**, 979–987.
- Selkoe, D.J. (2002) Alzheimer's disease is a synaptic failure. *Science*, **298**, 789–791.
- Selkoe, D.J. (2008) Soluble oligomers of the amyloid  $\beta$ -protein impair synaptic plasticity and behavior. *Behav. Brain Res.*, **192**, 106–113.
- Selkoe, D.J. & Schenk, D. (2003) Alzheimer's disease: molecular understanding predicts amyloid-based therapeutics. *Annu. Rev. Pharmacol. Toxicol.*, **43**, 545–584.
- Sharma, G. & Vijayaraghavan, S. (2003) Modulation of presynaptic store calcium induces release of glutamate and postsynaptic firing. *Neuron*, **38**, 929–939.
- Sharma, G., Grybko, M. & Vijayaraghavan, S. (2008) Action potential-independent and nicotinic receptor-mediated concerted release of multiple quanta at hippocampal CA3-Mossy fiber synapses. *J. Neurosci.*, **28**, 2563–2575.
- Small, D.H., Maksud, D., Kerr, M.L., Ng, J., Hou, X., Chu, C., Mehrani, H., Unabia, S., Azari, M.F., Loiacono, R., Aguilar, M.I. & Chebib, M. (2007) The  $\beta$ -amyloid protein of Alzheimer's disease binds to membrane lipids but does not bind to the  $\alpha 7$  nicotinic acetylcholine receptor. *J. Neurochem.*, **101**, 1527–1538.
- Tsui-Pierchala, B.A., Encinas, M., Millbrandt, J. & Johnson, E.M. (2002) Lipid rafts in neuronal signaling and function. *Trends Neurosci.*, **25**, 412–417.
- Umbriaco, D., Watkins, K.C., Descarries, L., Cozzari, C. & Hartman, B.K. (1994) Ultrastructural and morphometric features of the acetylcholine innervation in adult rat parietal cortex: an electron microscopic study in serial sections. *J. Comp. Neurol.*, **348**, 351–373.
- Walsh, D.M. & Selkoe, D.J. (2007) A $\beta$  oligomers – a decade of discovery. *J. Neurochem.*, **101**, 1172–1184.
- Wang, H.Y., Lee, D.H.S., D'Andrea, M.R., Peterson, P.A., Shank, R.P. & Reitz, A.B. (2000a)  $\beta$ -amyloid<sub>1–42</sub> binds to  $\alpha 7$  nicotinic acetylcholine receptor with high affinity. *J. Biol. Chem.*, **275**, 5626–5632.
- Wang, H.Y., Lee, D.H., Davis, C.B. & Shank, R.P. (2000b) Amyloid peptide A $\beta$ <sub>1–42</sub> binds selectively and with picomolar affinity to  $\alpha 7$  nicotinic acetylcholine receptors. *J. Neurochem.*, **75**, 1155–1161.
- Wang, H.Y., Li, W., Benedetti, N.J. & Lee, D.H.S. (2003)  $\alpha 7$  nicotinic acetylcholine receptors mediate  $\beta$ -amyloid peptide-induced tau protein phosphorylation. *J. Biol. Chem.*, **278**, 31547–31553.
- Wang, H.Y., Stucky, A., Liu, J., Shen, C., Trocme-Thibierge, C. & Morain, P. (2009) Dissociating  $\beta$ -amyloid from  $\alpha 7$  nicotinic acetylcholine receptor by a novel therapeutic agent, S 24795, normalizes  $\alpha 7$  nicotinic acetylcholine and NMDA receptor function in Alzheimer's disease brain. *J. Neurosci.*, **29**, 10961–10973.
- Warnock, D.E., Roberts, C., Lutz, M.S., Blackburn, W.A., Young, W.W. & Baezinger, J.U. (1993) Determination of plasma membrane lipid mass and composition in cultured Chinese hamster ovary cells using high gradient magnetic affinity chromatography. *J. Biol. Chem.*, **268**, 10145–10153.
- Wu, J., Kuo, Y.P., George, A.A., Xu, L., Hu, L. & Lukas, R.J. (2004)  $\beta$ -amyloid directly inhibits human  $\alpha 4\beta 2$ -nicotinic acetylcholine receptors heterologously expressed in hum SH-EP1 cells. *J. Biol. Chem.*, **279**, 37842–37851.
- Wu, J., Khan, G.M. & Nichols, R.A. (2007) Dopamine release in prefrontal cortex in response to  $\beta$ -amyloid activation of  $\alpha 7$  \* nicotinic receptors. *Brain Res.*, **1182**, 82–89.
- Yakel, J.L., Shao, X.M. & Jackson, M.B. (1991) Activation and desensitization of the 5-HT<sub>3</sub> receptor in a rat glioma X mouse neuroblastoma hybrid cell. *J. Physiol.*, **436**, 293–308.
- Zidovetzki, R. & Levitan, I. (2007) Use of cyclodextrins to manipulate membrane cholesterol: evidence, misconceptions and control strategies. *Biochem. Biophys. Acta*, **1768**, 1311–1324.