

## Apolipoprotein E and Alzheimer disease: risk, mechanisms and therapy

Chia-Chen Liu, Takahisa Kanekiyo, Huaxi Xu and Guojun Bu

**Abstract** | Apolipoprotein E (Apo-E) is a major cholesterol carrier that supports lipid transport and injury repair in the brain. *APOE* polymorphic alleles are the main genetic determinants of Alzheimer disease (AD) risk: individuals carrying the  $\epsilon 4$  allele are at increased risk of AD compared with those carrying the more common  $\epsilon 3$  allele, whereas the  $\epsilon 2$  allele decreases risk. Presence of the *APOE*  $\epsilon 4$  allele is also associated with increased risk of cerebral amyloid angiopathy and age-related cognitive decline during normal ageing. Apo-E-lipoproteins bind to several cell-surface receptors to deliver lipids, and also to hydrophobic amyloid- $\beta$  (A $\beta$ ) peptide, which is thought to initiate toxic events that lead to synaptic dysfunction and neurodegeneration in AD. Apo-E isoforms differentially regulate A $\beta$  aggregation and clearance in the brain, and have distinct functions in regulating brain lipid transport, glucose metabolism, neuronal signalling, neuroinflammation, and mitochondrial function. In this Review, we describe current knowledge on Apo-E in the CNS, with a particular emphasis on the clinical and pathological features associated with carriers of different Apo-E isoforms. We also discuss A $\beta$ -dependent and A $\beta$ -independent mechanisms that link Apo-E4 status with AD risk, and consider how to design effective strategies for AD therapy by targeting Apo-E.

Liu, C.-C. *et al.* *Nat. Rev. Neurol.* **9**, 106–118 (2013); published online 8 January 2013; corrected online 26 February 2013; doi:10.1038/nrneurol.2012.263

### Introduction

Alzheimer disease (AD) is a progressive neurodegenerative disease associated with cognitive decline, and is the most common form of dementia in the elderly. Approximately 13% of people over the age of 65 years and 45% over the age of 85 years are estimated to have AD.<sup>1</sup> Mounting evidence from genetic, pathological, and functional studies has shown that an imbalance between production and clearance of amyloid- $\beta$  (A $\beta$ ) peptides in the brain results in accumulation and aggregation of A $\beta$ . Aggregates of toxic A $\beta$  in the form of soluble A $\beta$  oligomers, intraneuronal A $\beta$ , and amyloid plaques injure synapses and ultimately cause neurodegeneration and dementia.<sup>2,3</sup> The toxicity of A $\beta$  seems to depend on the presence of microtubule-associated protein tau,<sup>4</sup> the hyperphosphorylated forms of which aggregate and deposit in AD brains as neurofibrillary tangles. A $\beta$  is composed of 40 or 42 amino acids and is generated through proteolytic cleavage of amyloid precursor protein (APP).<sup>5</sup>

Early-onset, familial AD—which typically develops before the age of 65 years and accounts for only a small proportion (<1%) of AD cases<sup>2,3</sup>—is primarily caused by overproduction of A $\beta$  owing to mutations in either the *APP* gene or in genes encoding presenilin 1 (*PSEN1*) or presenilin 2 (*PSEN2*), which are essential components of the  $\gamma$ -secretase complexes responsible for cleavage and release of A $\beta$ . The majority of AD cases occur late in life (>65 years) and are commonly referred to as late-onset

AD (LOAD). Although multiple genetic and environmental risk factors are involved in LOAD pathogenesis, overall impairment in A $\beta$  clearance is probably a major contributor to disease development.<sup>6</sup> Genetically, the  $\epsilon 4$  allele of the apolipoprotein E (*APOE*) gene is the strongest risk factor for LOAD.<sup>7–9</sup> The human *APOE* gene exists as three polymorphic alleles— $\epsilon 2$ ,  $\epsilon 3$  and  $\epsilon 4$ —which have a worldwide frequency of 8.4%, 77.9% and 13.7%, respectively.<sup>10</sup> However, the frequency of the  $\epsilon 4$  allele is dramatically increased, to ~40%, in patients with AD.<sup>10</sup>

Apo-E regulates lipid homeostasis by mediating lipid transport from one tissue or cell type to another.<sup>11</sup> In peripheral tissues, Apo-E is primarily produced by the liver and macrophages, and mediates cholesterol metabolism in an isoform-dependent manner. Apo-E4 is associated with hyperlipidaemia and hypercholesterolaemia, which lead to atherosclerosis, coronary heart disease and stroke.<sup>11,12</sup> In the CNS, Apo-E is mainly produced by astrocytes, and transports cholesterol to neurons via Apo-E receptors, which are members of the low-density lipoprotein receptor (LDLR) family.<sup>8</sup>

Apo-E is composed of 299 amino acids and has a molecular mass of ~34 kDa.<sup>11</sup> Differences between the three Apo-E isoforms are limited to amino acid residues 112 and 158, where either cysteine or arginine is present (Figure 1): Apo-E2 (Cys112, Cys158), Apo-E3 (Cys112, Arg158), and Apo-E4 (Arg112, Arg158).<sup>11</sup> The single amino acid differences at these two positions affect the structure of Apo-E isoforms and influence their ability to bind lipids, receptors and A $\beta$ .<sup>13–15</sup> Human and animal studies clearly

Fujian Provincial Key Laboratory of Neurodegenerative Disease and Aging Research, Institute of Neuroscience, College of Medicine, Xiamen University, Xiamen, Fujian 361005, China (C.-C. Liu, H. Xu, G. Bu). Department of Neuroscience, Mayo Clinic, 4500 San Pablo Road, Jacksonville, FL 32224, USA (T. Kanekiyo).

Correspondence to: G. Bu  
bu.guojun@mayo.edu

### Competing interests

The authors declare no competing interests.

indicate that Apo-E isoforms differentially affect A $\beta$  aggregation and clearance. Several A $\beta$ -independent functions are also associated with Apo-E isoforms.

In this Review, we provide an overview of clinical evidence for the association between *APOE* genotypes and the risk of cognitive decline in AD, mild cognitive impairment (MCI) and other CNS diseases with a cognitive component, and discuss our current understanding of the mechanisms underlying Apo-E actions and Apo-E-targeted therapies.

## APOE genotypes, AD and cognition

### APOE $\epsilon$ 4 as a strong risk factor for AD

Genome-wide association studies have confirmed that the  $\epsilon$ 4 allele of *APOE* is the strongest genetic risk factor for AD.<sup>16,17</sup> The presence of this allele is associated with increased risk of both early-onset AD and LOAD.<sup>18,19</sup> A meta-analysis of clinical and autopsy-based studies demonstrated that, in white individuals, risk of AD was increased in individuals with one copy of the  $\epsilon$ 4 allele ( $\epsilon$ 2/ $\epsilon$ 4, OR 2.6;  $\epsilon$ 3/ $\epsilon$ 4, OR 3.2) or two copies ( $\epsilon$ 4/ $\epsilon$ 4, OR 14.9) compared with those with an  $\epsilon$ 3/ $\epsilon$ 3 genotype.<sup>10</sup> Conversely, the  $\epsilon$ 2 allele of *APOE* has a protective effect against AD: the risk of AD in individuals carrying *APOE*  $\epsilon$ 2/ $\epsilon$ 2 (OR 0.6) or  $\epsilon$ 2/ $\epsilon$ 3 (OR 0.6) is lower than in those carrying  $\epsilon$ 3/ $\epsilon$ 3.<sup>10</sup>

In population-based studies, the *APOE*  $\epsilon$ 4–AD association was weaker among African American ( $\epsilon$ 4/ $\epsilon$ 4, OR 5.7) and Hispanic ( $\epsilon$ 4/ $\epsilon$ 4, OR 2.2) populations, and was stronger in Japanese people ( $\epsilon$ 4/ $\epsilon$ 4, OR 33.1) compared with white individuals ( $\epsilon$ 4/ $\epsilon$ 4, OR 12.5).<sup>10</sup> *APOE*  $\epsilon$ 4 is associated with increased prevalence of AD and lower age of onset.<sup>7,10,20</sup> The frequency of AD and mean age at clinical onset are 91% and 68 years in  $\epsilon$ 4 homozygotes, 47% and 76 years in  $\epsilon$ 4 heterozygotes, and 20% and 84 years in  $\epsilon$ 4 noncarriers,<sup>7,20</sup> indicating that *APOE*  $\epsilon$ 4 confers dramatically increased risk of development of AD with an earlier age of onset in a gene dose-dependent manner (Table 1).


Genetic variants in the *TOMM40* (translocase of outer mitochondrial membrane 40 homologue) gene, which lies adjacent to the *APOE* gene on chromosome 19, have been implicated as a modulator of AD age-of-onset in *APOE*  $\epsilon$ 3 carriers.<sup>21</sup> A subsequent study, however, cast doubt on the strength of this association.<sup>22</sup> Whether the effects of *APOE* and *TOMM40* on AD risk, both genetically and functionally, are synergistic requires further investigation.

### Apo-E and A $\beta$ deposition

Apo-E has an important role in A $\beta$  metabolism (Figure 2). Studies show that *APOE* genotypes strongly affect deposition of A $\beta$  to form senile plaques and cause cerebral amyloid angiopathy (CAA)—two major hallmarks of amyloid pathology in AD brains.<sup>23</sup> Immunohistological evidence demonstrates that Apo-E is co-deposited in senile plaques in the brains of patients with AD.<sup>24</sup> A $\beta$  deposition in the form of senile plaques is more abundant in *APOE*  $\epsilon$ 4 carriers than in noncarriers.<sup>25–27</sup> The difference was most evident among individuals aged 50–59 years: 40.7% of *APOE*  $\epsilon$ 4 carriers had senile plaques compared with 8.2% of noncarriers.<sup>25</sup>

## Key points

- The  $\epsilon$ 4 allele of the apolipoprotein E (*APOE*) gene is the main genetic risk factor for Alzheimer disease (AD)
- *APOE*  $\epsilon$ 4 carriers have enhanced AD pathology, accelerated age-dependent cognitive decline and worse memory performance than do noncarriers
- Numerous structural and functional brain changes associated with AD pathogenesis are detected in *APOE*  $\epsilon$ 4 carriers before clinical symptoms become evident
- Apo-E affects amyloid- $\beta$  (A $\beta$ ) clearance, aggregation and deposition in an isoform-dependent manner
- Apo-E4 also contributes to AD pathogenesis by A $\beta$ -independent mechanisms that involve synaptic plasticity, cholesterol homeostasis, neurovascular functions, and neuroinflammation
- Apo-E-targeted AD therapy should focus on restoration of the physiological function of Apo-E through increased expression and lipidation, and inhibition of the detrimental effects of Apo-E4



**Figure 1** | *APOE*  $\epsilon$ 4 is a major genetic risk factor for AD. The Apo-E2, E3, and E4 isoforms, which are encoded by the  $\epsilon$ 2,  $\epsilon$ 3 and  $\epsilon$ 4 alleles of the *APOE* gene, respectively, differ from one another at amino acid residues 112 and/or 158 (grey circles). Apo-E has two structural domains: the N-terminal domain, which contains the receptor-binding region (residues 136–150), and the C-terminal domain, which contains the lipid-binding region (residues 244–272); the two domains are joined by a hinge region. A meta-analysis demonstrated a significant association between the  $\epsilon$ 4 allele of *APOE* and AD.<sup>10</sup> Abbreviations: AD, Alzheimer disease; Apo-E, apolipoprotein E.

In individuals with positive Pittsburgh compound B (PiB)-PET images, which indicate fibrillar aggregates of A $\beta$ ,<sup>28</sup> *APOE*  $\epsilon$ 4 was more common than in those with negative scans (65% versus 22%) in patients with AD.<sup>29</sup>

Fibrillar A $\beta$  deposition is often detected in the brains of elderly, cognitively normal individuals in a manner that depends on the presence of *APOE*  $\epsilon$ 4, although such an association is weaker than that in patients with AD.<sup>30</sup> In addition, *APOE*  $\epsilon$ 4 carriers have lower cerebrospinal fluid (CSF) A $\beta$ <sub>42</sub> levels and show more PiB binding on PET than do noncarriers, which reflect the presence of cerebral amyloid deposition and serve as potential biomarkers for AD.<sup>31,32</sup> PiB-positivity can be detected in cognitively normal *APOE*  $\epsilon$ 4 carriers at about 56 years of age, compared with around 76 years of age in noncarriers.<sup>33</sup> This difference suggests that *APOE*  $\epsilon$ 4 probably increases the risk of AD by initiating and accelerating A $\beta$  accumulation, aggregation and deposition in the brain. Although *APOE*  $\epsilon$ 2 reduces the risk of dementia,<sup>34</sup> both the  $\epsilon$ 2 and  $\epsilon$ 4 alleles of *APOE* increase amyloid burden compared with *APOE*  $\epsilon$ 3 in individuals older than 90 years, suggesting that the protective effects of *APOE*  $\epsilon$ 2 against AD might not be associated with A $\beta$  deposition.

**Table 1** | The effect of *APOE*  $\epsilon$ 4 on AD frequency and age at onset<sup>7</sup>

| Characteristic                     | <i>APOE</i> $\epsilon$ 4 noncarrier | <i>APOE</i> $\epsilon$ 4 heterozygous | <i>APOE</i> $\epsilon$ 4 homozygous |
|------------------------------------|-------------------------------------|---------------------------------------|-------------------------------------|
| AD frequency (%)                   | 20                                  | 47                                    | 91                                  |
| Mean age of clinical onset (years) | 84                                  | 76                                    | 68                                  |

Abbreviations: AD, Alzheimer disease; *APOE*  $\epsilon$ 4,  $\epsilon$ 4 allele of the apolipoprotein E gene.

*APOE*  $\epsilon$ 4 also shows an association with CAA and CAA-related haemorrhages.<sup>35,36</sup> CAA refers to the pathological condition in which amyloid spreads and deposits throughout the cerebral blood vessel walls,<sup>37</sup> and is frequently detected in AD.<sup>23</sup> Interestingly, although *APOE*  $\epsilon$ 2 is protective against AD, it is a risk factor for CAA-related haemorrhage, independently of AD, possibly by predisposing vessels to vasculopathic complications of CAA.<sup>36</sup>

### Prediction of AD in MCI

MCI is a transitional stage between normal ageing and dementia, and is associated with increased risk of AD.<sup>38</sup> The rate at which patients with amnesic MCI (aMCI) progress to clinically diagnosable AD is 10–15% per year, in contrast to a rate of 1–2% per year among healthy elderly individuals.<sup>39</sup> The prevalence of *APOE*  $\epsilon$ 4 is substantially higher in both aMCI and dys-executive MCI than in control individuals.<sup>40</sup> Patients with MCI who harbour *APOE*  $\epsilon$ 4 exhibit distinct cognitive profiles, which seem to resemble those of patients in the early stages of AD.<sup>41</sup> A case-control study reported poorer memory performance among patients with MCI who were carriers of *APOE*  $\epsilon$ 4 compared with noncarriers.<sup>42</sup> *APOE*  $\epsilon$ 4 is associated with impaired memory performance and increased risk of memory decline in middle-aged (40–59 years) and elderly (60–85 years) people with MCI.<sup>43,44</sup> Furthermore, patients with MCI who are carriers of *APOE*  $\epsilon$ 4 experience more-rapid decline in several cognitive and functional domains, and severity of the deficits is strongly associated with the *APOE*  $\epsilon$ 4 gene dose.<sup>41,45,46</sup> Importantly, the presence of *APOE*  $\epsilon$ 4 is associated with increased risk of progression from MCI to AD-type dementia.<sup>47–49</sup> Among individuals with aMCI, *APOE*  $\epsilon$ 4 carriers tend to be younger than noncarriers, consistent with younger age of AD onset in individuals with *APOE*  $\epsilon$ 4.<sup>50</sup> These findings indicate that the *APOE*  $\epsilon$ 4 genotype in patients with MCI can serve as a predictive factor for determination of clinical outcome and the risk of conversion to AD.

In patients with MCI, the adverse effects of *APOE*  $\epsilon$ 4 on cognitive functions correlate with the severity of neuronal pathology. Those who are carriers of *APOE*  $\epsilon$ 4 have lower CSF A $\beta$ <sub>42</sub> levels, higher tau levels and greater brain atrophy than do noncarriers.<sup>50</sup> Furthermore, patients with MCI who are PiB-positive are more likely to be *APOE*  $\epsilon$ 4 carriers and exhibit worse memory performance than are PiB-negative patients.<sup>51</sup> Other findings suggest, although not without controversy,<sup>52</sup> that *APOE*  $\epsilon$ 4 has considerable deleterious effects on memory performance<sup>42</sup> and might be used to predict

disease progression in combination with AD biomarkers and neuroimaging approaches.<sup>53</sup>

### Predicting cognitive decline in healthy cases

Healthy *APOE*  $\epsilon$ 4 carriers not diagnosed with MCI or AD show accelerated longitudinal decline in memory tests, which starts around the age of 55–60 years, revealing a possible pre-MCI state in this genetic subset of individuals.<sup>54,55</sup> This memory decline, despite ongoing normal clinical status, suggests that pathological changes in AD might manifest in the brain as early as the sixth decade of life.<sup>56,57</sup> Thus, *APOE*  $\epsilon$ 4 is associated with cognitive decline many years before cognitive impairment becomes clinically apparent.<sup>56,58</sup> Interestingly, *APOE*  $\epsilon$ 4 has differential effects on memory performance depending on age. Some studies in young adults and children have found evidence of better cognitive performance in *APOE*  $\epsilon$ 4 carriers than in noncarriers, which could suggest antagonistic pleiotropy,<sup>59–61</sup> in which *APOE*  $\epsilon$ 4 might offer benefits during development and early adulthood at the expense of more-rapid decline in cognitive function with ageing.<sup>62</sup>


Similar to the situation in patients with MCI, *APOE*  $\epsilon$ 4 is associated with enhanced amyloid pathology in cognitively normal people. The proportion of PiB-positive individuals follows a strong *APOE* allele-dependent pattern ( $\epsilon$ 4 >  $\epsilon$ 3 >  $\epsilon$ 2),<sup>25,63,64</sup> and *APOE*  $\epsilon$ 4 increases the amount of amyloid deposition in a gene-dose-dependent manner.<sup>30</sup>

### *APOE* $\epsilon$ 4 and other AD risk factors

The *APOE*  $\epsilon$ 4 genotype combines synergistically with atherosclerosis, peripheral vascular disease, and type 2 diabetes in contributing to an increased risk of AD.<sup>65,66</sup> *APOE*  $\epsilon$ 4 is a risk factor for cardiovascular disease, suggesting that this allele and cerebrovascular disease might have compounding effects on cognitive decline in AD.<sup>67</sup> Diabetes also increases the risk of AD, and the association is particularly strong among *APOE*  $\epsilon$ 4 carriers.<sup>66,68,69</sup> Patients with diabetes who are carriers of *APOE*  $\epsilon$ 4 have more neuritic plaques, neurofibrillary tangles and CAA than do noncarriers.<sup>66</sup> The combination of a diabetes-related factor—that is, hyperglycaemia, hyperinsulinaemia, and insulin resistance—and the *APOE*  $\epsilon$ 4 allele promotes neuritic plaque formation.<sup>69</sup> *APOE*  $\epsilon$ 4 seems to modify the risk of AD in patients with diabetes—a disease that directly or indirectly causes vascular and neuronal damage and further exacerbates AD pathology. Furthermore, recent research demonstrated that, independently of A $\beta$ , Apo-E4 triggers inflammatory cascades that cause neurovascular dysfunction, including blood-brain barrier breakdown, leakage of blood-derived toxic proteins into the brain and reduction in the length of small vessels.<sup>70</sup> This result suggests that Apo-E4-associated damage to vascular systems in the brain could have a key role in AD pathogenesis.

### *APOE* and traumatic brain injury

Increasing evidence has shown that *APOE*  $\epsilon$ 4 is associated with poorer outcomes following traumatic brain injury (TBI) compared with *APOE*  $\epsilon$ 2 and  $\epsilon$ 3 alleles, regardless



**Figure 2** | Apolipoprotein E and amyloid- $\beta$  metabolism in the brain. Major A $\beta$  clearance pathways include receptor-mediated uptake by neurons and glia, drainage into interstitial fluid or through the BBB, and proteolytic degradation by IDE and neprilysin. Impaired A $\beta$  clearance can cause accumulation in brain parenchyma, triggering formation of A $\beta$  oligomers and amyloid plaques. Perivascular A $\beta$  accumulation leads to CAA, which disrupts blood vessel function. Apo-E is primarily synthesized by astrocytes and microglia, and is lipidated by the ABCA1 transporter, forming lipoprotein particles. Lipidated Apo-E binds soluble A $\beta$  and facilitates A $\beta$  uptake through cell-surface receptors, including LRP1, LDLR, and HSPG.<sup>175,177</sup> Apo-E facilitates binding and internalization of soluble A $\beta$  by glial cells, disrupts A $\beta$  clearance at the BBB in an isoform-dependent manner (Apo-E4 > Apo-E3 > Apo-E2) and influences CAA pathogenesis. Abbreviations: A $\beta$ , amyloid- $\beta$ ; ABCA1, ATP-binding cassette A1; Apo-E, apolipoprotein E; BBB, blood–brain barrier; CAA, cerebral amyloid angiopathy; HSPG, heparan sulphate proteoglycan; IDE, insulin-degrading enzyme; LDLR, low-density lipoprotein receptor; LRP1, low-density lipoprotein receptor-related protein 1; LXR, liver X receptor.

of the severity of initial injury.<sup>71</sup> A meta-analysis demonstrated that the outcome of TBI at 6 months after injury is worse in *APOE*  $\epsilon$ 4 carriers than in noncarriers.<sup>72</sup> TBI is associated with increased risk of AD,<sup>73</sup> and such a risk is more evident in patients with *APOE*  $\epsilon$ 4.<sup>74</sup> Only 10% of *APOE*  $\epsilon$ 4 noncarriers with TBI have A $\beta$  plaque pathology, whereas 35% and 100% of TBI patients with one or two *APOE*  $\epsilon$ 4 alleles, respectively, possess A $\beta$  pathology.<sup>75</sup> The poorer outcomes associated with Apo-E4 might relate to the reduced ability of this isoform to repair and remodel synapses and protect neurons upon injury compared with Apo-E3.<sup>8</sup> These possibilities are currently under investigation.

### ***APOE* and vascular diseases**

Vascular cognitive impairment, which comprises clinical conditions with cerebrovasculature-derived cognitive disturbances including vascular dementia, is observed in approximately 8–15% of aged individuals with cognitive dysfunction in Western clinic-based series.<sup>76</sup> A recent meta-analysis has shown evidence of increased risk of vascular dementia in individuals with *APOE*  $\epsilon$ 4 compared with *APOE*  $\epsilon$ 3 (OR 1.72).<sup>77</sup> Several studies suggest that the contribution of *APOE*  $\epsilon$ 4 to risk of vascular cognitive impairment is independent of other vascular risk factors including hypertension, dyslipidaemia and atherogenesis,<sup>78</sup> whereas another report shows that age-related cognitive decline among *APOE*  $\epsilon$ 4 carriers is induced by brain damage owing to increased blood pressure.<sup>79</sup> In addition, *APOE*  $\epsilon$ 4 is associated with poor


outcome after subarachnoid haemorrhage,<sup>80</sup> and is a strong risk factor for CAA-related intracranial haemorrhage.<sup>81</sup> These results suggest that *APOE*  $\epsilon$ 4 is closely associated with neurovascular dysfunctions.

### ***APOE* and other types of dementia**

Lewy body disease is thought to be the second most common form of dementia, and comprises a spectrum of diseases that includes Parkinson disease (PD), PD-associated dementia, and dementia with Lewy bodies (DLB). Clinical and pathological features of PD and AD frequently overlap. Most studies, however, have failed to report associations between *APOE*  $\epsilon$ 4 and susceptibility to PD and PD-associated dementia.<sup>82,83</sup>

DLB also shares clinical and pathological characteristics with AD and PD,<sup>84</sup> and several reports have shown that *APOE*  $\epsilon$ 4 increases risk of DLB.<sup>85</sup> Immunohistochemical analysis showed that deposition of Lewy bodies in patients with DLB who are *APOE*  $\epsilon$ 4 carriers is substantially more abundant than in those who are noncarriers.<sup>86</sup> As Lewy bodies are considerably increased in the cerebral cortex of DLB patients with A $\beta$  deposition,<sup>87</sup> the strong association between amyloid pathology and the pathology of Lewy body disease could explain why *APOE*  $\epsilon$ 4 increases risk of DLB. *APOE*  $\epsilon$ 4 might also be a risk factor for frontotemporal dementia,<sup>88</sup> although the pathophysiological role of Apo-E in this disease requires further investigation. The *APOE* genotypes do not seem to influence the risks of Huntington disease<sup>89</sup> nor amyotrophic lateral sclerosis.<sup>90</sup>





**Figure 3** | The role of apolipoprotein E4 in Alzheimer disease pathogenesis. Apo-E4 confers toxic gain of function, loss of neuroprotective function, or both, in the pathogenesis of Alzheimer disease. Key functional differences between Apo-E4 and Apo-E3 are illustrated. Abbreviations: Aβ, amyloid-β; Apo-E, apolipoprotein E.

### Mechanisms of Apo-E isoforms in AD

*APOE* ε4 confers a gain of toxic functions, a loss of neuroprotective functions or both in the pathogenesis of AD (Figure 3).

#### Aβ metabolism and aggregation

Studies in humans and transgenic mice showed that brain Aβ levels and amyloid plaque loads are Apo-E isoform-dependent (ε4 > ε3 > ε2),<sup>30,63,91</sup> suggesting an important role for Apo-E in modulating Aβ metabolism, aggregation, and deposition. Apo-E4 is less efficient in Aβ clearance than is Apo-E3 in young and old amyloid mouse models that express human Apo-E isoforms.<sup>63</sup> Additionally, Apo-E isoforms differentially regulate cholesterol levels, which have been shown to modulate γ-secretase activity and Aβ production.<sup>92</sup> Several studies reported an *APOE* genotype-dependent effect on CSF and brain Apo-E levels (ε4 < ε3 < ε2) in Apo-E-targeted-replacement (Apo-E-TR) mice, in which the mouse *Apoe* gene is replaced with human *APOE* isoforms.<sup>91,93,94</sup> This result suggests that lower levels of total Apo-E exhibited by *APOE* ε4 carriers might contribute to disease progression. However, whether human Apo-E isoform status affects CSF and brain Apo-E protein levels in healthy individuals and patients with AD remains to be established.<sup>95,96</sup>

Apo-E-knockout mice clear Aβ from the brain faster than do control mice.<sup>97</sup> Stimulation of liver X receptors (LXRs)<sup>98,99</sup> or the retinoid X receptor (RXR)<sup>100</sup> facilitates Aβ clearance, probably by increasing Apo-E levels and lipidation. Further investigation is needed to determine whether Apo-E levels are directly associated with Aβ clearance. In addition, a recent study showed that lack of one copy of ATP-binding cassette transporter A1 (ABCA1), which shuttles lipids to Apo-E, impairs Aβ clearance and exacerbates amyloid deposition and memory deficits in Apo-E4-TR mice, but not in Apo-E3-TR mice.<sup>101</sup> This result suggests that Apo-E isoforms exhibit differential

lipidation status, which affects Aβ clearance in an isoform-dependent manner. Alternatively, Apo-E-lipoprotein particles may sequester Aβ and promote cellular uptake and degradation of Apo-E-Aβ complexes.<sup>102</sup>

Apo-E4-lipoproteins bind Aβ with lower affinity than do Apo-E3-lipoproteins,<sup>103</sup> suggesting that Apo-E4 might be less efficient in mediating Aβ clearance. In addition, Apo-E might modulate Aβ removal from the brain to the systemic circulation by transporting Aβ across the blood-brain barrier. In this respect, Apo-E impedes Aβ clearance at the blood-brain barrier in an isoform-specific fashion (Apo-E4 > Apo-E3 and Apo-E2).<sup>104</sup> Finally, studies in microglia have shown that Apo-E3 promotes enzyme-mediated degradation of Aβ more efficiently than does Apo-E4.<sup>105</sup> Together, these studies suggest that Apo-E4 inhibits Aβ clearance and/or is less efficient in mediating Aβ clearance compared with Apo-E3 and Apo-E2.

Apo-E also seems to regulate Aβ aggregation and deposition. An important study showed that deletion of the mouse *Apoe* gene essentially eliminates deposition of fibrillar Aβ in amyloid mouse models.<sup>106</sup> Given that Apo-E is co-deposited with Aβ in human AD brains,<sup>24</sup> it is possible that Apo-E promotes Aβ aggregation and deposition in an isoform-dependent manner. The exact mechanisms by which Apo-E isoforms differentially regulate Aβ aggregation and deposition require further investigation.

#### Brain activity and atrophy

AD is associated with both functional abnormalities of the hippocampus and cortical atrophy in the memory network.<sup>107,108</sup> Patients with AD or MCI who are *APOE* ε4 carriers exhibit greater medial temporal lobe atrophy, particularly in the hippocampal area.<sup>41,109,110</sup> Structural MRI studies found that, compared with noncarriers, *APOE* ε4 carriers have accelerated age-related reduction in cortical thickness and hippocampal volume that are tightly coupled to decline in cognitive performance.<sup>111–113</sup>

Functional MRI (fMRI) studies reported that Apo-E4 disrupts resting-state fMRI connectivity and the balance between brain networks, in the absence of amyloid pathology.<sup>114,115</sup> Furthermore, cognitively normal *APOE* ε4 carriers have elevated resting-state activity in the default mode network—a network that is preferentially affected early in AD—and higher hippocampal activation during memory tasks.<sup>116–118</sup> Such changes have been hypothesized to represent a compensatory response by *APOE* ε4 carriers in which increased cognitive effort is required to achieve an equivalent level of performance to that of noncarriers.<sup>116,118</sup>

Elevated baseline activity in brain networks of *APOE* ε4 carriers could potentially contribute to increased Aβ production, as Aβ levels are regulated by neuronal activity.<sup>119,120</sup> Interestingly, in adults who do not have dementia, increased hippocampal activity was associated with reduced cortical thickness in the medial temporal lobe and brain regions that are vulnerable to AD pathology.<sup>121</sup> Studies suggested that hippocampal hyperactivity might represent impending synaptic dysfunction and incipient cognitive decline.<sup>122</sup> Interestingly, another study showed

a reduction of posterior default mode network connectivity in *APOE*  $\epsilon 4$  carriers in cognitively normal elderly people, implying that *APOE*  $\epsilon 4$  carriers exhibit more-rapid decline in connectivity of this network than do noncarriers as they age.<sup>115</sup>

<sup>18</sup>F-fluorodeoxyglucose PET imaging, which measures cerebral metabolic rates of glucose as a proxy for neuronal activity, correlates with disease progression and predicts histopathological diagnosis in AD.<sup>123</sup> Mounting evidence suggests that *APOE*  $\epsilon 4$  carriers exhibit lower cerebral glucose metabolism.<sup>124–126</sup> Healthy adults with *APOE*  $\epsilon 4$  show altered patterns of brain metabolism both at rest and during cognitive challenges compared with noncarriers.<sup>126,127</sup> Representative studies illustrating the association of Apo-E4 isoform with altered brain metabolism and activity, memory decline, and amyloid pathology in cognitively normal people are shown in Figure 4. Improved understanding of the mechanisms of Apo-E4-related brain activity changes, brain atrophy and reduced metabolism should help to explain why Apo-E4 is a risk factor for cognitive decline and AD.

### Tau phosphorylation and neurotoxicity

Apo-E is produced primarily by astrocytes and microglia. Neuronal Apo-E expression can, however, be induced in response to stress or injury, probably for the purpose of neuronal repair and remodelling.<sup>128,129</sup> A truncated fragment of Apo-E4, resulting from proteolytic cleavage of Apo-E following stress or injury, increases tau hyperphosphorylation, cytoskeletal disruption and mitochondrial dysfunction.<sup>128,130,131</sup> Apo-E4 also exacerbates neurotoxicity triggered by A $\beta$  and other insults.<sup>128,131</sup>

A recent study showed that neurons in patients with temporal lobe epilepsy who harbour *APOE*  $\epsilon 4$  are less resilient to the damaging hyperexcitability and more susceptible to A $\beta$  toxicity than are those in *APOE*  $\epsilon 3$  carriers,<sup>132</sup> suggesting that Apo-E3 might confer a neuroprotective advantage over Apo-E4 against neuronal stress. Interestingly, astrocyte-derived Apo-E4 has neuroprotective effects against excitotoxic injuries, whereas neuronal expression of Apo-E4 promotes excitotoxic cell death. This result suggests that Apo-E derived from various cellular sources might exhibit different physiological and pathological activity.<sup>133</sup>

### Lipid metabolism

Abnormal lipid metabolism is strongly related to the pathogenesis of AD. In the CNS, Apo-E mediates neuronal delivery of cholesterol, which is an essential component for axonal growth, synaptic formation and remodelling—events that are crucial for learning, memory formation and neuronal repair.<sup>134,135</sup> Brain cholesterol levels are substantially reduced in hippocampal and cortical areas in patients with AD compared with age-matched controls.<sup>136</sup> Preferential degradation of Apo-E4 relative to Apo-E3 in astrocytes in transgenic animals has been proposed to result in low levels of Apo-E in the brain and CSF and reduced capacity for neuronal delivery of cholesterol, suggesting that low levels of total Apo-E exhibited by *APOE*  $\epsilon 4$  carriers may directly contribute to disease

progression.<sup>93</sup> Apo-E4 is also less efficient than Apo-E3 in transporting brain cholesterol.<sup>137</sup> Moreover, Apo-E4-TR mice have abnormal cholesterol levels and impaired lipid metabolism.<sup>138</sup> Insufficient levels of Apo-E and/or impaired Apo-E function in carriers of the  $\epsilon 4$  allele might, therefore, lead to aberrant CNS cholesterol homeostasis and neuronal health, which contribute to AD risk.

### Synaptic plasticity and spine integrity

Synaptic failure is an early pathological feature of AD.<sup>139,140</sup> Increasing evidence demonstrates that Apo-E isoforms differentially regulate synaptic plasticity and repair.<sup>141,142</sup> In AD and healthy aged controls, *APOE*  $\epsilon 4$  gene dosage correlates inversely with dendritic spine density in the hippocampus.<sup>143</sup> Apo-E4-TR mice also have lower dendritic spine density and length compared with Apo-E3-TR mice.<sup>144,145</sup> Apo-E3, but not Apo-E4, prevents loss of synaptic networks induced by A $\beta$  oligomers.<sup>146</sup> Apo-E isoforms also differentially regulate dendritic spines during ageing.<sup>143,147</sup> The age-dependence of these differences implies that the effects of Apo-E isoforms on neuronal integrity might relate to increased risk of dementia in aged *APOE*  $\epsilon 4$  carriers.


Reduced synaptic transmission was observed in 1-month-old Apo-E4-TR mice compared with Apo-E3-TR mice, suggesting that Apo-E4 may also contribute to functional deficits early in development, which could account for alteration of neuronal circuitry that eventually results in cognitive disorders later in life.<sup>147</sup> In addition, Apo-E4 selectively impairs Apo-E receptor trafficking and signalling, as well as glutamate receptor function and synaptic plasticity.<sup>141</sup> Together, these findings suggest that the effect of *APOE*  $\epsilon 4$  genotype on risk of AD might be mediated, at least in part, through direct effects on synaptic function.

### Neuroinflammation

Neuroinflammation contributes to neuronal damage in the brain and is implicated in AD pathogenesis.<sup>148</sup> Apo-E colocalizes with plaque-associated amyloid and microglia, suggesting a role for Apo-E in the innate immune response in AD. Lack of Apo-E in mice is associated with increased inflammation in response to A $\beta$ ,<sup>149,150</sup> but Apo-E isoforms might differently regulate the innate immune response.<sup>151</sup> Apo-E4 seems to have pro-inflammatory and/or reduced anti-inflammatory functions, which could further exacerbate AD pathology. For example, Apo-E4-TR mice exhibit greater inflammatory responses to lipopolysaccharide compared with Apo-E3-TR mice.<sup>152</sup> In addition, young *APOE*  $\epsilon 4$  carriers show an increased inflammatory response that may relate to AD risk later in life.<sup>153</sup> Consistent with this notion, non-steroidal anti-inflammatory drugs were shown to reduce AD risk only in *APOE*  $\epsilon 4$  carriers,<sup>154</sup> suggesting that *APOE* genotype might determine the effect of anti-inflammatory medications for AD.

### Neurogenesis

Hippocampal neurogenesis has an important role in structural plasticity and maintenance of brain networks.



**Figure 4** | Brain function, neuropathology and memory in cognitively normal APOE  $\epsilon$ 4 carriers. **a** | On  $^{18}\text{F}$ -FDG PET imaging, cognitively normal APOE  $\epsilon$ 4 carriers have lower glucose metabolism than do noncarriers. **b** | APOE  $\epsilon$ 4 carriers exhibit greater increase in functional MRI signal in brain regions associated with task performance, and increases in additional regions compared with APOE  $\epsilon$ 3 carriers. **c** | Age-related memory decline occurs more rapidly in APOE  $\epsilon$ 4 carriers than noncarriers, starting from age 55–60 years. **d** | APOE  $\epsilon$ 4 carriers show increased cerebral A $\beta$  deposition, becoming more pronounced with age. Increased PiB binding and reduced CSF A $\beta$ <sub>42</sub> levels reflect cerebral amyloid deposition. Abbreviations: A $\beta$ , amyloid- $\beta$ ; APOE, apolipoprotein E; AVLT-LTM, long-term memory score of the Auditory Verbal Learning Test; CSF, cerebrospinal fluid; PiB, Pittsburgh compound B. Part a is modified, with permission from the National Academy of Science, USA © Small, G. W. *et al. Proc. Natl Acad. Sci. USA* **97**, 6037–6042 (2000). Part b is modified, with permission, from the Massachusetts Medical Society © Bookheimer, S. Y. *et al. N. Engl. J. Med.* **343**, 450–456 (2000). Part c is modified, with permission from the Massachusetts Medical Society © Caselli, R. J. *et al. N. Engl. J. Med.* **361**, 255–263 (2009). Part d is modified, with permission, from John Wiley and Sons © Morris, J. C. *et al. Ann. Neurol.* **67**, 122–131 (2010).

Dysfunctional neurogenesis resulting from early disease manifestations could, therefore, exacerbate neuronal vulnerability to AD and contribute to memory impairment.<sup>155</sup> Apo-E is required for maintenance of the neural stem or progenitor cell pool in the adult dentate gyrus region of the hippocampus.<sup>156</sup> In Apo-E-TR mice, Apo-E4 inhibits hippocampal neurogenesis by impairing maturation of hilar  $\gamma$ -aminobutyric acid-containing interneurons, which contributes to learning and memory deficits.<sup>157,158</sup> These results demonstrate an important pathological role of Apo-E4 in impairment of neurogenesis, which might contribute to AD pathogenesis.

### Apo-E as a therapeutic target in AD

Most therapeutic approaches for AD target the A $\beta$  pathway. With the recent failure of clinical trials of drugs targeting solely A $\beta$ , an urgent need exists to define new targets and develop alternative therapeutic strategies to treat AD. As APOE genotype determines AD risk, and

Apo-E has crucial roles in cognition, Apo-E might offer an attractive alternative target for AD therapy. APOE genotype status could be included in clinical trial enrolment criteria, as some therapies might be effective only in specific APOE genotypes. Here, we briefly discuss several approaches that are currently being explored (Table 2).

### APOE genotype and A $\beta$ immunotherapy

Recent phase III trials of immunotherapy have shown that bapineuzumab, an antibody that targets the N-terminus of A $\beta$ , prevents A $\beta$  deposition in the brains of APOE  $\epsilon$ 4 carriers with mild or moderate AD, but not noncarriers.<sup>159,160</sup> Bapineuzumab also lowers levels of phosphorylated tau in the CSF of both APOE  $\epsilon$ 4 carriers and noncarriers.<sup>159,160</sup> These reports suggest that A $\beta$  immunotherapy is useful to eliminate A $\beta$  from the brains of patients with AD and that its effect is likely to depend on Apo-E isoforms. Major adverse effects of bapineuzumab—namely, vasogenic cerebral oedema and microhaemorrhage—occur more

**Table 2** | Apo-E-targeted strategies for treatment of Alzheimer disease

| Strategy                                      | Rationale   | Examples  |
|---|---|---|
| <b>Pharmacological approaches</b>             |   |   |
| Modulate Apo-E levels                         | Promotes A $\beta$ clearance, lipid homeostasis and synaptic function                                 | LXR and RXR agonists, small molecules   |
| Increase ABCA1 expression                     | Promotes Apo-E lipidation and stabilizes Apo-E, thereby decreasing amyloid deposition                 | LXR and RXR agonists, small molecules   |
| Disrupt Apo-E–A $\beta$ interaction           | Reduces A $\beta$ aggregation and deposition  | A $\beta$ 12-28P, small molecule inhibitors, Apo-E-specific antibody  |
| Stimulate conversion of Apo-E4 to Apo-E3      | Increases Apo-E3-associated protective functions and decreases Apo-E4-related toxic effects           | Small molecules (for example, disulphonate and monosulphoalkyl)   |
| Restore Apo-E functions                       | Increases Apo-E protective functions and decreases neuroinflammation                                  | Apo-E-mimetic peptide   |
| Block Apo-E fragmentation                     | Decreases tau pathology and prevents toxicity to mitochondria   | Inhibitors of specific proteinases involved in Apo-E fragmentation  |
| Increase LRP1 and/or LDLR levels              | Enhances A $\beta$ clearance, cholesterol transport and synaptic plasticity                           | Small molecules   |
| Increase Apo-E receptor 2 and/or VLDLR levels | Increases Apo-E signalling and synaptic plasticity  | Small molecules   |
| Restore brain vascular integrity              | Eliminates Apo-E4-mediated blood–brain barrier breakdown and leakage of blood-derived toxic molecules | Cyclosporine A  |
| <b>Nonpharmacological approaches</b>          |   |   |
| APOE genotyping prior to immunotherapy        | Helps to predict clinical outcome of A $\beta$ -targeted or other therapies                           | A $\beta$ immunotherapy might be more effective in APOE $\epsilon$ 4 carriers or noncarriers  |
| Preventive care                               | Maintains healthy brain vasculature function  | Physical exercise, intellectual activities (for example, puzzles), social connections (for example, calling family and friends), healthy diet |

Abbreviations: A $\beta$ , amyloid- $\beta$ ; ABCA1, ATP-binding cassette transporter A1; AD, Alzheimer disease; Apo-E, apolipoprotein E; LDLR, low-density lipoprotein receptor; LRP1, low-density lipoprotein receptor-related protein 1; LXR, liver X receptor; RXR, retinoid X receptor; VLDLR, very-low-density lipoprotein receptor.

frequently in *APOE*  $\epsilon$ 4 carriers than in noncarriers.<sup>159,160</sup> Although bapineuzumab failed to prevent cognitive and functional decline in these clinical trials, a combination of A $\beta$  immunotherapy and an Apo-E-targeted approach might lead to more-effective therapeutic strategies.

### Prevention of cognitive decline in $\epsilon$ 4 carriers

A prospective study of a cognitively normal cohort showed that risk of dementia in *APOE*  $\epsilon$ 4 carriers is negatively associated with high education, high level of leisure activities, and absence of vascular risk factors.<sup>161</sup> A recent study demonstrated that physical exercise was strongly associated with reduced PiB-positivity in cognitively normal *APOE*  $\epsilon$ 4 carriers,<sup>31</sup> indicating that a sedentary lifestyle in *APOE*  $\epsilon$ 4 carriers might increase the risk of amyloid deposition. Such studies indicate that high education, active leisure activities and exercise, and maintenance of vascular health could be beneficial in reducing the risk of AD and cognitive decline, particularly in *APOE*  $\epsilon$ 4 carriers.

### Regulation of Apo-E expression

Apo-E levels in CSF and plasma tend to be lower in patients with AD than in healthy individuals, although such findings remain controversial.<sup>162,163</sup> Thus, increasing the expression of Apo-E in all *APOE* genotypes may prevent or slow progression of AD through acceleration of A $\beta$  metabolism and promotion of Apo-E functions in

lipid metabolism and synaptic support. Compounds that increase brain Apo-E expression can be identified through comprehensive drug screening. Given that expression of Apo-E is controlled by peroxisome proliferator-activated receptor- $\gamma$  and LXRs (which form complexes with RXRs),<sup>100,164</sup> agonists or antagonists of these nuclear receptors are potential candidates as Apo-E modulators. Indeed, recent work has demonstrated that oral administration of an RXR agonist, bexarotene, to an amyloid mouse model decreases A $\beta$  plaque deposition and improves cognitive function in an Apo-E-dependent manner.<sup>100</sup> The LXR agonist TO901317 also increases Apo-E levels in the brain, facilitates clearance of A $\beta$ <sub>42</sub>, and reverses contextual memory deficit in amyloid mouse models.<sup>98,99</sup>

In addition to Apo-E, LXRs also regulate ABCA1, which promotes cholesterol efflux.<sup>165</sup> Consequently, reduction of amyloid burden by the LXR agonist GW3965 depends on expression of ABCA1 in amyloid mouse models.<sup>166</sup> These results suggest that upregulation of lipidated Apo-E might be necessary to maximize therapeutic effects in AD. These studies did not, however, assess the effect of increasing human Apo-E3 or Apo-E4 specifically. Because A $\beta$  deposition is greater in *APP*-transgenic mice expressing mouse Apo-E than in those expressing human Apo-E isoforms,<sup>167</sup> further studies are needed to confirm the therapeutic effect of modulating the level of human Apo-E isoforms. In addition, whether increasing Apo-E4 is beneficial or harmful in



AD brains remains unclear, and the effects might depend on age and disease status. Toxic functions associated with Apo-E4 suggest that lowering Apo-E4 expression might be beneficial in *APOE*  $\epsilon$ 4 carriers with cognitive decline during MCI and AD. Additional preclinical studies are needed to test potential beneficial or harmful effects of increasing or decreasing Apo-E expression, particularly with regard to Apo-E isoforms.

### Blockade of Apo-E–A $\beta$ interaction

Apo-E is required for deposition of A $\beta$  fibrils in amyloid mouse models.<sup>106</sup> Recent studies have demonstrated that haploinsufficiency of human *APOE* results in significantly decreased amyloid plaque deposition in amyloid mouse models regardless of *APOE* isoform status.<sup>168,169</sup> Thus, disruption of the interaction between Apo-E and A $\beta$  might reduce A $\beta$  aggregation and deposition, and should be considered as a therapeutic approach. A $\beta$  interacts with Apo-E through amino acid residues 12–28. A synthetic peptide mimicking this sequence, A $\beta$ 12–28P, reduces A $\beta$  deposition and ameliorates memory deficits in amyloid mouse models.<sup>170</sup> Blocking the Apo-E–A $\beta$  interaction using A $\beta$ -mimicking peptides could, therefore, be an effective approach for treatment of AD. Screening assays can also be used to identify compounds or Apo-E-specific antibodies that block Apo-E–A $\beta$  interaction. These approaches should be assessed carefully because they could disrupt Apo-E–lipid interactions and the associated beneficial functions of Apo-E.

### Other Apo-E-based therapeutic approaches

Apo-E4 is structurally different from Apo-E2 and Apo-E3 owing to different domain interactions,<sup>131</sup> and this difference probably contributes to Apo-E4 isoform-specific harmful effects. Modification of the structure of Apo-E4 to form an Apo-E3-like molecule might, therefore, be an interesting approach to ameliorate these harmful effects. Indeed, several molecules that bind to Apo-E4 and interfere with domain interactions between the N-terminus and C-terminus have been found. GIND-25 (disulpho-nate) and GIND-105 (monosulphoalkyl) are good candidates because they decrease A $\beta$  production induced by Apo-E4 to a level similar to that induced by Apo-E3.<sup>131</sup> CB9032258 (a phthalazinone analogue) and its derivatives disrupt Apo-E4 domain interaction and restore functional activities of Apo-E4 in neurons.<sup>171</sup>

An Apo-E-mimetic peptide containing the receptor-binding region suppresses neuronal cell death and calcium influx associated with *N*-methyl-D-aspartate exposure *in vitro*.<sup>172</sup> COG112, a chimeric peptide containing the receptor-binding region, is also reported to improve symptoms in mouse models of multiple sclerosis<sup>173</sup> and sciatic nerve crush<sup>174</sup> through modulation of inflammatory responses. The effects of these peptides on AD pathogenesis are unknown, however, because they do not contain A $\beta$ -interacting or lipid-binding regions.<sup>13</sup>

Apo-E receptors are also potential targets for AD therapy. For example, low-density lipoprotein receptor-related protein 1 and low-density lipoprotein receptor have crucial roles in brain lipid metabolism and A $\beta$

clearance (Figure 2).<sup>175–177</sup> Apo-E receptor 2 and very low-density lipoprotein receptor are essential for reelin signaling, which is important for neuronal migration during development and synaptic plasticity in adult brains.<sup>178</sup> Modulation of expression of these Apo-E receptors in AD brains might, therefore, restore lipid homeostasis and synaptic plasticity, and augment A $\beta$  clearance.<sup>8,178</sup> Although Apo-E-based therapies are still in early stages of development, they offer great promise in the fight against AD. Clinical trials to further evaluate therapeutic potential of Apo-E-based strategies are needed, with an eventual goal to develop curative and/or protective treatments for AD.

### Conclusions

Work summarized in this Review highlights clinical evidence for the association between *APOE*  $\epsilon$ 4, AD and cognitive decline. Although the presence of *APOE*  $\epsilon$ 4 does not necessarily entail disease development, this genetic isoform probably accelerates the rate of disease conversion and progression. In particular, the effects of *APOE*  $\epsilon$ 4 on brain network connectivity, memory performance, and rate of cognitive decline are age-dependent in patients with AD and cognitively normal individuals. Thus, understanding the potential pathogenic link between *APOE*  $\epsilon$ 4 and cognitive function might enable earlier identification of people at increased risk of AD. In combination with other putative AD biomarkers—such as MRI scans, PiB-PET, and measurements of CSF A $\beta$  and tau—*APOE* allele status could add predictive value to clinical diagnosis and evaluation of treatment efficacy.

Mechanistically, Apo-E4 seems to increase risk of AD and cognitive decline through both A $\beta$ -dependent and A $\beta$ -independent pathways. Apo-E isoforms differentially regulate A $\beta$  production, aggregation and clearance. Independently of A $\beta$ , Apo-E4 might be less efficient than Apo-E3 and Apo-E2 in delivering cholesterol and essential lipids for maintenance of synaptic integrity and plasticity. In addition, Apo-E is a crucial regulator of the innate immune system, with Apo-E4 promoting proinflammatory responses that could exacerbate AD pathogenesis. Finally, Apo-E isoforms have differential roles in maintaining vascular health—roles that are crucial given that vascular pathology is strongly associated with AD. Elucidation of the contribution of Apo-E4 to AD pathogenesis is a considerable challenge, but one that affords the potential to assist in combating AD.

### Review criteria

This Review was based on searches of the PubMed database using the following terms: “apolipoprotein E”, “cognitive decline”, “Alzheimer disease”, “amyloid beta”, “synaptic plasticity”, “cerebral amyloid angiopathy”, “mild cognitive impairment”, “cholesterol”, “brain activity”, “cerebrovascular diseases”, “brain metabolism”, “neurogenesis”, “brain atrophy”, “neuroinflammation”, “tau” and “traumatic brain injury”. Only articles published in English were retrieved. Full-text papers were available for most of the articles that were chosen for review, and the references of these articles were searched for further relevant material.

1. Alzheimer's Association. 2012 Alzheimer's disease facts and figures. *Alzheimers Dement.* **8**, 131–168 (2012).
2. Hardy, J. & Selkoe, D. J. The amyloid hypothesis of Alzheimer's disease: progress and problems on the road to therapeutics. *Science* **297**, 353–356 (2002).
3. Blennow, K., de Leon, M. J. & Zetterberg, H. Alzheimer's disease. *Lancet* **368**, 387–403 (2006).
4. Roberson, E. D. et al. Reducing endogenous tau ameliorates amyloid  $\beta$ -induced deficits in an Alzheimer's disease mouse model. *Science* **316**, 750–754 (2007).
5. Zheng, H. & Koo, E. Biology and pathophysiology of the amyloid precursor protein. *Mol. Neurodegener.* **6**, 27 (2011).
6. Mawuenyega, K. G. et al. Decreased clearance of CNS  $\beta$ -amyloid in Alzheimer's disease. *Science* **330**, 1774 (2010).
7. Corder, E. H. et al. Gene dose of apolipoprotein E type 4 allele and the risk of Alzheimer's disease in late onset families. *Science* **261**, 921–923 (1993).
8. Bu, G. Apolipoprotein E and its receptors in Alzheimer's disease: pathways, pathogenesis and therapy. *Nat. Rev. Neurosci.* **10**, 333–344 (2009).
9. Huang, Y. & Mucke, L. Alzheimer mechanisms and therapeutic strategies. *Cell* **148**, 1204–1222 (2012).
10. Farrer, L. A. et al. Effects of age, sex, and ethnicity on the association between apolipoprotein E genotype and Alzheimer disease: a meta-analysis. *JAMA* **278**, 1349–1356 (1997).
11. Mahley, R. W. & Rall, S. C. Jr. Apolipoprotein E: far more than a lipid transport protein. *Annu. Rev. Genomics Hum. Genet.* **1**, 507–537 (2000).
12. Lahoz, C. et al. Apolipoprotein E genotype and cardiovascular disease in the Framingham Heart Study. *Atherosclerosis* **154**, 529–537 (2001).
13. Frieden, C. & Garai, K. Structural differences between apoE3 and apoE4 may be useful in developing therapeutic agents for Alzheimer's disease. *Proc. Natl Acad. Sci. USA* **109**, 8913–8918 (2012).
14. Chen, J., Li, Q. & Wang, J. Topology of human apolipoprotein E3 uniquely regulates its diverse biological functions. *Proc. Natl Acad. Sci. USA* **108**, 14813–14818 (2011).
15. Zhong, N. & Weisgraber, K. H. Understanding the association of apolipoprotein E4 with Alzheimer disease: clues from its structure. *J. Biol. Chem.* **284**, 6027–6031 (2009).
16. Harold, D. et al. Genome-wide association study identifies variants at *CLU* and *PICALM* associated with Alzheimer's disease. *Nat. Genet.* **41**, 1088–1093 (2009).
17. Lambert, J. C. et al. Genome-wide association study identifies variants at *CLU* and *CR1* associated with Alzheimer's disease. *Nat. Genet.* **41**, 1094–1099 (2009).
18. Chartier-Harlin, M. C. et al. Apolipoprotein E,  $\epsilon$ 4 allele as a major risk factor for sporadic early and late-onset forms of Alzheimer's disease: analysis of the 19q13.2 chromosomal region. *Hum. Mol. Genet.* **3**, 569–574 (1994).
19. Houlden, H. et al. ApoE genotype is a risk factor in nonpresenilin early-onset Alzheimer's disease families. *Am. J. Med. Genet.* **81**, 117–121 (1998).
20. Rebeck, G. W., Reiter, J. S., Strickland, D. K. & Hyman, B. T. Apolipoprotein E in sporadic Alzheimer's disease: allelic variation and receptor interactions. *Neuron* **11**, 575–580 (1993).
21. Roses, A. D. et al. A *TOMM40* variable-length polymorphism predicts the age of late-onset Alzheimer's disease. *Pharmacogenomics J.* **10**, 375–384 (2010).
22. Cruchaga, C. et al. Association and expression analyses with single-nucleotide polymorphisms in *TOMM40* in Alzheimer disease. *Arch. Neurol.* **68**, 1013–1019 (2011).
23. Ellis, R. J. et al. Cerebral amyloid angiopathy in the brains of patients with Alzheimer's disease: the CERAD experience, Part XV. *Neurology* **46**, 1592–1596 (1996).
24. Namba, Y., Tomonaga, M., Kawasaki, H., Otomo, E. & Ikeda, K. Apolipoprotein E immunoreactivity in cerebral amyloid deposits and neurofibrillary tangles in Alzheimer's disease and kuru plaque amyloid in Creutzfeldt–Jakob disease. *Brain Res.* **541**, 163–166 (1991).
25. Kok, E. et al. Apolipoprotein E-dependent accumulation of Alzheimer disease-related lesions begins in middle age. *Ann. Neurol.* **65**, 650–657 (2009).
26. Polvikoski, T. et al. Apolipoprotein E, dementia, and cortical deposition of  $\beta$ -amyloid protein. *N. Engl. J. Med.* **333**, 1242–1247 (1995).
27. Schmechel, D. E. et al. Increased amyloid  $\beta$ -peptide deposition in cerebral cortex as a consequence of apolipoprotein E genotype in late-onset Alzheimer disease. *Proc. Natl Acad. Sci. USA* **90**, 9649–9653 (1993).
28. Klunk, W. E. et al. Imaging brain amyloid in Alzheimer's disease with Pittsburgh compound-B. *Ann. Neurol.* **55**, 306–319 (2004).
29. Barthel, H. et al. Cerebral amyloid- $\beta$  PET with florbetaben ( $^{18}$ F) in patients with Alzheimer's disease and healthy controls: a multicentre phase 2 diagnostic study. *Lancet Neurol.* **10**, 424–35 (2011).
30. Reiman, E. M. et al. Fibrillar amyloid- $\beta$  burden in cognitively normal people at 3 levels of genetic risk for Alzheimer's disease. *Proc. Natl Acad. Sci. USA* **106**, 6820–6825 (2009).
31. Head, D. et al. Exercise engagement as a moderator of the effects of APOE genotype on amyloid deposition. *Arch. Neurol.* **69**, 636–643 (2012).
32. Prince, J. A., Zetterberg, H., Andreasen, N., Marcusson, J. & Blennow, K. APOE  $\epsilon$ 4 allele is associated with reduced cerebrospinal fluid levels of A $\beta$ 42. *Neurology* **62**, 2116–2118 (2004).
33. Fleisher, A. S. et al. Apolipoprotein E  $\epsilon$ 4 and age effects on florbetapir positron emission tomography in healthy aging and Alzheimer disease. *Neurobiol. Aging* **34**, 1–12 (2013).
34. Berlau, D. J., Corrada, M. M., Head, E. & Kawas, C. H. APOE  $\epsilon$ 2 is associated with intact cognition but increased Alzheimer pathology in the oldest old. *Neurology* **72**, 829–834 (2009).
35. Greenberg, S. M. et al. Apolipoprotein E  $\epsilon$ 4 is associated with the presence and earlier onset of hemorrhage in cerebral amyloid angiopathy. *Stroke* **27**, 1333–1337 (1996).
36. Biffi, A. et al. Variants at APOE influence risk of deep and lobar intracerebral hemorrhage. *Ann. Neurol.* **68**, 934–943 (2010).
37. Vinters, H. V. Cerebral amyloid angiopathy. A critical review. *Stroke* **18**, 311–324 (1987).
38. Morris, J. C. et al. Mild cognitive impairment represents early-stage Alzheimer disease. *Arch. Neurol.* **58**, 397–405 (2001).
39. Petersen, R. C. et al. Mild cognitive impairment: clinical characterization and outcome. *Arch. Neurol.* **56**, 303–308 (1999).
40. Pa, J. et al. Clinical-neuroimaging characteristics of dysexecutive mild cognitive impairment. *Ann. Neurol.* **65**, 414–423 (2009).
41. Farlow, M. R. et al. Impact of APOE in mild cognitive impairment. *Neurology* **63**, 1898–1901 (2004).
42. Smith, G. E. et al. Apolipoprotein E genotype influences cognitive 'phenotype' in patients with Alzheimer's disease but not in healthy control subjects. *Neurology* **50**, 355–362 (1998).
43. Ramakers, I. H. et al. The association between APOE genotype and memory dysfunction in subjects with mild cognitive impairment is related to age and Alzheimer pathology. *Dement. Geriatr. Cogn. Disord.* **26**, 101–108 (2008).
44. Dik, M. G. et al. APOE- $\epsilon$ 4 is associated with memory decline in cognitively impaired elderly. *Neurology* **54**, 1492–1497 (2000).
45. Whitehair, D. C. et al. Influence of apolipoprotein E  $\epsilon$ 4 on rates of cognitive and functional decline in mild cognitive impairment. *Alzheimers Dement.* **6**, 412–419 (2010).
46. Cosentino, S. et al. APOE  $\epsilon$ 4 allele predicts faster cognitive decline in mild Alzheimer disease. *Neurology* **70**, 1842–1849 (2008).
47. Fleisher, A. S. et al. Clinical predictors of progression to Alzheimer disease in amnesic mild cognitive impairment. *Neurology* **68**, 1588–1595 (2007).
48. Elias-Sonnenschein, L. S., Vechtbauer, W., Ramakers, I. H., Verhey, F. R. & Visser, P. J. Predictive value of APOE- $\epsilon$ 4 allele for progression from MCI to AD-type dementia: a meta-analysis. *J. Neurol. Neurosurg. Psychiatry* **82**, 1149–1156 (2011).
49. Petersen, R. C. et al. Apolipoprotein E status as a predictor of the development of Alzheimer's disease in memory-impaired individuals. *JAMA* **273**, 1274–1278 (1995).
50. Vemuri, P. et al. Effect of apolipoprotein E on biomarkers of amyloid load and neuronal pathology in Alzheimer disease. *Ann. Neurol.* **67**, 308–316 (2010).
51. Pike, K. E. et al.  $\beta$ -amyloid imaging and memory in non-demented individuals: evidence for preclinical Alzheimer's disease. *Brain* **130**, 2837–2844 (2007).
52. Bunce, D., Fratiglioni, L., Small, B. J., Winblad, B. & Bäckman, L. APOE and cognitive decline in preclinical Alzheimer disease and non-demented aging. *Neurology* **63**, 816–821 (2004).
53. Lin, A. L., Laird, A. R., Fox, P. T. & Gao, J. H. Multimodal MRI neuroimaging biomarkers for cognitive normal adults, amnesic mild cognitive impairment, and Alzheimer's disease. *Neurology Res. Int.* **2012**, 907409 (2012).
54. Caselli, R. J. et al. Cognitive domain decline in healthy apolipoprotein E  $\epsilon$ 4 homozygotes before the diagnosis of mild cognitive impairment. *Arch. Neurol.* **64**, 1306–1311 (2007).
55. Caselli, R. J. et al. Longitudinal changes in cognition and behavior in asymptomatic carriers of the APOE  $\epsilon$ 4 allele. *Neurology* **62**, 1990–1995 (2004).
56. Caselli, R. J. et al. Longitudinal modeling of age-related memory decline and the APOE  $\epsilon$ 4 effect. *N. Engl. J. Med.* **361**, 255–263 (2009).
57. Caselli, R. J. et al. Longitudinal modeling of frontal cognition in APOE  $\epsilon$ 4 homozygotes, heterozygotes, and noncarriers. *Neurology* **76**, 1383–1388 (2011).
58. Izaks, G. J. et al. The association of APOE genotype with cognitive function in persons aged 35 years or older. *PLoS ONE* **6**, e27415 (2011).
59. Mondadori, C. R. et al. Better memory and neural efficiency in young apolipoprotein E  $\epsilon$ 4 carriers. *Cereb. Cortex* **17**, 1934–1947 (2007).

60. Jochemsen, H. M., Muller, M., van der Graaf, Y. & Geerlings, M. I. APOE  $\epsilon 4$  differentially influences change in memory performance depending on age. The SMART-MR study. *Neurobiol. Aging* **33**, 832 e15–e22 (2012).
61. Bloss, C. S., Delis, D. C., Salmon, D. P. & Bondi, M. W. Decreased cognition in children with risk factors for Alzheimer's disease. *Biol. Psychiatry* **64**, 904–906 (2008).
62. Tuminello, E. R. & Han, S. D. The apolipoprotein E antagonistic pleiotropy hypothesis: review and recommendations. *Int. J. Alzheimers Dis.* **2011**, 126197 (2011).
63. Castellano, J. M. et al. Human apoE isoforms differentially regulate brain amyloid- $\beta$  peptide clearance. *Sci. Transl. Med.* **3**, 89a57 (2011).
64. Morris, J. C. et al. APOE predicts amyloid-beta but not tau Alzheimer pathology in cognitively normal aging. *Ann. Neurol.* **67**, 122–131 (2010).
65. Haan, M. N., Shemanski, L., Jagust, W. J., Manolio, T. A. & Kuller, L. The role of APOE  $\epsilon 4$  in modulating effects of other risk factors for cognitive decline in elderly persons. *JAMA* **282**, 40–46 (1999).
66. Peila, R., Rodriguez, B. L. & Launer, L. J. Type 2 diabetes, APOE gene, and the risk for dementia and related pathologies: the Honolulu-Asia Aging Study. *Diabetes* **51**, 1256–1262 (2002).
67. Kalmijn, S., Feskens, E. J., Launer, L. J. & Kromhout, D. Cerebrovascular disease, the apolipoprotein e4 allele, and cognitive decline in a community-based study of elderly men. *Stroke* **27**, 2230–2235 (1996).
68. Irie, F. et al. Enhanced risk for Alzheimer disease in persons with type 2 diabetes and APOE  $\epsilon 4$ : the Cardiovascular Health Study Cognition Study. *Arch. Neurol.* **65**, 89–93 (2008).
69. Matsuzaki, T. et al. Insulin resistance is associated with the pathology of Alzheimer disease: the Hisayama study. *Neurology* **75**, 764–770 (2010).
70. Bell, R. D. et al. Apolipoprotein E controls cerebrovascular integrity via cyclophilin A. *Nature* **485**, 512–516 (2012).
71. Verghese, P. B., Castellano, J. M. & Holtzman, D. M. Apolipoprotein E in Alzheimer's disease and other neurological disorders. *Lancet Neurol.* **10**, 241–252 (2011).
72. Zhou, W. et al. Meta-analysis of APOE4 allele and outcome after traumatic brain injury. *J. Neurotrauma* **25**, 279–290 (2008).
73. Fleminger, S., Oliver, D. L., Lovestone, S., Rabe-Hesketh, S. & Giora, A. Head injury as a risk factor for Alzheimer's disease: the evidence 10 years on; a partial replication. *J. Neurol. Neurosurg. Psychiatry* **74**, 857–862 (2003).
74. Magnoni, S. & Brody, D. L. New perspectives on amyloid- $\beta$  dynamics after acute brain injury: moving between experimental approaches and studies in the human brain. *Arch. Neurol.* **67**, 1068–1073 (2010).
75. Nicoll, J. A., Roberts, G. W. & Graham, D. I. Apolipoprotein E  $\epsilon 4$  allele is associated with deposition of amyloid  $\beta$ -protein following head injury. *Nat. Med.* **1**, 135–137 (1995).
76. Jellinger, K. A. Morphologic diagnosis of "vascular dementia"—a critical update. *J. Neurol. Sci.* **270**, 1–12 (2008).
77. Yin, Y. W. et al. Association between apolipoprotein E gene polymorphism and the risk of vascular dementia: a meta-analysis. *Neurosci. Lett.* **514**, 6–11 (2012).
78. Prince, M. et al. The association between APOE and dementia does not seem to be mediated by vascular factors. *Neurology* **54**, 397–402 (2000).
79. Bender, A. R. & Raz, N. Age-related differences in memory and executive functions in healthy APOE  $\epsilon 4$  carriers: the contribution of individual differences in prefrontal volumes and systolic blood pressure. *Neuropsychologia* **50**, 704–714 (2012).
80. Lanterna, L. A. et al. Meta-analysis of APOE genotype and subarachnoid hemorrhage. *Neurology* **69**, 766–775 (2007).
81. Biffi, A. et al. APOE genotype and extent of bleeding and outcome in lobar intracerebral haemorrhage: a genetic association study. *Lancet Neurol.* **10**, 702–709 (2011).
82. Federoff, M., Jimenez-Rolando, B., Nalls, M. A. & Singleton, A. B. A large study reveals no association between APOE and Parkinson's disease. *Neurobiol. Dis.* **46**, 389–392 (2012).
83. Ezquerra, M. et al. Lack of association of APOE and tau polymorphisms with dementia in Parkinson's disease. *Neurosci. Lett.* **448**, 20–23 (2008).
84. McKeith, I. G. et al. Dementia with Lewy bodies. *Semin. Clin. Neuropsychiatry* **8**, 46–57 (2003).
85. Harrington, C. R. et al. Influence of apolipoprotein E genotype on senile dementia of the Alzheimer and Lewy body types. Significance for etiological theories of Alzheimer's disease. *Am. J. Pathol.* **145**, 1472–1484 (1994).
86. Mann, D. M., Brown, S. M., Owen, F., Baba, M. & Iwatsubo, T. Amyloid  $\beta$  protein (A $\beta$ ) deposition in dementia with Lewy bodies: predominance of A $\beta$ <sub>42/43</sub> and paucity of A $\beta$ <sub>40</sub> compared with sporadic Alzheimer's disease. *Neuropathol. Appl. Neurobiol.* **24**, 187–194 (1998).
87. Pletnikova, O. et al. A $\beta$  deposition is associated with enhanced cortical  $\alpha$ -synuclein lesions in Lewy body diseases. *Neurobiol. Aging* **26**, 1183–1192 (2005).
88. Seripa, D. et al. The APOE gene locus in frontotemporal dementia and primary progressive aphasia. *Arch. Neurol.* **68**, 622–628 (2011).
89. Saft, C. et al. Apolipoprotein E genotypes do not influence the age of onset in Huntington's disease. *J. Neurol. Neurosurg. Psychiatry* **75**, 1692–1696 (2004).
90. Jawaide, A. et al. Does apolipoprotein E genotype modify the clinical expression of ALS? *Eur. J. Neurol.* **18**, 618–624 (2011).
91. Bales, K. R. et al. Human APOE isoform-dependent effects on brain  $\beta$ -amyloid levels in PDAPP transgenic mice. *J. Neurosci.* **29**, 6771–6779 (2009).
92. Osenkowski, P., Ye, W., Wang, R., Wolfe, M. S. & Selkoe, D. J. Direct and potent regulation of  $\gamma$ -secretase by its lipid microenvironment. *J. Biol. Chem.* **283**, 22529–22540 (2008).
93. Riddell, D. R. et al. Impact of apolipoprotein E (apoE) polymorphism on brain ApoE levels. *J. Neurosci.* **28**, 11445–11453 (2008).
94. Sullivan, P. M. et al. Reduced levels of human apoE4 protein in an animal model of cognitive impairment. *Neurobiol. Aging* **32**, 791–801 (2011).
95. Wahrle, S. E. et al. Apolipoprotein E levels in cerebrospinal fluid and the effects of ABCA1 polymorphisms. *Mol. Neurodegener.* **2**, 7 (2007).
96. Beffert, U. et al. Apolipoprotein E and  $\beta$ -amyloid levels in the hippocampus and frontal cortex of Alzheimer's disease subjects are disease-related and apolipoprotein E genotype dependent. *Brain Res.* **843**, 87–94 (1999).
97. DeMattos, R. B. et al. ApoE and clusterin cooperatively suppress A $\beta$  levels and deposition: evidence that ApoE regulates extracellular A $\beta$  metabolism in vivo. *Neuron* **41**, 193–202 (2004).
98. Riddell, D. R. et al. The LXR agonist T0901317 selectively lowers hippocampal A $\beta$ 42 and improves memory in the Tg2576 mouse model of Alzheimer's disease. *Mol. Cell Neurosci.* **34**, 621–628 (2007).
99. Vanmierlo, T. et al. Liver X receptor activation restores memory in aged AD mice without reducing amyloid. *Neurobiol. Aging* **32**, 1262–1272 (2011).
100. Cramer, P. E. et al. ApoE-directed therapeutics rapidly clear  $\beta$ -amyloid and reverse deficits in AD mouse models. *Science* **335**, 1503–1506 (2012).
101. Fitz, N. F. et al. Abca1 deficiency affects Alzheimer's disease-like phenotype in human ApoE4 but not in ApoE3-targeted replacement mice. *J. Neurosci.* **32**, 13125–13136 (2012).
102. Kim, J., Basak, J. M. & Holtzman, D. M. The role of apolipoprotein E in Alzheimer's disease. *Neuron* **63**, 287–303 (2009).
103. LaDu, M. J. et al. Isoform-specific binding of apolipoprotein E to  $\beta$ -amyloid. *J. Biol. Chem.* **269**, 23403–23406 (1994).
104. Deane, R. et al. ApoE isoform-specific disruption of amyloid  $\beta$  peptide clearance from mouse brain. *J. Clin. Invest.* **118**, 4002–4013 (2008).
105. Jiang, Q. et al. ApoE promotes the proteolytic degradation of A $\beta$ . *Neuron* **58**, 681–693 (2008).
106. Bales, K. R. et al. Lack of apolipoprotein E dramatically reduces amyloid  $\beta$ -peptide deposition. *Nat. Genet.* **17**, 263–264 (1997).
107. Ries, M. L. et al. Magnetic resonance imaging characterization of brain structure and function in mild cognitive impairment: a review. *J. Am. Geriatr. Soc.* **56**, 920–934 (2008).
108. Nestor, P. J., Scheltens, P. & Hodges, J. R. Advances in the early detection of Alzheimer's disease. *Nat. Med.* **10** (Suppl.), S34–S41 (2004).
109. Korf, E. S., Wahlund, L.-O., Visser, P. J. & Scheltens, P. Medial temporal lobe atrophy on MRI predicts dementia in patients with mild cognitive impairment. *Neurology* **63**, 94–100 (2004).
110. Hashimoto, M. et al. Apolipoprotein E  $\epsilon 4$  and the pattern of regional brain atrophy in Alzheimer's disease. *Neurology* **57**, 1461–1466 (2001).
111. Espeseth, T. et al. Accelerated age-related cortical thinning in healthy carriers of apolipoprotein E  $\epsilon 4$ . *Neurobiol. Aging* **29**, 329–340 (2008).
112. Reiman, E. M. et al. Hippocampal volumes in cognitively normal persons at genetic risk for Alzheimer's disease. *Ann. Neurol.* **44**, 288–291 (1998).
113. Fennema-Notestine, C. et al. Presence of APOE  $\epsilon 4$  allele associated with thinner frontal cortex in middle age. *J. Alzheimers Dis.* **26**, 49–60 (2011).
114. Sheline, Y. I. et al. APOE4 allele disrupts resting state fMRI connectivity in the absence of amyloid plaques or decreased CSF A $\beta$ 42. *J. Neurosci.* **30**, 17035–17040 (2010).
115. Machulda M. M. et al. Effect of APOE  $\epsilon 4$  status on intrinsic network connectivity in cognitively normal elderly subjects. *Arch. Neurol.* **68**, 1131–1136 (2011).
116. Bookheimer, S. Y. et al. Patterns of brain activation in people at risk for Alzheimer's disease. *N. Engl. J. Med.* **343**, 450–456 (2000).
117. Filippini, N. et al. Distinct patterns of brain activity in young carriers of the APOE- $\epsilon 4$  allele. *Proc. Natl Acad. Sci. USA* **106**, 7209–7214 (2009).
118. Bondi, M. W., Houston, W. S., Eyler, L. T. & Brown, G. G. fMRI evidence of compensatory mechanisms in older adults at genetic risk for Alzheimer disease. *Neurology* **64**, 501–508 (2005).
119. Bero, A. W. et al. Neuronal activity regulates the regional vulnerability to amyloid- $\beta$  deposition. *Nat. Neurosci.* **14**, 750–756 (2011).



120. Cirrito, J. R. *et al.* Synaptic activity regulates interstitial fluid amyloid- $\beta$  levels *in vivo*. *Neuron* **48**, 913–922 (2005).
121. Putcha, D. *et al.* Hippocampal hyperactivation associated with cortical thinning in Alzheimer's disease signature regions in non-demented elderly adults. *J. Neurosci.* **31**, 17680–17688 (2011).
122. O'Brien, J. L. *et al.* Longitudinal fMRI in elderly reveals loss of hippocampal activation with clinical decline. *Neurology* **74**, 1969–1976 (2010).
123. Mosconi, L. *et al.* Pre-clinical detection of Alzheimer's disease using FDG-PET, with or without amyloid imaging. *J. Alzheimers Dis.* **20**, 843–854 (2010).
124. Small, G. W. *et al.* Apolipoprotein E type 4 allele and cerebral glucose metabolism in relatives at risk for familial Alzheimer disease. *JAMA* **273**, 942–947 (1995).
125. Reiman, E. M. *et al.* Preclinical evidence of Alzheimer's disease in persons homozygous for the  $\epsilon$ 4 allele for apolipoprotein E. *N. Engl. J. Med.* **334**, 752–758 (1996).
126. Small, G. W. *et al.* Cerebral metabolic and cognitive decline in persons at genetic risk for Alzheimer's disease. *Proc. Natl Acad. Sci. USA* **97**, 6037–6042 (2000).
127. Reiman, E. M. *et al.* Correlations between apolipoprotein E  $\epsilon$ 4 gene dose and brain-imaging measurements of regional hypometabolism. *Proc. Natl Acad. Sci. USA* **102**, 8299–8302 (2005).
128. Huang, Y.  $\text{A}\beta$ -independent roles of apolipoprotein E4 in the pathogenesis of Alzheimer's disease. *Trends Mol. Med.* **16**, 287–294 (2010).
129. Aoki, K. *et al.* Increased expression of neuronal apolipoprotein E in human brain with cerebral infarction. *Stroke* **34**, 875–880 (2003).
130. Brecht, W. J. *et al.* Neuron-specific apolipoprotein E4 proteolysis is associated with increased tau phosphorylation in brains of transgenic mice. *J. Neurosci.* **24**, 2527–2534 (2004).
131. Mahley, R. W., Weisgraber, K. H. & Huang, Y. Apolipoprotein E4: a causative factor and therapeutic target in neuropathology, including Alzheimer's disease. *Proc. Natl Acad. Sci. USA* **103**, 5644–5651 (2006).
132. Aboud, O., Mrak, R. E., Boop, F. & Griffin, S. T. Apolipoprotein epsilon 3 alleles are associated with indicators of neuronal resilience. *BMC Med.* **10**, 35 (2012).
133. Buttini, M. *et al.* Cellular source of apolipoprotein E4 determines neuronal susceptibility to excitotoxic injury in transgenic mice. *Am. J. Pathol.* **177**, 563–569 (2010).
134. Mauch, D. H. *et al.* CNS synaptogenesis promoted by glia-derived cholesterol. *Science* **294**, 1354–1357 (2001).
135. Pfriger, F. W. Cholesterol homeostasis and function in neurons of the central nervous system. *Cell Mol. Life Sci.* **60**, 1158–1171 (2003).
136. Svennerholm, L. & Gottfries, C.-G. Membrane lipids, selectively diminished in Alzheimer brains, suggest synapse loss as a primary event in early-onset form (Type I) and demyelination in late-onset form (Type II). *J. Neurochem.* **62**, 1039–1047 (1994).
137. Rapp, A., Gmeiner, B. & Hüttinger, M. Implication of apoE isoforms in cholesterol metabolism by primary rat hippocampal neurons and astrocytes. *Biochimie* **88**, 473–483 (2006).
138. Hamanaka, H. *et al.* Altered cholesterol metabolism in human apolipoprotein E4 knock-in mice. *Hum. Mol. Genet.* **9**, 353–361 (2000).
139. Selkoe, D. J. Alzheimer's disease is a synaptic failure. *Science* **298**, 789–791 (2002).
140. Koffie, R., Hyman, B. & Spires-Jones, T. Alzheimer's disease: synapses gone cold. *Mol. Neurodegener.* **6**, 63 (2011).
141. Chen, Y., Durakoglugil, M. S., Xian, X. & Herz, J. ApoE4 reduces glutamate receptor function and synaptic plasticity by selectively impairing ApoE receptor recycling. *Proc. Natl Acad. Sci. USA* <http://dx.doi.org/10.1073/pnas.0914984107>.
142. Buttini, M. *et al.* Modulation of Alzheimer-like synaptic and cholinergic deficits in transgenic mice by human apolipoprotein E depends on isoform, aging, and overexpression of amyloid  $\beta$  peptides but not on plaque formation. *J. Neurosci.* **22**, 10539–10548 (2002).
143. Ji, Y. *et al.* Apolipoprotein E isoform-specific regulation of dendritic spine morphology in apolipoprotein E transgenic mice and Alzheimer's disease patients. *Neuroscience* **122**, 305–315 (2003).
144. Dumanis, S. B. *et al.* ApoE4 decreases spine density and dendritic complexity in cortical neurons *in vivo*. *J. Neurosci.* **29**, 15317–15322 (2009).
145. Wang, C. *et al.* Human apoE4-targeted replacement mice display synaptic deficits in the absence of neuropathology. *Neurobiol. Dis.* **18**, 390–398 (2005).
146. Sen, A., Alkon, D. L. & Nelson, T. J. Apolipoprotein E3 (ApoE3) but not ApoE4 protects against synaptic loss through increased expression of protein kinase C $\alpha$ . *J. Biol. Chem.* **287**, 15947–15958 (2012).
147. Klein, R. C., Mace, B. E., Moore, S. D. & Sullivan, P. M. Progressive loss of synaptic integrity in human apolipoprotein E4 targeted replacement mice and attenuation by apolipoprotein E2. *Neuroscience* **171**, 1265–1272 (2010).
148. Bales, K. R., Du, Y., Holtzman, D., Cordell, B. & Paul, S. M. Neuroinflammation and Alzheimer's disease: critical roles for cytokine/ $\text{A}\beta$ -induced glial activation, NF- $\kappa$ B, and apolipoprotein E. *Neurobiol. Aging* **21**, 427–432 (2000).
149. LaDu, M. J. *et al.* Apolipoprotein E and apolipoprotein E receptors modulate  $\text{A}\beta$ -induced glial neuroinflammatory responses. *Neurochem. Int.* **39**, 427–434 (2001).
150. Lynch, J. R., Morgan, D., Mance, J., Matthew, W. D. & Laskowitz, D. T. Apolipoprotein E modulates glial activation and the endogenous central nervous system inflammatory response. *J. Neuroimmunol.* **114**, 107–113 (2001).
151. Keene, C. D., Cudaback, E., Li, X., Montine, K. S. & Montine, T. J. Apolipoprotein E isoforms and regulation of the innate immune response in brain of patients with Alzheimer's disease. *Curr. Opin. Neurobiol.* **21**, 920–928 (2011).
152. Lynch, J. R. *et al.* APOE genotype and an ApoE-mimetic peptide modify the systemic and central nervous system inflammatory response. *J. Biol. Chem.* **278**, 48529–48533 (2003).
153. Ringman, J. M. *et al.* Plasma signaling proteins in persons at genetic risk for Alzheimer disease: influence of APOE genotype. *Arch. Neurol.* **69**, 757–764 (2012).
154. Szekely, C. A. *et al.* NSAID use and dementia risk in the Cardiovascular Health Study: role of APOE and NSAID type. *Neurology* **70**, 17–24 (2008).
155. Mu, Y. & Gage, F. Adult hippocampal neurogenesis and its role in Alzheimer's disease. *Mol. Neurodegener.* **6**, 85 (2011).
156. Yang, C.-P., Gilley, J. A., Zhang, G. & Kernie, S. G. ApoE is required for maintenance of the dentate gyrus neural progenitor pool. *Development* **138**, 4351–4362 (2011).
157. Andrews-Zwilling, Y. *et al.* Apolipoprotein E4 causes age- and Tau-dependent impairment of GABAergic interneurons, leading to learning and memory deficits in mice. *J. Neurosci.* **30**, 13707–13717 (2010).
158. Li, G. *et al.* GABAergic interneuron dysfunction impairs hippocampal neurogenesis in adult apolipoprotein E4 knockin mice. *Cell Stem Cell* **5**, 634–645 (2009).
159. Salloway, S. *et al.* A randomized, double-blind, placebo-controlled clinical trial of intravenous bapineuzumab in patients with Alzheimer's disease who are apolipoprotein E  $\epsilon$ 4 non-carriers. Presented at the 16<sup>th</sup> meeting of the European Federation of Neurological Societies (Stockholm, Sweden, 2012).
160. Sperling, R. *et al.* A randomized, double-blind, placebo-controlled clinical trial of intravenous bapineuzumab in patients with Alzheimer's disease who are apolipoprotein E  $\epsilon$ 4 carriers. Presented at the 16<sup>th</sup> meeting of the European Federation of Neurological Societies (Stockholm, Sweden, 2012).
161. Ferrari, C. *et al.* How can elderly apolipoprotein E  $\epsilon$ 4 carriers remain free from dementia? *Neurobiol. Aging* <http://dx.doi.org/10.1016/j.neurobiolaging.2012.03.003>.
162. Hesse, C. *et al.* Measurement of apolipoprotein E (apoE) in cerebrospinal fluid. *Neurochem. Res.* **25**, 511–517 (2000).
163. Gupta, V. B. *et al.* Plasma apolipoprotein E and Alzheimer disease risk. *Neurology* **76**, 1091–1098 (2011).
164. Chawla, A. *et al.* A PPAR $\gamma$ -LXR-ABCA1 pathway in macrophages is involved in cholesterol efflux and atherogenesis. *Mol. Cell* **7**, 161–171 (2001).
165. Vaya, J. & Schipper, H. M. Oxysterols, cholesterol homeostasis, and Alzheimer disease. *J. Neurochem.* **102**, 1727–1737 (2007).
166. Donkin, J. J. *et al.* ATP-binding cassette transporter A1 mediates the beneficial effects of the liver X receptor agonist GW3965 on object recognition memory and amyloid burden in amyloid precursor protein/presenilin 1 mice. *J. Biol. Chem.* **285**, 34144–34154 (2010).
167. Fryer, J. D. *et al.* Human apolipoprotein E4 alters the amyloid- $\beta$  40:42 ratio and promotes the formation of cerebral amyloid angiopathy in an amyloid precursor protein transgenic model. *J. Neurosci.* **25**, 2803–2810 (2005).
168. Kim, J. *et al.* Haploinsufficiency of human APOE reduces amyloid deposition in a mouse model of amyloid- $\beta$  amyloidosis. *J. Neurosci.* **31**, 18007–18012 (2011).
169. Bien-Ly, N., Gillespie, A. K., Walker, D., Yoon, S. Y. & Huang, Y. Reducing human apolipoprotein E levels attenuates age-dependent  $\text{A}\beta$  accumulation in mutant human amyloid precursor protein transgenic mice. *J. Neurosci.* **32**, 4803–4811 (2012).
170. Sadowski, M. J. *et al.* Blocking the apolipoprotein E/amyloid- $\beta$  interaction as a potential therapeutic approach for Alzheimer's disease. *Proc. Natl Acad. Sci. USA* **103**, 18787–18792 (2006).
171. Chen, H. K. *et al.* Small molecule structure correctors abolish detrimental effects of apolipoprotein E4 in cultured neurons. *J. Biol. Chem.* **287**, 5253–5266 (2012).
172. Aono, M. *et al.* Protective effect of apolipoprotein E-mimetic peptides on N-methyl-D-aspartate excitotoxicity in primary rat neuronal-glial cell cultures. *Neuroscience* **116**, 437–445 (2003).
173. Li, F. Q. *et al.* Apolipoprotein E-derived peptides ameliorate clinical disability and inflammatory infiltrates into the spinal cord in a murine model of multiple sclerosis. *J. Pharmacol. Exp. Ther.* **318**, 956–965 (2006).



174. Li, F. Q., Fowler, K. A., Neil, J. E., Colton, C. A. & Vitek, M. P. An apolipoprotein E-mimetic stimulates axonal regeneration and remyelination after peripheral nerve injury. *J. Pharmacol. Exp. Ther.* **334**, 106–115 (2010).
175. Kanekiyo, T. et al. Heparan sulphate proteoglycan and the low-density lipoprotein receptor-related protein 1 constitute major pathways for neuronal amyloid- $\beta$  uptake. *J. Neurosci.* **31**, 1644–1651 (2011).
176. Liu, Q. et al. Neuronal LRP1 knockout in adult mice leads to impaired brain lipid metabolism and progressive, age-dependent synapse loss and neurodegeneration. *J. Neurosci.* **30**, 17068–17078 (2010).
177. Kim, J. et al. Overexpression of low-density lipoprotein receptor in the brain markedly inhibits amyloid deposition and increases extracellular A $\beta$  clearance. *Neuron* **64**, 632–644 (2009).
178. Herz, J. & Chen, Y. Reelin, lipoprotein receptors and synaptic plasticity. *Nat. Rev. Neurosci.* **7**, 850–859 (2006).

## Acknowledgements

Work in authors' laboratories is supported by the NIH, the Alzheimer's Association, the American Health Assistance Foundation, and Xiamen University Research Funds. We thank C. Stetler and O. Ross for critical reading of the manuscript before submission.

## Author contributions

All authors contributed to researching data for the article, discussion of the content, writing the article, and to review and/or editing of the manuscript before submission.

## **CORRECTION**

### **Apolipoprotein E and Alzheimer disease: risk, mechanisms and therapy**

Chia-Chen Liu, Takahisa Kanekiyo, Huaxi Xu & Guojun Bu

*Nat. Rev. Neurol.* **9**, 106–118 (2013); doi:10.1038/nrneurol.2012.263

In the version of this article initially published, in the author list, Chia-Chen Liu's name was misspelt. The error has been corrected for the HTML and PDF versions of the article.