

Ch 27: Magnetism

Season 3 Episode 3 - preparing for boss fight

In this episode of LARC Physics 3B, we're going to . . .

- Create a foundation for solving "Magnetism" problems by
 - Determining how (moving) charges behave in magnetic fields
 - Using Ampere's Law

Lecture Review

Big Ideas: Force due to Magnetic Field and Ampere's Law

Key Words: moving charge, current, magnetic field (i.e. B-field), Ampere's Law.

Here are some important equations/concepts:


- $F_B = q(\vec{v} \times \vec{B})$, magnetic force on moving point charge
- $F_B = I(\vec{\ell} \times \vec{B})$, magnetic force on segment of current where $\vec{\ell}$ is the direction of the current.
- Ampere's Law: $\int \vec{B} \cdot d\vec{\ell} = \mu_0 I_{\text{enc}}$
 - Similar to Gauss' Law, we primarily rely on this equation to find the B-field.
 - Involves drawing an "Amperian loop" (like a "Gaussian surface")
 - Often reduces such that

$$\int \vec{B} \cdot d\vec{\ell} = \mu_0 I_{\text{enc}} \implies B(2\pi r) = \mu_0 I_{\text{enc}}, \quad \text{where } r \text{ is the radius of the Amperian loop}$$


Guided Practice

Consider the following scenarios with a given magnetic field \vec{B} and a charge moving with some velocity \vec{v} . Determine the direction of the magnetic force acting on the charge.


\vec{B} pointing out of the page




\vec{B} pointing to the right



\vec{B} pointing into the page



\vec{B} pointing up



Answers:

(a) \vec{F}_B points to the right

(b) \vec{F}_B points into the page

(c) \vec{F}_B points downwards (Left-Hand Rule for negative charges!)

(d) \vec{F}_B is zero since \vec{v} and \vec{B} are parallel and $\sin \theta = 0$ for $\theta = 0^\circ$

Breakout-Room Activity


Determine the magnitude and direction of the force on an electron traveling 8.75×10^5 m/s horizontally to the East in a vertically upward magnetic field of strength 0.45 T.

Useful info: $q = -1.6 \times 10^{-19}$ C for the charge of an electron.

Answer: $\vec{F}_B = 6.3 \times 10^{-14} \text{ N}(-\hat{z})$, into the page (i.e. away from you).

Breakout-Room Activity

Just as magnetic fields push moving charges; moving charges can create magnetic fields. In other words, a current may produce a magnetic field. Let's consider the scenario of current moving through a straight wire:



Derive an expression for the **magnitude of the magnetic field** $B = |\tilde{\mathbf{B}}|$ at a certain radius r away from the wire with current I running vertically upwards as shown in the figure.

Answer: $B = \frac{\mu_0}{2\pi} \frac{I}{r}$

Here is an application of this concept in the case of two current-carrying wires running parallel to each other. Their individual magnetic fields cause the two wires to attract each other via the magnetic force. https://www.youtube.com/watch?v=8T_dP70q620