# Bounds on the Number of Divisors

0xHaru

April 30, 2026

## Preliminary Lemma

**Lemma 1.** *Let $d \in \mathbb{N}$ be a divisor of $n \in \mathbb{N}$. Then at least one of $d$ or $k = n/d$ is less than or equal to $\sqrt{n}$.*

*Proof 1 (by contradiction).* Since $d$ divides $n$, there exists an integer $k$ such that $d \cdot k = n$.

Suppose for contradiction that *both* $d$ and $k$ are strictly greater than $\sqrt{n}$:

$$d > \sqrt{n} \quad \text{and} \quad k > \sqrt{n}.$$

Multiplying these two inequalities gives

$$d \cdot k > \sqrt{n} \cdot \sqrt{n} = n,$$

which contradicts $d \cdot k = n$. Therefore, at least one of $d \leq \sqrt{n}$ or $k \leq \sqrt{n}$ must hold. □

*Proof 2 (direct).* Since $d$ divides $n$, there exists an integer $k = n/d$ such that $d \cdot k = n$. We consider two cases.

*Case $d \leq \sqrt{n}$.* The thesis holds immediately, since $d$ itself is at most $\sqrt{n}$.

*Case $d > \sqrt{n}$.* We show that $k < \sqrt{n}$:

$$\begin{aligned}
d &> \sqrt{n} && \text{(assumption of this case)} \\
\frac{1}{d} &< \frac{1}{\sqrt{n}} && \text{(taking reciprocals of positive quantities reverses the inequality)} \\
\frac{n}{d} &< \frac{n}{\sqrt{n}} && \text{(multiplying both sides by } n > 0\text{)} \\
k = \frac{n}{d} &< \sqrt{n} && \text{(simplifying } n/\sqrt{n} = \sqrt{n}\text{)}
\end{aligned}$$

In both cases at least one of $d$ or $k$ is less than or equal to $\sqrt{n}$. □

*Remark.* This lemma tells us that every divisor $d$ of $n$ either is at most $\sqrt{n}$, or its complementary divisor $n/d$ is at most $\sqrt{n}$. In other words, to enumerate *all* divisors of $n$ it suffices to check only the natural numbers up to $\sqrt{n}$.

# Main Theorem

**Theorem 1.** *For every $n \in \mathbb{N}$, the number of divisors $d(n)$ satisfies*

$$d(n) \;\leq\; 2\sqrt{n}.$$

*Proof.* Let $D$ denote the set of all positive divisors of $n$. By definition $|D| = d(n)$.
**Partition of $D$.** We partition $D$ into three pairwise disjoint subsets:

$$\begin{aligned} S_1 &= \{\, d \in D \;:\; d < \sqrt{n} \,\}, \\ S_2 &= \{\, d \in D \;:\; d = \sqrt{n} \,\}, \\ S_3 &= \{\, d \in D \;:\; d > \sqrt{n} \,\}. \end{aligned}$$

Since $S_1, S_2, S_3$ cover $D$ without overlap:

$$d(n) = |S_1| + |S_2| + |S_3|.$$

**Step 1: A bijection between $S_1$ and $S_3$.**
We claim $|S_1| = |S_3|$. We prove this by exhibiting a bijection

$$\varphi \;:\; S_1 \;\longrightarrow\; S_3, \qquad \varphi(d) = \frac{n}{d}.$$

We verify in turn that $\varphi$ is well-defined, injective, and surjective.[1]

***Well-definedness.***
Let $d \in S_1$ be arbitrary; we must show $\varphi(d) = n/d \in S_3$. Since $d \in S_1$, $d$ divides $n$, so $n/d \in \mathbb{N}$ and $n/d$ divides $n$. It remains to check $n/d > \sqrt{n}$. Since $d < \sqrt{n}$,

$$\begin{aligned} &\frac{1}{d} > \frac{1}{\sqrt{n}} && \text{(taking reciprocals of positive quantities reverses the inequality)} \\ &\frac{n}{d} > \frac{n}{\sqrt{n}} = \sqrt{n} && \text{(multiplying both sides by } n > 0\text{)} \end{aligned}$$

Hence $\varphi(d) \in S_3$ for all $d \in S_1$, and $\varphi$ is well-defined.

***Injectivity.***
Suppose $\varphi(a) = \varphi(b)$ for some $a, b \in S_1$, i.e. $n/a = n/b$.

$$\frac{n}{a} = \frac{n}{b} \;\Longrightarrow\; n = \frac{na}{b} \;\Longrightarrow\; bn = an \;\Longrightarrow\; b = a.$$

Hence $\varphi$ is injective.

[1] Let $f\colon A \to B$ be a function. $f$ is *injective* if $f(a) = f(b) \Rightarrow a = b$. $f$ is *surjective* if $\forall\, b \in B \;\; \exists\, a \in A$ such that $f(a) = b$.

***Surjectivity.***

*Scratch work.*

$$\varphi(a) = b \implies \frac{n}{a} = b \implies n = ab \implies a = \frac{n}{b}.$$

Let $b \in S_3$ be arbitrary. Set $a := n/b$. Since $b \in S_3$, $b$ divides $n$, so $a \in \mathbb{N}$ and $a$ divides $n$. Moreover, since $b > \sqrt{n}$,

$$a = \frac{n}{b} < \frac{n}{\sqrt{n}} = \sqrt{n},$$

so $a \in S_1$. Then

$$\varphi(a) = \frac{n}{a} = \frac{n}{\frac{n}{b}} = n \cdot \frac{b}{n} = b.$$

Since $b \in S_3$ was arbitrary, $\varphi$ is surjective.

Having verified that $\varphi$ is well-defined, injective, and surjective, we conclude that $\varphi$ is a bijection from $S_1$ to $S_3$, and therefore $|S_1| = |S_3|$.

**Step 2: Bounding $|S_1|$.**

Let $A(n) := |S_1|$. Using $|S_1| = |S_3|$, we rewrite the divisor count as

$$d(n) = A(n) + |S_2| + A(n).$$

By definition, $A(n)$ counts all naturals in $S_1$, each of which satisfies $d < \sqrt{n}$. The number of positive integers strictly less than $\sqrt{n}$ is at most $\lfloor \sqrt{n} \rfloor$, and $\lfloor \sqrt{n} \rfloor < \sqrt{n}$ always holds.[2] Therefore

$$A(n) < \sqrt{n}.$$

**Step 3: Final bounds.**

We distinguish two cases.

*Case A: $n$ is not a perfect square.*

$\sqrt{n} \notin \mathbb{N}$, so no divisor of $n$ can equal $\sqrt{n}$; hence $S_2 = \varnothing$ and $|S_2| = 0$. Therefore

$$d(n) = A(n) + 0 + A(n) = 2A(n) < 2\sqrt{n}.$$

*Case B: $n$ is a perfect square.*

$\sqrt{n} \in \mathbb{N}$ and $\sqrt{n}$ divides $n$ (since $\sqrt{n} \cdot \sqrt{n} = n$). Thus $S_2$ contains exactly one element, namely $\sqrt{n}$, so $|S_2| = 1$. Therefore

$$d(n) = A(n) + 1 + A(n) = 2A(n) + 1 < 2\sqrt{n} + 1.$$

Since $n$ is a perfect square both $d(n)$ and $2\sqrt{n}$ are integers, so the strict inequality with $2\sqrt{n} + 1$ is equivalent to

$$d(n) \leq 2\sqrt{n}.$$

In both cases we obtain $d(n) \leq 2\sqrt{n}$, which completes the proof. $\square$

[2] If $n$ is not a perfect square then $\sqrt{n}$ is irrational, so $\lfloor \sqrt{n} \rfloor < \sqrt{n}$ directly. If $n$ is a perfect square, the positive integers strictly less than $\sqrt{n}$ number at most $\sqrt{n} - 1 < \sqrt{n}$.