

# Problem F

## A Grouping Problem

**Input:** Standard Input  
**Output:** Standard Output

You are given a set of  $N$  integers. You can take  $K$  different elements from them to make a group. Two groups will be different if there is at least one element which is not common to both. For example, if there are 4 elements –  $a, b, c, d$  and you are asked to take two elements then  $ab, ad, bc, cd$  are all valid and different groups. A grouping system is complete if for a particular  $K$ , number of different groups is the maximum. In the former case,  $\{ab, bc, cd, bd, ad, ac\}$  is a complete grouping system.

For a particular complete grouping system, the **fitness** is calculated in the following way –

1. Each group of a grouping system contributes a part – the multiplication of all numbers of that group
2. Contribution from all groups are added
3. The fitness is equivalent to *Total Contribution mod M*,  $M$  is the bounding parameter

In our example, for  $K = 2$ , the fitness is  $F_2 = (ab+bc+cd+bd+ad+ac) \bmod M$ . If  $K = 1$ , then fitness is  $F_1 = (a+b+c+d) \bmod M$ .

Here, in this problem you have to find the complete grouping system with maximum fitness.

### Input

Each test case starts with two positive integer  $N$  ( $2 \leq N \leq 1000$ ) and  $M$  ( $1 \leq M < 2^{31}$ ). In next few lines there will be  $N$  positive integers. Each integer will be at best 1000. Input will be terminated by a case where  $N=M=0$ .

### Output

For each test case, print in a line the maximum fitness possible for a grouping system.

### Sample Input

```
4 10
1 2 3 4
4 100
1 2 3 4
4 6
1 2 3 4
0 0
```

### Output for Sample Input

```
5
50
5
```

---

**Problem setter: Md. Kamruzzaman, EPS**