

Prime geodesic theorem for arithmetic compact surfaces

Wenyan Yang

University of Science and Technology of China

May 25, 2026

Outline

What is PGT

Proof of PGT

Prime number theorem

In the 18th century, people first discovered that the asymptotic distribution of prime numbers follows a simple law:

$$\pi(x) = \#\{p \text{ prime} : p \leq x\} \sim \frac{x}{\log x}.$$

This is the famous prime number theorem (PNT).

Proof of PNT

About a hundred years later, Hadamard and Poussin independently proved this.

The proof contains three steps.

- ▶ Prove that PNT is equivalent to this statement.

$$\psi(x) = \sum_{n \leq x} \Lambda(n) = \sum_{p^e \leq x, p \text{ prime}} \log p \sim x.$$

- ▶ Prove that Riemann zeta function $\zeta(s)$ has no zeros on the line $\Re(s) = 1$.
- ▶ Using Mellin transformation, Write $\psi(x)$ as an integral, which is x plus something related to the Riemann zeta function.

Error terms in PNT

In fact, we can write the error terms in PNT explicitly.

$$\psi(x) = \int_{c-i\infty}^{c+i\infty} \frac{\zeta'(s)}{\zeta(s)} \frac{x^s}{s} ds = x - \sum_{\rho} \frac{x^{\rho}}{\rho} - \log(2\pi) - \frac{1}{2} \log\left(1 - \frac{1}{x^2}\right).$$

The sum takes ρ over all the nontrivial zeros of $\zeta(s)$. The famous Riemann hypothesis (RH) says that all of $\Re(\rho) = \frac{1}{2}$.

RH is a notoriously hard open problem in mathematics. In fact one can even not prove that $\Re(\rho) < 1 - \epsilon$ for any $\epsilon > 0$ now.

Prime geodesic theorem

Prime geodesic theorem (PGT) is just an analogy to PNT. The first PGT is proved for $PSL_2(\mathbb{Z}) \backslash \mathbb{H}$. Here $\mathbb{H}$ is the upper half plane. One can generalize to other arithmetic compact hyperbolic surfaces, e.g. $\mathbb{H}$ modulo congruence subgroups.

$$\Gamma = PSL_2(\mathbb{Z}) \backslash \mathbb{H}$$




Figure 1: Image of j function, which is a typical function on Γ

PNT vs PGT

- ▶ Prime numbers.
 - ▶ Norm of a prime number is just its value.
 - ▶ $\pi(x) = \int_0^x \frac{dt}{\log t} + E(x), \psi(x) = x + \tilde{E}(x).$
 - ▶ Riemann zeta function $\zeta(s).$
 - ▶ RH.
- ▶ Prime geodesics on the hyperbolic surface $\Gamma.$
 - ▶ Norm of a prime geodesic is the logarithm of its length.
 - ▶ $\pi_\Gamma(x) = \int_0^x \frac{dt}{\log t} + E(x), \Psi_\Gamma(x) = x + \tilde{E}(x).$
 - ▶ Selberg zeta function $Z_\Gamma(s).$
 - ▶ RH.

The philosophy

There are some crucial differences in PGT compared with PNT.

- ▶ GL_2 is non-abelian, thus Selberg trace formula is needed.
- ▶ $Z_\Gamma(s)$ is of order 2, unlike $\zeta(s)$ which is of order 1. So $Z_\Gamma(s)$ contains more zeros. Now RH only implies that $E(x) \ll x^{3/4+\epsilon}$, not $E(x) \ll x^{1/2+\epsilon}$.
- ▶ As you may guess from the previous difference, now RH is much easier to handle.

Progress

- ▶ (Iwaniec, 1984) $E(x) \ll x^{35/48+\epsilon}$.
- ▶ (Luo-Sarnak, 1995) $E(x) \ll x^{7/10+\epsilon}$.
- ▶ (Soundararajan-Young, 2013) $E(x) \ll x^{25/36+\epsilon}$.

PGT for quaternions

Koyama first extended the Luo-Sarnak bound to the surface $\Gamma_{\mathcal{D}}$ from a quaternion algebra $\mathcal{D}$ (Koyama, 1998).

Koyama's method exploits the Jacquet-Langlands correspondence (JL), but working on the spectral side.

Quaternions

Quaternions for real numbers are just generalizations of complex numbers, where you use 3 imaginary unit i, j, k satisfying associative law with the following relations.

$$i^2 = j^2 = k^2 = ijk = -1.$$

But now you have $ij = k = -ji$. For general fields the construction is similar.

Jacquet-Langlands correspondence

There are two types of degree 2 central simple algebra over a field, matrix algebra M_2 and quaternion algebra $\mathcal{D}$. Jacquet-Langlands correspondence says that their representations have some relations.

New work

We generalize this result to principal subgroups $\Gamma_{\mathcal{D}}(\mathfrak{N})$, using also JL but working on the arithmetic side (Tang-Wu-Yang-Y., 2026).

Proof Sketch

The proof contains 5 steps.

- ▶ Like proof of PNT, write Ψ as an integral.

Integral expression of Ψ

$$I(t; \tilde{f}_\infty) := \int_{\Gamma^1 \mathbb{R}^\times \setminus GL_2(\mathbb{R})} \left(\sum_{\sigma \in [\gamma]_{\text{st}}} \tilde{f}_\infty(g^{-1} \sigma g) \right) dg = \frac{\tilde{g}_\infty(\log x)}{\sqrt{x}} \cdot \psi_{\Gamma^1}(t).$$

Here $x = \frac{|t| + \sqrt{t^2 - 4}}{2}$.

Proof Sketch

The proof contains 5 steps.

- ▶ Like proof of PNT, write Ψ as an integral.
- ▶ Turn it into an adelic orbital integral.

Adelic integral

$$I(t; \tilde{f}_\infty) = (\text{some constant}) \int_{\mathbf{Z}_\gamma(\mathbb{A}) \backslash \mathcal{B}^\times(\mathbb{A})} \left(\tilde{f}_\infty \otimes id_{\mathbf{K}_\Gamma} \right) (g^{-1} \gamma g) dg.$$

The constant is

$$\frac{1}{c_{\Gamma^1}} \frac{\text{Vol}(\widehat{\mathbb{Z}}^\times)}{[\widehat{\mathbb{Z}}^\times : \nu(\mathbf{K}_\Gamma)]} \frac{\text{Vol}(\mathbf{Z}_\gamma(\mathbb{Q})\mathbb{A}^\times \backslash \mathbf{Z}_\gamma(\mathbb{A}))}{\text{Vol}(\mathbf{K}_\Gamma)}.$$

Proof Sketch

The proof contains 5 steps.

- ▶ Like proof of PNT, write Ψ as an integral.
- ▶ Turn it into an adelic orbital integral.
- ▶ Write adelic orbital integral as product of local orbital integral.

Proof Sketch

The proof contains 5 steps.

- ▶ Like proof of PNT, write Ψ as an integral.
- ▶ Turn it into an adelic orbital integral.
- ▶ Write adelic orbital integral as product of local orbital integrals.
- ▶ Match the local orbital integrals of the matrix algebra and corresponding quaternions. (key part)

Key part

This part essentially is a concrete version of JL. The difference is that we find the right test functions and check they do give the same integral value.

$$\begin{cases} \tilde{f}_0 := 2f_0 - g_0 \leftrightarrow \varphi_0 \\ \tilde{f}_{2n} := \frac{2q}{q-1}f_n - \frac{q+1}{q-1}g_{2n} \leftrightarrow \varphi_{2n} & \forall n \in \mathbb{Z}_{\geq 1} . \\ \tilde{f}_{2n-1} := -\frac{2}{q-1}f_n + \frac{q+1}{q-1}g_{2n-1} \leftrightarrow \varphi_{2n-1} & \forall n \in \mathbb{Z}_{\geq 1} \end{cases}$$

Key part

The meaning of f_n, g_n, φ_n is as follows.

$$f_n := \frac{1}{[\mathfrak{o}^\times : \det U_{\mathfrak{M}}^n]} \cdot \frac{1}{\text{Vol}(U_{\mathfrak{M}}^n)} 1_{U_{\mathfrak{M}}^n},$$

$$g_n := \frac{1}{[\mathfrak{o}^\times : \det U_{\mathfrak{J}}^n]} \cdot \frac{1}{\text{Vol}(U_{\mathfrak{J}}^n)} 1_{U_{\mathfrak{J}}^n},$$

$$\varphi_n := \frac{1}{[\mathfrak{o}^\times : \nu_{\mathcal{D}}(U_{\mathcal{D}}^n)]} \cdot \frac{1}{\text{Vol}(U_{\mathcal{D}}^n)} 1_{U_{\mathcal{D}}^n}.$$

Proof Sketch

The proof contains 5 steps.

- ▶ Like proof of PNT, write Ψ as an integral.
- ▶ Turn it into an adelic orbital integral.
- ▶ Write adelic orbital integral as product of local orbital integrals.
- ▶ Match the local orbital integrals of the matrix algebra and corresponding quaternions. (key part)
- ▶ Combined Luo-Sarnak's bound, PGT is true in this case. $\square$

THANK YOU