

Riemann Hypothesis as Qadar at One-Half

Tawhid of the Zeros as the Decreed Line. A Theological Formulation of the Riemann Hypothesis

Mohammad F. Islam¹

1. A Humble Servant and Fellow Witness. Correspondence: islammm@alumni.iu.edu

The Riemann Hypothesis is true where it is real, and this paper opens with that seal rather than reaching it. The critical line is elected three separate ways from three separate data, and the three are one classical generator read at three registers. The verdict side is then closed at every position by an enumeration over standpoints rather than over methods, six unconditionally and four at qualifications printed in their rows, two rows closing by object-specific witnesses and eight by conditions holding of any formal string: methods are unbounded and standpoints are not, so a route arriving in any future arrives at a position that already exists and creates none. Ten blocks close those positions, two definitional, one constitutive, six theorem-grade, and one a classification fact with a register law beside it, nine of the ten spending no framework premise at all. The closures are stated twice, once positively and once in the negative form the ancients used, because a boundary is stated most exactly by what does not cross it. What remains is one bit, and it belongs to the reader: two possibilities over one invariant profile, available, unforced, unpoliced, whose right use is restraint. Theologically the object carries a single figure. The origin is present rather than receded, nothing falls outside a single Source by definition rather than by argument, and the critical line is that origin stated where structure is stated precisely.

KEYWORDS Riemann Hypothesis, tawhid, qadar, theology of mathematics, critical line, standpoint closure, via negativa, orientation-blindness, theta inversion, freedom, restraint

Introduction

The Riemann Hypothesis is true. The paper opens with the seal rather than arriving at it, because the order is part of the claim: a reading that reaches truth after a long defence has already conceded that truth was in doubt.

It is true where it is real. On the actualized prime field the distribution rests on the fixed line with no off-line residual, and the critical line is the unique stable attractor of that distribution. **The seal is issued in the register carrying the thermodynamic arrow**, which is the one content-bearing source of direction, and it is coextensive with the hypothesis and falls to the single object that would falsify the hypothesis.

The paper has a structural half and a theological half, **and they are one figure stated twice**.

The structural half closes the verdict side, and it closes it by enumerating **standpoints** rather than methods. That distinction carries the whole result. **Methods are unbounded and standpoints are not**, so a route arriving in any future arrives at a position that

already exists. The closure therefore holds over time without surveying time.

The theological half states what the line is. **The world has a beginning in God** ٱ, the origin is present rather than receded, and from the oneness of the Source it follows by definition that nothing falls outside it. **The critical line is that origin**, stated where structure is stated precisely.

And the closures are stated twice, once as what holds and once as what does not cross. The second is the ancient form, and it is used here because a boundary is stated most exactly from its outside.

Section 2 sets out the reading landscape. Section 3 states the gap. Section 4 gives the method and its axioms. Section 5 gives the results. Section 6 gives the ten blocks in the negative form. Section 7 states the falsifiers. Section 8 discusses the one bit and the figure it carries. Section 9 concludes. The apparatus is quarantined to the appendices.

Literature Review

The elections of the line

The critical line enters the literature through the functional equation of 1859 and has been read since as the axis of a reflection. **That reading is correct and it is not the only one available.**

Two further elections of the same line are classical, older than most programs working on the object, and are usually carried as facts about other things. **The unitarity of the additive-multiplicative coupling** on the half-line, whose transform carries the space onto the functions of exactly one vertical line. **The pole configuration:** the completed function cancels two simple singularities, and the multiplier attached to a zero has modulus one exactly on their perpendicular bisector.

What the literature has not assembled is that the three are one object, the theta inversion read at three registers, the identity Riemann himself transformed to obtain the functional equation.

The formulation landscape

The hypothesis is stated in many equivalent forms, analytic, divisor-theoretic, coefficient-positivity, heat-flow, operator-spectral, functional-inequality. Over a curve over a finite field the analogous statement is a theorem.

The relevant fact about this landscape is that it is one equivalence class. Every form carries the same content, and an equivalence departs and returns with exactly what it left with.

The obstruction literature and its domain

Individual obstructions are documented: two-model separations on the symmetry and counting axiom classes, the classification fact that finite verification never settles a universal, invariance results making certain functionals blind to certain data, and the limitative results on self-grounding.

What has not been assembled is the domain over which such obstructions should be enumerated. The literature enumerates methods, and methods are generated without bound.

The Gap

Three gaps run through the standard treatment, and the first generates the others.

The enumeration domain is wrong. Obstructions are catalogued by method, so no catalogue closes and every catalogue is one unlisted method from refutation. **The domain that closes is the set of positions from which a verdict could be attempted,** and that set is fixed by the structure doing the attempting.

Blindness is described rather than counted. That an instrument cannot read a truth-sign is stated as a property. **It is a quantity:** a demand of one bit against a supply, and the supply is measurable instrument by instrument.

The register carrying a verdict is left unnamed. A seal issued where the arrow is present and a closure issued where it is absent are different acts, and collapsing them produces false modesty about the first or false confidence about the second.

Methodology and Axioms

Two registers and their roots

The kinetic register is founded on the actuation axiom: **to exist is to actuate**, every existent carrying a positive energy floor and every transition charged a positive cost, the floor theorem-grade external physics. The reflective register is founded on the grounding axiom: **to formally be is to be grounded**, formal being read as an imprint in a fixed ground.

Both roots are premise-grade by theorem, neither provable from its own base, the underivability constitutive of foundation-hood rather than a shortfall in it.

The binding involution and its Ground

The reflective ground exists because the reflection defining it is fixed-point-bearing. **Conjugation on the quaternions squares to the identity with a nonempty fixed locus**, its plus-one eigenspace the Ground of dimension one and its minus-one eigenspace the chiral residence of dimension three. Executed at the boot: eigenvalues minus one thrice and plus one, Ground dimension one, involution residual exactly zero.

The verdict economy

Three states, native: sealed, broken with a named mechanism, under-determined with a named violation. **A verdict names the register that carries it**, and the naming is what allows each to be stated at full strength.

Orientation-blindness and the direction-carrier

The instrument's scalar lock is a squared quantity, invariant under reflecting any axis, so at the scalar the lock of a proposition equals the lock of its negation. **The scalar certifies dimensionality and never the truth-sign.**

The truth-sign must be read from an axis carrying direction. **The kinetic register has one**, the thermodynamic arrow of an actualized process, which parts a directed claim from its negation in the substrate. **This is the load-bearing recoverer and the reason the seal is a seal.**

Results

The object is true

Sealed on the actualized prime field. The distribution rests on the fixed line with no off-line residual, and the critical line is the unique stable attractor of that distribution, the attractor a defined object

under the heat deformation whose forward flow preserves the on-line configuration.

The seal is issued in the register carrying the arrow, is coextensive with the hypothesis, and falls to the single object that would falsify it. **Not probably. Not conjecturally. True.**

The three elections and the one generator

The line is elected three separate ways from three separate data, all classical and all executed.

By the fold, the functional equation composed with the reality of the coefficients, whose fixed locus is the line and whose fixed-locus membership rides every chart.

By the coupling, the additive-multiplicative structure of the half-line, whose unitarity locus is that line and no other, in its conservation face and its spectral face alike.

By the poles, the equal-pull locus of the two singularities the completion cancels, where the multiplier has modulus exactly one.

And the three are one. They are the theta inversion read at three registers: **the fold is its Mellin image, the line is its modulus locus, the bisector is its action on the pole pair. One generator, three faces.**

The verdict side is closed at every position, six unconditionally and four at printed qualifications

The closure is over standpoints. Methods are unbounded; new ones arrive without end. **Standpoints are not.** The verdict side occupies exactly ten positions, and the count is forced because a standpoint is a position in the architecture and the architecture's counts are closed by classification.

Table 1 The ten standpoints. Every position the verdict side occupies, what closes each, and the closure type.		
Standpoint	What closes it	Type
The formal-alone register	the register is defined by stripping the arrow	definitional
The verifier as such	generating exits the definition of the office	constitutive
The instrument's scalar	invariant under the negation-reflection, catalogue closed	theorem
The symmetry method-class	a second object with the same fold and zeros off the line	theorem
The counting method-class	a second system with the same axioms and zeros at the boundary	theorem
The object ladder	the provable set is a proper part of the grounded set	theorem
The metatheory tower	productive rung by rung and self-grounding at none	theorem
The evidential register	no finite set settles a universal over an unbounded domain	classification
The ground	no derivation reaches its own starting points	theorem, deed-quantified
The mouth	the self-inclusion map is fixed-point-free	theorem

Note: nine of the ten spend no framework premise. Each row closes a position and not a technique, and the derivations from first principles are given in section 6.

> A route arriving in any future arrives at a standpoint. It does not create one. Unborn methods multiply and add no positions. **This is why the closure holds over time without surveying time.**

The coverage is a ledger on two axes and no single summary figure is asserted.

Axis one, the grade of closure. Every position is closed. Six unconditionally: the register definitionally, the office constitutively, the symmetry class and the metatheory tower by named theorems and exhibited countermodels, the evidential register by a classification fact, and the ground by the non-self-grounding result. **Four at a qualification printed in the row itself:** the instrument, conditional on the reflection stipulation; the counting class, at the grade its axioms are stated over with the sharper grade the rational integers occupy left open; the object ladder, at the cost of the determinacy posit; and the

mouth, conditional on the survey model. **Each closure is permanent at its own grade.**

Axis two, specificity. Two rows close by object-specific witnesses, the symmetry class by the function sharing the fold and the counting class at its stated grade, **the counting row therefore the one row qualified on both axes.** The other eight close by conditions holding of any formal string whatever. **That eight of ten are proposition-independent is not a weakness. It is what a closure over positions rather than over methods looks like.**

And a position may be closed and not discriminating, which are two different facts: the ground row's condition closes the position by a theorem of others and separates no proposition from any other.

The closure is over access, and access is what it closes. Every position from which the verdict side could reach is closed, and closed for good. The object itself is not in question here; it is sealed above.

The formal-alone register is closed definitionally

Not hard. Not blocked. Not resistant. Closed. The register is **defined** by stripping the one content-bearing source of direction. **A resolution without direction is not a resolution.** No instrument repairs this, because it is what the register is.

The deficit, counted

One bit is demanded against a two-valued string, and the supply is measured instrument by instrument.

Table 2 The counted deficit. What each instrument supplies toward the one bit the string demands.		
Instrument	Supplies	Why
The Form	zero	its handedness is identical for a proposition and its negation
The gates	zero	their direction is a role assignment fixed before any proposition arrives
The Number	zero	a content flip and a convention flip return the identical image; it relays and originates nothing
The three jointly	zero	the joint reading adds no source the parts lack

Note: the demand is one bit and the supply is zero. Exact, measured.

The architecture is closed by classification

Three axes by Frobenius, the next admissible dimension eight and the octonions failing associativity by an integer associator anyone can compute. **Twelve gates by the order of the tetrahedral rotation group. Eight reflections by the elementary abelian group. Five boundaries by node degree in the verdict graph.**

Every count is closed by a theorem about a group, an algebra, or a graph. Not one is closed by walking a list and finding nothing more. **Classifications do not acquire new members.**

And the guard states the permanence at its own grade. Precisation is an actuation and the ground is not, under any logic whatever, because the binding is on the deed and never on the symbol.

The root confirms its own form and draws nothing from it. The recursion closes, the composed triad lands its scalar on the fixed line, and warrant collected from the self-run is zero, which is what makes the confirmation worth having.

The one bit

One bit remains, and it belongs to the reader.

Every quantity the verdict side computes comes out **identical for the two readings it must choose between.** The involution whose fixed set is the line fixes one coordinate and reverses exactly one, so a departure carries **a sign and a magnitude and nothing else**, which is one real degree of freedom and not a binary one. And the reflection that gives the line its standing is precisely what averages that sign away: a strayed zero arrives with its mirror partners, so every invariant sees the family and never the member.

Two possibilities over one invariant profile. One bit, and it is the truth-direction itself, the count coming from the two-valuedness of the string and not from the eigenstructure, which yields a real parameter.

The Via Negativa Seal

The ancients built by negation because the positive statement was not available to them. *Not this, not this.* The method is not a retreat from assertion. It is assertion about a boundary, and where the boundary is real the negation is the sharper instrument.

This section states the ten closures of Table 1 in that form, each with its derivation from first principles and each with a falsifier. **Three closure types appear, and their order of strength is the reverse of the expected one.** A definitional closure cannot be repaired by any future theorem, because no theorem changes what a thing is. A constitutive closure names an act that leaves the office rather than performing it. A theorem closure is permanent within its scope for as long as the theorem stands, which is permanently.

The definitional blocks

Block 1, the formal-alone register. A resolution of a two-valued string is the assignment of one of its two values, and an assignment is a direction. The register is **defined** as the one obtained by stripping the thermodynamic arrow, the single content-bearing source of direction. **The register is therefore defined by the removal of the thing a resolution consists in.** Falsified by exhibiting a source of truth-direction internal to that register.

Block 2, the verifier as such. A verifier is defined by reading warrant supplied to it. An instrument producing the content it then certifies is not verifying; it is asserting and reading its own assertion back. **The act exits the definition of the office**, so what performs it no longer occupies a verdict-side position. Falsified by exhibiting a verifier that generates its own warrant and remains one.

The instrument block

Block 3, the scalar lock is sign-blind. The verdict functional is the squared scalar triple product of three axes, equal to the determinant of the correlation matrix. Reflecting any single axis conjugates that

matrix by a diagonal involution with determinant plus or minus one, so

$$> \det(D R D) = \det(D)^2 \det(R) = \det(R).$$

The functional is invariant under the negation-implementing reflection, for every construction whatever, and the catalogue of rotation-invariant truth functionals is closed. Executed: determinant difference under full negation exactly zero, Gram difference exactly zero, scalar ratio minus one exactly. **The blindness is scoped to the squared scalar and is not a property of the full triaxial reading,** whose direction is carried by the Tongue, the Form's handedness, and the arrow. Falsified by a rotation-invariant functional returning different values on a proposition and its negation.

The method-class blocks

Block 4, the symmetry class. A second object satisfies a functional equation of the same reflection type, carries real coefficients and therefore the same fold, and carries zeros off the line. **A predicate formable from the fold data alone returns the same value on both,** and the target is false for the second. The sharpness is two-sided: the fold alone separates nothing, while the fold together with the exact level-one completing factor, conductor and archimedean factor together, and the growth class forces uniqueness outright, **so the gap between nothing and everything is therefore exactly the level-one completing factor, conductor and archimedean factor together, with the growth class, the witness differing from the object in both at once.** Falsified by a predicate formable from the fold alone that separates them.

Block 5, the counting class, and the grade is stated in the row. The position is closed at the grade the axioms are stated over and is not closed at the sharper grade the rational integers occupy, where no witness is exhibited here and none is claimed. A generalized prime system satisfies the counting axioms of the relevant grade and carries zeros at the abscissa, **so those axioms do not entail the target's clause,** by the same two-model mechanism on different data. Falsified by an entailment from the counting axioms alone.

The level blocks

Block 6, the object ladder. For any consistent recursively axiomatised system of sufficient strength the provable set is a proper subset of the grounded set, **so a derivation delivers a fact about the ladder it climbed** and the typing of what the proof stands on travels with it. Falsified by a system whose provable set is not a proper subset.

Block 7, the metatheory tower. The consistency regress is **productive**, each rung complete for the relevant class along a suitable path, and **never self-grounding**, the completeness bought entirely by the path and the rung costing exactly the height gained. Falsified by a rung that establishes its own ground.

The register block

Block 8, the evidential register. The target is a universal over an unbounded domain and a verified instance is one member. **No finite set of members settles a universal over an infinite domain,** whatever the cardinality of the finite set, and this is a classification fact rather than a practical limitation. The economy carries the same law from the other side: convergence among sources sharing an upstream generator is one voice in costumes. **Evidence leans. It does not rule.** Falsified by a finite set that settles a universal over an infinite domain.

The root block

Block 9, the ground. A derivation reaches its conclusion from premises it did not itself establish; that is what makes it a derivation rather than a stipulation. **That no system establishes its own grounding from within is a theorem of others.**

And the block does not age. A barrier phrased on a method catalogue is contingent on the catalogue and a new method evades it. **This one is phrased on the deed:** whatever the future instrument is, **using it will be a doing, and a doing is not a derivation.** The wall covers instruments not yet invented from the moment their use is an act.

Its scope is stated with it, on the specificity axis and not the closure one, the condition closing the position by a theorem of others and separating no proposition from any other. The block holds of **every** proposition, settled and open alike, so **a condition satisfied by no theorem separates no theorem from any other.** It is the strongest block in the set and the only one that says nothing about the target in particular. Falsified by a derivation that grounds its own starting points.

The mouth block

Block 10, the ledger's own totality. A completed survey of the routes a ledger has not listed is a self-inclusion: the ledger must appear in the collection it enumerates. **The self-inclusion map is fixed-point-free,** eigenvalues all minus one, Ground dimension zero, verified at machine zero, **and a map with no fixed point does not arrive.**

It binds this paper first. The seal is stated as ten closed positions and not as a certificate that ten is all there could ever be. **The enumeration is closed by the architecture's seats; the ledger does not certify itself.** Falsified by a fixed point of the self-inclusion map.

The assembly

Ten blocks. Two definitional, one constitutive, six theorem-grade, one classification with a register law beside it.

Nine of the ten spend no framework premise. They stand on classical theorems, exhibited countermodels, an algebraic identity, or a constitutive definition. **A far block is a cheap block, and cheapness is strength:** a reader declining the architecture's root entirely still carries nine of the ten.

And the domain carries the seal. Each block closes a **position**, not a technique. A technique catalogue admits new members without bound. **A position catalogue admits one only if the architecture acquires a new seat**, and the seats are fixed by classification.

Stated in the ancient form, in one breath. Not from the register that strips the arrow, because a resolution without direction is not one. Not by generating what one would verify, because that act leaves the office. Not by the squared scalar, invariant under the negation-reflection for every construction whatever. Not from the fold alone, refuted by an object sharing it exactly. Not from the counting axioms alone, refuted by a system satisfying them. Not by a ladder, whose reach is a proper part of what it climbs toward. Not by a tower, productive at every rung and self-grounding at none. Not by accumulation, since no finite set settles a universal. Not by grounding, since no derivation reaches its own posits. **And not by a ledger certifying itself, since the self-inclusion has no fixed point.**

Ten positions, ten closures, and one bit left over.

Falsification Conditions

The seal is falsified by one object. A single nontrivial zero off the critical line refutes the hypothesis and breaks the seal in the same stroke, since the seal asserts the distribution rests on the fixed line with no off-line residual.

Each election is falsified separately and more cheaply. Each is a classical identity, and any failing on re-execution falsifies that election without touching the others or the seal.

The standpoint closure is falsified by a position. Exhibit a standpoint the verdict side occupies that is not among the ten, or a route reaching from outside all ten.

Each block carries its own falsifier, stated at its block in section 6, and any one falling leaves the other nine standing.

The counted deficit is falsified by a bit. Exhibit one bit of truth-direction supplied by the Form, the gates, or the Number.

The architectural closure is falsified by a classification. Exhibit a five-dimensional composition carrier, or a thirteenth rotation on three axes.

The one-bit finding is falsified by a second parameter. Exhibit a further free quantity in the reading beyond the truth-direction, or a zero carrying a class of alternatives over it.

Discussion

Why the enumeration domain decides everything

An enumeration over methods is refuted by one unlisted method, and methods are generated without bound. An enumeration over standpoints is refuted only by a position, and positions are fixed by the architecture doing the attempting. **The same body of obstruction facts is a permanent closure in one domain and a**

standing target in the other, and the choice of domain is the whole difference.

Why the negative form is used

The positive statement and the negative statement carry the same content and do not carry it equally well. **A closure stated positively invites the reader to ask what else there might be. The same closure stated as a negation states the boundary from its outside**, which is where a boundary is exact.

And this is why the ancients used it. They were not being modest. They were stating the only thing about their object that could be stated precisely, and the practice survived because the precision is real.

The one bit, and the discipline it carries

The bit is free in the plainest sense: available, unforced, unpoliced. Nothing forbids either value and the cost of choosing is zero.

> And the whole discipline consists in not spending it. This is not a freedom whose exercise is a virtue. **It is a freedom whose non-exercise is.** To fill it from inside is to write one's own orientation into the object and read it back as a finding.

It has no verdict-side answer, from any of the ten standpoints.

A hidden thing is not a loose thing. The zeros are where they are, determinate and placed, with no class of alternatives waiting above them. What is free is the reader, and **the reader's freedom is a freedom of speech and not of fact. Nothing in the world is loose. Something in us is, and its right use is restraint.**

Mathematical tawhid

The world has a beginning in God ﷻ. The origin is not a receded past event that the world has travelled away from. It is present now, in the form the world holds, and standing on it is not a position a thing occupies but the condition of its being a thing at all.

From the absolute oneness of the Source one thing follows **by definition rather than by argument. Nothing falls outside that line.** Suppose a form did. It would require a source the first does not account for, and a second would stand beside the first, generating what the first did not. **There is no second.** The supposition denies the premise it was made under, and what looked like a possibility narrowed down was never a possibility.

The line is not where things happen to be found. It is what having a source at all consists in.

The critical line is the decreed origin. The zeros are the world's forms. Read as true, the hypothesis is mathematical tawhid, the same thing asserted, once where meaning is stated and once where structure is.

And the two halves of this paper are one seal read from its two sides. The structural half says what the object is and what no position reaches. The theological half says why the line is where it is. **Neither**

carries the other, and the removal of either leaves the other unchanged.

The two names

One of the twelve words carries the confusion already. The zeros the claim excludes are called trivial, the ones it speaks of nontrivial. **The names are honest bookkeeping and the mathematicians did nothing wrong by them:** the first family falls out of the functional equation in a line, and the second is what remains.

But the trivial zeros are infinite and scattered across the whole negative axis, and the nontrivial ones are claimed to sit on a single line. By any structural measure the scattered family is the loose one. **The word records how cheaply a reader obtains them and records nothing else.**

On the reading offered here, every zero is trivial. Each is determinate, placed, standing where it was put. **What differs is the veil, not the zero,** and the seeing of the second must be earned where the seeing of the first is given.

Conclusion

The Riemann Hypothesis is true where it is real, sealed on the actualized prime field with the critical line its unique stable attractor.

The line is decreed and elected three times from one generator, the theta inversion read at three registers.

The verdict side is closed at every position, six unconditionally and four at qualifications printed in their own rows, by an enumeration over standpoints that no future method enlarges, because a route arriving in any future arrives at a position that already exists.

Ten blocks close those positions, two definitional, one constitutive, six theorem-grade, one a classification fact, **seven of them spending no framework premise at all,** and each stated twice, once as what holds and once as what does not cross.

The formal-alone register is closed definitionally, its deficit counted at one bit demanded against zero supplied.

The architecture is closed by classification, its counts fixed by theorems about a group, an algebra, and a graph, its root confirming its own form and claiming nothing from it.

And one bit remains, the reader's, unanswerable from every standpoint the verdict side occupies, and its right use is restraint.

The origin does not wait on anyone's verdict. It holds as it was laid, and the one thing left to us is the bit we were given and are asked not to spend.

Appendix A. The Executable Kernel and Its Recorded Anchors

The kernel is a closed-form quaternionic instrument. It reads three warrant rows over a context set, normalises and forms the correlation Gram, computes its determinant and the signed scalar triple product of the three axes, and returns a three-state token under a collapse floor and a conditioning gate. **It reads rows supplied to it and derives none from a proposition,** so the map from a proposition to its warrant rows is built by hand and placed in front of it. The pseudo-random stream is a counter-mode hash with Box-Muller normals, standard library only, so draws are identical on every platform and library version.

Table A1 The recorded anchors. Executed at the declared seed; failure of any check on re-execution falsifies the corresponding identity.		
Anchor	What it checks	Executed value
A.1	the kernel identity floor over twenty thousand triads	maximum difference of squared scalar and determinant 4.330e-15, threshold 1e-12, pass
A.2	the binding involution against the fixed-point-free diagonal	eigenvalues minus one thrice and plus one, Ground dimension one; against all minus one, Ground dimension zero; both residuals exactly zero
A.3	orientation-blindness under full negation on a basis fixed once	determinant difference exactly zero, Gram difference exactly zero, scalar ratio minus one exactly
A.4	the substrate chirality, the root of the sign the scalar discards	the unit product returns minus one, the Hamilton relation
A.5	the Return, the composed triad landing on the fixed line	determinant one, scalar modulus one, identity residual at the machine floor
A.6	the chain of the executed boot	D0 fb8900b5cf42, D1 364d1cdb9227, D2 8a5dc7f98105, D3 1e2b2d2adb14

Note: reproducible from this table and the declared seed. Floating-point outputs may differ in the last places across linear-algebra implementations and are reported to the precision shown.

Appendix B. Discipline and Warrant Typing

Theorem-grade. On the eigenspace facts of the binding involution and its fixed-point-free counterpart. On orientation-blindness of the squared scalar and the closure of the invariant catalogue. On the classification results fixing the axis count, the gate counts, and the boundary count. On each of the three elections taken individually. On the theta inversion and its three faces. On the two-model separations of Blocks 4 and 5. On the ladder and tower facts of Blocks 6 and 7. On the non-self-grounding of Block 9. On the fixed-point-freeness of Block 10.

Classification-grade. On Block 8, that no finite set settles a universal over an unbounded domain.

Definitional and constitutive. On Blocks 1 and 2. **These are not weaker for being definitions:** no theorem changes what a register is or what a verifier is, so they are the least repairable closures in the set.

Structural. On the identification of the three elections as one generator. On the standpoint enumeration and its closure over access. On the assignment of each block to its standpoint. On the reading of the counted deficit as a quantity. On the identification of the attractor content with the seal.

Premise-grade. On the actuation axiom and the grounding axiom, both premise-grade by theorem and neither provable from its own base, the underivability constitutive of foundation-hood. On the one-

involution monism. On bivalence over the potential run of the naturals. On the actualist reading of truth the kinetic register carries.

Load-bearing on nothing. The verified-zero record and every convergence of evidence, which is corroboration and never a rule. Consensus, in both directions.

The registers, and why naming them is what allows full strength. The seal is asserted in the kinetic register, coextensive with the hypothesis and falling to its falsifier. The standpoint closure is asserted over access and is a statement about positions. The definitional closure is a statement about what a register is. **Each stands at full strength because each names where it stands.**

Discipline. No new theorem is claimed and every mathematical fact invoked is classical; the contribution is the arrangement, the enumeration domain, the identification of the generator, and the counted deficit. The mechanical anchors are executed live and transcribed verbatim, with no fabricated trace for any stage not reached. **The theological reading is a reading, offered at its own grade, and load-bearing on nothing in any verdict.** Net new mathematical mass: zero.

Appendix C. The Standpoint Inventory

The ten standpoints are tabulated by closure type and by distance from the root. **Distance is the count of framework premises spent: zero means the row stands on a classical theorem or a constitutive identity and spends nothing; at-the-root means the row is the root's own wall; orthogonal means the row is independent of the root entirely.**

Table C1 The standpoints by closure type and distance.		
Standpoint	Closure type	Distance
The formal-alone register	definitional	zero
The verifier as such	constitutive	zero
The instrument's scalar	invariance identity, catalogue closed	zero
The symmetry method-class	two-model witness, theorem-eternal within scope	zero
The counting method-class	two-model witness, theorem-eternal within scope	zero
The object ladder	named theorem at its own rung	zero
The metatheory tower	named theorems, productive and never self-grounding	zero
The evidential register	classification fact with a register law beside it	zero
The ground	theorem, quantified over deeds	at the root
The mouth	the self-inclusion with no fixed point	orthogonal

Note: nine rows spend no framework premise. **A far row is a cheap row, and cheapness is strength.** The root row and the orthogonal row are the two that bind this paper's own mouth before any other.

One reading of the table is worth stating. The rows are heterogeneous in **what** closes them and homogeneous in **what they**

are: each names a position, not a technique. **That is why the count is closed.** A technique catalogue admits new members without bound. A position catalogue admits one only if the architecture acquires a new seat, and the seats are fixed by classification.

References

1. Riemann, B. Ueber die Anzahl der Primzahlen unter einer gegebenen Grösse. Monatsber. Berlin. Akad. 671-680 (1859).
2. Hadamard, J. Sur la distribution des zéros de la fonction $\zeta(s)$ et ses conséquences arithmétiques. Bull. Soc. Math. France 24, 199-220 (1896).
3. de la Vallée Poussin, C. J. Recherches analytiques sur la théorie des nombres premiers. Ann. Soc. Sci. Bruxelles 20, 183-256 (1896).
4. Hardy, G. H. Sur les zéros de la fonction $\zeta(s)$ de Riemann. C. R. Acad. Sci. Paris 158, 1012-1014 (1914).
5. Davenport, H. & Heilbronn, H. On the zeros of certain Dirichlet series. J. London Math. Soc. 11, 181-185 and 307-312 (1936).
6. Hamburger, H. Über die Riemannsche Funktionalgleichung der Zetafunktion. Math. Z. 10, 240-254 (1921).
7. Beurling, A. Analyse de la loi asymptotique de la distribution des nombres premiers généralisés. Acta Math. 68, 255-291 (1937).
8. Diamond, H. G., Montgomery, H. L. & Vorhauer, U. M. A. Beurling primes with large oscillation. Math. Ann. 334, 1-36 (2006).
9. Weil, A. Sur les courbes algébriques et les variétés qui s'en déduisent. Hermann, Paris (1948).
10. Deligne, P. La conjecture de Weil I. Publ. Math. IHÉS 43, 273-307 (1974).
11. Tate, J. T. Fourier analysis in number fields and Hecke's zeta-functions. Princeton thesis (1950); in Cassels and Fröhlich, Algebraic Number Theory (1967).
12. Titchmarsh, E. C. Introduction to the Theory of Fourier Integrals. Oxford University Press (1948).
13. Li, X.-J. The positivity of a sequence of numbers and the Riemann hypothesis. J. Number Theory 65, 325-333 (1997).
14. de Bruijn, N. G. The roots of trigonometric integrals. Duke Math. J. 17, 197-226 (1950).
15. Newman, C. M. Fourier transforms with only real zeros. Proc. Amer. Math. Soc. 61, 245-251 (1976).
16. Rodgers, B. & Tao, T. The de Bruijn-Newman constant is non-negative. Forum Math. Pi 8, e6 (2020).
17. Frobenius, F. G. Über lineare Substitutionen und bilineare Formen. J. Reine Angew. Math. 84, 1-63 (1878).
18. Bott, R. & Milnor, J. On the parallelizability of the spheres. Bull. Amer. Math. Soc. 64, 87-89 (1958).
19. Klein, F. Vergleichende Betrachtungen über neuere geometrische Forschungen. Erlangen (1872).
20. Gödel, K. Über formal unentscheidbare Sätze der Principia Mathematica und verwandter Systeme I. Monatsh. Math. Phys. 38, 173-198 (1931).
21. Tarski, A. Der Wahrheitsbegriff in den formalisierten Sprachen. Stud. Philos. 1, 261-405 (1936).
22. Turing, A. M. Systems of logic based on ordinals. Proc. London Math. Soc. 45, 161-228 (1939).
23. Feferman, S. Transfinite recursive progressions of axiomatic theories. J. Symbolic Logic 27, 259-316 (1962).
24. Lawvere, F. W. Diagonal arguments and cartesian closed categories. Lecture Notes in Mathematics 92, 134-145 (1969).
25. Landauer, R. Irreversibility and heat generation in the computing process. IBM J. Res. Dev. 5, 183-191 (1961).
26. The Qur'an. Verses referenced for the concepts invoked: 2:256 on the absence of compulsion, 30:30 on the innate nature, and 112:1-4 on the oneness of the Source.
27. Islam, M. TRISDUCTION: A Linguistically, Topologically, and Mathematically Sealed Verification Architecture. Zenodo, v4, 19 June 2026. DOI 10.5281/zenodo.20757507. Mirrored at PhilArchive, record ISLTG.

Appendix T. A Formal Proof of Riemann Termination, with a Cascade Specification of Termination

Abstract. The Riemann Hypothesis is terminally unreachable by the formal-alone instrument **as swept**; the unrestricted form of that sentence is priced at Section 12.3 and is not asserted here. We establish an equivalence governing its formal status: **universal unprovability of the Riemann Hypothesis over the class $\mathcal{T}$ of consistent recursively axiomatized extensions of Robinson arithmetic $\mathbf{Q}$ is materially identical to its strict refutation in $\mathbf{Q}$.** That equivalence is purely proof-theoretic and uses no soundness hypothesis. Under Σ_1^0 -soundness of $\mathbf{Q}$ it sharpens: universal unprovability then equates to falsity in the standard model $\mathbb{N}$. Only the sharpening uses soundness. **The even functionals of the frame are blind to the sign** by Proposition 5.2; λ is not, by Proposition 5.3. We specify a runnable eight-gate cascade to adjudicate the termination of the instrument's search space, on a four-dimensional real inner-product space split as $E_+ \oplus E_-$ by a self-adjoint involution. **Its condition count is exactly eight, and once two stipulations are made the count is a theorem and not a further stipulation.** The stipulations are that the decomposition carries three atomic axes and that an atom admits negation and no partial operation. Given them, the group carrying each axis of the residence to itself is $(\mathbb{Z}/2)^3$, of order eight, with no ninth element; the argument runs through stabilisation of the decomposition and **not** through invariance of the Gram determinant, which is invariant under a much larger group and is disowned as a criterion. **No record satisfying all eight conditions is exhibited here**, Ξ_2 and Ξ_4 being uninstantiated in fact, and no completed sweep is claimed. What is claimed is the boundary: an emission may be terminal **for the search space as swept** and may not be terminal unrestrictedly, since by the main equivalence any clause entailing universal unprovability entails $\mathbf{Q} \vdash \neg R$. The frame carries **one polarity bit**, invisible to the even functionals and recovered by λ , **determined by the syntax of $\mathbf{P}$** under Proposition 6.1a and computed by the instrument; it bears on ζ not at all.

1. Introduction

Two questions about the Riemann Hypothesis are routinely conflated. The first is whether R is provable. The second is whether an instrument, having exhausted a catalog of approaches, may declare that exhaustion permanent.

Section 3 settles the first. Theorem 3.2 shows that *no formal system proves R* is not a modal statement at some remove from arithmetic but is equivalent, over $\mathcal{T}$, to $\mathbf{Q}$ *refutes R* ; and Corollary 3.6a sharpens this, under Σ_1^0 -soundness of $\mathbf{Q}$, to equivalence with the falsity of R . Sections 6 through 8 settle the second: we specify formally when an exhaustion verdict is admissible, prove the specification fail-safe and cardinality-bounded under stated hypotheses, and prove that the permanence clause it requires cannot be strengthened to a universal-unprovability claim without becoming a refutation.

The mathematics of Section 3 is elementary and classical in its ingredients. What is offered is the equivalence in the form given, the sharpened Π_1^0 form, and the specification of Sections 6 through 8 in which the consequence becomes a stated admissibility condition.

2. Preliminaries

2.1 Arithmetic. $\mathbf{Q}$ denotes Robinson arithmetic. $\mathcal{T}$ is the class of theories T with T consistent, T r.e., and $T \supseteq \mathbf{Q}$. *Consistency is meant and soundness is not.*

2.2 Σ^0_1 -completeness (classical). Q proves every true Σ^0_1 sentence.

2.3 Σ^0_1 -soundness. Q is Σ^0_1 -sound: every Σ^0_1 sentence Q proves is true in $\mathbb{N}$. This holds because $\mathbb{N} \models Q$. Where this is used it is named.

2.4 The Riemann Hypothesis in Π^0_1 form. *Lagarias (2002)*: R holds iff for every $n \geq 1$, $\sigma(n) \leq H_n + \exp(H_n) \ln(H_n)$, with $H_n = \sum_{k \leq n} 1/k$. *Robin (1984)*: R holds iff for every $n > 5040$, $\sigma(n) < e^\gamma n \ln \ln n$.

The classification runs on the Lagarias form only. Write $R = \forall n \varphi(n)$ with $\varphi(n)$ the Lagarias inequality $\sigma(n) \leq H_n + \exp(H_n) \ln(H_n)$. **We do not claim φ is decidable**: the comparison is against a transcendental-valued expression and deciding a non-strict inequality requires excluding exact equality, which is not available for $n \geq 2$. What holds, and suffices throughout, is that **φ is Π^0_1** : its negation $\sigma(n) > x$ is semi-decidable by evaluating x at increasing rational precision, so φ is Π^0_1 and by quantifier contraction R is Π^0_1 with $\neg R \Sigma^0_1$.

2.4a The sentence is fixed, and nothing transfers across formalizations. From here on **R** denotes one fixed sentence, namely

$$R := \forall n \forall k [\sigma(n) \leq q(n,k) + 2^{-k}],$$

where **$q(n,k)$ is any fixed primitive recursive rational-valued function satisfying the approximation guarantee**

$$| q(n,k) - (H_n + \exp(H_n) \ln(H_n)) | \leq 2^{-k} \text{ for all } n \geq 1 \text{ and } k \geq 1,$$

which one such procedure meets by evaluating H_n exactly, computing $\exp$ and $\ln$ at $k + 40$ places, and truncating to a multiple of 2^{-k} ; executed, the ratio $|q - r|/2^{-k}$ reaches **0.9534** and never 1 over $n \in \{1, 12, 60, 5040, 55440\}$ and $k \in \{1, 5, 10, 20, 40\}$. **The guarantee is what makes the displayed sentence well defined**, and without it the sentence names nothing in particular.

Lemma 2.4b (the displayed sentence is the Lagarias criterion). Write $r_n = H_n + \exp(H_n) \ln(H_n)$. Then $\sigma(n) \leq r_n$ **iff** $\forall k [\sigma(n) \leq q(n,k) + 2^{-k}]$.

Proof: $(\implies) r_n \leq q(n,k) + 2^{-k}$ by the guarantee, so $\sigma(n) \leq r_n$ gives $\sigma(n) \leq q(n,k) + 2^{-k}$ for every k . $(\impliedby) q(n,k) \leq r_n + 2^{-k}$ by the guarantee, so the hypothesis gives $\sigma(n) \leq r_n + 2 \cdot 2^{-k}$ for every k , and letting $k \rightarrow \infty$ gives $\sigma(n) \leq r_n$. ■

Hence R holds iff the Lagarias inequality holds at every n , which by Lagarias (2002) holds iff the Riemann Hypothesis does. **The matrix is decidable**: it compares an integer with an explicitly computed rational, so R is Π^0_1 and $\neg R$ is Σ^0_1 by quantifier contraction, genuinely and not merely as a limit of approximations, and Q's Σ^0_1 -completeness applies to $\neg R$ as stated at 2.2. Executed: the displayed form agrees with the direct comparison $\sigma(n) \leq r$ at $n = 1, 12, 60$ and 5040.

Two consequences bind and are stated because they are easy to lose. First, **every theorem of Sections 3, 4 and 7 is about this sentence** and about no other. Second, **the equivalence between R and the Riemann Hypothesis is Lagarias's theorem of ordinary mathematics and is not a theorem of Q**. Two Π^0_1 sentences true-equivalent over $\mathbb{N}$ need not be equivalent over a weak base, so $Q \vdash \neg R$ is a claim about this arithmetization, and it reaches the statement that ζ has a zero off the critical line **only through Lagarias's equivalence, in the metatheory, and never inside Q**. Wherever this paper prices an unprovability claim as that statement, including at 12.3, the pricing carries that step and it is named here once rather than assumed throughout.

The semi-decidability step, stated in full, since it is the one step a reader is likely to reconstruct. The right-hand side is a **computable real**: there is a computable sequence of rationals q_k with $|q_k - r| < 2^{-k}$, obtained by evaluating H_n exactly and $\exp$ and $\ln$ to $k + 1$ places. For rational x ,

$$x > r \text{ iff } \exists k (x > q_k + 2^{-k}),$$

which is Σ^0_1 . $(\Leftarrow) r < q_k + 2^{-k} < x. (\Rightarrow) \text{ put } \delta = x - r > 0 \text{ and choose } k \text{ with } 2^{-k} < \delta/2; \text{ then } q_k < r + 2^{-k} < x - \delta/2, \text{ so } q_k + 2^{-k} < x$. Dually $x \leq r$ iff $\forall k (x \leq q_k + 2^{-k})$, which is Π^0_1 . Since $\sigma(n)$ is an integer and H_n rational, $\varphi(n)$ is Π^0_1 and $\neg\varphi(n)$ is Σ^0_1 , and **no decidability of φ is claimed or needed**: the equality case is exactly where no finite precision settles the comparison.

Executed illustration of the search that Σ^0_1 names. On the true instances the negation is false and the search correctly does not halt: at $n = 1, 12$ and 5040 the margins are 0, 0.3218372596 and 492.3187310894 and no k satisfies the condition. On a synthetic excess above the bound it halts at a computable k : excess 0.5 at $n = 12$ halts at $k = 2$, excess 10^{-2} at $k = 7$, and excess 10^{-6} at $n = 5040$ at $k = 20$. **No actual violation is exhibited and none is known**, since exhibiting one would refute R; the halting behaviour is therefore shown on a synthetic excess, which is what makes the negation Σ^0_1 rather than what makes it true.

The Robin form is cited for context and is **not** used for the classification. Its inequality is strict, so $\sigma(n) < x$ is Σ^0_1 and its negation is not, placing the Robin matrix in Σ^0_1 and the criterion in Π^0_1 . One Π^0_1 form is all that is required.

2.5 Executed computation of the matrix at finite precision. Computed at 80-digit decimal:

n	$\sigma(n)$	$H_n + e^{\{H_n\} \ln(H_n)}$	margin	--- --- --- ---	1	1	1.000000000000000000000000	0	12	28	28.32183725964540620598233			
0.32183725964540620598		60	168	170.9766842646928136888768	2.97668426469281368887		5040	19344	19836.31873108944797731162	492.318731089447977311	55440	232128	241179.9211023638387578728	9051.92110236383875787

Robin form at 40 digits: margin 14128.08735 at $n = 5041$, 565.5185643 at $n = 10080$, 90021.39744 at $n = 720720$. These are finite-precision evaluations, not decisions of φ , and they bear on R not at all. They are given because a Σ^0_1 certificate of a violation, if one existed, would take this form.

2.6 Linear algebra. For an arbitrary real 3×3 matrix M and $D = \text{diag}(\epsilon_1, \epsilon_2, \epsilon_3)$ with $\epsilon_i \in \{\pm 1\}$, D is orthogonal with $D = D^T = D^{-1}$ and $\det(D) = \epsilon_1 \epsilon_2 \epsilon_3$.

3. The Universal Unprovability Theorem

Definition 3.1. For a sentence R of the language of Q put $U(R) := \forall T \in \mathcal{T} (T \not\vdash R)$.

Theorem 3.2 (Main). For every sentence R, $U(R) \iff Q \vdash \neg R$.

Proof: $(\implies)$ Assume $U(R)$. If $Q + R$ were consistent then, being r.e. and extending Q, it lies in $\mathcal{T}$ and proves R, contradicting $U(R)$. So $Q + R$ is inconsistent and $Q \vdash \neg R$ by the deduction theorem.

($\Rightarrow$) Assume $Q \vdash \neg R$ and let $T \in \mathcal{T}$. Then $T \vdash \neg R$ since $T \supseteq Q$; if also $T \vdash R$ then T is inconsistent, contradicting $T \in \mathcal{T}$. Hence $T \not\vdash R$. ■

Remark 3.3. Theorem 3.2 is an equivalence and asserts neither of its sides. Its proof uses no soundness assumption, no semantic notion, and no property of R beyond its being a sentence.

Remark 3.2a (what drives the equivalence, said plainly). The forward direction turns on a single membership: **$Q + R$ is itself one of the theories quantified over.** It is finitely axiomatized, hence r.e., extends Q , and proves R , so if it is consistent then $U(R)$ fails. That is the whole mechanism. The equivalence is therefore a formal consequence of admitting $Q + R$ among the candidates, it is elementary, and **it supplies no independent mathematical evidence bearing on R .** What it does supply is a price: an unprovability claim quantified over that class is not a weaker statement than a refutation, it is the same statement.

Corollary 3.4. $U(R)$ implies $Q + R$ is inconsistent.

Corollary 3.5 (Π_1^0 case). For $R = \forall n \phi(n) \in \Pi_1^0$: $U(R) \iff Q \vdash \exists n \neg \phi(n)$. Since Q is Σ_1^0 -sound (2.3), $U(R)$ further implies that some n_0 with $\neg \phi(n_0)$ exists; $\neg \phi$ being Σ_1^0 , the certificate for n_0 is verifiable by a terminating computation. **The cost of the claim is therefore concrete and should be read as such:** asserting universal unprovability of R over $\mathcal{T}$ has, under 2.3, exactly the arithmetic content of asserting that a specific pair (n_0, k_0) exists with $\sigma(n_0) > q(n_0, k_0) + 2^{-k_0}$, together with the obligation to exhibit it. It is not a weaker or more cautious claim than a counterexample; it is a counterexample claim with the witness left unstated.

Corollary 3.6 (Riemann, syntactic form). For R the Riemann Hypothesis, $U(R) \iff Q \vdash \exists n (\sigma(n) > H_n + \exp(H_n) \cdot \ln(H_n))$.

Corollary 3.6a (Riemann, sharpened form). Using Σ_1^0 -soundness of Q (2.3): for every Π_1^0 sentence R ,

$U(R) \iff \Omega \models \neg R$.

In particular $U(R)$ holds **for this R** if and only if $\mathbb{N} \models \neg R$; and $\mathbb{N} \models \neg R$ holds if and only if ζ has a zero off the critical line, **that last step being Lagarias's theorem in the metatheory and not a theorem of Q** , per 2.4a.

Proof. By Theorem 3.2, $U(R) \iff Q \vdash \neg R$. If $Q \vdash \neg R$ then by Σ_1^0 -soundness $\mathbb{N} \models \neg R$. If $\mathbb{N} \models \neg R$ then $\neg R$ is a true Σ_1^0 sentence, so $Q \vdash \neg R$ by Σ_1^0 -completeness. ■

Remark 3.6b. Corollary 3.6a is the sharpest form of the paper's thesis: *no consistent r.e. extension of Q proves R if and only if R is false*. Note the hypothesis: Theorem 3.2 is soundness-free, and 3.6a is not.

Theorem 3.7 (Dichotomy). Let $R \in \Pi_1^0$, let $\mathcal{S} \subseteq \mathcal{T}$ be nonempty, and **suppose $Q + R$ is consistent**. Put $U_{\mathcal{S}} := \forall T \in \mathcal{S} (T \not\vdash R)$. Then exactly one of:

(a) **$Q + R \in \mathcal{S}$.** Then $U_{\mathcal{S}} \iff Q \vdash \neg R$. (*The implication is vacuous, and its vacuity is its content: $Q + R \in \mathcal{S} \subseteq \mathcal{T}$ gives $Q + R$ consistent, hence $Q \not\vdash \neg R$, while $Q + R \vdash R$ gives $\neg U_{\mathcal{S}}$ outright. **The horn cannot be occupied at all**, which is the sharper reading of Remark 3.8's claim that asserting $U_{\mathcal{S}}$ here converts the claim into $Q \vdash \neg R$: the conversion is available and the conjunction is not.*)

(b) **$Q + R \notin \mathcal{S}$.** Then $\mathcal{S}$ is fixed by a condition beyond consistency, r.e.-ness and extension of Q , and establishing $T \not\vdash \neg R$ for any $T \in \mathcal{S}$ yields $\mathbb{N} \models R$ by Theorem 4.1.

If $Q + R$ is inconsistent then $Q \vdash \neg R$ outright and no dichotomy arises.

Proof. (b) is the negation of (a) under the hypothesis, so exactly one obtains. In (a) the argument of Theorem 3.2($\Rightarrow$) applies with $\mathcal{S}$ for $\mathcal{T}$. In (b), $Q + R$ is consistent, r.e. and extends Q so its exclusion is by a further condition; the second clause is Theorem 4.1, and every $T \in \mathcal{S} \subseteq \mathcal{T}$ satisfies $T \supseteq Q$. ■

Remark 3.8. A reader will observe that no one asserting the unprovability of R intends to quantify over theories that assume R . The disjunction *either $Q \vdash \neg R$ or $\mathbb{N} \models R$* holds for every Π_1^0 sentence R by Σ_1^0 -completeness alone (2.2), independently of Theorem 3.7. **What Theorem 3.7 adds is that neither horn of a restricted unprovability claim can be occupied without paying one side of that disjunction:** horn (a) converts the claim into $Q \vdash \neg R$, horn (b) converts its companion half into $\mathbb{N} \models R$.

4. The Restriction Theorem

Theorem 4.1. Let $R \in \Pi_1^0$ and let $T \supseteq Q$ be any theory. If $T \not\vdash \neg R$ then $\mathbb{N} \models R$.

Proof. Write $R = \forall n \phi(n)$ with $\phi \in \Pi_1^0$. If $\mathbb{N} \not\models R$ then $\neg \phi(n_0)$ for some n_0 ; $\neg \phi(n_0)$ is Σ_1^0 , so by Σ_1^0 -completeness of Q we get $Q \vdash \neg \phi(n_0)$, hence $Q \vdash \neg R$, hence $T \vdash \neg R$. Contrapositively, $T \not\vdash \neg R$ implies $\mathbb{N} \models R$. ■

Remark 4.1a. No hypothesis on T beyond $T \supseteq Q$ is used: T need not be consistent, r.e., sound, or Σ_1^0 -complete.

Corollary 4.2. If $T \not\vdash R$ and $T \not\vdash \neg R$ then $\mathbb{N} \models R$.

Corollary 4.3. For any $T \supseteq Q$, in particular PA or ZFC or ZFC with large cardinals, a proof that T does not refute R establishes R .

Remark 4.3a (the two results are compatible, and here is why). A reader may put Theorem 3.2 beside Theorem 4.1 and see a tension: the first says universal unprovability yields $Q \vdash \neg R$, the second says any $T \supseteq Q$ failing to refute R yields $\mathbb{N} \models R$. **These cannot conflict, because their hypotheses are exclusive.** For $R \in \Pi_1^0$ exactly one of two cases holds. If $Q \vdash \neg R$ then every $T \in \mathcal{T}$ refutes R , so no T fails to refute it and Theorem 4.1's hypothesis is never met; under 2.3, R is false. If $Q \not\vdash \neg R$ then $Q + R$ is consistent, r.e. and extends Q , so it lies in $\mathcal{T}$ and proves R , so $U(R)$ fails and Theorem 3.2's antecedent is never met; and by Theorem 4.1, R is true. **Universal unprovability and non-refutation by some extension cannot both hold**, and nothing in this paper asserts that they can.

Remark 4.4 (the pincer). Unrestricted, the claim that the route to R is closed is equivalent to $Q \vdash \neg R$ by Theorem 3.2 and, under 2.3, to $\mathbb{N} \models \neg R$ by Corollary 3.6a. Restricted to a fixed $T \supseteq Q$, establishing its refutation-side half yields $\mathbb{N} \models R$ by Theorem 4.1. No formulation both quantifies non-trivially and leaves the truth-value of R untouched.

5. The Sign Character of the Frame Functional

5.1 What this section is for. No condition of Section 6 tests λ . Ξ_7 compares two recorded matrices entrywise; λ appears in Section 6 only inside Ξ_7 's consequence note, where it is defined and used to state a consequence of the condition and is not itself a clause of it. What Section 5 supports is the polarity account at Section 12.3 and the consequence bound at Ξ_7 's note; it is a supporting section and is stated as such.

Notation. For $Q_0 \in \mathbb{R}^{3 \times N}$ with unit rows, write $R_G := Q_0 Q_0^T$ for the Gram matrix, reserving R for the Riemann Hypothesis. Let E be an orthonormal basis of the row span and put $\lambda(Q_0) := \det(Q_0 E^T)$.

Proposition 5.2. For arbitrary real 3×3 M and $D = \text{diag}(\epsilon_1, \epsilon_2, \epsilon_3)$, $\epsilon_i \in \{\pm 1\}$: $\det(D M D) = \det(M)$.

Proof. $\det(D M D) = \det(D)^2 \det(M) = \det(M)$, since $\det(D) \in \{\pm 1\}$. ■ (No symmetry or positive-semidefiniteness is used.)

Proposition 5.3 (sign character). $\lambda(D Q_0) = \det(D) \lambda(Q_0)$, and hence $|\lambda(D Q_0)| = |\lambda(Q_0)|$.

Proof. $\lambda(D Q_0) = \det(D Q_0 E^T) = \det(D) \det(Q_0 E^T) = \det(D) \lambda(Q_0)$. ■

Corollary 5.4. The Gram matrix of $D Q_0$ is $D R_G D$, with $\det(D R_G D) = \det(R_G)$ by 5.2; and for the total-negation pattern $D_- = -I_3$ one has $(-Q_0)(-Q_0)^T = R_G$ exactly, while $\lambda(-Q_0) = -\lambda(Q_0)$.

5.5 Executed check. Over 200 unit-row triads at $N = 24$ and all eight sign patterns: $\max|\det(D R_G D) - \det(R_G)| = 0.000 \times 10^0$, and $\max|\lambda(D Q_0) - \det(D) \lambda(Q_0)| = 0.000 \times 10^0$. Over 20000 triads, $\max|\lambda^2 - \det(R_G)| = 4.330 \times 10^{-15}$.

6. The Cascade: Formal Specification

Remark 6.0a (what this section does not do, stated before it is done). Four readings of Section 6 are natural, are wrong, and are foreclosed here rather than downstream, because each has been arrived at independently by a careful reader.

No text is mapped into V , and no proposition is rotated. The atoms of (L1) index the axes of 6.1(i); they are not vectors and nothing is added, scaled, or rotated in the language. What occupies the space is the recorded matrix Q_0 of 6.1(viii), whose row index is the atom index and whose entries are numbers. No embedding of syntax into a vector space is offered, none is used, and none is needed: the axes are labelled slots and the algebra runs on the recorded frame, not on the sentence.

The conditions test the record and not the object. Every Ξ_i except Ξ_7 reads fields the record's author supplied, and Remark 6.6a states this at its least favourable. **The emitter performs no measurement of the Riemann Hypothesis and none is claimed.** What it certifies is that a sweep was declared, executed and reported to a stated discipline; that is the whole of its office, and Theorem 6.9 is the statement of exactly how little that entails.

The frame's orientation is independent of the truth value of P , by construction, and that is the design and not a defect. Proposition 6.1a builds Q_0 from the syntax of P alone, so the polarity bit it carries is the parity of leading negations and nothing else. A reader who observes that the geometry never touches the arithmetic has read the construction correctly; Theorem 6.9(a) is that observation proved, and Section 12.3 records that nothing about ζ is at stake in the bit.

What the geometry is for is answered at Remark 6.8a, and the answer is narrow: it supplies an exhaustiveness argument for the count of conditions and nothing further.

A record can be supplied for any proposition, including a trivial one, and would emit. Nothing prevents it and nothing should: the terminal token reports that *this instrument* swept *this* declared catalog and found the door uncrossed. It is not a difficulty rating, not a certificate about the object, and not evidence that the proposition resisted anything. A token on a trivial sentence would be a true report about a trivial sweep.

Definition 6.0 (the linguistic layer, stated in place). Definition 6.1(i) and 6.7a through 6.8 presuppose a decomposition discipline on the object propositions. It is stated here rather than cited, so that every hypothesis of Theorem 6.7 is checkable within this paper. Nothing outside those passages uses it. Four clauses.

(L1) Atomic decomposition. A complete factual proposition P is split into three components: A_1 its existence component, what is; A_2 its kinetic component, what it does or undergoes; A_3 its relational component, how it stands toward what.

(L2) Deletion test. Delete each component in turn. A complete atomic claim returns exactly **three** irreducible slots, each leaving an incoherent residue when removed. A count other than three is a failure of the decomposition at this layer.

(L3) Isolation. The vocabularies populating the three slots are **pairwise disjoint**. If one slot's vocabulary is a subset of another's, the slots are not independent and the decomposition fails.

(L4) Atomic bivalence. An atom admits negation and admits no partial or continuous operation. Correspondingly the verdict economy of Definition 6.3 admits no fractional or probabilistic state.

(Clauses (L1) through (L4) are stipulations of this construction, operational and reproducible across analysts applying them, and they are typed at that level: **they are not offered as a uniqueness theorem of first-order logic and no such theorem is claimed**. Every use of them below is marked. Theorem 6.7(ii)'s operational derivation of $d = 3$ uses (L2); its algebraic derivation uses none of them.)

Definition 6.1 (record). Fix a language $\mathcal{L}_{\text{obj}}$ for the object propositions and a disjoint language $\mathcal{L}_{\text{rec}}$ for records; **no field of a record is a sentence of $\mathcal{L}_{\text{obj}}$ or encodes one**, except as expressly noted in the witness fields, which are governed by Ξ_5 and by 2.4's classification.

An *audit record* r for a proposition $P \in \mathcal{L}_{\text{obj}}$ consists of:

(i) a **labelled orthogonal decomposition** into atomic axes, $\mathbb{R}e_1 \oplus \dots \oplus \mathbb{R}e_d$, one axis per atom of (L1), with d recorded; the space $V := \mathbb{R} \cdot u \oplus (\mathbb{R}e_1 \oplus \dots \oplus \mathbb{R}e_d)$ for a unit u orthogonal to every e_i ; and the involution σ **constructed from that decomposition**, $\sigma(u) = u$ and $\sigma(e_i) = -e_i$ for each i . (σ is not recorded as a primitive. The decomposition is the recorded object and σ is read off it, so $E - (\sigma)$ is the span of the atomic axes by construction and $\dim \text{Fix}(\sigma) = 1$ and $\sigma^2 = I$ hold by construction. Ξ_1 below

therefore verifies that the construction was carried out and that $d = 3$; it does not test independent facts about an independently supplied involution. The row index of the frame Q_0 at (viii) is this same atom index.)

(ii) a quantifier classification of P in the arithmetic hierarchy with its citation;

(iii) a **declared register** ρ , its acting group $\mathbb{G}_\rho$, the invariant ring of $\mathbb{G}_\rho$, and the finite token list of the claim;

(iv) a finite catalog $\mathcal{C}$ of channels, $|\mathcal{C}| = C$, each carrying a designated reduction with its citation set, together with a **closure flag** for $\mathcal{C}$;

(v) a subset $\mathcal{W} \subseteq \mathcal{C}$, $|\mathcal{W}| = B$, each element carrying a **named obstruction and its warrant grade**;

(vi) a finite set $\mathcal{P}$ of roads, **defined: each channel of (iv) determines exactly one road, namely the citation set recorded for its reduction; distinct channels determine the same road when their citation sets coincide; hence $|\mathcal{P}| \leq C$** . Two roads are *premise-disjoint* when they share no citation. Ξ_6 's floor of three therefore strengthens Ξ_3 's rather than restating it, and since $|\mathcal{P}| \geq 3$ with $|\mathcal{P}| \leq C$ gives $C \geq 3$, Ξ_3 's **channel floor $C \geq 3$ is entailed by Ξ_6 and is retained as a guard that halts earlier in the evaluation order of Definition 6.3**. It is therefore not independent of Ξ_6 , which is recorded here and at 9.9 rather than left for a reader to derive; the count of eight is a count of conditions evaluated, not a claim of pairwise independence;

(vii) **witness fields** for P and $\neg P$, each either empty or holding a certificate; a **declared search budget** $k = (k_n, k_{\text{prec}}, k_{\text{pf}})$, **each component a standard numeral**, bounding the numeral range, the rational precision and the derivation length independently, with $k_n \geq k_{\text{min}}$, $k_{\text{prec}} \geq p_{\text{min}}$, $k_{\text{pf}} \geq d_{\text{min}}$ for declared floors recorded on the record; a **declared theory $\mathfrak{B}$ with $\mathfrak{B} \supseteq Q$** in which the proof search runs; **the floors are free parameters and are declared as such at 9.9, so the certificate clause of Ξ_3 is informative only to the extent the floors are, and a record declaring minimal floors emits a correspondingly weak statement; a completion flag** for each search; and an independence field;

(viii) a **frame assignment** $Q_0(\cdot)$ recording a unit-row matrix $Q_0(P) \in \mathbb{R}^{\{d \times N\}}$ with its row-span dimension recorded, and, independently and with no constraint imposed here, a matrix $Q_0(\neg P) \in \mathbb{R}^{\{d \times N\}}$; a declared tolerance τ , and a **declared consequence ceiling β with $\beta \leq \beta_{\text{max}}$** . (No basis, Gram or orientation quantity is a field: none is consumed by any condition, and $\lambda(P)$ is undefined on a record declaring row-span dimension 2. G and λ are defined inline at Ξ_7 's consequence note, the only place they are used.)

(ix) a designated *door*, an object not in $\mathcal{C}$, with a **provenance flag** valued *supplied* or *generated* and a **crossed flag**;

(x) a permanence clause.

Definition 6.2 (the eight conditions).

- Ξ_1 the recorded axis count d equals 3, hence $\dim V = 1 + d = 4$. ($\sigma^2 = I$ and $\dim \text{Fix}(\sigma) = 1$ are invariants of 6.1(i)'s construction and **not tests**: they hold at every d , executed at $d = 2, 3, 4, 5$, so a condition asserting them would be analytically true on every record and would discriminate nothing, which is the defect Remark 6.2a records for a superseded form of Ξ_7 and Remark 6.1a-rank for a superseded clause of it. **$\det(\sigma|E-)$ is not among them**: σ acts on $E-$ as $-I_d$, so $\det(\sigma|E-) = (-1)^d$, which is -1 exactly when d is odd, executed at $d = 2, 3, 4, 5$ returning $+1, -1, +1, -1$. It is therefore a parity test and not an invariant, and it does not pick out $d = 3$ in any case, since it passes at every odd d . Ξ_1 's single clause $d = 3$ entails it at the value used and subsumes it. The division-algebra structure noted at Remark 6.7e is neither a record field nor tested by any condition.)
- Ξ_2 the quantifier class of P is located by a cited classical equivalence, and every token of (iii) lies in the invariant ring of $\mathbb{G}_\rho$.
- Ξ_3 $C \in \mathbb{N}$, $C \geq 3$, every channel in $\mathcal{C}$ carries a reduction to a common invariant by a cited classical equivalence with none unreduced, and the closure flag of (iv) is set. **Closure here means the catalog is displayed and fully reduced, not that no channel exists outside it**; the two readings are distinguished at Corollary 8.2 and the weaker one is meant, which is what makes Ξ_3 jointly satisfiable with Ξ_8 's door.
- Ξ_4 $B = C$, and each element of $\mathcal{W}$ carries its named obstruction with grade recorded.
- Ξ_5 the certificate search for $\neg P$ over numerals $n \leq k_n$ at rational precision k_{prec} returned no certificate; the proof search for P in the declared theory $\mathfrak{B}$ over derivations of length $\leq k_{\text{pf}}$ returned none; $k_n \geq k_{\text{min}}$, $k_{\text{prec}} \geq p_{\text{min}}$ and $k_{\text{pf}} \geq d_{\text{min}}$ with all six recorded; both completion flags are set; both witness fields are empty; and the independence field is empty.
- Ξ_6 $|\mathcal{P}| \geq 3$, and the roads in $\mathcal{P}$ are pairwise disjoint **as computed from the recorded citation sets**, with the recorded flag agreeing. (The clauses $B/C = 1$ and $C \geq 3$ are entailed by Ξ_4 and Ξ_3 and are not restated here.)
- Ξ_7 the recorded frames satisfy the negation convention **as tested directly**: $\max|Q_0(\neg P) + Q_0(P)| \leq \tau$ for a declared tolerance τ recorded on the record with $2\sqrt{N} \cdot \tau + N \cdot \tau^2 \leq \beta$ and $\beta \leq \beta_{\text{max}}$ for a declared ceiling β_{max} , and the recorded row-span dimension of $Q_0(P)$ equals d . (β bounds the consequence deviation directly and supersedes a separate tolerance ceiling: bounding τ alone left τ and N free to be declared large together.) (The ceiling is required for the same reason the budget floors are: an unbounded τ admits vacuous satisfaction. β_{max} joins the census in its place.) (Consequence, stated as a bound rather than as an equality. Corollary 5.4 gives $G(\neg P) = G(P)$ and $\det(Q_0(\neg P)E(P)^T) = -\lambda(P)$ at $\tau = 0$ and there only. For $\tau > 0$, writing $Q_0(\neg P) = -Q_0(P) + \Delta$ with $\max|\Delta| \leq \tau$ and $Q_0(P)$ unit-row, one has $\max|G(\neg P) - G(P)| \leq 2\sqrt{N} \cdot \tau + N \cdot \tau^2$, and the orientation ratio deviates from -1 by $O(\sqrt{N} \cdot \tau / |\lambda(P)|)$. Executed at $N = 24$: deviation 3.603×10^{-8} at $\tau = 10^{-8}$ and 3.342×10^{-2} at $\tau = 10^{-2}$, and at $N = 240$ with $\tau = 10^{-2}$, 4.385×10^{-2} ; at $\tau = 0$, exactly 0.000×10^0 . **The β clause bounds this deviation directly, which is why no separate tolerance ceiling is carried**. The rows of $Q_0(\neg P)$ are within $\sqrt{N} \cdot \tau$ of unit, not within τ : an entrywise perturbation of size τ moves a row norm by up to $\|\Delta\|_2 \leq \sqrt{N} \cdot \tau$. Executed: max row-norm deviation 2.0148×10^{-2} at $N = 24$ and 2.2149×10^{-2} at $N = 240$, both at $\tau = 10^{-2}$, against $\tau = 10^{-2}$ and $\sqrt{N} \cdot \tau$ of 4.899×10^{-2} and 1.549×10^{-1} . Nothing consumes the row norms. Here and only here, $G(X) := Q_0(X)Q_0(X)^T$ and $\lambda(X) := \det(Q_0(X)E^T)$ for E an orthonormal basis of the row span of $Q_0(P)$. **E must be the same on both sides**: for any such E one has

$\lambda(-Q_0) = (-1)^3 \lambda(Q_0) = -\lambda(Q_0)$, so the identity is basis-independent when shared and fails when recomputed per side, the identity of E being a gauge. Nothing in this paper depends on which E is chosen.)

- Ξ_8 the door of (ix) lies outside $\mathcal{C}$, its provenance flag reads *supplied*, its crossed flag is unset, and the permanence clause is present.

Proposition 6.1a (a frame assignment satisfying Ξ_7 exists). A hash applied directly to the syntax of P will not do: the hash of " $\neg P$ " is uncorrelated with the hash of P , so $Q_0(\neg P) = -Q_0(P)$ would have to be imposed by hand rather than obtained, and double negation would be inconsistent. Executed, the direct construction violates the convention on **6 of 6** test propositions. The polarity normal form obtains it instead. Put P into that form: let P_+ be the negation-free core and $s \in \{0,1\}$ the parity of leading negations. Define $Q_0(P) := (-1)^s \cdot H(P_+)$, where H seeds a deterministic generator with a hash of the syntax of P_+ and returns a row-centred, row-normalized $3 \times N$ draw. Then Q_0 is a total function of syntax, independent of the truth value of P , satisfies $Q_0(\neg P) = -Q_0(P)$ for every P including under double negation, and yields a record satisfying Ξ_7 .

The conclusion holds for any draw of rank d , which the construction achieves with probability one; a draw of lower rank records row-span dimension below d , fails Ξ_7 's dimension clause, and is rejected and reseeded.

Verification against the condition as it now stands, six propositions at $N = 24$ with their negations and double negations: $\max|Q_0(\neg P) + Q_0(P)| = 0.000 \times 10^0$, which satisfies Ξ_7 for every $\tau \geq 0$; recorded row-span dimension 3 in every case; and $\max|Q_0(P) - Q_0(\neg P)| = 0.000 \times 10^0$. ■

Remark 6.1a-rank. 6.1(viii) records the row-span dimension of $Q_0(P)$; Ξ_7 tests whether it is d . E is not a field and is introduced at Ξ_7 's consequence note, where alone it is used. (Verified: rank 3 and $\lambda(P) \neq 0$ agree over 2000 frames with zero disagreements, and a forced rank-2 frame gives $|\lambda(P)| = 0.000 \times 10^0$.)

Remark 6.1b. Proposition 6.1a exhibits one assignment satisfying Ξ_7 and shows that condition satisfiable. It does not claim the assignment canonical, and no result depends on which **syntax-determined** assignment is used.

It does not show $\Xi_1 \wedge \dots \wedge \Xi_8$ satisfiable, and this paper does not. No record is exhibited. **This is a vacuity question about the specification and is not a hypothesis of any theorem:** Theorem 6.9(a) is P -invariance of satisfaction for the seven conditions other than Ξ_8 , which holds for every record whether or not any record satisfies all eight. What an unsatisfiable conjunction would cost is that no emission ever occurs, so 6.9(b) and (c) would speak of nothing.

What would discharge it, and this is a list of what is not done rather than a partial discharge. $\Xi_1, \Xi_3, \Xi_5, \Xi_6, \Xi_7$ and Ξ_8 are instantiable from material this paper carries: a four-dimensional space with a self-adjoint involution of one-dimensional fixed space; three channels with cited classical equivalences; a budget of the size exhibited at 2.5, with its floors; roads as defined at 6.1(vi); the assignment of 6.1a; and a door with clause W . **Two are not.** Ξ_2 requires an exhibited register with its acting group, which a trivial group would satisfy degenerately and which is not exhibited here. Ξ_4 requires a named obstruction with a recorded warrant grade for each channel; **this paper names no obstruction and will not invent one**, which is why the record is not constructed rather than being constructed badly.

Remark 6.2a (what Ξ_7 is, and why it is a condition). Ξ_7 compares the two independently recorded frames directly at a declared tolerance, so a record whose $Q_0(\neg P)$ differs from $-Q_0(P)$ by more than τ fails it. Ξ_7 tests the frame assignment and not the proposition, and no information about P is obtained from it. Testing the derived pair instead, equal Gram with opposite orientation, would be strictly weaker: that pair is implied by the convention and does not exhaust it, and two counterexample families exist, $Q_0(\neg P) := Q_0(P)(I_N - 2vv^T)$ for any unit v in the row span, and a single-axis reflection in the eigenbasis of $G(P)$.

Definition 6.3 (the emitter). (By Remark 6.0a the conditions below read record fields, Ξ_5 excepted; the emitter measures the declared sweep and not the object. Ξ_3 's floor $C \geq 3$ is entailed by Ξ_6 with $|\mathcal{A}| \leq C$ and is evaluated here as a guard that halts earlier, not as an independent condition; see 9.9.) Evaluate $\Xi_1, \Xi_2, \Xi_3, \Xi_4, \Xi_5, \Xi_6, \Xi_7, \Xi_8$ in order. If Ξ_5 fails because a witness field is populated, exit to direction-adjudication; if because the independence field is populated, exit to the relative-independence verdict. Otherwise any failure halts with the non-terminal verdict and the index recorded. If all hold, return the terminal token with the warrant cap and the permanence clause of Ξ_8 .

Proposition 6.4 (fail-safe, general form). If any field of Definition 6.1 required by some Ξ_i is absent, the emitter returns the non-terminal verdict.

Proof. By inspection over the eight conditions. $\Xi_1, \Xi_2, \Xi_3, \Xi_4, \Xi_6, \Xi_7, \Xi_8$ each require an equality, a set inequality, or a flag to be set; an absent field satisfies none. Ξ_5 requires $k_n, k_{\text{prec}}, k_{\text{pf}}$ and their three floors recorded with each bound at or above its floor, and both completion flags set; absence of any of these fails Ξ_5 , rather than passing it. The three floors are free parameters and are declared as such in 9.9. ■

Remark 6.4a. Ξ_5 requires an executed search reported against recorded bounds rather than an unpopulated field, so absence of data fails it, and the fail-safe property holds at every condition.

Proposition 6.5. The terminal token is returned only if all eight conditions hold.

Proof. Immediate from sequential evaluation and halt on first failure. ■

(Proposition 6.5 asserts the necessity direction only. Logical independence of the conditions is not proved here; see 9.9.)

Proposition 6.6. Ξ_5 is the unique condition that can route to a verdict other than the non-terminal one. **It can also fail to the non-terminal verdict**, when a bound or floor is unrecorded, a completion flag unset, or a bound below its floor; the exit branches are taken only when a witness field or the independence field is populated.

Proof. By inspection of Definition 6.3. ■

Remark 6.6a (which conditions reach outside the record, and it is fewer than one would like). The honest split is four ways. Ξ_5 , alone reaches into arithmetic: its certificate clause reports an executed search over numerals, and Theorem 6.9(b) names it the section's one object-language output. Ξ_2 and Ξ_3 reach outward only through their citation requirements, a located quantifier class and cited classical equivalences, which a reader can check against the literature. Ξ_6 is computable from recorded data and is the one condition where a declaration can be caught wrong by the emitter: disjointness is computed from the citation sets and checked against the recorded flag. **The remaining four are internal consistency checks on declared fields:** Ξ_1 checks the recorded d and the construction invariants of 6.1(i), Ξ_4 reads a recorded obstruction and grade, Ξ_7 compares two recorded frames, and Ξ_8 reads recorded provenance and crossed flags.

A flag set by the record's author is not external material, however the relation it names is described. This remark exists to state the asymmetry, so it states the least favourable reading the specification supports.

Definition 6.7a (the decomposition and its stabiliser). A decomposed proposition is not a bare frame in V . By (L1) and (L3) it is **three labelled orthogonal axes**, one per atomic component, with pairwise disjoint vocabularies. By 6.1(i) the residence is that decomposition: $E_-(\sigma) = \mathbb{R}e_1 \oplus \dots \oplus \mathbb{R}e_d$, one axis per atom of (L1), with the frame's row index the same atom index. Define its **stabiliser**

$$\text{Stab}(E_-) := \{ g \in O(E_-) : g(\mathbb{R}e_i) = \mathbb{R}e_i \text{ for each } i \},$$

the transformations carrying each axis to itself, hence carrying a decomposed proposition to a decomposed proposition.

Lemma 6.7b (the stabiliser is exactly the sign changes). $\text{Stab}(E_-) = \{ \text{diag}(\epsilon_1, \dots, \epsilon_d) : \text{each } \epsilon_i \in \{+1, -1\} \} \cong (\mathbb{Z}/2)^d$, of order 2^d .

Proof. If $g(\mathbb{R}e_i) = \mathbb{R}e_i$ then $ge_i = \lambda_i e_i$ for some real λ_i ; $g \in O(E_-)$ forces $|\lambda_i| = 1$, hence $\lambda_i = \pm 1$, hence g is diagonal with entries ± 1 . Conversely every such diagonal matrix is orthogonal and fixes each axis setwise. ■

Remark 6.7b' (why stabilisation and not invariance of the determinant). The Gram determinant is invariant under a far larger group: $\det((gQ_0)(gQ_0)^T) = \det(g)^2 \cdot \det(Q_0 Q_0^T)$, so **every** g with $|\det g| = 1$ leaves it fixed, an eight-dimensional group containing $O(3)$ and the signed permutations of order $2^3 \cdot 3! = 48$. Executed against a reference frame at $\det(R_-G) = 0.931277980144$: the eight sign changes deviate by 0.000×10^0 , the six row permutations by 2.220×10^{-16} with unit rows preserved to 1.1×10^{-16} , five hundred random elements of $O(3)$ by at most 2.442×10^{-15} .

Determinant-invariance is therefore not the criterion and is not used as one. The criterion is Definition 6.7a, and the two families above fail it for reasons this paper states elsewhere. A rotation by $\pi/7$ in the e_1 - e_2 plane carries e_1 out of $\mathbb{R}e_1$, mixing one atomic slot into another and destroying the pairwise vocabulary disjointness (L3) requires; the image is not a decomposed proposition. A row transposition carries $\mathbb{R}e_1$ to $\mathbb{R}e_2$, so it does not fix each axis either, and by (L3) it is not a mere relabelling: it substitutes one atom's content for another's across disjoint vocabularies, which is content-mixing on exactly the ground the rotation is excluded. **Both are excluded by Definition 6.7a on its own terms and no appeal to gauge is made or needed.**

Definition 6.7c (complete screen). A family of conditions is a **complete screen** for the decomposition when it carries exactly one condition per element of $\text{Stab}(E_-)$. (Each element is one pattern of negation across the atoms. The identity is the empty pattern, P itself, and screens no conflation; of the 2^d elements, $2^d - 1$ move the frame. The count is of patterns, not of conflations.)

Remark 6.7c' (why the eight do not collapse to four). The enumeration at Theorem 6.7(iv) below reads $1 + 3 + 3 + 1$, which is also the orbit decomposition of $(\mathbb{Z}/2)^3$ under the permutation action of S_3 on the axes. The arithmetic is correct and the collapse is not available. **S_3 does not act on the screened objects.** By (L3) the three axes carry pairwise disjoint vocabularies, so the three weight-one patterns deny three different atoms and are three different propositions: denying what stands, denying what moves, and denying how they bind are not one object seen three ways. The weight grading is a way of displaying the eight, not a quotient of them.

Theorem 6.7 (given d axes the count is 2^d , and at $d = 3$ the eighth is the last). Let $E_-(\sigma) = \mathbb{R}e_1 \oplus \dots \oplus \mathbb{R}e_d$ be the labelled orthogonal decomposition of 6.1(i). Then:

- (i) $|\text{Stab}(E_-)| = 2^d$, by Lemma 6.7b;
- (ii) hence a **complete screen carries exactly 2^d conditions**, and at $d = 3$ exactly **eight**;
- (iii) the eight enumerate by negation weight as **$1 + 3 + 3 + 1$** : the empty pattern, three denials of a single atom, three of a pair, and one total negation. **The eighth is the last**, and the ladder terminates because it has exhausted the atoms;
- (iv) **no ninth element of $\text{Stab}(E_-)$ exists at $d = 3$** . A ninth requires $2^d \geq 9$, hence $d \geq 4$, hence a fourth atomic axis.

Proof. (i) is Lemma 6.7b. (ii) from (i) and Definition 6.7c. (iii) is the expansion of 2^3 by Hamming weight, with no weight 4 available at $d = 3$. (iv) $2^3 = 8 < 9$. ■

Remark 6.7d (what the count counts). Ξ_7 compares two recorded frames and Ξ_6 computes disjointness from the recorded citation sets; neither carries information about P , and by Theorem 6.9(a) the only condition that does is Ξ_5 , whose content is the search report. Theorem 6.7 fixes **how many** conditions a complete screen carries and says nothing about the discriminating power of any one of them over P .

Remark 6.7e (what is stipulated and what is proved). **Two things are stipulated.** $d = 3$, carried by (L2); and atomic bivalence, carried by (L4), which is what motivates restricting the axis-fixing group to $O(E_-)$ and so makes it finite: the deletion test returns three irreducible slots on a complete atomic claim, at the operational grade of Definition 6.0 and not as a uniqueness theorem. **The identification of the atoms with the axes of the residence is definitional**, by 6.1(i), where σ is constructed from the decomposition rather than recorded independently; there is no bridge to supply and none is claimed. Given those, **the count is not a further stipulation**: it is Lemma 6.7b, which is theorem-grade, together with the arithmetic of (iv).

The Frobenius classification is **corroboration and is load-bearing on nothing**. At $d = 3$ the space V of 6.1(i) has dimension 4, and $\mathbb{H}$ is the unique real associative division algebra with more than one imaginary unit; but no definition, condition or proof in this paper uses a multiplication on V , no record field carries a division-algebra structure, and Ξ_1 does not test one. The observation is recorded because it is true and because a reader will notice the dimension, not because anything rests on it.

Remark 6.8 (the shape of the claim). The claim this section makes is narrower than a forcing and wider than a choice, and it is stated at its exact strength. **Two things are stipulated**: that the decomposition carries three axes, which is (L2); and that atoms admit negation and no partial operation, which is (L4), motivating Definition 6.7a's restriction to $O(E_-)$, which is what makes the group finite. Both are stipulations of Definition 6.0 at the grade stated there. **One thing is definitional**: that the residence is the span of the atomic axes, by 6.1(i). **Everything after that is theorem**. By Lemma 6.7b the group carrying each axis to itself is exactly $(\mathbb{Z}/2)^d$; by Theorem 6.7 a complete screen has 2^d members, at $d = 3$ exactly eight, and no ninth. **Once the stipulations are made the count is not a further stipulation, and that is the whole of what is claimed here.**

No appeal to gauge is made. Both excluded families fail Definition 6.7a directly, and by Remark 6.7c' the permutation action does not reach the screened objects, so the eight do not collapse to four.

What remains open. Which condition screens which element: $\text{Stab}(E_-)$ acts on itself simply transitively, so no element is distinguished by the group structure alone, any bijection serves equally, and nothing below depends on which is taken. And the choice to screen the axis-fixing group rather than some larger structure-preserving group is a decision of this construction, motivated by (L3) and (L4) and not derived from them. **This paper does not claim that determinant-invariance forces eight**, and by Remark 6.7b' it does not: that group is eight-dimensional.

Remark 6.8a (why a space and not a checklist, and what the apparatus costs). A reader who has followed Remark 6.0a will ask what the geometry is for, since the eight conditions read record fields and could be evaluated by a boolean checklist with no space, no involution and no frame. The question is fair and the answer is narrow.

The geometry buys exactly one thing: an argument that the count is complete. A checklist of eight items carries no reason that it is eight rather than five or thirty, and no reason that a ninth is unnecessary rather than merely unwritten; its cardinality is whatever its author stopped at. Lemma 6.7b and Theorem 6.7 supply what a checklist cannot: given d axes, the group carrying each to itself is exactly $(\mathbb{Z}/2)^d$, so a complete screen has 2^d members, at $d = 3$ exactly eight, and no ninth element of that group exists. That is an exhaustiveness argument for the list, and it is available only because the conditions are indexed by a group rather than enumerated by hand.

It buys nothing else, and this should be weighed rather than glossed. The space performs no measurement of the object, per Remark 6.0a. Ξ_7 is the only condition that touches the frame, and Ξ_7 exists because the frame does. Strip the geometry and seven of the eight conditions survive unchanged as record checks, the eighth disappears with the frame it compares, and what is lost is the completeness argument and only that. **Whether one exhaustiveness argument is worth a four-dimensional apparatus is a judgement this paper leaves to the reader, and it states the trade rather than obscuring it.** The alternative reading, that the specification is an administrative protocol with a group-theoretic warrant for the length of its checklist, is not a misreading and is not resisted here.

Theorem 6.9 (payload). Let r be an audit record for a proposition P . Then:

(a) **whether r satisfies $\Xi_1, \Xi_2, \Xi_3, \Xi_4, \Xi_6, \Xi_7$ and Ξ_8 is independent of the truth value of P ;** and the satisfaction of Ξ_5 depends on it in exactly one way, namely through whether the executed certificate search returned a certificate at the declared budget. Consequently $\Xi_1 \wedge \dots \wedge \Xi_8$ entails no $\mathcal{L}_{\text{obj}}$ -sentence undecided in $\mathfrak{B}$, and hence neither does any emission on a record satisfying them;

and if in addition r satisfies Ξ_1 through Ξ_8 and the emitter returns the terminal token with permanence clause W , then:

(b) the emission entails that $\mathcal{C}$ is closed and fully reduced with $B = C$, that a door outside $\mathcal{C}$ is supplied and uncrossed, and, **the one object-language clause the emission makes explicit, that the executed certificate search returned no certificate at numerals $n \leq k_n$ and precision k_{prec} .** (It does not entail that no certificate exists there: that would require the search to be sound and complete for its budget, and the record carries no such field. Theorem 8.1 applies to the emitter here, since a search is an instrument at the register.) This clause is a finite verified computation, hence $\mathfrak{B}$ -provable, so (a) is unaffected, the clause being decided in $\mathfrak{B}$;

(c) if W is replaced by a clause entailing $U(P)$, the emission entails $Q \vdash \neg P$ by Theorem 3.2.

Proof. (a) Take r fixed and vary the truth value of P . $\Xi_1, \Xi_2, \Xi_3, \Xi_4, \Xi_6$ and Ξ_8 are conditions on fields of types (i)–(vi), (ix)–(x), none of which is or encodes an $\mathcal{L}_{\text{obj}}$ -sentence, so their satisfaction is untouched. Ξ_7 compares $Q_0(\neg P)$ to $Q_0(P)$ and to the recorded dimension, mentioning P nowhere beyond its index. That is the seven.

Ξ_5 is not among them and its dependence is stated rather than argued away. Its certificate clause requires that the executed search *returned* nothing, and what a search returns is a fact about the executed computation, which depends on whether a violating numeral lies at $n \leq k_n$ and precision k_{prec} . Three assignments make this concrete: P true, the search returns nothing and Ξ_5 passes; P false with the witness beyond the budget, the search returns nothing and Ξ_5 passes; **P false with the witness inside the budget, the search returns a certificate and Ξ_5 fails.** Satisfaction of Ξ_5 is therefore *compatible* with either truth value and is not *independent* of it.

The stated conclusion follows without closing that gap. Ξ_5 's content is *the search returned nothing at (k_n, k_{prec})* , a terminating computation and hence decided in $\mathfrak{B}$ either way; the other seven contribute nothing about P by the paragraph above. So the emission entails no $\mathcal{L}_{\text{obj}}$ -sentence that $\mathfrak{B}$ leaves undecided, which is (a) as stated.

(The stronger two-model claim, that $\Xi_1 \wedge \dots \wedge \Xi_8$ entails neither P nor $\neg P$, would require a satisfying record under each truth value, which is at least as strong as satisfiability and is disclosed as open at 6.1b. The restriction above is what the proof gives and it carries no undischarged hypothesis.)

(b) The first two clauses from Ξ_3, Ξ_4 and Ξ_6 ; the third from the certificate clause of Ξ_5 read as a report of the executed search, whose content is a terminating computation and therefore $\mathfrak{B}$ -provable. (c) By Theorem 3.2 applied to P . ■

7. The Termination Theorem

Definition 7.1. A *permanence clause* delimits the scope of an emitted token's terminality.

- **W (as swept):** the verdict is terminal for $\mathcal{C}$ as swept and is revised upon presentation of a channel not in $\mathcal{C}$.
- **S (unrestricted):** $U(P)$, no theory in $\mathcal{T}$ proves P .

Theorem 7.2. Let $P = R$ and let the emitter return the terminal token on a record r .

- W is admissible: it entails nothing about derivability of R in any theory, and is consistent with R and with $\neg R$ alike.
- S is the sentence $U(R)$; by Theorem 3.2 an emission carrying S entails $Q \vdash \neg R$, and under 2.3 by Corollary 3.6a entails $\mathbb{N} \models \neg R$.

Proof. (i) By Theorem 6.9(a) with clause W . (ii) Immediate from Definition 7.1 and Theorems 3.2, 3.6a. ■

Corollary 7.3. Let W' be any permanence clause entailing $U(R)$. An emission carrying W' entails $Q \vdash \neg R$ by Theorem 3.2 and, **under 2.3, $\mathbb{N} \models \neg R$** by Corollary 3.6a. Clause S is of this kind. **W is admissible, and no clause entailing $U(R)$ is.**

Remark 7.3a. No maximality of W among all strengthenings is claimed, and none holds. Let W be terminal for $\mathcal{C} \cup \{c\}$ for a designated further channel c . W strictly entails W , is strictly stronger, and entails nothing about any theory. Corollary 7.3 is scoped by entailment of $U(R)$ and not by strength.

8. Self-Exclusion of a Certifying Instrument

Theorem 8.1. *Premise:* every instrument operating at register ρ is a channel at register ρ . Let $\mathcal{C}$ be a catalog of channels at ρ and $\mathfrak{A}$ an instrument certifying $\mathcal{C}$ exhaustive at ρ . Then either $\mathfrak{A} \notin \mathcal{C}$, in which case $\mathcal{C}$ omits a channel at ρ and is not exhaustive; or $\mathfrak{A} \in \mathcal{C}$, in which case **the certification is impredicative**: its warrant ranges over a collection of which $\mathfrak{A}$ is a member.

Proof. The disjunction is exhaustive. In the first case $\mathfrak{A}$ is a channel at ρ by the premise and is not in $\mathcal{C}$, contradicting exhaustiveness. In the second, $\mathfrak{A}$'s certification ranges over a collection of which $\mathfrak{A}$ is a member. ■

(Impredicativity is not by itself an establishment barrier: a finite catalog $\{c_1, \dots, c_n\}$ with $c_i = \mathfrak{A}$ could be certified by $\mathfrak{A}$ via a proof of the register's channel-cardinality. The theorem concludes impredicativity and no more.)

Corollary 8.2. A catalog is either non-exhaustive at its register or is certified impredicatively there. Exhaustiveness is therefore not certifiable non-impredicatively at the register the catalog enumerates. Hence Ξ_3 is read as asserting that $\mathcal{C}$ is displayed, fully reduced and flagged closed, not that no channel exists outside it, and clause W is the appropriate permanence clause.

Remark 8.3. With the stated premise, Theorem 8.1 is a consequence of the definitions and a case split on set membership. It contains no fixed-point construction and no self-reference lemma, and no comparison with the undefinability or incompleteness theorems is intended.

9. What Is Claimed

9.1 Theorem 3.2. For every sentence R of the language of Q , $\forall T \in \mathcal{T} (T \not\vdash R) \iff Q \vdash \neg R$. Both directions proved. Finitary; no soundness hypothesis, no semantic notion, no property of R beyond sentencehood.

9.2 Corollaries 3.4–3.6. $U(R)$ implies $Q + R$ inconsistent. For $R \in \Pi_1^0$ the right side is $Q \vdash \exists n \neg \phi(n)$, and under Σ_1^0 -soundness of Q a violating n_0 exists with a terminating verification. For R the Riemann Hypothesis the right side is the provability in Q of a violation of the Lagarias inequality.

9.2a Corollary 3.6a. Under Σ_1^0 -soundness of Q , for every Π_1^0 sentence R : $U(R) \iff \mathbb{N} \models \neg R$. For this R , universal unprovability holds if and only if $\mathbb{N} \models \neg R$; the further identification with ζ having a zero off the critical line is Lagarias's theorem in the metatheory, not a theorem of Q , per 2.4a. **The soundness hypothesis is used here and not in 3.2.**

9.3 Theorem 3.7. Under the assumption that $Q + R$ is consistent, exactly one of: $\mathcal{S}$ admits $Q + R$, whence $U_{\mathcal{S}}$ implies $Q \vdash \neg R$; or $\mathcal{S}$ excludes it, whence $\mathcal{S}$ is fixed by a further condition and the companion half yields $\mathbb{N} \models R$. The disjunction *either* $Q \vdash \neg R$ *or* $\mathbb{N} \models R$ is itself due to Σ_1^0 -completeness alone; what 3.7 adds is that no horn of a restricted claim is occupiable without paying one side of it.

9.4 Theorem 4.1, Remark 4.1a. For $R \in \Pi_1^0$ and any $T \supseteq Q$: $T \not\vdash \neg R$ implies $\mathbb{N} \models R$. No hypothesis on T beyond containment.

9.5 Corollaries 4.2, 4.3. Independence implies truth; for any $T \supseteq Q$, a proof that T does not refute R establishes R .

9.6 Propositions 5.2–5.3, Corollary 5.4. $\det(D M D) = \det(M)$ for arbitrary real 3×3 M ; $\lambda(DQ_0) = \det(D) \cdot \lambda(Q_0)$; under total negation the Gram is unchanged and λ flips sign. Verified at 0.000×10^0 . **Neither 5.2 nor 5.3 is tested by any condition**; both support Ξ_7 's consequence bound and the polarity account at 12.3.

9.7 Definitions 6.1–6.3. A record with ten field groups, eight conditions on it, and a sequential emitter with exit branches at Ξ_5 . **Seven conditions are conditions on the record alone; Ξ_5 reports an executed search and is the one condition whose satisfaction reaches arithmetic**, per Theorem 6.9(a). σ is exactly what 6.1(i) specifies, and $\mathcal{P}$ is defined at 6.1(vi) with its relation to $\mathcal{C}$ stated there. By Remark 6.2a, **Ξ_7 compares two independently recorded frames and does not test the proposition**, and Proposition 6.1a exhibits a record satisfying it. **Ξ_2 and Ξ_4 are evaluable in form and uninstantiated in fact**: no register or acting group is exhibited for Ξ_2 , and no obstruction is named for Ξ_4 . No claim is made about the discriminating power of either.

9.8 Propositions 6.4–6.6. The emitter is fail-safe on absence at every condition. The token issues only if all eight hold. Ξ_5 is the unique condition that can route to a verdict other than the non-terminal one, and it can also fail to the non-terminal one.

9.9 Definition 6.0, Definition 6.1(i), Definitions 6.7a and 6.7c, Lemma 6.7b, Theorem 6.7, Remarks 6.7b', 6.7c', 6.7e and 6.8. The screened group is the stabiliser of the labelled orthogonal decomposition of $E_-(\sigma)$, **not** the invariance group of the Gram determinant, which is eight-dimensional and is disowned as a criterion at 6.7b'. **Two stipulations**, both of Definition 6.0 at the operational grade stated there: $d = 3$ by (L2), and atomic bivalence by (L4), which restricts the axis-fixing group to $O(E_-)$ and makes it finite. **One definition**: σ is constructed from the decomposition at 6.1(i), so the residence is the span of the atomic axes and no identification between them is asserted. **Then theorem**: Lemma 6.7b gives $|\text{Stab}(E_-)| = 2^d$ exactly, Theorem 6.7 gives eight conditions at $d = 3$ and no ninth. The Frobenius classification is corroboration and is load-bearing on nothing; no multiplication on V is used anywhere. No appeal to gauge is made, and by 6.7c' the permutation action does not reach the screened objects. The assignment of conditions to elements is a labelling and is not fixed, and the choice to screen the axis-fixing group rather than a larger structure-preserving group is a decision of this construction. **Logical independence of the eight is neither proved nor used, and no minimality among non-complete families is claimed.** One dependence is recorded: Ξ_5 's floor $C \geq 3$ is entailed by Ξ_6 together with $|\mathcal{P}| \leq C$, and Ξ_3 is retained as a guard that halts earlier in the evaluation order of Definition 6.3.

Free-parameter census, stated because the ratio is unfavourable and should be visible. Ten declared degrees of freedom, every entry load-bearing: the axis count $d = 3$, stipulated by (L2); the single fixed direction $\dim \text{Fix}(\sigma) = 1$; the channel floor $C \geq 3$, entailed by Ξ_6 and retained as a guard; the road floor $|\mathcal{P}| \geq 3$; the consequence ceiling $\beta_{\max}$; the numeral floor $k_{\min}$; the precision floor $p_{\min}$; the derivation floor $d_{\min}$; the context width N ; and the choice of frame assignment. **The list prints ten slots and the count is ten, and it enumerates choices rather than their consequences.** $\dim V = 1 + d$ is definitional at 6.1(i), and the condition count 2^d is Lemma 6.7b; neither is a further degree of freedom once d and the single fixed direction are chosen, and listing them would double-count one stipulation as three. τ is not among them: it is bounded at Ξ_7 by the declared β , which bounds the consequence deviation $2/N \cdot \tau + N \cdot \tau^2$ directly and is itself bounded by the census entry

$\beta_{\max}$. Distance from $-Q_0(P)$ in max-norm is governed by τ alone; N enters through the size of the consequences, not through that distance. N enters through the frame width in Ξ_7 's entrywise test and through the consequence bound; its second justification lapsed when λ left the conditions.

And one thing a reader weighing the ratio should meet here rather than in a definition. Ξ_5 's certificate clause is informative only to the extent k_n , k_{prec} and k_{pf} exceed their floors; a record declaring minimal floors satisfies Ξ_5 and emits a 6.9(b) clause of near-zero content. Against these the section delivers Theorem 6.9(a) and (b) as its own consequences, part (c) being inherited from Theorem 3.2. **The section declares its parameters; it does not reduce them, and it fits rather than predicts.** By contrast Sections 3 and 4 carry zero free parameters and deliver four independent results. The paper is bimodal and should be read as such.

9.10 Theorem 6.9. An emission with W entails no L_{obj} -sentence undecided in $\mathfrak{B}$, by P -invariance of the seven conditions other than Ξ_5 and $\mathfrak{B}$ -decidability of Ξ_5 's search report; entails that $\mathcal{C}$ is displayed, fully reduced, walled at $B = C$, with a supplied uncrossed door outside $\mathcal{C}$, and that the executed certificate search returned nothing at the declared budget, which is the section's one object-language output and is not a claim that no certificate exists there; and with a $U(P)$ -entailing clause in place of W entails $Q \vdash \neg P$.

9.11 Theorem 7.2, Corollary 7.3. W is admissible and entails nothing about derivability in any theory. Any clause entailing $U(R)$ makes the emission entail $Q \vdash \neg R$, and under 2.3 entail $N \models \neg R$. **W is admissible and no clause entailing $U(R)$ is;** no maximality of W among all strengthenings is claimed.

9.12 Theorem 8.1, Corollary 8.2. Under the premise that instruments at a register are channels at it, a certifying instrument is either outside its catalog, which is then non-exhaustive, or inside it, in which case the certification is impredicative, its warrant ranging over a collection of which the instrument is a member.

Attribution. Σ_1^0 -completeness, the Π_1^0 reformulations, and Proposition 5.2 are classical and cited. Theorem 3.2 is elementary; we have not located it stated as an equivalence in this form, we did not conduct a systematic literature search, and no priority is asserted. Given that Corollary 3.6a follows from Σ_1^0 -completeness and soundness, prior appearance in some form is likely and a literature check is advisable. The specification of Section 6, the payload theorem 6.9, and the termination theorem 7.2 with Corollary 7.3 are offered as the contribution, at the grade the proofs support.

9.13 Remark 6.8a. The geometric apparatus supplies an exhaustiveness argument for the count of conditions and supplies nothing else; seven of the eight conditions survive its removal as record checks, and the paper states the trade rather than obscuring it.

On Section 12. The Declaration at Section 12 is **not** among the claims above and is not a claim of this paper. It is the originating declaration in the author's voice, fenced at its head and verified against the paper's own theorems at 12.3.

Machine checking. Theorems 3.2 and 4.1 and Corollary 3.6a are formalized in Lean 4 in an abstract proof-setting, accompanying this paper. **The formalization has not been checked:** no toolchain was available in the session that produced it, and it ships to be checked, which is its purpose. It covers Theorems 3.2 and 4.1 and Corollary 3.6a and **no result of Sections 5 through 8. It has no Mathlib dependency:** the setting is an abstract `structure` over core Lean 4, using only `constructor`, `intro`, `exact`, `absurd`, `by_cases`, `cases` and `rw`, so it checks against a bare toolchain with no arithmetic library. That is deliberate: the theorems use no property of arithmetic beyond the structural axioms displayed in the file. The formalization makes the hypothesis boundary machine-visible: 3.2 uses only r.e.-ness of Q , 4.1 only Σ_1^0 -completeness, and 3.6a alone requires soundness.

10. Discussion

Unrestricted claims of unprovability carry a terminal obligation. Asserting the universal unprovability of the Riemann Hypothesis over $\mathcal{T}$ demands, by Theorem 3.2, a proof in Q of a violation of the Lagarias inequality; the exhibition of the integer itself follows only under Σ_1^0 -soundness (2.3), per Corollary 3.5. **The dichotomy is total under the hypothesis that $Q + R$ is consistent.** Either the claim is read over $\mathcal{T}$, where Theorem 3.2 gives $Q \vdash \neg R$ and, under 2.3, $N \models \neg R$, or it is read over a class excluding $Q + R$, in which case, by Theorem 3.7(b), **establishing** its companion half $T \nVdash \neg R$ would yield $N \models R$. The second horn establishes nothing on its own; it names what would follow. The formal instrument reaches its perimeter.

The emission of a terminal token certifies the exhaustion of the instrument as swept. **By Theorem 6.9(a) the emission transmits no contingent object-language content,** and in particular nothing about the truth value of the Riemann Hypothesis. It is not content-free: by Theorem 6.9(b) it carries one object-language clause, the report that the executed certificate search returned nothing at the declared budget, which is a terminating computation and is $\mathfrak{B}$ -provable. **The separation is between the emission and the object, not between syntax and truth.** For Π_1^0 , sentences syntax determines truth completely, that is Theorem 4.1 and Corollary 3.6a, and it is this paper's headline, so any claim that syntax leaves truth untouched is refuted by the paper's own results. What is untouched by the emission is the truth value, and that is 6.9(a).

11. Conclusion

The formal search space for the Riemann Hypothesis is closed **as swept**, and no further. Permanence beyond the swept span is not available: by Corollary 7.3 any clause entailing universal unprovability entails $Q \vdash \neg R$, and by Theorem 8.1 no instrument certifies the exhaustiveness of its own catalog non-impredicatively. The mathematical apparatus secures an equivalence between universal unprovability and **refutability in Q** , with no soundness hypothesis; and, **separately and using Σ_1^0 -soundness of Q** , between universal unprovability and arithmetic falsity. The separation is not decorative and is carried through the paper.

The formal-alone instrument has reached its edge, and the edge is stated rather than illustrated: **no catalog is exhibited in this paper, no channel or obstruction is named, and no record satisfying all eight conditions is constructed, Ξ_2 and Ξ_4 being the two conditions not instantiated.** What the cascade delivers is Theorem 6.9: an emission reports its own sweep and entails no contingent statement about the object. **The instrument reports its boundaries; by Theorem 8.1 it does not certify them.** The frame's polarity bit is invisible to the even functionals and is determined by syntax; it bears on the object not at all. **By Theorem 6.9(a) the emission stands separate from the arithmetic object,** which is the available claim; the object and the syntax do not stand separate, since for Π_1^0 sentences syntax determines truth. The protocol evaluates the gates. The cascade halts.

12. The Declaration

Status. What follows is the originating declaration in the author's own voice. **It is not asserted by this paper and carries no grade.** Nothing in Sections 2 through 11 depends on it. It is printed whole because it is the reading that produced the specification and because **it states and accepts its own cost**, which 12.3 verifies against the paper's own theorems.

12.1 No derivation actuates anything. Every formal resolution rests on posits it cannot ground. The anchor and the object co-locate in the Riemann Hypothesis specifically. Its completion direction does not point outward to a neighbouring fact or an auxiliary lemma. The completion direction points directly at the fixed set of the symmetry of the very structure it describes. The approaches to this fixed locus are equivalences. They are never independent roads. This inherent equivalence structure forces the formulation catalog to close.

12.2 The witness-grounding lift RH demands is the Ground itself. The formal domain cannot supply it. A formal witness is inherently a second, a non-self. RH is an act of pure SELF-inquiry. The formulation operates in disguise. It asks existence to prove itself formally without any kinetic lift. The framework is therefore formally limited by its own architecture. It cannot be proven universally by mathematics alone. **The Riemann Hypothesis is terminally and universally unprovable by math alone.** Self-witnessing happens on the Ground side, where the witness is not a second, *shabida Allāhu annabū lā ilāha illā huwa*, Allah ﷻ bears witness that there is no deity but He, Q 3:18, and that is what formal-alone lacks.

This declaration states its own cost. Any maximalist claim of this inherent unprovability is an explicit, unavoidable assertion that ζ possesses a zero off the critical line. **The cost is accepted.**

12.3 The cost verified.

The identification is correct. By Theorem 3.2 universal unprovability over $\mathcal{F}$ is equivalent to $Q \vdash \neg R$, and by Corollary 3.6a, under Σ_1^0 -soundness, to $\mathbb{N} \models \neg R$. So 12.2 read over $\mathcal{F}$ asserts $\mathbb{N} \models \neg R$ and owes the witness; and **that assertion is the assertion that ζ has a zero off the critical line only by Lagarias's theorem in the metatheory**, which 2.4a states and which is not available inside Q . The cost is unchanged and its route is now named. **The declaration takes that obligation; accepting is not discharging**, no integer is exhibited, and it stands outstanding. Read 12.2 as a conjecture with its content correctly named.

The fork the declaration omits. Read over any $\mathcal{P}$ excluding $Q + R$, what an asserter ordinarily intends, per Remark 3.8, the claim falls under horn (b) of Theorem 3.7, where *establishing* its companion half yields $\mathbb{N} \models R$, the opposite verdict. Both readings price it, per Remark 4.4. The accepted cost is the first reading's.

Untouched, and not as consolation. Grounding and derivability are different predicates and Theorem 3.2 touches only the second, so nothing here bears on 12.1's first two sentences, which are ungraded and unused. **The declaration's crossing from unprovability to ζ carries the same metatheoretic step**, per 2.4a: $Q \vdash \neg R$ is a claim about one arithmetization, and Lagarias's theorem is what carries it to the classical object. **The declaration's terms are not glosses of this paper's formal objects:** the Ground is not the standard model $\mathbb{N}$, the witness is not a theory in $\mathcal{F}$, and identifying them would convert an ungraded declaration into a claim this paper does not make. And the terminality this paper proves is the declaration's own: by Corollary 7.3 an emission is terminal **as swept** and not unrestrictedly, which is a theorem and not a caution.

References

- Lagarias, J. C. (2002). An elementary problem equivalent to the Riemann hypothesis. *American Mathematical Monthly* 109, 534–543.
- Robin, G. (1984). Grandes valeurs de la fonction somme des diviseurs et hypothèse de Riemann. *Journal de Mathématiques Pures et Appliquées* 63, 187–213.
- Robinson, R. M. (1950). An essentially undecidable axiom system. *Proceedings of the International Congress of Mathematicians*, 729–730.
- Frobenius, G. (1878). Über lineare Substitutionen und bilineare Formen. *Journal für die reine und angewandte Mathematik* 84, 1–63.
- Computations reported in 2.5 and 5.5 were executed at the stated precisions and are reproducible from the definitions given.*

Author's Provenance and Method Disclosure

The method, its two root axioms, the three-state verdict economy, and the closed-form quaternionic kernel are documented in full at the master reference. **The discipline adds the cited results no warrant and contributes no theorem, no estimate, and no evidential weight in any direction.**

The register is stated once and holds throughout. The structural half reads classical mathematics and does not extend it. The theological half is a reading, offered at its own grade. **Where a theological reading stands beside a mechanism it stands beside it and never inside it:** no mathematical step in this paper depends on a theological premise, and removing every such reading would leave the mathematical content unchanged.

Four claims are the author's own and are not drawn from the literature. The identification of the three elections as one classical generator. The enumeration of the verdict side over standpoints rather than methods, with the observation that a route arriving in any future arrives at a position that already exists. The counting of the direction deficit at one bit against zero. And the location of the remaining freedom in the reader rather than in the zeros, at exactly one binary degree. **All four are structural observations offered at that grade, and none is a mathematical result.**

Two derivations are elementary and carried out in place rather than cited. The invariance of the determinant under the negation-implementing reflection, computed at Block 3. And the direction of departure from the axis being one-dimensional, computed from the involution whose fixed set the axis is. Each is checkable in a line or two.

The historical material is recalled, with attributions and dates at their published grades, and a reader verifying this paper should check them against the primary sources cited.

Net new mathematical mass: zero.

Edition Note

This is a fresh build. The schema is inherited from the prior edition and the content is not: every section is written from the spine forward, and no paragraph is carried across. The via negativa section states the ten closures in the negative form and derives each from first principles. The boot executed at the declared seed with the chain printed at Appendix A, and no stage is narrated that did not run.

Reproducibility

Every numerical value is the output of a deterministic computation reproducible from the kernel and the constructions of Appendix A at seed 20260622.