

Summary: Improper integrals

Consider $f, g > 0$. We say that g grows faster than f as x tends towards ∞ , and write this as

$$0 < f(x) \ll g(x),$$

$$\text{if } f, g \longrightarrow \infty \quad \text{and} \quad \frac{f(x)}{g(x)} \longrightarrow 0 \quad \text{as } x \rightarrow \infty.$$

Examples

For $p > 0$,

$$\ln x \ll x^p \ll e^x \ll e^{x^2}.$$

Note that $e^{x^2} = e^{(x^2)}$, not $(e^x)^2 = e^{2x}$.

Rate of Decay (as $x \rightarrow \infty$)

Consider $f, g > 0$. We say that f decays faster than g as x tends towards ∞ , and write this as

$$g(x) \gg f(x) > 0,$$

$$\text{if } f, g \longrightarrow 0 \quad \text{and} \quad \frac{f(x)}{g(x)} \longrightarrow 0 \quad \text{as } x \rightarrow \infty.$$

Examples

For $p > 0$,

$$\frac{1}{\ln x} \gg \frac{1}{x^p} \gg e^{-x} \gg e^{-x^2}.$$

Improper integrals definition

An **improper integral** is defined by $\int_a^\infty f(x) dx = \lim_{N \rightarrow \infty} \int_a^N f(x) dx$.

This improper integral **converges** if the limit exists and is finite.

This improper integral **diverges** if the limit does not exist (this includes when the limit is $\pm\infty$).

Conclusion of example: powers of x

$$\int_a^\infty \frac{dx}{x^p} \quad \begin{cases} \text{diverges} & \text{if } p \leq 1 \\ \text{converges to } \frac{a^{-p+1}}{p-1} & \text{if } p > 1 \end{cases}$$

Notation

We say that $f(x) \sim g(x)$ as $x \rightarrow \infty$ if $\frac{f(x)}{g(x)} \xrightarrow{x \rightarrow \infty} 1$.

In words, we say that $f(x)$ and $g(x)$ are **similar** as $x \rightarrow \infty$. (The idea is that $f(x)$ and $g(x)$ have the same asymptotic behavior as x tends to infinity.)

Limit comparison

The idea behind limit comparison is that if the asymptotic behavior is the same, then the improper integrals have the same behavior.

If $f(x) \sim g(x)$ as $x \rightarrow \infty$, then the two integrals $\int_a^\infty f(x) dx$ and $\int_a^\infty g(x) dx$ (for large a) either **both converge** or **both diverge**.

This also works in the case where one function decays faster than the other as x tends towards infinity.

Suppose that $g(x)$ decays faster than $f(x)$ as $x \rightarrow \infty$. That is $f(x) \gg g(x)$ as $x \rightarrow \infty$.

- If $\int_a^\infty f(x) dx$ converges, then $\int_a^\infty g(x) dx$ converges.
- If $\int_a^\infty g(x) dx$ diverges, then $\int_a^\infty f(x) dx$ diverges.

Comparison

Suppose $f(x) \geq g(x) > 0$ for $x \geq a$.

If $\int_a^\infty f(x) dx$ converges, then $\int_a^\infty g(x) dx$ converges also.

If $\int_a^\infty g(x) dx$ diverges, then $\int_a^\infty f(x) dx$ diverges also.

Definition of singularity

A **singularity** of a function $f(x)$ is a point $x = s$ such that the function $f(x)$ does not exist at $x = s$.

There are three main ways that the function can fail to exist at a point:

- $\lim_{x \rightarrow s^+} |f(x)|$ and/or $\lim_{x \rightarrow s^-} |f(x)|$ tends to ∞ . This is the case of most interest in this section.
- $\lim_{x \rightarrow s^\pm} f(x)$ does not exist. In this case the function f may oscillate, or have a jump discontinuity.
- $\lim_{x \rightarrow s^\pm} f(x)$ exists and is finite. In this case, the function f has a removable discontinuity.

Definition of improper integrals of the 2nd type

An **improper integral of the 2nd type** is an integral $\int_a^b f(x) dx$ such that the function $f(x)$ has a singularity at $x = s$ for some s with $a \leq s \leq b$.

For example, if $f(x)$ has a singularity at $x = b$, then

$$\int_a^b f(x) dx = \lim_{C \rightarrow b^-} \int_a^C f(x) dx.$$

We say

- the integral **converges** if the limit exists and is finite.
- the integral **diverges** if the limit does not exist (which includes the case that the limit is $\pm\infty$.)

Overview of improper integrals

The improper integral $\int_a^\infty \frac{dx}{x^p}$ $\begin{cases} \text{diverges} & \text{if } p \leq 1 \\ \text{converges to } \frac{a^{1-p}}{p-1} & \text{if } p > 1 \end{cases}$.

The improper integral $\int_0^a \frac{dx}{x^p}$ $\begin{cases} \text{diverges} & \text{if } p \geq 1 \\ \text{converges to } \frac{a^{1-p}}{1-p} & \text{if } p < 1 \end{cases}$.

Comparison tests for improper integrals of 2nd type

Suppose that $f(x)$ and $g(x)$ both have a singularity at $x = s$.

Suppose $f(x) > g(x) \geq 0$ for all $a \leq x \leq b$ except at $x = s$.

If $\int_a^b f(x) dx$ converges, then $\int_a^b g(x) dx$ converges also.

If $\int_a^b g(x) dx$ diverges, then $\int_a^b f(x) dx$ diverges also.




Figure 1: From left to right, we see the areas for $0 \leq x \leq 1$ and $1 \leq x < \infty$ under the graphs of $\frac{1}{x}$, $\frac{1}{x^2}$, and $\frac{1}{\sqrt{x}}$. The areas shaded in pink are infinite. The areas shaded in green are finite.

More notation

Suppose that $f(x)$ and $g(x)$ have a singularity at $x = s$. ($f, g \rightarrow \pm\infty$ as $x \rightarrow s^+$ and/or as $x \rightarrow s^-$.)

We say that $f(x)$ is **similar** to $g(x)$, and write $f(x) \sim g(x)$ as $x \rightarrow s^+$ or $x \rightarrow s^-$ if

$$\frac{f(x)}{g(x)} \rightarrow 1 \quad \text{as } x \rightarrow s^\pm. \quad (1)$$

We say that $f(x)$ grows faster than $g(x)$ as x tends towards s , and write

$$f(x) \gg g(x) \text{ as } x \rightarrow s^\pm, \text{ if } \begin{cases} f(x) \rightarrow \infty \\ g(x) \rightarrow \infty \\ \frac{g(x)}{f(x)} \rightarrow 0 \end{cases} \quad \text{as } x \rightarrow s^\pm.$$

Limit comparison tests for improper integrals of 2nd type

Suppose that $f(x)$ and $g(x)$ both have a singularity at $x = s$.

Suppose $f(x), g(x) \geq 0$ for all $a \leq x \leq b$ except at $x = s$.

1. If $f(x) \sim g(x)$ as $x \rightarrow s^\pm$,
then the two integrals $\int_a^b f(x) dx$ and $\int_a^b g(x) dx$ either **both converge** or **both diverge**.

2. Suppose that $f(x)$ grows faster than $g(x)$ as x tends towards s^\pm . In other words, $f(x) \gg g(x)$ as $x \rightarrow s^\pm$.

- If $\int_a^b f(x) dx$ converges, then $\int_a^b g(x) dx$ converges.

- If $\int_a^b g(x) dx$ diverges, then $\int_a^b f(x) dx$ diverges.

Note that this notation is exactly the same notation that we had before. The only difference is that instead of having $x \rightarrow \infty$, we have $x \rightarrow s^+$ or $x \rightarrow s^-$, where s is a finite number that is a singularity of the function of interest.