

(ii) where the compound of Formula (I) contains an alcohol functionality (-OH), an ester thereof, such as a compound wherein the hydrogen of the alcohol functionality of the compound of Formula (I) is replaced by $-\text{CO}(\text{C}_{1-8}\text{alkyl})$ (e.g. methylcarbonyl) or the alcohol is esterified with an amino acid;

(iii) where the compound of Formula (I) contains an alcohol functionality (-OH), an ether thereof, such as a compound wherein the hydrogen of the alcohol functionality of the compound of Formula (I) is replaced by $(\text{C}_{1-8}\text{alkyl})\text{C}(=\text{O})\text{OCH}_2-$ or $-\text{CH}_2\text{OP}(=\text{O})(\text{OH})_2$;

(iv) where the compound of Formula (I) contains an alcohol functionality (-OH), a phosphate thereof, such as a compound wherein the hydrogen of the alcohol functionality of the compound of Formula (I) is replaced by $-\text{P}(=\text{O})(\text{OH})_2$ or $-\text{P}(=\text{O})(\text{ONa})_2$ or $-\text{P}(=\text{O})(\text{O}-)_2\text{Ca}^{2+}$;

(v) where the compound of Formula (I) contains a primary or secondary amino functionality ($-\text{NH}_2$ or $-\text{NHR}$ where $\text{R} \neq \text{H}$), an amide thereof, for example, a compound wherein, as the case may be, one or both hydrogens of the amino functionality of the compound of Formula (I) is/are replaced by $(\text{C}_{1-10})\text{alkanoyl}$, $-\text{COCH}_2\text{NH}_2$ or the amino group is derivatised with an amino acid;

(vi) where the compound of Formula (I) contains a primary or secondary amino functionality ($-\text{NH}_2$ or $-\text{NHR}$ where $\text{R} \neq \text{H}$), an amine thereof, for example, a compound wherein, as the case may be, one or both hydrogens of the amino functionality of the compound of Formula (I) is/are replaced by $-\text{CH}_2\text{OP}(=\text{O})(\text{OH})_2$.

Certain compounds of Formula (I) may themselves act as prodrugs of other compounds of Formula (I). It is also possible for two compounds of Formula (I) to be joined together in the form of a prodrug. In certain circumstances, a prodrug of a compound of Formula (I) may be created by internally linking two functional groups in a compound of Formula (I), for instance by forming a lactone.

As used herein, the term "Formula I" may be referred to as a "compound(s) of the invention," "compound(s) of the present invention," "the invention," and "compound of Formula I." Such terms are used interchangeably. Furthermore, it is intended that the embodiments discussed herein with reference to Formula I also concern compounds of Formula I(a) or Formula I(b). Such terms are also defined to include all forms of the compound of Formula I, including hydrates, solvates, clathrates, stereoisomers, geometric isomers, crystalline (including co-crystals) and non-crystalline forms, isomorphs, polymorphs and tautomers thereof. For example, the compounds of the invention, or pharmaceutically acceptable salts thereof, may exist in unsolvated and solvated forms. When the solvent or water is tightly bound, the complex will have a well-defined stoichiometry independent of humidity. When, however, the solvent or water is weakly bound, as in channel solvates and hygroscopic compounds, the water/solvent