

Claims

1. A combination comprising (1) a GLP-1R agonist and (2) 4-(4-(1-isopropyl-7-oxo-1,4,6,7-tetrahydrospiro[indazole-5,4'-piperidine]-1'-carbonyl)-6-methoxypyridin-2-yl)benzoic acid or a pharmaceutically acceptable salt thereof, for use in a method for treating a disease or condition in a patient in need thereof, wherein:

the disease or condition is selected from nonalcoholic steatohepatitis with liver fibrosis, nonalcoholic steatohepatitis with cirrhosis, and nonalcoholic steatohepatitis with cirrhosis and with hepatocellular carcinoma or with a metabolic-related disease; and

the GLP-1R agonist is selected from:

2-({4-[2-(4-chloro-2-fluorophenyl)-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(2S)-oxetan-2-ylmethyl]-1H-benzimidazole-6-carboxylic acid;

2-({4-[2-(4-chloro-2-fluorophenyl)-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-7-fluoro-1-[(2S)-oxetan-2-ylmethyl]-1H-benzimidazole-6-carboxylic acid;

2-({4-[(2S)-2-(4-chloro-2-fluorophenyl)-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(2S)-oxetan-2-ylmethyl]-1H-benzimidazole-6-carboxylic acid;

2-({4-[(2S)-2-(4-chloro-2-fluorophenyl)-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-7-fluoro-1-[(2S)-oxetan-2-ylmethyl]-1H-benzimidazole-6-carboxylic acid;

2-({4-[2-(4-chloro-2-fluorophenyl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(2S)-oxetan-2-ylmethyl]-1H-benzimidazole-6-carboxylic acid;

2-({4-[2-(4-cyano-2-fluorophenyl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(2S)-oxetan-2-ylmethyl]-1H-benzimidazole-6-carboxylic acid;

2-({4-[2-(5-chloropyridin-2-yl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(2S)-oxetan-2-ylmethyl]-1H-benzimidazole-6-carboxylic acid;

2-({4-[2-(4-chloro-2-fluorophenyl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-3-(1,3-oxazol-2-ylmethyl)-3H-imidazo[4,5-b]pyridine-5-carboxylic acid;

2-({4-[2-(4-chloro-2-fluorophenyl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(1-ethyl-1H-imidazol-5-yl)methyl]-1H-benzimidazole-6-carboxylic acid;

2-({4-[2-(4-chloro-2-fluorophenyl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-(1,3-oxazol-4-ylmethyl)-1H-benzimidazole-6-carboxylic acid;

2-({4-[2-(4-chloro-2-fluorophenyl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-(pyridin-3-ylmethyl)-1H-benzimidazole-6-carboxylic acid;

2-({4-[2-(4-chloro-2-fluorophenyl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-(1,3-oxazol-5-ylmethyl)-1H-benzimidazole-6-carboxylic acid;

2-({4-[2-(4-chloro-2-fluorophenyl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(1-ethyl-1*H*-1,2,3-triazol-5-yl)methyl]-1*H*-benzimidazole-6-carboxylic acid;

2-({4-[2-(4-chloro-2-fluorophenyl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-(1,3-oxazol-2-ylmethyl)-1*H*-benzimidazole-6-carboxylic acid;

2-({4-[2-(4-chloro-2-fluorophenyl)-7-fluoro-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(2*S*)-oxetan-2-ylmethyl]-1*H*-benzimidazole-6-carboxylic acid;

2-({4-[2-(4-cyano-2-fluorophenyl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-(1,3-oxazol-2-ylmethyl)-1*H*-benzimidazole-6-carboxylic acid;

2-({4-[(2*S*)-2-(4-chloro-2-fluorophenyl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-7-fluoro-1-[(2*S*)-oxetan-2-ylmethyl]-1*H*-benzimidazole-6-carboxylic acid;

2-({4-[(2*S*)-2-(4-chloro-2-fluorophenyl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(2*S*)-oxetan-2-ylmethyl]-1*H*-benzimidazole-6-carboxylic acid;

2-({4-[(2*S*)-2-(4-chloro-2-fluorophenyl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-7-fluoro-1-[(2*S*)-oxetan-2-ylmethyl]-1*H*-benzimidazole-6-carboxylic acid;

2-({4-[(2*S*)-2-(4-cyano-2-fluorophenyl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(2*S*)-oxetan-2-ylmethyl]-1*H*-benzimidazole-6-carboxylic acid;

2-({4-[(2*S*)-2-(5-chloropyridin-2-yl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(2*S*)-oxetan-2-ylmethyl]-1*H*-benzimidazole-6-carboxylic acid;

2-({4-[(2*S*)-2-(4-chloro-2-fluorophenyl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(1-ethyl-1*H*-imidazol-5-yl)methyl]-1*H*-benzimidazole-6-carboxylic acid;

2-({4-[(2*R*)-2-(4-cyano-2-fluorophenyl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(2*S*)-oxetan-2-ylmethyl]-1*H*-benzimidazole-6-carboxylic acid;

2-({4-[(2*R*)-2-(5-chloropyridin-2-yl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(2*S*)-oxetan-2-ylmethyl]-1*H*-benzimidazole-6-carboxylic acid;

2-({4-[(2*R*)-2-(4-chloro-2-fluorophenyl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(1-ethyl-1*H*-imidazol-5-yl)methyl]-1*H*-benzimidazole-6-carboxylic acid;

2-({4-[2-(5-chloropyridin-2-yl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(2*S*)-oxetan-2-ylmethyl]-1*H*-benzimidazole-6-carboxylic acid;

2-({4-[(2*S*)-2-(5-chloropyridin-2-yl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(2*S*)-oxetan-2-ylmethyl]-1*H*-benzimidazole-6-carboxylic acid;

2-({4-[(2*R*)-2-(5-chloropyridin-2-yl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(2*S*)-oxetan-2-ylmethyl]-1*H*-benzimidazole-6-carboxylic acid; and

2-({4-[2-(5-chloropyridin-2-yl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(2*S*)-oxetan-2-ylmethyl]-1*H*-benzimidazole-6-carboxylic acid, DIAST-X2;

or pharmaceutically acceptable salt thereof.

2. A combination for use in a method of treating, as claimed in claim 1, wherein the GLP-1R agonist is selected from:

2-({4-[(2S)-2-(4-chloro-2-fluorophenyl)-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(2S)-oxetan-2-ylmethyl]-1H-benzimidazole-6-carboxylic acid;

2-({4-[2-(4-chloro-2-fluorophenyl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(2S)-oxetan-2-ylmethyl]-1H-benzimidazole-6-carboxylic acid;

2-({4-[2-(4-cyano-2-fluorophenyl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(2S)-oxetan-2-ylmethyl]-1H-benzimidazole-6-carboxylic acid;

2-({4-[2-(5-chloropyridin-2-yl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(2S)-oxetan-2-ylmethyl]-1H-benzimidazole-6-carboxylic acid;

2-({4-[2-(4-chloro-2-fluorophenyl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-3-(1,3-oxazol-2-ylmethyl)-3H-imidazo[4,5-b]pyridine-5-carboxylic acid;

2-({4-[2-(4-chloro-2-fluorophenyl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(1-ethyl-1H-imidazol-5-yl)methyl]-1H-benzimidazole-6-carboxylic acid;

2-({4-[(2S)-2-(4-chloro-2-fluorophenyl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(2S)-oxetan-2-ylmethyl]-1H-benzimidazole-6-carboxylic acid;

2-({4-[(2S)-2-(4-cyano-2-fluorophenyl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(2S)-oxetan-2-ylmethyl]-1H-benzimidazole-6-carboxylic acid;

2-({4-[(2S)-2-(5-chloropyridin-2-yl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(2S)-oxetan-2-ylmethyl]-1H-benzimidazole-6-carboxylic acid;

2-({4-[(2R)-2-(5-chloropyridin-2-yl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(2S)-oxetan-2-ylmethyl]-1H-benzimidazole-6-carboxylic acid; and

2-({4-[2-(5-chloropyridin-2-yl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(2S)-oxetan-2-ylmethyl]-1H-benzimidazole-6-carboxylic acid, DIAST-X2;

or a pharmaceutically acceptable salt thereof.

3. A combination for use in a method of treating, as claimed in claim 2, wherein the GLP-1R agonist is 2-({4-[2-(5-chloropyridin-2-yl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(2S)-oxetan-2-ylmethyl]-1H-benzimidazole-6-carboxylic acid, DIAST-X2, or a pharmaceutically acceptable salt thereof.

4. A combination for use in a method of treating, as claimed in any one of claims 1 to 3, wherein the 4-(4-(1-isopropyl-7-oxo-1,4,6,7-tetrahydrospiro[indazole-5,4'-piperidine]-1'-carbonyl)-6-methoxypyridin-2-yl)benzoic acid or pharmaceutically acceptable salt thereof is the 2-amino-2-(hydroxymethyl)propane-1,3-diol salt of 4-(4-(1-Isopropyl-7-oxo-1,4,6,7-tetrahydrospiro[indazole-5,4'-piperidine]-1'-carbonyl)-6-methoxypyridin-2-yl)benzoic acid.

5. A combination comprising (1) GLP-1R agonist and (2) [(1R,5S,6R)-3-{2-[(2S)-2-methylazetidin-1-yl]-6-(trifluoromethyl)pyrimidin-4-yl}-3-azabicyclo[3.1.0]hex-6-yl]acetic acid or a pharmaceutically acceptable salt thereof, for use in a method of treating a disease or condition, wherein:

the disease or condition is selected from nonalcoholic steatohepatitis with liver fibrosis, nonalcoholic steatohepatitis with cirrhosis, and nonalcoholic steatohepatitis with cirrhosis and with hepatocellular carcinoma or with a metabolic-related disease; and

the GLP-1R agonist is selected from:

2-({4-[2-(4-chloro-2-fluorophenyl)-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(2S)-oxetan-2-ylmethyl]-1H-benzimidazole-6-carboxylic acid;

2-({4-[2-(4-chloro-2-fluorophenyl)-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-7-fluoro-1-[(2S)-oxetan-2-ylmethyl]-1H-benzimidazole-6-carboxylic acid;

2-({4-[(2S)-2-(4-chloro-2-fluorophenyl)-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(2S)-oxetan-2-ylmethyl]-1H-benzimidazole-6-carboxylic acid;

2-({4-[(2S)-2-(4-chloro-2-fluorophenyl)-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-7-fluoro-1-[(2S)-oxetan-2-ylmethyl]-1H-benzimidazole-6-carboxylic acid;

2-({4-[2-(4-chloro-2-fluorophenyl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(2S)-oxetan-2-ylmethyl]-1H-benzimidazole-6-carboxylic acid;

2-({4-[2-(4-cyano-2-fluorophenyl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(2S)-oxetan-2-ylmethyl]-1H-benzimidazole-6-carboxylic acid;

2-({4-[2-(5-chloropyridin-2-yl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(2S)-oxetan-2-ylmethyl]-1H-benzimidazole-6-carboxylic acid;

2-({4-[2-(4-chloro-2-fluorophenyl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-3-(1,3-oxazol-2-ylmethyl)-3H-imidazo[4,5-b]pyridine-5-carboxylic acid;

2-({4-[2-(4-chloro-2-fluorophenyl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(1-ethyl-1H-imidazol-5-yl)methyl]-1H-benzimidazole-6-carboxylic acid;

2-({4-[2-(4-chloro-2-fluorophenyl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-(1,3-oxazol-4-ylmethyl)-1H-benzimidazole-6-carboxylic acid;

2-({4-[2-(4-chloro-2-fluorophenyl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-(pyridin-3-ylmethyl)-1*H*-benzimidazole-6-carboxylic acid;

2-({4-[2-(4-chloro-2-fluorophenyl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-(1,3-oxazol-5-ylmethyl)-1*H*-benzimidazole-6-carboxylic acid;

2-({4-[2-(4-chloro-2-fluorophenyl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(1-ethyl-1*H*-1,2,3-triazol-5-yl)methyl]-1*H*-benzimidazole-6-carboxylic acid;

2-({4-[2-(4-chloro-2-fluorophenyl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-(1,3-oxazol-2-ylmethyl)-1*H*-benzimidazole-6-carboxylic acid;

2-({4-[2-(4-chloro-2-fluorophenyl)-7-fluoro-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(2*S*)-oxetan-2-ylmethyl]-1*H*-benzimidazole-6-carboxylic acid;

2-({4-[2-(4-cyano-2-fluorophenyl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-(1,3-oxazol-2-ylmethyl)-1*H*-benzimidazole-6-carboxylic acid;

2-({4-[(2*S*)-2-(4-chloro-2-fluorophenyl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-7-fluoro-1-[(2*S*)-oxetan-2-ylmethyl]-1*H*-benzimidazole-6-carboxylic acid;

2-({4-[(2*S*)-2-(4-chloro-2-fluorophenyl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(2*S*)-oxetan-2-ylmethyl]-1*H*-benzimidazole-6-carboxylic acid;

2-({4-[(2*S*)-2-(4-chloro-2-fluorophenyl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-7-fluoro-1-[(2*S*)-oxetan-2-ylmethyl]-1*H*-benzimidazole-6-carboxylic acid;

2-({4-[(2*S*)-2-(4-cyano-2-fluorophenyl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(2*S*)-oxetan-2-ylmethyl]-1*H*-benzimidazole-6-carboxylic acid;

2-({4-[(2*S*)-2-(5-chloropyridin-2-yl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(2*S*)-oxetan-2-ylmethyl]-1*H*-benzimidazole-6-carboxylic acid;

2-({4-[(2*S*)-2-(4-chloro-2-fluorophenyl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(1-ethyl-1*H*-imidazol-5-yl)methyl]-1*H*-benzimidazole-6-carboxylic acid;

2-({4-[(2*R*)-2-(4-cyano-2-fluorophenyl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(2*S*)-oxetan-2-ylmethyl]-1*H*-benzimidazole-6-carboxylic acid;

2-({4-[(2*R*)-2-(5-chloropyridin-2-yl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(2*S*)-oxetan-2-ylmethyl]-1*H*-benzimidazole-6-carboxylic acid;

2-({4-[(2*R*)-2-(4-chloro-2-fluorophenyl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(1-ethyl-1*H*-imidazol-5-yl)methyl]-1*H*-benzimidazole-6-carboxylic acid;

2-({4-[2-(5-chloropyridin-2-yl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(2*S*)-oxetan-2-ylmethyl]-1*H*-benzimidazole-6-carboxylic acid;

2-({4-[(2*S*)-2-(5-chloropyridin-2-yl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(2*S*)-oxetan-2-ylmethyl]-1*H*-benzimidazole-6-carboxylic acid;

2-({4-[(2R)-2-(5-chloropyridin-2-yl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(2S)-oxetan-2-ylmethyl]-1H-benzimidazole-6-carboxylic acid; and

2-({4-[2-(5-chloropyridin-2-yl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(2S)-oxetan-2-ylmethyl]-1H-benzimidazole-6-carboxylic acid, DIAST-X2;
or pharmaceutically acceptable salt thereof.

6. A combination for use in a method of treating, as claimed in claim 5, wherein the GLP-1R agonist is selected from:

2-({4-[(2S)-2-(4-chloro-2-fluorophenyl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(2S)-oxetan-2-ylmethyl]-1H-benzimidazole-6-carboxylic acid;

2-({4-[2-(4-chloro-2-fluorophenyl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(2S)-oxetan-2-ylmethyl]-1H-benzimidazole-6-carboxylic acid;

2-({4-[2-(4-cyano-2-fluorophenyl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(2S)-oxetan-2-ylmethyl]-1H-benzimidazole-6-carboxylic acid;

2-({4-[2-(5-chloropyridin-2-yl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(2S)-oxetan-2-ylmethyl]-1H-benzimidazole-6-carboxylic acid;

2-({4-[2-(4-chloro-2-fluorophenyl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-3-(1,3-oxazol-2-ylmethyl)-3H-imidazo[4,5-b]pyridine-5-carboxylic acid;

2-({4-[2-(4-chloro-2-fluorophenyl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(1-ethyl-1H-imidazol-5-yl)methyl]-1H-benzimidazole-6-carboxylic acid;

2-({4-[(2S)-2-(4-chloro-2-fluorophenyl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(2S)-oxetan-2-ylmethyl]-1H-benzimidazole-6-carboxylic acid;

2-({4-[(2S)-2-(4-cyano-2-fluorophenyl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(2S)-oxetan-2-ylmethyl]-1H-benzimidazole-6-carboxylic acid;

2-({4-[(2S)-2-(5-chloropyridin-2-yl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(2S)-oxetan-2-ylmethyl]-1H-benzimidazole-6-carboxylic acid;

2-({4-[(2R)-2-(5-chloropyridin-2-yl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(2S)-oxetan-2-ylmethyl]-1H-benzimidazole-6-carboxylic acid; and

2-({4-[2-(5-chloropyridin-2-yl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(2S)-oxetan-2-ylmethyl]-1H-benzimidazole-6-carboxylic acid, DIAST-X2;
or a pharmaceutically acceptable salt thereof.

7. A combination for use in a method of treating, as claimed in claim 6, wherein the GLP-1R agonist is 2-({4-[2-(5-chloropyridin-2-yl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-

1-[(2S)-oxetan-2-ylmethyl]-1H-benzimidazole-6-carboxylic acid, DIAST-X2, or a pharmaceutically acceptable salt thereof.

8. A combination for use in a method of treating, as claimed in any one of claims 5 to 7, wherein the [(1R,5S,6R)-3-{2-[(2S)-2-methylazetidin-1-yl]-6-(trifluoromethyl)pyrimidin-4-yl}-3-azabicyclo[3.1.0]hex-6-yl]acetic acid or salt thereof is a crystal form of [(1R,5S,6R)-3-{2-[(2S)-2-methylazetidin-1-yl]-6-(trifluoromethyl)pyrimidin-4-yl}-3-azabicyclo[3.1.0]hex-6-yl]acetic acid.

9. A combination comprising (1) a GLP-1R agonist and (2) (S)-2-(5-((3-ethoxypyridin-2-yl)oxy)pyridin-3-yl)-N-(tetrahydrofuran-3-yl)pyrimidine-5-carboxamide or a pharmaceutically acceptable salt thereof, for use in a method for treating a disease or condition in a patient in need thereof, wherein:

the disease or condition is selected from nonalcoholic steatohepatitis with liver fibrosis, nonalcoholic steatohepatitis with cirrhosis, and nonalcoholic steatohepatitis with cirrhosis and with hepatocellular carcinoma or with a metabolic-related disease; and

the GLP-1R agonist is selected from:

2-({4-[2-(4-chloro-2-fluorophenyl)-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(2S)-oxetan-2-ylmethyl]-1H-benzimidazole-6-carboxylic acid;

2-({4-[2-(4-chloro-2-fluorophenyl)-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-7-fluoro-1-[(2S)-oxetan-2-ylmethyl]-1H-benzimidazole-6-carboxylic acid;

2-({4-[(2S)-2-(4-chloro-2-fluorophenyl)-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(2S)-oxetan-2-ylmethyl]-1H-benzimidazole-6-carboxylic acid;

2-({4-[(2S)-2-(4-chloro-2-fluorophenyl)-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-7-fluoro-1-[(2S)-oxetan-2-ylmethyl]-1H-benzimidazole-6-carboxylic acid;

2-({4-[2-(4-chloro-2-fluorophenyl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(2S)-oxetan-2-ylmethyl]-1H-benzimidazole-6-carboxylic acid;

2-({4-[2-(4-cyano-2-fluorophenyl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(2S)-oxetan-2-ylmethyl]-1H-benzimidazole-6-carboxylic acid;

2-({4-[2-(5-chloropyridin-2-yl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(2S)-oxetan-2-ylmethyl]-1H-benzimidazole-6-carboxylic acid;

2-({4-[2-(4-chloro-2-fluorophenyl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-3-(1,3-oxazol-2-ylmethyl)-3H-imidazo[4,5-b]pyridine-5-carboxylic acid;

2-({4-[2-(4-chloro-2-fluorophenyl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(1-ethyl-1H-imidazol-5-yl)methyl]-1H-benzimidazole-6-carboxylic acid;

2-({4-[2-(4-chloro-2-fluorophenyl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-(1,3-oxazol-4-ylmethyl)-1*H*-benzimidazole-6-carboxylic acid;

2-({4-[2-(4-chloro-2-fluorophenyl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-(pyridin-3-ylmethyl)-1*H*-benzimidazole-6-carboxylic acid;

2-({4-[2-(4-chloro-2-fluorophenyl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-(1,3-oxazol-5-ylmethyl)-1*H*-benzimidazole-6-carboxylic acid;

2-({4-[2-(4-chloro-2-fluorophenyl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(1-ethyl-1*H*-1,2,3-triazol-5-yl)methyl]-1*H*-benzimidazole-6-carboxylic acid;

2-({4-[2-(4-chloro-2-fluorophenyl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-(1,3-oxazol-2-ylmethyl)-1*H*-benzimidazole-6-carboxylic acid;

2-({4-[2-(4-chloro-2-fluorophenyl)-7-fluoro-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(2*S*)-oxetan-2-ylmethyl]-1*H*-benzimidazole-6-carboxylic acid;

2-({4-[2-(4-cyano-2-fluorophenyl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-(1,3-oxazol-2-ylmethyl)-1*H*-benzimidazole-6-carboxylic acid;

2-({4-[(2*S*)-2-(4-chloro-2-fluorophenyl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-7-fluoro-1-[(2*S*)-oxetan-2-ylmethyl]-1*H*-benzimidazole-6-carboxylic acid;

2-({4-[(2*S*)-2-(4-chloro-2-fluorophenyl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(2*S*)-oxetan-2-ylmethyl]-1*H*-benzimidazole-6-carboxylic acid;

2-({4-[(2*S*)-2-(4-chloro-2-fluorophenyl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-7-fluoro-1-[(2*S*)-oxetan-2-ylmethyl]-1*H*-benzimidazole-6-carboxylic acid;

2-({4-[(2*S*)-2-(4-cyano-2-fluorophenyl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(2*S*)-oxetan-2-ylmethyl]-1*H*-benzimidazole-6-carboxylic acid;

2-({4-[(2*S*)-2-(5-chloropyridin-2-yl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(2*S*)-oxetan-2-ylmethyl]-1*H*-benzimidazole-6-carboxylic acid;

2-({4-[(2*S*)-2-(4-chloro-2-fluorophenyl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(1-ethyl-1*H*-imidazol-5-yl)methyl]-1*H*-benzimidazole-6-carboxylic acid;

2-({4-[(2*R*)-2-(4-cyano-2-fluorophenyl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(2*S*)-oxetan-2-ylmethyl]-1*H*-benzimidazole-6-carboxylic acid;

2-({4-[(2*R*)-2-(5-chloropyridin-2-yl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(2*S*)-oxetan-2-ylmethyl]-1*H*-benzimidazole-6-carboxylic acid;

2-({4-[(2*R*)-2-(4-chloro-2-fluorophenyl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(1-ethyl-1*H*-imidazol-5-yl)methyl]-1*H*-benzimidazole-6-carboxylic acid;

2-({4-[2-(5-chloropyridin-2-yl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(2*S*)-oxetan-2-ylmethyl]-1*H*-benzimidazole-6-carboxylic acid;

2-({4-[(2S)-2-(5-chloropyridin-2-yl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(2S)-oxetan-2-ylmethyl]-1H-benzimidazole-6-carboxylic acid;

2-({4-[(2R)-2-(5-chloropyridin-2-yl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(2S)-oxetan-2-ylmethyl]-1H-benzimidazole-6-carboxylic acid, and

2-({4-[2-(5-chloropyridin-2-yl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(2S)-oxetan-2-ylmethyl]-1H-benzimidazole-6-carboxylic acid, DIAST-X2;
or pharmaceutically acceptable salt thereof.

10. A combination for use in a method of treating, as claimed in claim 9, wherein the disease or condition is nonalcoholic steatohepatitis with liver fibrosis.

11. A combination for use in a method of treating, as claimed in claim 9, wherein the disease or condition is nonalcoholic steatohepatitis with cirrhosis.

12. A combination for use in a method of treating, as claimed in any one of claims 9 to 11, wherein the GLP-1R agonist is selected from:

2-({4-[(2S)-2-(4-chloro-2-fluorophenyl)-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(2S)-oxetan-2-ylmethyl]-1H-benzimidazole-6-carboxylic acid;

2-({4-[2-(4-chloro-2-fluorophenyl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(2S)-oxetan-2-ylmethyl]-1H-benzimidazole-6-carboxylic acid;

2-({4-[2-(4-cyano-2-fluorophenyl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(2S)-oxetan-2-ylmethyl]-1H-benzimidazole-6-carboxylic acid;

2-({4-[2-(5-chloropyridin-2-yl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(2S)-oxetan-2-ylmethyl]-1H-benzimidazole-6-carboxylic acid;

2-({4-[2-(4-chloro-2-fluorophenyl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-3-(1,3-oxazol-2-ylmethyl)-3H-imidazo[4,5-b]pyridine-5-carboxylic acid;

2-({4-[2-(4-chloro-2-fluorophenyl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(1-ethyl-1H-imidazol-5-yl)methyl]-1H-benzimidazole-6-carboxylic acid;

2-({4-[(2S)-2-(4-chloro-2-fluorophenyl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(2S)-oxetan-2-ylmethyl]-1H-benzimidazole-6-carboxylic acid;

2-({4-[(2S)-2-(4-cyano-2-fluorophenyl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(2S)-oxetan-2-ylmethyl]-1H-benzimidazole-6-carboxylic acid;

2-({4-[(2S)-2-(5-chloropyridin-2-yl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(2S)-oxetan-2-ylmethyl]-1H-benzimidazole-6-carboxylic acid;

2-({4-[(2R)-2-(5-chloropyridin-2-yl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(2S)-oxetan-2-ylmethyl]-1H-benzimidazole-6-carboxylic acid; and

2-({4-[2-(5-chloropyridin-2-yl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(2S)-oxetan-2-ylmethyl]-1H-benzimidazole-6-carboxylic acid, DIAST-X2;
or a pharmaceutically acceptable salt thereof.

13. A combination for use in a method of treating, as claimed in claim 12, wherein the GLP-1R agonist is 2-({4-[2-(5-chloropyridin-2-yl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(2S)-oxetan-2-ylmethyl]-1H-benzimidazole-6-carboxylic acid, DIAST-X2, or a pharmaceutically acceptable salt thereof.

14. A combination for use in a method of treating, as claimed in claim 13, wherein the pharmaceutically acceptable salt is a tris salt.

15. A combination for use in a method of treating, as claimed in any one of claims 9 to 14, wherein the (S)-2-(5-((3-ethoxypyridin-2-yl)oxy)pyridin-3-yl)-N-(tetrahydrofuran-3-yl)pyrimidine-5-carboxamide or salt thereof is a crystal form of (S)-2-(5-((3-ethoxypyridin-2-yl)oxy)pyridin-3-yl)-N-(tetrahydrofuran-3-yl)pyrimidine-5-carboxamide.

16. A combination comprising (1) GLP-1R agonist and (2) 2-[(1R,3R,5S)-3-({5-cyclopropyl-3-[2-(trifluoromethoxy)phenyl]-1,2-oxazol-4-yl}methoxy)-8-azabicyclo[3.2.1]octan-8-yl]-4-fluoro-1,3-benzothiazole-6-carboxylic acid or a pharmaceutically acceptable salt thereof, for use in a method for treating a disease or condition in a patient in need thereof, wherein:

the disease or condition selected from nonalcoholic steatohepatitis with liver fibrosis, nonalcoholic steatohepatitis with cirrhosis, and nonalcoholic steatohepatitis with cirrhosis and with hepatocellular carcinoma or with a metabolic-related disease; and

the GLP-1R agonist is selected from:

2-({4-[2-(4-chloro-2-fluorophenyl)-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(2S)-oxetan-2-ylmethyl]-1H-benzimidazole-6-carboxylic acid;

2-({4-[2-(4-chloro-2-fluorophenyl)-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-7-fluoro-1-[(2S)-oxetan-2-ylmethyl]-1H-benzimidazole-6-carboxylic acid;

2-({4-[(2S)-2-(4-chloro-2-fluorophenyl)-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(2S)-oxetan-2-ylmethyl]-1H-benzimidazole-6-carboxylic acid;

2-({4-[(2S)-2-(4-chloro-2-fluorophenyl)-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-7-fluoro-1-[(2S)-oxetan-2-ylmethyl]-1H-benzimidazole-6-carboxylic acid;

2-({4-[2-(4-chloro-2-fluorophenyl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(2S)-oxetan-2-ylmethyl]-1H-benzimidazole-6-carboxylic acid;

2-({4-[2-(4-cyano-2-fluorophenyl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(2S)-oxetan-2-ylmethyl]-1H-benzimidazole-6-carboxylic acid;

2-({4-[2-(5-chloropyridin-2-yl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(2S)-oxetan-2-ylmethyl]-1H-benzimidazole-6-carboxylic acid;

2-({4-[2-(4-chloro-2-fluorophenyl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-3-(1,3-oxazol-2-ylmethyl)-3H-imidazo[4,5-b]pyridine-5-carboxylic acid;

2-({4-[2-(4-chloro-2-fluorophenyl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(1-ethyl-1H-imidazol-5-yl)methyl]-1H-benzimidazole-6-carboxylic acid;

2-({4-[2-(4-chloro-2-fluorophenyl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-(1,3-oxazol-4-ylmethyl)-1H-benzimidazole-6-carboxylic acid;

2-({4-[2-(4-chloro-2-fluorophenyl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-(pyridin-3-ylmethyl)-1H-benzimidazole-6-carboxylic acid;

2-({4-[2-(4-chloro-2-fluorophenyl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-(1,3-oxazol-5-ylmethyl)-1H-benzimidazole-6-carboxylic acid;

2-({4-[2-(4-chloro-2-fluorophenyl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(1-ethyl-1H-1,2,3-triazol-5-yl)methyl]-1H-benzimidazole-6-carboxylic acid;

2-({4-[2-(4-chloro-2-fluorophenyl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-(1,3-oxazol-2-ylmethyl)-1H-benzimidazole-6-carboxylic acid;

2-({4-[2-(4-chloro-2-fluorophenyl)-7-fluoro-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(2S)-oxetan-2-ylmethyl]-1H-benzimidazole-6-carboxylic acid;

2-({4-[2-(4-cyano-2-fluorophenyl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-(1,3-oxazol-2-ylmethyl)-1H-benzimidazole-6-carboxylic acid;

2-({4-[(2S)-2-(4-chloro-2-fluorophenyl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-7-fluoro-1-[(2S)-oxetan-2-ylmethyl]-1H-benzimidazole-6-carboxylic acid;

2-({4-[(2S)-2-(4-chloro-2-fluorophenyl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(2S)-oxetan-2-ylmethyl]-1H-benzimidazole-6-carboxylic acid;

2-({4-[(2S)-2-(4-chloro-2-fluorophenyl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-7-fluoro-1-[(2S)-oxetan-2-ylmethyl]-1H-benzimidazole-6-carboxylic acid;

2-({4-[(2S)-2-(4-cyano-2-fluorophenyl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(2S)-oxetan-2-ylmethyl]-1H-benzimidazole-6-carboxylic acid;

2-({4-[(2S)-2-(5-chloropyridin-2-yl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(2S)-oxetan-2-ylmethyl]-1H-benzimidazole-6-carboxylic acid;

2-({4-[(2S)-2-(4-chloro-2-fluorophenyl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(1-ethyl-1H-imidazol-5-yl)methyl]-1H-benzimidazole-6-carboxylic acid;

2-({4-[(2R)-2-(4-cyano-2-fluorophenyl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(2S)-oxetan-2-ylmethyl]-1H-benzimidazole-6-carboxylic acid;

2-({4-[(2R)-2-(5-chloropyridin-2-yl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(2S)-oxetan-2-ylmethyl]-1H-benzimidazole-6-carboxylic acid;

2-({4-[(2R)-2-(4-chloro-2-fluorophenyl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(1-ethyl-1H-imidazol-5-yl)methyl]-1H-benzimidazole-6-carboxylic acid;

2-({4-[2-(5-chloropyridin-2-yl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(2S)-oxetan-2-ylmethyl]-1H-benzimidazole-6-carboxylic acid;

2-({4-[(2S)-2-(5-chloropyridin-2-yl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(2S)-oxetan-2-ylmethyl]-1H-benzimidazole-6-carboxylic acid;

2-({4-[(2R)-2-(5-chloropyridin-2-yl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(2S)-oxetan-2-ylmethyl]-1H-benzimidazole-6-carboxylic acid, and

2-({4-[2-(5-chloropyridin-2-yl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(2S)-oxetan-2-ylmethyl]-1H-benzimidazole-6-carboxylic acid, DIAST-X2;

or pharmaceutically acceptable salt thereof.

17. A combination for use in a method of treating, as claimed in claim 16, wherein the GLP-1R agonist is selected from:

2-({4-[(2S)-2-(4-chloro-2-fluorophenyl)-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(2S)-oxetan-2-ylmethyl]-1H-benzimidazole-6-carboxylic acid;

2-({4-[2-(4-chloro-2-fluorophenyl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(2S)-oxetan-2-ylmethyl]-1H-benzimidazole-6-carboxylic acid;

2-({4-[2-(4-cyano-2-fluorophenyl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(2S)-oxetan-2-ylmethyl]-1H-benzimidazole-6-carboxylic acid;

2-({4-[2-(5-chloropyridin-2-yl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(2S)-oxetan-2-ylmethyl]-1H-benzimidazole-6-carboxylic acid;

2-({4-[2-(4-chloro-2-fluorophenyl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-3-(1,3-oxazol-2-ylmethyl)-3H-imidazo[4,5-b]pyridine-5-carboxylic acid;

2-({4-[2-(4-chloro-2-fluorophenyl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(1-ethyl-1H-imidazol-5-yl)methyl]-1H-benzimidazole-6-carboxylic acid;

2-({4-[(2S)-2-(4-chloro-2-fluorophenyl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(2S)-oxetan-2-ylmethyl]-1H-benzimidazole-6-carboxylic acid;
2-({4-[(2S)-2-(4-cyano-2-fluorophenyl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(2S)-oxetan-2-ylmethyl]-1H-benzimidazole-6-carboxylic acid;
2-({4-[(2S)-2-(5-chloropyridin-2-yl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(2S)-oxetan-2-ylmethyl]-1H-benzimidazole-6-carboxylic acid;
2-({4-[(2R)-2-(5-chloropyridin-2-yl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(2S)-oxetan-2-ylmethyl]-1H-benzimidazole-6-carboxylic acid; and
2-({4-[2-(5-chloropyridin-2-yl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(2S)-oxetan-2-ylmethyl]-1H-benzimidazole-6-carboxylic acid, DIAST-X2;
or a pharmaceutically acceptable salt thereof.

18. A combination for use in a method of treating, as claimed in claim 17, wherein the GLP-1R agonist is 2-({4-[2-(5-chloropyridin-2-yl)-2-methyl-1,3-benzodioxol-4-yl]piperidin-1-yl}methyl)-1-[(2S)-oxetan-2-ylmethyl]-1H-benzimidazole-6-carboxylic acid, DIAST-X2, or a pharmaceutically acceptable salt thereof.

19. A combination for use in a method of treating, as claimed in any one of claims 16 to 18, wherein the 2-[(1R,3R,5S)-3-({5-cyclopropyl-3-[2-(trifluoromethoxy)phenyl]-1,2-oxazol-4-yl}methoxy)-8-azabicyclo[3.2.1]octan-8-yl]-4-fluoro-1,3-benzothiazole-6-carboxylic acid or salt thereof is 2-[(1R,3R,5S)-3-({5-cyclopropyl-3-[2-(trifluoromethoxy)phenyl]-1,2-oxazol-4-yl}methoxy)-8-azabicyclo[3.2.1]octan-8-yl]-4-fluoro-1,3-benzothiazole-6-carboxylic acid.