

EP 16802532.8 – 8 November 2021

USE OF PTERIDINONE DERIVATIVE SERVING AS EGFR INHIBITOR

Technical field

5 The present invention relates to the field of medicinal chemistry; and in particular, the present invention relates to novel pteridine derivatives, pharmaceutical compositions comprising these compounds and compounds for use as inhibitors of epidermal growth factor receptor tyrosine kinase (EGFR) for treating EGFR-mediated diseases (tumors). Synthesis methods for the compounds are disclosed, but not covered by the claimed
10 invention.

Background

Cancer, also known as malignant tumor, is a group of diseases characterized by abnormal cell proliferation, metastasis, high incidence and high mortality, and is one of
15 the malignant diseases that threaten human health and cause death. Research data show that in 2008, there were 12.7 million cancer patients, in which the number of death was up to 700 million. And 20% of new cases of tumor in the world occur in China, and 24% of tumor-caused death occurs in China. Without effective prevention or better treatment options, it is estimated that by 2030 there will be 26 million new cases of cancer
20 worldwide each year and the number of cancer-caused deaths will reach 17 million. Among the existing cancers, lung cancer is the malignant tumor with the highest morbidity and mortality in the world, wherein non-small cell lung cancer (NSCLC) accounts for more than 80% of lung cancer patients. According to World Health Organization (WHO), it is predicted that by 2025, annual increase of new cases of lung
25 cancer will be more than 1 million in China. Once diagnosed as having lung cancer, patients have only little prospect for survival, and 5-year survival rate of less than 15%.

Since the 1980s, molecular mechanisms of tumorigenesis and development have been clear with deep researches on tumor molecular biology. Among many cancer-inducing factors, some of highly expressed protein kinases in cancer cells caused
30 by gene mutations are one of major factors leading to abnormal signal transduction pathways. Protein, tyrosine kinase is an important factor in signal transmission process, involved in a series of cell activities, and closely related to cell growth, differentiation, and proliferation. It catalyzes the transfer of γ -phosphate group of ATP to tyrosine residues of many important proteins and phosphorylates phenolic hydroxyl, thereby
35 transferring signals. Therefore, for developing anti-tumor drugs, it is an effective research strategy to develop selective protein kinase inhibitors to block or regulate diseases caused by these abnormal signaling pathways. Among many tyrosine kinases, epidermal growth factor receptor tyrosine kinase (EGFR) is an indispensable and important part. EGFR consists of 1186 amino acids and encodes a transmembrane glycoprotein with a molecular

EP 16802532.8 – 8 November 2021

weight of 170-kDa. EGFR can mediate multiple signal transduction pathways and transmit extracellular signals into cells, which plays an important regulatory role in the proliferation, differentiation and apoptosis of normal and tumor cells (Cell, 2000, 100, 113-127). EGFR is a constitutively expressed component in many normal epithelial tissues, such as the skin and hair follicles, while overexpressed or highly expressed in most solid tumors. For example, in lung cancer, the expression rate of EGFR reaches 40 to 80%. Therefore, the purpose of treating lung cancer can be achieved by selectively inhibiting EGFR and interfering with its signal transduction pathways, which will open up a feasible way for targeted treatment of lung cancer.

In clinical practice, EGFR-targeting drugs such as Iressa, Tarceva, etc in combination with traditional radiotherapy and chemotherapy were proved to be very effective in the treatment of lung cancer. However, clinical practice shows that most patients with non-small cell lung cancer develop acquired resistance within 6-12 months after treatment with Gefitinib or Erlotinib. Approximately 50% of cases of resistance are related to mutation in one amino acid residue in EGFR kinase domain (mutation of threonine residue at position 790 to methionine, T790M) (The New England Journal of Medicine, 2005, 352, 786-792). T790M mutation results in steric hindrance when an inhibitor binds to EGFR or increases the affinity of EGFR to ATP, so that the anticancer effects of such reversibly-binding competitive inhibitors are greatly reduced. Drug resistance not only reduces the patient's sensitivity to drugs, but also greatly reduces the quality of lives of cancer patients. For overcoming the resistance induced by T790M mutation, a series of irreversible ATP competitive inhibitors (such as CI-1033, HKI-272, PF00299804, etc.) have entered the clinical research stage. The irreversible inhibitors contain a Michael acceptor fragment that forms a covalent bond with a conserved amino acid residue (Cys797) at the ATP binding site of EGFR, thereby resulting in stronger EGFR-binding affinity as compared with the reversible inhibitor. However, this type of drugs exhibit poor selectivity to both wild-type and mutant EGFR, therefore maximal tolerated dose (MTD) thereof is low, and clinical effects are less effective. In addition, some compounds exhibit poor druggability despite good activity, which limits their clinical application.

Therefore, it is of great clinical significance and application prospects for studying and developing EGFR-targeted drugs that can selectively inhibit T790M mutation and overcome clinical resistance.

Wei Zhou et al. (2013) *J. Med. Chem.*, 56, 7821-7837 discloses a pteridine-7(8H)-one scaffold which was transformed into irreversible inhibitors of the EGFR T790M mutant.

CN 1 373 763 A relates to 8H-pteridine-7-ones, tetrahydropteridin-7-ones, 5H,

EP 16802532.8 – 8 November 2021

8H-pteridine-6,7-diones and pteridine-7-ureas that inhibit cyclin-dependent serine/threonine kinase, Wee 1 tyrosine kinase, and growth factor-mediated tyrosine kinase enzymes, and as such are useful to treat cell proliferation diseases and disorders.

WO 2006/002367 A1 discloses compounds, pharmaceutical compositions comprising such compounds and methods of using such compounds to treat or prevent diseases or disorders associated with abnormal or deregulated kinase activity, particularly diseases or disorders that involve abnormal activation of the Abl, Bcr-abl, Bmx, c-RAF, CSK, Fes, FGFR3, Flt3, GSK3 β , IR, JNK1 α 1, JNK2 α 2, Lck, MKK4, MKK6, p70S6K, PDGFR α , Rsk1, SAPK2 α , SAPK2 β , Syk and TrkB kinases.


WO 2013/170671 A1 relates to pteridine ketones synthesis and to pteridine ketones as EGFR, BLK and FLT3 inhibitors in the preparation of a medicament for tumor-related diseases.

Summary of the invention

The invention is defined in the claims. Aspects not falling within the scope of the claims are disclosed and for information only.

An object of the present invention is to provide a derivative of pteridinone with novel structure, which is capable of selectively inhibiting EGFR T790M mutation and possessing good druggability.

In the first aspect of the present invention, a compound of formula I or a pharmaceutical acceptable salt thereof is provided:



I

wherein

A is a divalent radical $-(CR_aR_b)_n-A'$, wherein R_a and R_b are independently selected from a group consisting of H, a C_{1-3} alkyl, a halogen, and n is an integer from 0 to 3;

m is an integer from 0 to 3;


R^1 is independently selected from substituted or unsubstituted phenyl, optionally substituted C_3 - C_8 cycloalkyl;

A' is a benzene ring, a five- or six-membered nitrogen-containing heterocycle;

R^2 is independently selected from a C_2 - C_4 alkenyl substituted acylamino, substituted or unsubstituted C_2 - C_4 alkenyl substituted acyl;


R^7 is selected from a group consisting of:

EP 16802532.8 – 8 November 2021



R^3 , R^4 , R^5 and R^6 are independently selected from a group consisting of H, a substituted or unsubstituted C_1 - C_6 (preferably C_1 - C_3) alkoxy, a substituted or unsubstituted C_1 - C_6 (preferably C_1 - C_3) alkyl,

- 5 or a compound or a pharmaceutical acceptable salt thereof selected from the following group:



In a particular embodiment,

- 10 R^1 is independently selected from substituted or unsubstituted phenyl;


$n = 0$;

A' is a benzene ring, or a five-membered nitrogen-containing heterocycle;

R^2 is independently selected from a C_2 - C_4 alkenyl substituted acylamino, or C_2 - C_4 alkenyl substituted acyl;

- 15 m is an integer from 0 to 2;

R^7 is selected from:









R^3 , R^4 , R^5 and R^6 are independently selected from a group consisting of H, a substituted or unsubstituted C_1 - C_3 alkoxy, a substituted or unsubstituted C_1 - C_3 alkyl.










20

In a particular embodiment of the present invention, a compound of formula I or a pharmaceutical acceptable salt thereof selected from the following group is provided:


EP 16802532.8 – 8 November 2021




EP 16802532.8 – 8 November 2021




EP 16802532.8 – 8 November 2021



In a particular embodiment of the present invention, a compound of formula I or a pharmaceutical acceptable salt thereof selected from the following group is provided:



EP 16802532.8 – 8 November 2021

In the second aspect, a pharmaceutical composition is provided in the present invention, comprising a compound or a pharmaceutical acceptable salt thereof of the first aspect and a pharmaceutically acceptable carrier or excipient.

In a preferred embodiment, the pharmaceutical composition is in a form suitable for oral administration, including but not limited to tablets, solutions, suspensions, capsules, granules, powders.

In the third aspect, a compound of the first aspect for use in treating or preventing EGFR-mediated diseases, or inhibiting EGFR is provided in the present invention, wherein

the EGFR-mediated disease is cancer.

In a particular embodiment, the cancer is selected from a group consisting of non-small cell lung cancer, small cell lung cancer, lung adenocarcinoma, lung squamous cell carcinoma, breast cancer, prostate cancer, glioblastoma, ovarian cancer, squamous cell carcinoma of head and neck, cervical cancer, esophageal cancer, liver cancer, kidney cancer, pancreas cancer, colon cancer, skin cancer, leukemia, lymphoma, stomach cancer, multiple myeloma and solid tumors.

Also disclosed herein but not covered by the claimed invention is a method for treating or preventing EGFR-mediated diseases by using the compound of the first aspect of the invention.

It is disclosed that the EGFR-mediated disease is cancer; preferably, the cancer is selected from a group consisting of non-small cell lung cancer, small cell lung cancer, lung adenocarcinoma, lung squamous cell carcinoma, breast cancer, prostate cancer, glioblastoma, ovarian cancer, squamous cell carcinoma of head and neck, cervical cancer, esophageal cancer, liver cancer, kidney cancer, pancreas cancer, colon cancer, skin cancer, leukemia, lymphoma, stomach cancer, multiple myeloma and solid tumors.

It should be understood that in the present invention, the technical features specifically mentioned above and below (such as in the Examples) can be combined

EP 16802532.8 – 8 November 2021

with each other, thereby constituting a new or preferred technical solution which needs not be individually described.

Description of drawings

5 Figure 1 shows pharmacodynamic evaluation of a series of compounds: after orally administered to nude mice of H1975 cell xenograft model for 14 days, Compound 14 exhibited superior efficacy over CO-1686.

Figure 2 shows pharmacodynamics comparison between Compound 14 and CO-1686 and AZD9291: Compound 14 (10 mg/kg) exhibited better efficacy over CO-1686 (50
10 mg/kg), Compound 14 (25 mg/kg) exhibited comparable efficacy to AZD9291 (25 mg/kg), both of which can effectively reduce the size of a tumor, and the mice still live and the tumor did not recur after administration was stopped for 1 month.

Figure 3 shows that compound 14 exhibited better selectivity on wild-type EGFR. Nude mice of EGFR wild-type A431 cell xenograft model were administered for 14 days
15 by oral administration, compound 14 (25 mg/kg) exhibited performance comparable to the same dose of AZD9291, that is, the size of the tumor did not substantially change, and the selectivity was significantly better than afatinib (a medicament of 2nd generation).

Figure 4 shows the toxicity assessment of Compound 14 and AZD9291.

Figure 5 shows the pathological analysis of tumor-bearing mice after 14 consecutive
20 days of administration.

Modes for carrying out the invention

Through comprehensive and intensive research, the inventors have unexpectedly found derivatives of pteridinone with novel structure, which are capable of selectively
25 inhibiting EGFR T790M mutation and possessing good druggability. For these compounds, IC₅₀ values of inhibitory activity on EGFR-T790M/L858R kinase and IC₅₀ values of inhibiting proliferation of H1975 cells (non-small cell lung cancer cells, EGFR-T790M / L858R) reached nM levels and these compounds exhibited excellent solubility. Based on the above findings, the present invention is completed.

30

Definition on terms

The terms mentioned herein are further defined as follows:

As used herein, “alkyl” refers to a saturated straight chain or branched chain alkyl having 1 to 10 carbon atoms, and preferably alkyl includes an alkyl with 2-8 carbon atoms,
35 1-6 carbon atoms, 1-4 carbon atoms, 1-3 carbon atoms in length. Examples of alkyl include, but not limited to, methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl, heptyl, and the like. Alkyl can be substituted by one or more substituents, for example substituted

EP 16802532.8 – 8 November 2021

by a halogen or a haloalkyl. For example, alkyl may be an alkyl substituted by 1-4 fluorine atoms, or an alkyl substituted by fluorinated alkyl.

As used herein, "cycloalkyl" refers to a saturated cyclic alkyl having 3-10, preferably 3-8 ring carbon atoms. Exemplary cycloalkyl includes, but not limited to, cyclopropyl, cyclobutyl, cyclohexyl, cycloheptyl, and the like. Cycloalkyl can be substituted by one or more substituents, for example substituted by a halogen or a haloalkyl. For example, cycloalkyl may be substituted by 1-4 fluorine atoms. In a preferred embodiment, the cycloalkyl is cyclohexyl.

As used herein, "alkoxyl" refers to an oxy substituted by alkyl. A preferred alkoxyl is an alkoxyl with 1-6 carbon atoms in length, more preferably an alkoxyl with 1-4 carbon atoms in length. Examples of alkoxyl include, but not limited to, methoxyl, ethoxyl, propoxyl and the like. In a particular embodiment, alkoxy can be a substituted alkoxy group, for example, an alkoxy-substituted alkoxy. In a particular embodiment, a C₁-C₃ alkoxy-substituted C₁-C₃ alkoxy is preferable.

As used herein, "alkenyl" generally means a monovalent hydrocarbon group with at least one double bond, generally comprises 2-8 carbon atoms, preferably 2-6 carbon atoms and may be of straight or branched chain. Examples of alkenyl include, but not limited to, ethenyl, propenyl, iso-propenyl, butenyl, iso-butenyl, hexenyl, and the like.

As used herein, "halogen" refers to fluorine, chlorine, bromine and iodine.

As used herein, "aryl" means a monocyclic, bicyclic or tricyclic aromatic group with 6 to 14 carbon atoms, and includes phenyl, naphthyl, phenanthryl, anthryl, indenyl, fluorenyl, tetralin, indanyl and the like. Aryl can be optionally substituted with 1 - 5 (e.g., 1, 2, 3, 4, or 5) substituents selected from: a halogen, a C₁₋₄ aldehyde group, a C₁₋₆ alkyl, a cyano, a nitro, an amino, a hydroxyl, a hydroxymethyl, a halogen-substituted alkyl (e.g., trifluoromethyl), halogen-substituted alkoxyl (e.g., trifluoromethoxyl), a carboxyl, a C₁₋₄ alkoxyl, a ethoxyformyl, N(CH₃) and a C₁₋₄ acyl, a heterocyclyl or a heteroaryl, and the like.

As used herein, "heterocycle group" includes, but not limited to, 5- or 6- member heterocyclic groups comprising 1-3 heteroatoms selected from O, S or N, including (but not limited to) a furyl, a thienyl, a pyrrolyl, a pyrrolidinyl, a pyrazolyl, an imidazolyl, a triazolyl, an oxazolyl, a pyranlyl, a pyridyl, a pyrimidinyl, a pyrazinyl, a piperidinyl, a morpholinyl and the like.

As used herein, "aromatic heterocycle group" means that the group comprises 5 to 14 ring atoms, and 6, 10, or 14 electrons are shared in the ring system. And the contained ring atoms are carbon atoms and 1-3 heteroatoms optionally selected from O, N, S. Useful aromatic heterocycle group includes a piperazinyl, a morpholinyl, a piperidinyl, a pyrrolidinyl, a thienyl, a furyl, a pyranlyl, a pyrrolyl, an imidazolyl, a pyrazolyl, a pyridyl,

EP 16802532.8 – 8 November 2021

including, but not limited to, 2-pyridyl, 3-pyridyl and 4-pyridyl, a pyrazinyl, a pyrimidinyl and the like.

Aromatic heterocycle group may be optionally substituted by 1-5 (e.g., 1, 2, 3, 4 or 5) substituents selected from: a halogen, a C₁₋₄ aldehyde group, a C₁₋₆ a straight chain or
5 branched chain alkyl, a cyano, a nitro, an amino, a hydroxyl, a hydroxymethyl, a halogen-substituted alkyl (e.g., trifluoromethyl), a halogen-substituted alkoxy (e.g., trifluoromethoxy), a carboxyl, a C₁₋₄ alkoxy, an ethoxyformyl, N(CH₃) and a C₁₋₄ acyl.

As used herein, "acyl" refers to a group with the structure of formula "-C(O)-R", wherein R can be selected from an alkyl, an alkenyl or an alkynyl. And R can be
10 optionally substituted. In a particular embodiment, R in "acyl" described herein is substituted with an optionally substituted alkenyl. In a preferred embodiment, R in "acyl" described herein is substituted with optionally substituted C₂-C₄ alkenyl. In a preferred embodiment, "acyl" according to the present invention is formyl substituted with optionally substituted alkenyl, for example, formyl substituted with vinyl, formyl
15 substituted with NR_xR_y-substituted propenyl, wherein R_x and R_y can independently selected from alkyl or H.

As used herein, "acylamino" refers to a group with the structure of formula "-NH-C(O)-R", wherein R can be selected from an alkyl, an alkenyl, an alkynyl, a NR_xR_y-substituted alkyl, a NR_xR_y-substituted alkenyl, a NR_xR_y-substituted alkynyl, a
20 halogen-substituted alkyl, a cyano-substituted alkenyl, wherein R_x and R_y are independently selected from an alkyl or an alkenyl. In a particular embodiment, R in "acylamino" described herein is substituted with an alkenyl. In a preferred embodiment, R in "acylamino" described herein is substituted with C₂-C₄ alkenyl. In a preferred embodiment, "acylamino" according to the present invention is formylamino substituted
25 with vinyl.

As used herein, "optionally substituted" means that the group modified by the term can be optionally substituted by 1-5 (e.g., 1, 2, 3, 4, or 5) substituents selected from: a halogen, a C₁₋₄ aldehyde group, a C₁₋₆ straight chain or branched chain alkyl, a cyano, a nitro, an amino, a hydroxyl, a hydroxymethyl, a halogen-substituted alkyl (e.g.,
30 trifluoromethyl), a halogen-substituted alkoxy (e.g., trifluoromethoxy), a carboxyl, a C₁₋₄ alkoxy, an ethoxyformyl, N(CH₃) and a C₁₋₄ acyl.

Compounds of the present invention


The inventors have synthesized a series of candidates possessing EGFR inhibitory
35 activity. A series of 7(8*H*)-pteridinone compounds with novel structure were designed and synthesized through optimization of structure of the obtained candidate compounds, and structurally characterized. Biological activities and physical-chemical properties of this

EP 16802532.8 – 8 November 2021

series of compounds were tested, and a series of compounds capable of selectively inhibiting EGFR T790M mutation and possessing good druggability were obtained. Among these compounds, IC₅₀ values of inhibitory activity of compound ZW-39 on EGFR-T790M/L858R kinase was 3.9 nM, IC₅₀ values of inhibiting proliferation of H1975
5 cells (non-small cell lung cancer cells, EGFR-T790M / L858R) was 9 nM, and its solubility in water (20 mM phosphate buffer, pH 6.8) was 1367 µg/mL.

After further research, the inventors found that when judging a candidate compound, not only the absolute activity but also the solubility of the candidate compound should be taken into consideration, that is, the druggability of a candidate compound. For evaluating
10 a candidate compounds as a whole, ratio of the solubility of a candidate compound to IC₅₀ value of EGFR-T790M/L858R kinase inhibitory activity (ratio 1) and ratio of the solubility of the candidate compound to IC₅₀ value of inhibiting proliferation of H1975 cell (ratio 2) was creatively adopted by the inventors as criteria, and selective EGFR inhibitor AZD9291 of the third generation, which is currently in Phase II clinical trials,
15 was used as a positive control, thereby identifying a series of excellent candidate compounds.

In a particular embodiment, the compound of the present invention is a compound of the following general formula I or a pharmaceutically acceptable salt thereof:




20 wherein

A is a divalent radical $-(CR_aR_b)_n-A'$, wherein R_a and R_b are independently selected from a group consisting of H, a C₁₋₃ alkyl, a halogen, and n is an integer from 0 to 3;

EP 16802532.8 – 8 November 2021


m is an integer from 0 to 3;

R¹ is independently selected from substituted or unsubstituted phenyl, optionally substituted C₃-C₈ cycloalkyl; A' is a benzene ring, a five- or six-membered nitrogen-containing heterocycle; R² is independently selected from a C₂-C₄ alkenyl substituted acylamino, substituted or unsubstituted C₂-C₄ alkenyl substituted acyl; R⁷ is selected from a group consisting of:




R³, R⁴, R⁵ and R⁶ are independently selected from a group consisting of H, a substituted or unsubstituted C₁-C₆ (preferably C₁-C₃) alkoxy, a substituted or unsubstituted C₁-C₆ (preferably C₁-C₃) alkyl,

or a compound or a pharmaceutical acceptable salt thereof selected from the following group:









12.











In a particular embodiment, following compounds are provided in the present invention:




EP 16802532.8 – 8 November 2021




EP 16802532.8 – 8 November 2021



EP 16802532.8 – 8 November 2021





In a further preferred embodiment, R^1 is independently selected from substituted or unsubstituted phenyl; $n = 0$; A' is a benzene ring, or a five-membered nitrogen-containing heterocycle; R^2 is independently selected from a C_2 - C_4 alkenyl substituted acylamino, or C_2 - C_4 alkenyl substituted acyl; m is an integer from 0 to 2; R^7 is selected from:



R^3 , R^4 , R^5 and R^6 are independently selected from a group consisting of H, a substituted or unsubstituted C_1 - C_3 alkoxy, a substituted or unsubstituted C_1 - C_3 alkyl.

In a preferred embodiment, following compounds are provided in the present invention:



EP 16802532.8 – 8 November 2021

Based on the above compounds, a pharmaceutical composition is provided in the present invention, wherein the composition comprises a compound of formula I or a pharmaceutical acceptable salt thereof of the present invention and a pharmaceutically acceptable carrier or excipient.

5 The example of the pharmaceutically acceptable salt of the compound of the invention includes, but not limited to, inorganic salts and organic salts, for example, hydrochloride, hydrobromide, sulfate, citrate, lactate, tartrate, maleate, fumarate, mandelate and oxalate; and inorganic and organic salts formed with base, such as sodium hydroxyl, tri(hydroxymethyl) aminomethane (TRIS, tromethamine) and N-methyl
10 glucamine.

The skilled in the art can determine the optimal dosage of each active ingredient in the pharmaceutical compositions of the invention, although the individual's need differs. Generally, the compounds of the present invention or a pharmaceutically acceptable salt thereof are administered orally to a mammal in a daily dose of about 0.0025 to 50 mg / kg
15 body weight, preferably, administered orally about 0.01 to 10 mg per kg. For example, a unit oral dose may comprise about 0.01 to 50 mg, preferably about 0.1 to 10 mg of compounds of the invention. A unit dose may be administered once or more times, one or more tablets a day, with each tablet containing about 0.1 to 50 mg, preferably about 0.25 to 10 mg of the compounds of the invention or a solvate thereof.

20 The pharmaceutical composition of the invention may be formulated into forms

EP 16802532.8 – 8 November 2021

suitably for various administration routes, including, but not limited to, parenteral, subcutaneous, intravenous, intramuscular, intraperitoneal, transdermal, buccal, intrathecal, intracranial, intranasal or topical administration, for treating cancer and other diseases. The administration amount is the amount effective to ameliorate or eliminate one or more diseases. For treating a specific disease, the effective amount is the amount sufficient for ameliorating or alleviating disease relevant symptoms. Such amount can be administered in a single dose or according to the effective therapeutic schedule. The administration amount may be effective in curing diseases, but usually the purpose is to ameliorate symptoms of diseases. Generally, administration needs to be repeated for achieving the amelioration of symptoms. The dose can be determined according to the age, healthy status and weight of the patient, concurrent treatment, frequency of treatment, and the desired therapeutic benefit.

The pharmaceutical formulations of the invention may be administered to any mammal, as long as the compounds of the present invention are effective to them. In the mammals, human being is the most important.

The compound or pharmaceutical composition of the present invention may be useful in the treatment or prevention of various diseases mediated by epidermal growth factor receptor kinase (EGFR). Herein, the EGFR-mediated disease is various cancers. The cancers include, but are not limited to, non-small cell lung cancer, breast cancer, prostate cancer, glial cell tumors, ovarian cancer, head and neck squamous cell carcinoma, cervical cancer, esophageal cancer, liver cancer, kidney cancer, pancreatic cancer, colon cancer, skin cancer, leukemia, lymphoma, gastric cancer, multiple myeloma, and solid tumors.

The pharmaceutical formulations of the invention can be prepared in a known manner, e.g. by conventional mixing, granulating, tableting, dissolving, or lyophilizing processes. When manufacturing oral formulation, a solid excipient and an active compound can be combined, and the mixture can be ground optionally. If necessary, appropriate additives can be added, and the mixture can be processed into particles for obtaining tablets or troche cores.

Suitable excipients, particularly fillers, include for example, sugars, such as lactose or sucrose, mannitol or sorbitol; cellulose preparations or calcium phosphates, such as tricalcium phosphate or calcium hydrogen phosphate; and a binder, such as starch paste, including corn starch, wheat starch, rice starch, potato starch, gelatin, tragacanth, methyl cellulose, hydroxypropyl methyl cellulose, sodium carboxymethyl cellulose, or polyvinyl pyrrolidone. If desired, disintegrating agent, such as, the above-mentioned starches as well as carboxymethyl starch, cross-linked polyvinyl pyrrolidone, agar, or alginic acid or a salt thereof such as sodium alginate, can be added. Adjuvants especially includes flow

EP 16802532.8 – 8 November 2021

modifier and lubricants, such as silica, talc, stearates such as magnesium and calcium stearate, stearic acid or polyethylene glycol. If necessary, the troche core can be coated suitably to be resistant to gastric juices. For this purpose, concentrated saccharide solutions may be applied, which may contain gum arabic, talc, polyvinylpyrrolidone, polyethylene glycol and / or titanium dioxide, lacquer solutions, and suitable organic solvents or solvent mixtures. For preparing gastric juice-resistant coatings, suitable cellulose solutions may be used, such as cellulose acetate phthalate or hydroxypropylmethyl cellulose phthalate. Dyestuffs or pigments may be added to the tablets or troche cores coating, for example, for identifying or characterizing the dosage combinations of active ingredients.

Accordingly, a method for treating EGFR mediated diseases is disclosed herein, but not covered by the claimed invention, comprising administering the compound of the invention or the pharmaceutical composition comprising the compound of the invention to a object in need thereof.


The administration methods include, but not limited to, those methods known in the art, which can be determined according to the condition of the patient, including, but not limited to, parenteral, subcutaneous, intravenous, intramuscular, intraperitoneal, transdermal, buccal, intrathecal, intracranial, intranasal or topical administration.

A compound of the present invention for use in treating EGFR mediated diseases, wherein the EGFR mediated disease is cancer, is also included in the present invention.

The technical solutions of the present invention are further described below with reference to specific implementation examples. However, the following embodiments are not intended to limit the present invention. All the application methods adopted according to the principles and technical solutions of the present invention fall within the scope of the present invention. In the following examples, experimental methods with no specific conditions are specified, usually according to the usual conditions or according to the manufacturer's suggested conditions. Unless otherwise indicated, percentages and parts are by weight.

Material and method

Synthesis of 7(8H)-pteridinone compounds of the present invention



EP 16802532.8 – 8 November 2021

Reagents and conditions: (a) amine, DIPEA, 1,4-dioxane, r.t.; (b) ArNH₂, DIPEA, 1,4-dioxane, r.t.; (c) Pd/C, H₂, EtOH; (d) R²COCOOEt, HOAc, EtOH, reflux; (e) trifluoroacetic acid, CH₂Cl₂, 0°C - r.t.; (f) acid chloride, Et₃N, CH₂Cl₂, 0°C - r.t. or acid chloride, 1-methyl-2-pyrrolidone, CH₃CN, 0°C - r.t.


5 In the above preparation procedure, R¹, R², R³, R⁴ are defined as described above. Various starting compounds routinely obtained in the art as raw material can be used by a skilled person in the art according to actual needs to prepare the compound of the present invention.

10 Example

Example 1


The particular method for steps a-f as said above is shown as follows:

Synthesis of *tert*-butyl (3-(2-chloro-5-nitropyrimidyl-4-amino)phenyl)carbamate



15 2,4-dichloro-5-nitro-pyrimidine (3.80 g, 19.59 mmol) was placed into a 100 mL round bottom flask, 80 mL of 1,4-dioxane was added, and stirred at room temperature. *Tert*-butyl (3-aminophenyl) carbamate (4 g, 19.21 mmol) and N,N-diisopropylethylamine (2.98 g, 23.05 mmol) were dissolved in 20 mL of 1,4-dioxane. The resulting solution was added dropwise into the reaction solution as said above. Upon completion of addition, the
20 resulting mixture was stirred at room temperature for 0.5 h, and TLC showed that the raw material was completely conversed. The solvent was removed by rotary evaporation, and the crude product was separated through silica gel column chromatography (petroleum ether / ethyl acetate = 10: 1, v/v) to obtain *tert*-butyl (3-(2-chloro-5-nitropyrimidyl-4-amino)phenyl)carbamate as orange solids (5.9 g, yield
25 84%). ¹H NMR (400 MHz, DMSO-*d*₆): δ10.37(s, 1H), 9.48(s, 1H), 9.13(s, 1H), 7.67 (s, 1H), 7.31 (m, 2H), 7.15(m, 1H), 1.48(s, 9H).


Synthesis of *tert*-butyl (3-((2-(methoxymethoxy)-4-(4-methyl-1-piperazinyl)phenyl)amino)-5-nitro-4-pyrimidinyl)amino)phenyl)carbamate



EP 16802532.8 – 8 November 2021

tert-butyl-3-(2-chloro-5-nitropyrimidin-4-amino)phenylcarbamate (902 mg, 2.47 mmol), 2-(methoxymethoxy)-4-(4-methyl-1-piperazinyl)-aniline (775 mg, 2.47 mmol) and DIPEA (956 mg, 7.40 mmol) were weighed into a 100 ml of round bottom flask. 50 ml of tetrahydrofuran was added, heated to reflux and mixed overnight. After the reaction was completed, the solvent was removed by rotary evaporation, diluted with water and extracted with ethyl acetate (100 mL × 2). The organic phase was washed with saturated ammonium chloride solution (50 mL × 2) and saturated sodium chloride solution, and dried over sodium sulfate. The solvent was removed by rotary evaporation and the crude product was purified by column chromatography on silica gel (dichloromethane / methanol = 30: 1, v / v) to give *tert*-butyl (3-((2-(methoxymethoxy)-4-(4-methyl-1-piperazinyl)phenyl)amino)-5-nitro-4-pyrimidinylamino)phenyl)carbamate as orange-red solid, 945 mg, yield 66%. ¹H NMR (400 MHz, DMSO-*d*₆): δ 10.25 (s, 1H), 9.44 (s, 1H), 9.28 (s, 1H), 9.04 (s, 1H), 7.54 (s, 1H), 7.42 (d, *J*=8.8 Hz, 1H), 7.23-7.20 (m, 2H), 7.08 (t, *J*=8.0 Hz, 1H), 6.70 (s, 1H), 6.43 (d, *J*=8.8 Hz, 1H), 5.15 (s, 2H), 3.32 (s, 3H), 3.11 (br, 4H), 2.46 (br, 4H), 2.23 (s, 3H), 1.46 (s, 9H).

Synthesis of *tert*-butyl (3-(2-((2-(methoxymethoxy)-4-(4-methyl-1-piperazinyl)phenyl)amino)-7-oxo-8(7*H*)-pteridinyl)phenyl)carbamate




tert-butyl (3-((2-((2-(methoxymethoxy)-4-(4-methyl-1-piperazinyl)phenyl)amino)-5-nitro-4-pyrimidinylamino)phenyl)carbamate (940 mg, 1.62 mmol) was weighed into a 400 mL round bottom flask. 120 mL of ethanol, 100 mL of methylene chloride and 250 mg of palladium on carbon (10% Pd) were added. Hydrogen was ventilated and the reaction system was stirred for 8 hours at room temperature. After the reaction was completed, the reaction mixture was suction-filtered, the filtrate was spin-dried, and the product was directly used in the next reaction without purification.

The product from the previous reaction was placed in a 50 mL round bottom flask and 1.45 mL of glacial acetic acid and 25 mL of absolute ethanol were added followed by ethyl glyoxylate (50% in toluene) (0.495 mL, 2.43 mmol), heated to reflux and stirred overnight. After the reaction was completed, the solvent was removed by rotary evaporation and 15 mL of ethanol was added for recrystallization. Solids precipitated and were suction-filtered. The filter cake was washed with ethanol, aqueous ammonia and deionized water and dried to give *tert*-butyl (3-(2-((2-(methoxymethoxy)-4-(4-methyl-1-piperazinyl)phenyl)amino)-7-oxo-8(7*H*)-pteridinyl)phenyl)carbamate as orange-red

EP 16802532.8 – 8 November 2021

solids 452 mg, yield 47%. ¹H NMR (400 MHz, DMSO-*d*₆): δ 9.65 (s, 1H), 8.80 (s, 1H), 8.55 (br, 1H), 8.02 (s, 1H), 7.59 (s, 1H), 7.52 (d, *J*=8.0 Hz, 1H), 7.43 (t, *J*=8.0 Hz, 1H), 7.36-7.33 (m, 1H), 6.97 (d, *J*=8.0 Hz, 1H), 6.65 (d, *J*=2.0 Hz, 1H), 6.16 (br, 1H), 5.17 (s, 2H), 3.36 (s, 3H), 3.03 (br, 4H), 2.44 (t, *J*=3.6 Hz, 4H), 2.22 (s, 3H), 1.46 (s, 9H).


5 **Synthesis of 8-(3-aminophenyl)-2-((2-(methoxymethoxy)-4-(4-methyl-1-piperazinyl)phenyl)amino)-7(8H)pteridinone**



tert-butyl

10 (3-((2-((2-(methoxymethoxy)-4-(4-methyl-1-piperazinyl)phenyl)amino)-7-oxo-8(7H)-pteridinyl)phenyl)carbamate (452 mg, 0.77 mmol) was weighed into a 25 mL round bottom flask. 7 mL of methylene chloride was added and 2 mL of trifluoroacetic acid was added in an ice-bath with stirring. The reaction system was stirred for another 1 hour in an ice bath and for 1 hour at room temperature. After the reaction was completed, saturated sodium bicarbonate solution was added to neutralize the solution to pH = 9. The filtrate
15 was extracted with dichloromethane. The organic phase was washed with deionized water and saturated sodium chloride solution, and dried, and the solvent was spin-dried. The crude product was purified by column chromatography on silica gel (dichloromethane / methanol = 20: 1, v/v) to obtain 8-(3-aminophenyl)-2-((2-(methoxymethoxy)-4-(4-methyl-1-piperazinyl)phenyl)amino)-7(8H)pteridinone, orange-red solids 132 mg, yield 35%. MS
20 (ESI) *m/z* 489.2 [M + H]⁺.

Synthesis of N-(3-(2-((2-(methoxymethoxy)-4-(4-methyl-1-piperazinyl)phenyl)amino)-7-oxo-8(7H)pteridinyl)phenyl)acrylamide (Compound 9) (Comparative)




25 8-(3-aminophenyl)-2-((2-(methoxymethoxy)-4-(4-methyl-1-piperazinyl)phenyl)amino)-7(8H)pteridinone (132 mg, 0.270 mmol) was weighed into a 25 mL round bottom flask. 4 mL of N-methylpyrrolidone was added and stirred in an ice bath. Acryloyl chloride (25 μL, 0.297 mmol) was dissolved in 2 mL of acetonitrile and added dropwise to the reaction

EP 16802532.8 – 8 November 2021

mixture with a constant pressure dropping funnel. Upon addition, the mixture was stirred for half an hour in an ice bath and then for 2 hours at room temperature. After the reaction was completed, saturated sodium bicarbonate solution was added to the reaction liquid and stirred for another 15 minutes. The reaction mixture was diluted with water and extracted with ethyl acetate. The organic phase was washed with deionized water and saturated sodium chloride solution. After concentration, the residue was purified by column chromatography on silica gel (methylene chloride / methanol / aqueous ammonia = 100: 5: 0.5, v/v) to obtain *N*-(3-(2-((2-(methoxymethoxy)-4-(4-methyl-1-piperazinyl)phenyl)amino)-7-oxo-8(7*H*)pteridiny)phenyl)acrylamide, orange solids 101 mg, yield 69%. ¹H NMR (400 MHz, DMSO-*d*₆): δ 10.40 (s, 1H), 8.81 (s, 1H), 8.53 (br, 1H), 8.03 (s, 1H), 7.85 (d, *J*=6.4 Hz, 1H), 7.72 (s, 1H), 7.53 (t, *J*=8.0 Hz, 1H), 7.34-7.33 (m, 1H), 7.10 (d, *J*=8.0 Hz, 1H), 6.64 (s, 1H), 6.46 (dd, *J* = 16.8, 10.0 Hz, 1H), 6.27 (dd, *J* = 16.8, 1.6 Hz, 1H), 6.13-6.09 (m, 1H), 5.78 (dd, *J* = 10.0, 1.6 Hz, 1H), 5.16 (s, 2H), 3.35 (s, 3H), 3.00 (br, 4H), 2.43 (t, *J*=4.4 Hz, 4H), 2.22 (s, 3H). HRMS (ESI) (*m/z*): [M + H]⁺calcd for C₂₈H₃₁N₈O₄, 543.2468; found, 543.2476.

Following compounds were synthesized using the same or similar synthesis route as described above:


(*S*)-8-(1-acryloyl-3-pyrrolidinyl)-2-((2-methoxy)-4-(4-methyl-1-piperazinyl)phenyl)amino)-7(8*H*)pteridinone(Compound 1) (Comparative)



orange-red solids, yield 40%. ¹H NMR (400 MHz, DMSO-*d*₆): δ 8.96 (s, 1H), 8.73 (s, 0.5 H), 8.72 (s, 0.5H), 7.86 (s, 0.5 H), 7.84 (s, 0.5 H), 7.32 (m, 1H), 6.57 (s, 1H), 6.43-6.42 (m, 1H), 6.16 (ddd, *J* = 16.8, 10.0, 2.4 Hz, 1H), 5.72 (dd, *J* = 10.0, 2.4 Hz, 1H), 5.65 (dd, *J* = 10.0, 2.4 Hz, 1H), 4.08 (t, *J* = 10.2 Hz, 0.5H), 3.88-3.82 (m, 0.5 H), 3.76 (s, 3H), 3.67-3.62 (m, 1H), 3.58-3.55 (m, 1H), 3.21-3.17 (m, 4H), 2.81-2.71 (m, 1H), 2.68-2.61 (m, 4H), 2.38 (s, 1.4H), 2.34 (s, 1.6H), 2.04-1.96 (m, 2H). HRMS (ESI) (*m/z*): [M + H]⁺calcd for C₂₅H₃₁N₈O₃, 491.2519; found, 491.2520. HPLC purity: 95.7%, retention time = 9.43 min.


(*R*)-8-(1-acryloyl-3-pyrrolidinyl)-2-((2-methoxy)-4-(4-methyl-1-piperazinyl)phenyl)amino)-7(8*H*)pteridinone(Compound 2) (Comparative)

EP 16802532.8 – 8 November 2021




orange-red solids, yield 40%. ^1H NMR (400 MHz, $\text{DMSO}-d_6$): δ 8.94 (s, 1H), 8.72-8.71 (m, 1H), 7.86-7.84 (m, 1H), 7.33 (t, $J = 8.4$ Hz, 1H), 6.56 (s, 1H), 6.42-6.40 (m, 1H), 6.16 (ddd, $J = 16.8, 10.4, 2.4$ Hz, 1H), 5.70 (dd, $J = 10.4, 2.4$ Hz, 1H), 5.65 (dd, $J = 10.4, 2.4$ Hz, 1H), 4.10 (t, $J = 10.2$ Hz, 0.5H), 3.88-3.83 (m, 0.6H), 3.76 (s, 3H), 3.67-3.62 (m, 1H), 3.58-3.55 (m, 1H), 3.15-3.10 (m, 4H), 2.82-2.64 (m, 1H), 2.45 (br, 4H), 2.23 (s, 3H), 2.06-1.96 (m, 1H). HRMS (ESI) (m/z): $[\text{M} + \text{H}]^+$ calcd for $\text{C}_{25}\text{H}_{31}\text{N}_8\text{O}_3$, 491.2519; found, 491.2473.

(S)-8-(1-acryloyl-3-piperidiny)-2-((2-methoxy)-4-(4-methyl-1-piperazinyl)phenyl)amino-7(8H)pteridinone (Compound 3) (Comparative)



orange-red solids, yield 38%. ^1H NMR (400 MHz, $\text{DMSO}-d_6$): δ 9.14 (br, 1H), 8.71 (s, 1H), 7.83 (s, 1H), 7.22-7.21 (m, 1H), 6.80-6.62 (m, 2H), 6.48-6.47 (m, 1H), 6.15-6.08 (m, 1H), 5.71-5.60 (m, 1H), 4.82-4.76 (m, 0.6H), 4.34-4.24 (m, 1H), 3.95-3.92 (m, 1H), 3.74 (s, 3H), 3.14 (br, 4H), 2.46 (br, 4H), 1.64 (br, 2H), 1.34-1.30 (m, 1H). HRMS (ESI) (m/z): $[\text{M} + \text{H}]^+$ calcd for $\text{C}_{26}\text{H}_{33}\text{N}_8\text{O}_3$, 505.2676; found, 505.2676.

N-(3-(2-((3-methoxy-4-(methyl-1-piperazinyl)phenyl)amino)-7-oxo-8(7H)-pteridinyl)phenyl)acrylamide (Compound 4)




red solids, yield 55%. ^1H NMR (400 MHz, $\text{DMSO}-d_6$): δ 10.41 (s, 1H), 10.00 (br,

EP 16802532.8 – 8 November 2021


1H), 8.86 (s, 1H), 8.04 (s, 1H), 7.87 (d, $J = 8.0$ Hz, 1H), 7.73 (s, 1H), 7.54 (t, $J = 8.0$ Hz, 1H), 7.11 (d, $J = 8.0$ Hz, 1H), 7.04-6.95 (m, 2H), 6.48-6.42 (m, 2H), 6.27 (dd, $J = 17.2$, 2.0 Hz, 1H), 5.77 (dd, $J = 10.0$, 2.0 Hz, 1H), 3.55 (s, 3H), 2.83 (br, 4H), 2.43 (br, 4H), 2.22 (s, 3H). HRMS (ESI) (m/z): $[M + H]^+$ calcd for $C_{27}H_{29}N_8O_3$, 513.2363; found, 513.2362.

***N*-(3-(2-((3-methoxy-4-(4-methyl-1-piperazinyl)phenyl)amino)-7-oxo)-8(7H)pteridiny)phenyl)acrylamide (Compound 5)**



yellow solids, yield 63%. 1H NMR (400 MHz, DMSO- d_6): δ 10.41 (s, 1H), 10.04 (s, 1H), 8.86 (s, 1H), 8.04 (s, 1H), 7.87 (d, $J = 8.0$ Hz, 1H), 7.76 (s, 1H), 7.55 (t, $J = 8.0$ Hz, 1H), 7.22 (s, 1H), 7.13-7.10 (m, 2H), 6.70-6.69 (m, 1H), 6.45 (dd, $J = 17.0$, 10.2 Hz, 1H), 6.26 (dd, $J = 17.0$, 2.0 Hz, 1H), 5.76 (dd, $J = 10.2$, 2.0 Hz, 1H), 2.70 (br, 4H), 2.44 (br, 4H), 2.23 (s, 3H), 1.97 (s, 3H). HRMS (ESI) (m/z): $[M + H]^+$ calcd for $C_{27}H_{29}N_8O_2$, 497.2413; found, 497.2428.


***N*-(3-(2-((2-methoxy-5-methyl-4-(4-methyl-1-piperazinyl)phenyl)amino)-7-oxo)-8(7H)-pteridiny)phenyl)acrylamide (Compound 6) (Comparative)**




red solids, yield 50%. 1H NMR (400 MHz, DMSO- d_6): δ 10.38 (s, 1H), 8.82 (s, 1H), 8.37 (s, 1H), 8.04 (s, 1H), 7.82 (d, $J = 8.0$ Hz, 1H), 7.74 (s, 1H), 7.53 (t, $J = 8.0$ Hz, 1H), 7.29 (s, 1H), 7.10 (d, $J = 8.8$ Hz, 1H), 6.62 (s, 1H), 6.45 (dd, $J = 16.8$, 2.0 Hz, 1H), 6.25 (dd, $J = 16.8$, 2.0 Hz, 1H), 5.77-5.76 (m, 1H), 3.78 (s, 1H), 2.76 (br, 4H), 2.46 (br, 4H), 2.24 (s, 3H), 1.85 (s, 3H). HRMS (ESI) (m/z): $[M + H]^+$ calcd for $C_{28}H_{31}N_8O_3$, 527.2519; found, 527.2518.

***N*-(3-(2-((4-(4-acetyl-1-piperazinyl)-2-methoxyphenyl)amino)-7-oxo)-8(7H)-pteridiny)phenyl)acrylamide (Compound 7) (Comparative)**

EP 16802532.8 – 8 November 2021

dinyl)phenyl)acrylamide (Compound 7) (Comparative)


red solids, yield 50%. ^1H NMR (400 MHz, DMSO- d_6): δ 10.36 (s, 1H), 8.80 (s, 1H), 8.44 (s, 1H), 8.03 (s, 1H), 7.86 (d, J = 8.0 Hz, 1H), 7.69 (t, J = 2.0 Hz, 1H), 7.52 (t, J = 8.0 Hz, 1H), 7.32 (d, J = 8.8 Hz, 1H), 7.09 (d, J = 8.0 Hz, 1H), 6.57 (d, J = 2.4 Hz, 1H), 6.45 (dd, J = 17.0, 2.0 Hz, 1H), 6.27 (dd, J = 17.0, 10.0 Hz, 1H), 6.09 (br, 1H), 5.77 (dd, J = 10.0, 2.0 Hz, 1H), 3.77 (s, 1H), 3.58-3.54 (m, 4H), 3.06-3.00 (m, 4H), 2.05 (s, 3H). HRMS (ESI) (m/z): $[\text{M} + \text{H}]^+$ calcd for $\text{C}_{28}\text{H}_{29}\text{N}_8\text{O}_4$, 541.2312; found, 541.2312.

 N -(3-(2-((4-(2-(4-methyl-1-piperazinyl)ethyl)phenyl)amino)-7-oxo-8(7H)-pteridinyl)phenyl)acrylamide (Compound 8) (Comparative)

yellow solids, yield 50%. ^1H NMR (400 MHz, DMSO- d_6): δ 10.41 (s, 1H), 10.13 (s, 1H), 8.88 (s, 1H), 8.07 (s, 1H), 7.88 (d, J = 8.0 Hz, 1H), 7.77 (s, 1H), 7.57 (t, J = 8.0 Hz, 1H), 7.31-7.30 (m, 2H), 7.14 (d, J = 8.0 Hz, 1H), 6.86 (br, 2H), 6.45 (dd, J = 16.8, 10.0 Hz, 1H), 6.26 (dd, J = 16.8, 2.0 Hz, 1H), 5.77 (dd, J = 10.0, 2.0 Hz, 1H), 2.56 (t, J = 7.6 Hz, 2H), 2.39-2.31 (m, 9H), 2.15 (s, 3H). HRMS (ESI) (m/z): $[\text{M} + \text{H}]^+$ calcd for $\text{C}_{28}\text{H}_{31}\text{N}_8\text{O}_2$, 511.2570; found, 511.2571.


 N -(3-(2-((2-(methoxymethoxy)-4-(4-methyl-1-piperazinyl)phenyl)amino)-7-oxo-8(7H)-pteridinyl)phenyl)acrylamide (Compound 9) (Comparative)

EP 16802532.8 – 8 November 2021




orange solids, yield 69%. mp 187.3-188.1 °C. ¹H NMR (400 MHz, DMSO-*d*₆): δ 10.40 (s, 1H), 8.81 (s, 1H), 8.53 (br, 1H), 8.03 (s, 1H), 7.85 (d, *J*=6.4 Hz, 1H), 7.72 (s, 1H), 7.53 (t, *J*=8.0 Hz, 1H), 7.34-7.33 (m, 1H), 7.10 (d, *J*=8.0 Hz, 1H), 6.64 (s, 1H), 6.46 (dd, *J* = 16.8, 10.0 Hz, 1H), 6.27 (dd, *J* = 16.8, 1.6 Hz, 1H), 6.13-6.09 (m, 1H), 5.78 (dd, *J* = 10.0, 1.6 Hz, 1H), 5.16 (s, 2H), 3.35 (s, 3H), 3.00 (br, 4H), 2.43 (t, *J*=4.4 Hz, 4H), 2.22 (s, 3H). HRMS (ESI) (*m/z*): [*M* + *H*]⁺calcd for C₂₈H₃₁N₈O₄, 543.2468; found, 543.2476.

***N*-(3-(2-((2-methoxyethoxy)-4-(4-methyl-1-piperazinyl)phenyl)amino)-7-oxo-8(*H*)-pteridin-2-yl)phenylacrylamide (Compound 10) (Comparative)**



orange-red solids, yield 61%. ¹H NMR (400 MHz, CDCl₃): δ 8.78 (s, 1H), 8.65-8.62 (m, 1H), 8.24 (s, 1H), 8.07 (s, 1H), 7.85 (s, 1H), 7.41 (br, 3H), 6.93 (d, *J* = 5.6 Hz, 1H), 6.44 (s, 1H), 6.33 (d, *J* = 17.0 Hz, 1H), 6.10 (dd, *J* = 17.0, 10.4 Hz, 1H), 5.78 (d, *J* = 10.4 Hz, 1H), 4.10 (br, 2H), 3.69 (br, 2H), 3.43 (s, 3H), 3.08 (br, 4H), 2.54 (br, 4H), 2.34 (s, 3H). HRMS (ESI) (*m/z*): [*M* + *H*]⁺calcd for C₂₉H₃₃N₈O₄, 557.2625; found, 557.2621.

***N*-(3-(2-((4-(4-(2-hydroxyethyl)-1-piperazinyl)-2-methoxyphenyl)amino)-7-oxo-8(*H*)-pteridin-2-yl)phenylacrylamide (Compound 11) (Comparative)**




orange solids, yield 53%. ¹H NMR (400 MHz, DMSO-*d*₆): δ 10.39(s, 1H), 8.79 (s,

EP 16802532.8 – 8 November 2021


1H), 8.43 (s, 1H), 8.02 (s, 1H), 7.85 (d, $J = 7.6$ Hz, 1H), 7.71 (s, 1H), 7.52 (t, $J = 8.0$ Hz, 1H), 7.30 (d, $J = 7.6$ Hz, 1H), 7.09 (d, $J = 8.0$ Hz, 1H), 6.52 (s, 1H), 6.45 (dd, $J = 16.8$, 10.0 Hz, 1H), 6.27 (dd, $J = 16.8$, 2.0 Hz, 1H), 6.02 (br, 1H), 5.78 (dd, $J = 10.0$, 2.0 Hz, 1H), 4.43 (t, $J = 5.2$ Hz, 1H), 3.76 (s, 3H), 3.56-3.52 (m, 2H), 3.04 (br, 4H), 2.53 (t, $J =$
5 4.8 Hz, 4H), 2.44 (t, $J = 6.0$ Hz, 1H). HRMS (ESI) (m/z): $[M + H]^+$ calcd for $C_{29}H_{33}N_8O_3$, 543.2468; found, 543.2466.

***N*-(3-(2-((4-(2-(4-methyl-1-piperazinyl)ethyl)phenyl)amino)-7-oxo-8(7H)-pteridinyl)phenyl)acrylamide(Compound 12)**



10 red solids, yield 50%. 1H NMR (400 MHz, DMSO- d_6): δ 10.41 (s, 1H), 8.80 (s, 1H), 8.43 (s, 1H), 8.03 (s, 1H), 7.88-7.87 (m, 1H), 7.70 (s, 1H), 7.52 (t, $J = 8.0$ Hz, 1H), 7.29 (d, $J = 8.0$ Hz, 1H), 7.09 (d, $J = 8.0$ Hz, 1H), 6.52 (s, 1H), 6.46 (dd, $J = 17.2$, 10.0 Hz, 1H), 6.27 (dd, $J = 17.2$, 2.0 Hz, 1H), 6.04 (br, 1H), 5.78 (dd, $J = 10.0$, 2.0 Hz, 1H), 3.76 (s, 3H), 3.60-3.58 (m, 2H), 2.58 (t, $J = 10.8$ Hz, 2H), 2.20-2.18 (m, 7H), 1.82-1.79 (d, $J =$
15 12 Hz, 2H), 1.48-1.40 (m, 2H). HRMS (ESI) (m/z): $[M + H]^+$ calcd for $C_{29}H_{33}N_8O_3$, 541.2676; found, 541.2674.


***N*-(3-(6-isopropyl-2-((4-(4-methyl-1-piperazinyl)phenyl)amino)-7-oxo-8(7H)-pteridinyl)phenyl)acrylamide(Compound 13) (Comparative)**



20 Red solids 163 mg, yield 75%. 1H NMR (400 MHz, DMSO- d_6): δ 10.42 (s, 1H), 9.85 (br, 1H), 8.79 (s, 1H), 7.91 (d, $J = 6.4$ Hz, 1H), 7.71 (s, 1H), 7.55 (t, $J = 8.0$ Hz, 1H), 7.25 (s, 2H), 7.11 (d, $J = 8.0$ Hz, 1H), 6.59-6.58 (m, 2H), 6.45 (dd, $J = 17.0$, 10.2 Hz, 1H), 6.26 (dd, $J = 17.0$, 2.0 Hz, 1H), 5.78 (dd, $J = 10.2$, 2.0 Hz, 1H), 3.45-3.38 (m, 1H), 2.97 (br, 4H), 2.42 (br, 4H), 2.21 (s, 3H), 1.25 (d, $J = 6.8$ Hz, 6H). HRMS (ESI) (m/z): $[M +$
25 $H]^+$ calcd for $C_{29}H_{33}N_8O_2$, 525.2726; found, 525.2728.


EP 16802532.8 – 8 November 2021

***N*-(3-(2-((2-methoxy-4-(4-methyl-1-piperazinyl)phenyl)amino)-7-oxo-6-phenyl-8(7*H*)-pteridiny)phenyl)acrylamide(Compound 14)**



red solids, yield 75%. ¹H NMR (400 MHz, DMSO-*d*₆): δ 10.41 (s, 1H), 8.87 (s, 1H), 8.41 (s, 1H), 8.21-8.19 (m, 2H), 7.89-7.88 (m, 1H), 7.75 (t, *J* = 2.0 Hz, 1H), 7.54 (t, *J* = 8.0 Hz, 1H), 7.50-7.48 (m, 3H), 7.35 (d, *J* = 8.8 Hz, 1H), 7.17-7.14 (m, 1H), 6.54 (d, *J* = 2.0 Hz, 1H), 6.46 (dd, *J* = 17.0, 10.2 Hz, 1H), 6.27 (dd, *J* = 17.0, 2.0 Hz, 1H), 6.05 (br, 1H), 5.78 (dd, *J* = 10.2, 2.0 Hz, 1H), 3.78 (s, 3H), 3.05 (br, 4H), 2.45 (br, 4H), 2.23 (s, 3H). HRMS (ESI) (*m/z*): [*M* + *H*]⁺ calcd for C₃₃H₃₃N₈O₃, 589.2676; found, 589.2676.


8-(1-acryloyl-4-piperidiny)l-2-((2-methoxy-4-(4-methyl-1-piperazinyl)phenyl)amino)-7(8*H*)-pteridinone(Compound 15) (Comparative)



orange solids, yield 53%. ¹H NMR (400 MHz, DMSO-*d*₆): δ 8.83 (s, 1H), 8.71 (s, 1H), 7.83 (s, 1H), 7.53 (br, 1H), 6.85 (dd, *J* = 17.2, 10.4 Hz, 1H), 6.61 (d, *J* = 2.4 Hz, 1H), 6.47 (d, *J* = 8.0 Hz, 1H), 6.13 (dd, *J* = 17.2, 2.4 Hz, 1H), 5.70 (dd, *J* = 10.4, 2.4 Hz, 1H), 5.29 (br, 1H), 4.60 (d, *J* = 10.4 Hz, 1H), 4.20 (d, *J* = 12.8 Hz, 1H), 3.80 (s, 3H), 3.13 (t, *J* = 4.8 Hz, 4H), 2.59-2.53 (m, 3H), 2.45 (t, *J* = 4.8 Hz, 4H), 2.22 (s, 3H), 1.67 (d, *J* = 10.0 Hz, 2H), 1.23 (s, 1H).


(*R*)-8-((1-acryloyl-3-piperidiny)methyl)-2-((2-methoxy-4-(4-methyl-1-piperazinyl)phenyl)amino)-7(8*H*)-pteridinone(Compound 16) (Comparative)

EP 16802532.8 – 8 November 2021




orange solids, yield 61%. ^1H NMR (400 MHz, $\text{DMSO}-d_6$): δ 8.92 (s, 1H), 8.74 (s, 1H), 7.91-7.89 (m, 1H), 7.50-7.44 (m, 1H), 6.75 (dd, $J = 16.8, 10.4$ Hz, 1H), 6.64 (s, 1H), 6.56-6.51 (m, 1H), 6.02 (d, $J = 16.8$ Hz, 1H), 5.64-5.54 (m, 1H), 4.13-4.10 (m, 1H), 3.94-3.83 (m, 3H), 3.76 (s, 3H), 3.16 (br, 4H), 3.03-2.98 (m, 1H), 2.84-2.74 (m, 1H), 2.47 (t, $J = 4.8$ Hz, 4H), 1.92-1.91 (m, 1H), 1.64-1.57 (m, 2H), 1.30-1.23 (m, 3H).

(S)-8-((1-acryloyl-3-piperidinyl)methyl)-2-((2-methoxy-4-(4-methyl-1-piperazinyl)phenyl)amino)-7(8H)-pteridinone(Compound 17) (Comparative)



orange solids, yield 54%. ^1H NMR (400 MHz, $\text{DMSO}-d_6$): δ 8.92 (s, 1H), 8.74 (s, 1H), 7.91-7.89 (m, 1H), 7.49-7.43 (m, 1H), 6.75 (dd, $J = 16.4, 10.4$ Hz, 1H), 6.64 (s, 1H), 6.56-6.51 (m, 1H), 6.02 (d, $J = 16.4$ Hz, 1H), 5.64-5.54 (m, 1H), 4.13-4.08 (m, 1H), 3.94-3.86 (m, 3H), 3.76 (s, 3H), 3.16 (br, 4H), 3.03-2.97 (m, 1H), 2.84-2.74 (m, 1H), 2.47 (t, $J = 4.8$ Hz, 4H), 1.92 (br, 1H), 1.64-1.57 (m, 2H), 1.30-1.20 (m, 3H).

N-(3-(2-((4-((2-(dimethylamino)ethyl)(methyl)amino)-2-methoxyphenyl)amino)-7-oxo-8(7H)-pteridinyl)phenyl)acrylamide(Compound 18) (Comparative)




red solids, yield 61%. ^1H NMR (400 MHz, $\text{DMSO}-d_6$): δ 10.37(s, 1H), 8.76 (s, 1H), 8.41 (br, 1H), 7.99 (s, 1H), 7.84-7.83 (m, 1H), 7.69 (t, $J = 2.0$ Hz, 1H), 7.51 (t, $J = 8.0$ Hz,

EP 16802532.8 – 8 November 2021


1H), 7.26-7.19 (m, 1H), 7.09 (d, $J = 8.0$ Hz, 1H), 6.45 (dd, $J = 16.8, 10.4$ Hz, 1H), 6.29-6.24 (m, 2H), 5.77 (dd, $J = 10.4, 2.0$ Hz, 1H), 3.74 (s, 3H), 2.84 (s, 3H), 2.35-2.33 (m, 2H), 2.19 (s, 6H).

5 ***N*-(3-(2-((4-((2-(dimethylamino)ethyl)(methyl)amino)-2-methoxyphenyl)amino)-6-isopropyl-7-oxo-8(7*H*)-pteridinyl)phenyl)acrylamide(Compound 19) (Comparative)**



orange solids, yield 62%. ^1H NMR (400 MHz, $\text{DMSO-}d_6$): δ 10.37 (s, 1H), 8.72 (s, 1H), 8.21 (s, 1H), 7.84 (d, $J = 8.0$ Hz, 1H), 7.69 (s, 1H), 7.51 (t, $J = 8.0$ Hz, 1H), 7.26 (d, $J = 7.6$ Hz, 1H), 7.09 (d, $J = 8.4$ Hz, 1H), 6.45 (dd, $J = 16.8, 10.4$ Hz, 1H), 6.29-6.24 (m, 2H), 5.87 (br, 1H), 5.76 (dd, $J = 10.4, 2.0$ Hz, 1H), 3.75 (s, 3H), 3.43-3.38 (m, 1H), 2.84 (s, 3H), 2.34 (t, $J = 6.8$ Hz, 2H), 2.19 (s, 6H), 1.24 (d, $J = 6.8$ Hz, 6H).


***N*-(3-(2-((4-((2-(dimethylamino)ethyl)(methyl)amino)-2-methoxyphenyl)amino)-7-oxo-6-phenyl-8(7*H*)-pteridinyl)phenyl)acrylamide(Compound 20)**



15 red solids, yield 64%. ^1H NMR (400 MHz, $\text{DMSO-}d_6$): δ 10.39 (s, 1H), 8.83 (s, 1H), 8.38 (br, 1H), 8.21-8.18 (m, 2H), 7.87 (d, $J = 7.2$ Hz, 1H), 7.75 (s, 1H), 7.53 (t, $J = 8.0$ Hz, 1H), 7.49-7.47 (m, 3H), 7.28 (br, 1H), 7.15 (d, $J = 8.4$ Hz, 1H), 6.46 (dd, $J = 16.8, 10.0$ Hz, 1H), 6.29-6.25 (m, 2H), 5.83 (br, 1H), 5.76 (dd, $J = 10.0, 2.0$ Hz, 1H), 3.76 (s, 3H), 3.35 (br, 2H), 2.85 (s, 3H), 2.35 (t, $J = 5.6$ Hz, 2H), 2.20 (s, 6H).


20 ***(R)*-8-((1-acryloyl-3-piperidinyl)methyl)-6-isopropyl-2-((2-methoxy-4-(4-methyl-1-piperazinyl)phenyl)amino)-7(8*H*)-pteridinone(Compound 21) (Comparative)**

EP 16802532.8 – 8 November 2021




yellow solids, yield 55%. ^1H NMR (400 MHz, $\text{DMSO}-d_6$): δ 8.70(br, 2H), 7.57-7.52 (m, 1H), 6.76 (dd, $J = 16.4, 10.4\text{Hz}$, 1H), 6.64 (s, 1H), 6.57-6.51 (m, 1H), 6.05-5.98 (m, 1H), 5.64-5.53 (m, 1H), 4.15-3.96 (m, 3H), 3.90-3.87 (m, 1H), 3.78 (s, 3H), 3.16 (br, 4H), 3.04-2.75 (m, 2H), 2.47 (t, $J = 3.6\text{ Hz}$, 4H), 2.24 (s, 3H), 1.94 (br, 1H), 1.66-1.61 (m, 2H), 1.20 (d, $J = 6.8\text{ Hz}$, 6H).

(R)-8-(1-acryloyl-3-piperidinyl)-2-((2-methoxy)-4-(4-methyl-1-piperazinyl)phenyl)amino-7(8H)pteridinone (Compound 22) (Comparative)



orange solids, yield 46%. ^1H NMR (400 MHz, $\text{DMSO}-d_6$): δ 9.17 (s, 1H), 8.71 (s, 1H), 7.83 (s, 1H), 7.20 (br, 1H), 6.81 (br, 1H), 6.68-6.62 (m, 2H), 6.47 (br, 1H), 6.14-6.08 (m, 1H), 5.72-5.61 (m, 1H), 4.78-4.72 (m, 1H), 4.35-4.24 (m, 1H), 3.95-3.82 (m, 1H), 3.73 (s, 3H), 3.14 (br, 4H), 2.46 (br, 4H), 2.23 (s, 3H), 1.62 (br, 2H), 1.34-1.26 (m, 1H).

N-(2-((2-(dimethylamino)ethyl)(methyl)amino)-5-((6-isopropyl-8-methyl)-7-oxo-7,8-dihydrogen-2-pteridinyl)amino)-4-methoxyphenyl)acrylamide (Compound 23) (Comparative)




yellow solids, yield 43%. ^1H NMR (400 MHz, $\text{DMSO}-d_6$): δ 10.08 (s, 1H), 8.96 (s,

EP 16802532.8 – 8 November 2021


1H), 8.73 (s, 1H), 8.64 (s, 1H), 7.02 (s, 1H), 6.42 (dd, $J = 16.4, 10.0$ Hz, 1H), 6.24 (d, $J = 16.4$ Hz, 1H), 5.75 (d, $J = 10.4$ Hz, 1H), 3.85 (s, 3H), 3.59 (s, 3H), 3.43-3.36 (m, 1H), 2.89 (br, 2H), 2.71 (s, 3H), 2.35 (br, 2H), 2.23 (s, 6H), 1.20 (d, $J = 6.8$ Hz, 6H).

(S)-8-(1-acryloyl-3-pyrrolidinyl)-6-isopropyl-2-((3-methyl-4-(4-methyl-1-piperazinyl)phenyl)amino)-7(8H)-pteridinone(Compound 24) (Comparative)



yellow solids, yield 81%. ^1H NMR (400 MHz, DMSO- d_6): δ 9.90 (s, 1H), 8.77 (s, 0.6H), 8.76 (s, 0.4H), 7.49-7.41 (m, 2H), 6.93 (t, $J = 7.2$ Hz, 1H), 6.73-6.47 (m, 1H), 6.23-6.15 (m, 1H), 6.03-5.94 (m, 1H), 5.76-5.65 (m, 1H), 4.22 (t, $J = 8.8$ Hz, 0.6H), 4.02-3.86 (m, 1.7H), 3.83-3.65 (m, 1.7H), 3.47-3.40 (m, 1H), 2.92-2.85 (m, 1H), 2.78 (br, 4H), 2.46 (br, 4H), 2.23 (s, 3H), 2.20 (s, 3H), 2.14-2.07 (m, 1H), 1.20 (d, $J = 6.4$ Hz, 6H). HRMS (ESI) (m/z): $[\text{M} + \text{H}]^+$ calcd for $\text{C}_{28}\text{H}_{37}\text{N}_8\text{O}_2$ $[\text{M} + \text{H}]^+$ 517.3039; found, 517.3038.


(S)-8-(1-acryloyl-3-pyrrolidinyl)-2-((3-methyl-4-(4-methyl-1-piperazinyl)phenyl)amino)-6-phenyl-7(8H)-pteridinone(Compound 25)




orange solids, yield 77%. ^1H NMR (400 MHz, DMSO- d_6): δ 10.09 (s, 1H), 8.88 (s, 1H), 8.13 (br, 2H), 7.52-7.49 (m, 5H), 6.98-6.96 (m, 1H), 6.74-6.48 (m, 1H), 6.23-6.15 (m, 1H), 6.13-6.02 (m, 1H), 5.73-5.65 (m, 1H), 4.28-4.24 (m, 0.5H), 4.12-3.90 (m, 1.7H), 3.85-3.67 (m, 1.8H), 3.52-3.44 (m, 1H), 2.94-2.89 (m, 1H), 2.82 (br, 4H), 2.28 (s, 3H), 2.22 (s, 3H). HRMS (ESI) (m/z): $[\text{M} + \text{H}]^+$ calcd for $\text{C}_{31}\text{H}_{35}\text{N}_8\text{O}_2$ $[\text{M} + \text{H}]^+$ 551.2883; found, 551.2880.

(S)-8-(1-acryloyl-3-pyrrolidinyl)-6-isopropyl-2-((2-methoxy-4-(4-methyl-1-piperazinyl)phenyl)amino)-7(8H)-pteridinone(Compound 26) (Comparative)

EP 16802532.8 – 8 November 2021

azinyl)phenyl)amino)-7(8*H*)-pteridinone(Compound 26) (Comparative)


orange solids, yield 54%. ¹H NMR (400 MHz, DMSO-*d*₆): δ 8.71 (s, 1H), 8.69 (s, 0.5H), 8.68 (s, 0.5H), 7.39 (t, *J* = 7.6 Hz, 1H), 6.57 (s, 1H), 6.41 (d, *J* = 8.8 Hz, 1H), 6.17 (dd, *J* = 16.4 Hz, 10.8 Hz, 1H), 5.86-5.80 (m, 1H), 5.72-5.63 (m, 1H), 4.12 (t, *J* = 8.8 Hz, 0.6H), 3.91-3.86 (m, 0.6H), 3.77 (s, 3H), 3.69-3.64 (m, 1H), 3.59 (br, 1H), 3.40-3.38 (m, 0.8H), 3.12 (br, 4H), 2.81-2.70 (m, 1H), 2.45 (br, 4H), 2.23 (s, 3H), 2.09-1.98 (m, 1H), 1.18 (d, *J* = 6.4 Hz, 6H). HRMS (ESI) (*m/z*): [*M* + *H*]⁺ calcd for 533.2989; found, 533.2994.

(*S*)-8-(1-acryloyl-3-pyrrolidinyl)-2-((2-methoxy-4-(4-methyl-1-piperazinyl)phenyl)amino)-6-phenyl-7(8*H*)-pteridinone(Compound 27)

red solids, yield 63%. ¹H NMR (400 MHz, DMSO-*d*₆): δ 8.90 (s, 1H), 8.79 (s, 1H), 8.10 (br, 2H), 7.46 (br, 3H), 7.41-3.39 (m, 1H), 6.58-6.42 (m, 3H), 6.17 (dd, *J* = 16.4 Hz, 10.4 Hz, 1H), 5.98-5.82 (m, 1H), 5.72-5.64 (m, 1H), 4.16 (t, *J* = 8.8 Hz, 0.6H), 3.95-3.90 (m, 0.6H), 3.78 (s, 3H), 3.71 (t, *J* = 9.6 Hz, 0.8H), 3.60 (br, 1H), 3.40-3.34 (m, 1H), 3.13 (br, 4H), 2.83-2.71 (m, 1H), 2.45 (br, 4H), 2.23 (s, 3H), 2.12-2.02 (m, 1H). HRMS (ESI) (*m/z*): [*M* + *H*]⁺ calcd for C₃₁H₃₅N₈O₃ [*M* + *H*]⁺ 567.2832; found, 567.2831.


(*S,E*)-8-(1-(4-(dimethylamino)-2-butenoyl)-3-pyrrolidinyl)-6-isopropyl-2-((3-methyl-4-(4-methyl-1-piperazinyl)phenyl)amino)-7(8*H*)-pteridinone (Compound 28) (Comparative)

EP 16802532.8 – 8 November 2021



orange solids, yield 89%. ^1H NMR (400 MHz, CDCl_3) δ 8.75 (s, 1H), 8.00 (s, 1H), 7.73 (s, 0.6H), 7.65 (s, 0.4H), 7.02-6.89 (m, 1H), 6.55-6.50 (m, 2H), 6.42-6.20 (m, 1H), 6.17-6.07 (m, 1H), 4.48-4.45 (m, 0.5H), 4.18-3.99 (m, 2H), 3.90 (s, 3H), 3.83 (t, $J = 8.8$ Hz, 0.8H), 3.76-3.69 (m, 0.6H), 3.65-3.56 (m, 0.7H), 3.52-3.44 (m, 1H), 3.22-3.18 (m, 4H), 3.15 (t, $J = 6.0$ Hz, 1H), 3.04 (d, $J = 5.6$ Hz, 2H), 2.62-2.60 (br, 4H), 2.38 (s, 3H), 2.30 (s, 3H), 2.21 (s, 3H), 2.14-2.10 (m, 1H), 1.99 (s, 2H), 1.26 (t, $J = 6.0$ Hz, 6H).


(S,E)-8-(1-(4-(dimethylamino)-2-butenoyl)-3-pyrrolidinyl)-2-((3-methyl-4-(4-methyl-1-piperazinyl)phenyl)amino)-6-phenyl-7(8H)-pteridinone (Compound 29)



orange solids, yield 82%. ^1H NMR (400 MHz, $\text{DMSO}-d_6$): δ 10.10 (s, 1H), 8.88 (s, 1H), 8.12 (s, 2H), 7.49 (br, 5H), 6.97 (d, $J = 8.4$ Hz, 1H), 6.72-6.61 (m, 1H), 6.54-6.31 (m, 1H), 6.10-6.00 (m, 1H), 4.28-4.24 (m, 0.6H), 4.05-4.00 (m, 0.7H), 3.93-3.86 (m, 1.5H), 3.81-3.76 (m, 1.5H), 3.73-3.67 (m, 1H), 3.55 (br, 4H), 3.18 (d, $J = 6.8$ Hz, 2H), 3.08 (d, $J = 5.2$ Hz, 1H), 2.83 (br, 4H), 2.57 (s, 3H), 2.32 (s, 1.5H), 2.30 (s, 1.5H), 2.57 (s, 3H), 2.22 (s, 3H), 2.17 (s, 3H). HRMS (ESI) (m/z): $[\text{M} + \text{H}]^+$ calcd for $\text{C}_{34}\text{H}_{42}\text{N}_9\text{O}_2$ $[\text{M} + \text{H}]^+$ 608.3461; found, 608.3466.


(E)-8-(1-(4-(dimethylamino)-2-butenoyl)-4-piperidinyl)-2-((2-methoxy-4-(4-methyl-1-piperazinyl)phenyl)amino)-7(8H)-pteridinone (Compound 30) (Comparative)

EP 16802532.8 – 8 November 2021



Red solids (yield 71%). mp 157.3-157,6 °C. ¹H NMR (400 MHz, DMSO-*d*₆): δ 8.86 (s, 1H), 8.72 (s, 1H), 7.83 (s, 1H), 7.54 (br, 1H), 6.64-6.62(m, 3H), 6.47 (d, *J* = 7.6 Hz, 1H), 5.27 (br, 1H), 4.61-4.58 (m, 1H), 4.21-4.18 (m, 1H), 3.79 (s, 3H), 3.14(br, 4H),
 5 3.05-3.04 (br, 2H), 2.57 (br, 4H), 2.46 (t, *J* = 4.0 Hz, 4H), 2.23 (s, 3H), 2.16 (s, 6H), 1.67 (br, 2H). HRMS (ESI) (*m/z*): [M + H]⁺ calcd for C₂₉H₄₀N₉O₃ , 562.3254; found, 562.3259.


(S,E)-8-(1-(4-(dimethylamino)-2-butenoyl)3-pyrrolidinyl)-2-((2-methoxy-4-(4-methyl-1-piperazinyl)phenyl)amino)-6-phenyl-7(8H)pteridinone (Compound 31)



orange solids, yield 89%. mp 253.1-254.0 °C. ¹H NMR (400 MHz, DMSO-*d*₆): δ 9.92 (d, *J* = 6.8 Hz, 1H), 8.80 (d, *J* = 3.2 Hz, 1H), 8.12-8.11 (m, 2H), 7.31-7.67 (m, 1H), 7.47 (m, 3H), 7.42 (t, *J* = 8.0 Hz, 1H), 7.71-6.62 (m, 1H), 6.59 (br, 1H), 6.46-6.42 (m, 1H),
 15 4.19-4.10 (m, 2H), 3.94-3.89 (m, 1H), 3.71 (t, *J* = 10.8 Hz, 1H), 4.60-3.59 (m, 1H), 3.79 (s, 3H), 3.14 (br, 4H), 3.09-3.03 (m, 1H), 3.00 (d, *J* = 5.6, 1H), 2.87-2.74 (m, 1H), 2.47 (br, 4H), 2.24 (s, 3H), 2.19 (s, 3H), 2.13 (s, 3H). HPLC purity: 98.1%, Retention time = 9.63 min. HRMS (ESI) (*m/z*): [M+H]⁺ calcd for C₃₄H₄₂N₉O₃ [M+H]⁺ 624.3411; found, 624.3413.


N-(3-(6-isopropyl-2-((3-methoxy-4-(4-methyl-1-piperazinyl)phenyl)amino)-7-oxo-8(7H)pteridinyl)phenyl)acrylamide(Compound 32) (Comparative)

EP 16802532.8 – 8 November 2021



Yellow solids (yield 69%). mp 249.0-249.5 °C. ¹H NMR (400 MHz, DMSO-*d*₆): δ 10.42 (s, 1H), 9.83 (br, 1H), 8.82 (s, 1H), 7.87 (d, *J* = 8.0 Hz, 1H), 7.73 (s, 1H), 7.53 (t, *J* = 8.0 Hz, 1H), 7.12 (d, *J* = 8.0 Hz, 1H), 7.04 (br, 1H), 6.96 (br, 1H), 6.49-6.42 (m, 2H), 6.26 (dd, *J* = 16.8 Hz, 2.0 Hz, 1H), 5.77 (dd, *J* = 10.0 Hz, 2.0 Hz, 1H), 3.55 (s, 3H), 3.44-3.39 (m, 1H), 2.82 (br, 4H), 2.42 (br, 4H), 2.21 (s, 3H), 1.25 (d, *J* = 6.8 Hz, 6H). HRMS (ESI) (*m/z*): [M + H]⁺ calcd for C₃₀H₃₅N₈O₃, 555.2832; found, 555.2838.


N-(3-(2-((3-methoxy-4-(4-methyl-1-piperazinyl)phenyl)amino)-7-oxo-6-(trifluoromethyl)-8(7H)-pteridinyl)phenyl)acrylamide (Compound 33) (Comparative)



Brown solids (yield 59%). mp >300 °C. ¹H NMR (400 MHz, DMSO-*d*₆): δ 10.44 (s, 1H), 10.40 (s, 1H), 9.00 (s, 1H), 7.88 (d, *J* = 7.2 Hz, 1H), 7.81 (s, 1H), 7.56 (t, *J* = 8.0 Hz, 1H), 7.14 (d, *J* = 8.0 Hz, 1H), 7.01 (d, *J* = 7.2 Hz, 1H), 6.95 (s, 1H), 6.49-6.42 (m, 2H), 6.27 (d, *J* = 16.4 Hz, 1H), 5.77 (d, *J* = 10.4 Hz, 1H), 3.53 (s, 3H), 2.83 (br, 4H), 2.41 (br, 4H), 2.21 (s, 3H). HRMS (ESI) (*m/z*): [M+H]⁺ calcd for C₂₈H₂₈N₈O₃F₃, 581.2236; found, 581.2241.


(S)-8-(1-acryloyl-3-pyrrolidinyl)-2-((3-methyl-4-(4-methyl-1-piperazinyl)phenyl)amino)-7(8H)pteridinone (Compound 34) (Comparative)

EP 16802532.8 – 8 November 2021



Yellow solids (yield 45%). mp 215.5-215.7 °C. ¹H NMR (400 MHz, DMSO-*d*₆): δ 10.08 (s, 1H), 8.81 (d, *J* = 3.2 Hz, 1H), 7.89 (d, *J* = 6.8 Hz, 1H), 7.47-7.42 (m, 2H), 6.93 (t, *J* = 6.8 Hz, 1H), 6.72-6.46 (m, 1H), 6.22-6.6.14 (m, 1H), 5.98-5.88 (m, 1H), 5.75-5.64 (m, 1H), 4.20 (t, *J* = 8.8, 1H), 4.00-3.64 (m, 4H), 3.47-3.40 (m, 1H), 2.78 (br, 4H), 2.45 (s, 3H), 2.23 (s, 3H), 2.20 (br, 4H). HRMS (ESI) (*m/z*): [M + H]⁺ calcd for C₂₅H₃₁N₈O₂, 475.2570; found, 475.2550.


(S)-8-(1-acryloyl-30pyrrolidinyl)-2-((3-methoxy-4(4-methyl-1-piperazinyl)phenyl)amino)-6-phenyl-7(8H)pteridinone (Compound 35)



Orange solids (yield 40%). mp 208.3-208.8 °C. ¹H NMR (400 MHz, DMSO-*d*₆): δ 10.10 (s, 1H), 8.88 (d, *J* = 3.2, 1H), 8.13-8.12 (m, 2H), 7.48 (d, *J* = 2.4Hz, 1H), 7.47 (s, 1H), 7.34-7.27 (m, 2H), 6.81 (t, *J* = 8.4Hz, 1H), 6.74-6.48 (m, 1H), 6.23-6.15 (m, 1H), 6.12-6.04 (m, 1H), 5.76-5.65 (m, 1H), 4.25 (t, *J* = 9.2, 1H), 4.13-3.76 (m, 3H), 3.76 (s, 3H), 3.74-3.67 (m, 1H), 3.51-3.44 (m, 1H), 2.91 (br, 4H), 2.44 (br, 4H), 2.21 (s, 3H). HRMS (ESI) (*m/z*): [M + H]⁺ calcd for C₃₁H₃₅N₈O₃, 567.2832; found, 567.2835.


(S,E)-8-(1-(4-(dimethylamino)-2-butenoyl)3-pyrrolidinyl)-2-((3-methyl-4-(4-methyl-1-piperazinyl)phenyl)amino)-6-isopropyl-7(8H)pteridinone (Compound 36) (Comparative)

EP 16802532.8 – 8 November 2021



Red solids (yield 62%). mp 159.9-160.7 °C. ¹H NMR (400 MHz, DMSO-*d*₆): δ 9.94 (s, 1H), 8.77 (d, *J* = 3.6 Hz, 1H), 7.49-7.40 (m, 2H), 6.94 (d, *J* = 8.4, 1H), 6.72-6.61 (m, 1H), 6.53 (d, *J* = 15.2 Hz, 0.5H), 6.34 (d, *J* = 15.2 Hz, 0.5H), 6.04-5.91 (m, 1H), 4.22 (t, *J* = 8.4 Hz, 1H), 4.00-3.64 (m, 4H), 3.40-3.35 (m, 2H), 3.22 (t, *J* = 6.4 Hz, 1H), 3.10 (d, *J* = 6.0 Hz, 1H), 2.82 (br, 4H), 2.61 (br, 4H), 2.34 (d, 11.2 Hz, 3H), 2.28 (s, 3H), 2.20 (br, 4H), 2.19 (s, 3H), 1.20 (d, *J* = 6.8 Hz, 6H). HRMS (ESI) (*m/z*): [M + H]⁺ calcd for C₃₁H₄₄N₉O₂, 574.3618; found, 574.3584.


10 **N-(3-(6-isopropyl-2-((2-methoxy-4-(4-methyl-1-piperazinyl)phenyl)amino)-7-oxo-8(7H)pteridinyloxy)phenyl)acrylamide (Compound 37) (Comparative)**



Orange solids (yield 65%). mp 242.9-243.9 °C. ¹H NMR (400 MHz, DMSO-*d*₆): δ 10.42 (s, 1H), 8.75 (s, 1H), 8.26 (s, 1H), 7.87 (d, *J* = 7.2 Hz, 1H), 7.71 (s, 1H), 7.53 (t, *J* = 8.0 Hz, 1H), 7.34 (d, *J* = 8.8 Hz, 1H), 7.10 (d, *J* = 8.8 Hz, 1H), 6.53 (d, *J* = 2.0 Hz, 1H), 6.46 (dd, *J* = 16.8 Hz, 10.0 Hz, 1H), 6.27 (dd, *J* = 16.8 Hz, 2.0 Hz, 1H), 6.12-6.00 (m, 1H), 5.78 (dd, *J* = 10.0 Hz, 2.0 Hz), 3.76 (s, 3H), 3.44-3.41 (m, 1H), 3.04 (br, 4H), 2.43 (t, *J* = 4.8 Hz, 4H), 2.22 (s, 3H), 1.25 (d, *J* = 6.8 Hz, 6H). HRMS (ESI) (*m/z*): [M + H]⁺ calcd for C₃₀H₃₅N₈O₃, 555.2832; found, 555.2821.


20 **N-(3-(2-((2-methoxy-5-methyl-4-(4-methyl-1-piperazinyl)phenyl)amino)-7-oxo-6-phenyl-8(7H)pteridinonyloxy)phenyl)acrylamide (Compound 38)**

EP 16802532.8 – 8 November 2021



Yellow solids (yield 62%). mp 292.9-293.4 °C. ^1H NMR (400 MHz, $\text{DMSO}-d_6$): δ 10.41 (s, 1H), 8.90 (s, 1H), 8.36 (br, 1H), 8.21-8.18 (m, 2H), 7.84 (d, $J = 8.0\text{Hz}$, 1H), 7.80 (s, 1H), 7.55 (t, $J = 8.0\text{Hz}$, 1H), 7.50-7.48 (m, 3H), 7.33 (s, 1H), 7.16 (d, $J = 7.6\text{Hz}$, 1H), 6.63 (s, 1H), 6.45 (dd, $J = 16.8\text{Hz}$, 10.0Hz, 1H), 6.26 (dd, $J = 16.8\text{Hz}$, 1.6Hz, 1H), 5.77 (dd, $J = 10.0\text{Hz}$, 1.6Hz, 1H), 3.79 (s, 3H), 2.77 (br, 4H), 2.48 (br, 4H), 2.25 (s, 3H), 1.85 (br, 3H). HRMS (ESI) (m/z): $[\text{M} + \text{H}]^+$ calcd for $\text{C}_{34}\text{H}_{35}\text{N}_8\text{O}_3$, 603.2832; found, 603.2834.


N-(3-(6-isopropyl-2-((2-methoxy-5-methyl-4-(4-methyl-1-piperazinyl)phenyl)amino)-7-oxo-8(7H)pteridiny)phenyl)acrylamide (Compound 39) (Comparative)



orange solids (yield 60%). mp 227.5-228.5 °C. ^1H NMR (400 MHz, $\text{DMSO}-d_6$): δ 10.40 (s, 1H), 8.79 (s, 1H), 8.22 (s, 1H), 7.82 (d, $J = 8.0\text{Hz}$, 1H), 7.73 (s, 1H), 7.53 (t, $J = 8.0\text{Hz}$, 1H), 7.31 (s, 1H), 7.11 (d, $J = 7.6\text{Hz}$, 1H), 6.62 (s, 1H), 6.44 (dd, $J = 16.8\text{Hz}$, 10.0Hz, 1H), 6.25 (dd, $J = 16.8\text{Hz}$, 0.8Hz, 1H), 5.75 (d, $J = 10.0\text{Hz}$, 1H), 3.78 (s, 3H), 3.43-3.39 (m, 1H), 2.77 (br, 4H), 2.26 (s, 3H), 1.85 (br, 3H), 1.24 (d, $J = 6.8\text{Hz}$, 6H). HRMS (ESI) (m/z): $[\text{M} + \text{H}]^+$ calcd for $\text{C}_{31}\text{H}_{37}\text{N}_8\text{O}_3$, 569.2989; found, 569.2989.


(S)-8-(1-acryloyl-3-pyrrolidinyl)-2-((4-((2-(dimethylamino)ethyl)(methyl)amino)-3-methylphenyl)amino)-6-phenyl-7(8H)pteridinone (Compound 40)

EP 16802532.8 – 8 November 2021



red solids (yield 40%). mp 256.1-256.4 °C. ^1H NMR (400 MHz, $\text{DMSO}-d_6$): δ 10.08 (s, 1H), 8.88 (d, $J = 3.2\text{Hz}$, 1H), 8.13-8.12 (m, 2H), 7.52-7.45 (m, 5H), 7.02 (dd, $J = 8.4\text{Hz}$, 5.6Hz, 1H), 6.74-6.49 (m, 1H), 6.23-6.15 (m, 1H), 6.12-6.02 (m, 1H), 5.75-5.65 (m, 1H), 4.28-4.23 (m, 1H), 4.15-4.12 (m, 1H), 3.94-3.68 (m, 3H), 3.52-3.44 (m, 1H), 2.90 (t, $J = 6.8\text{Hz}$, 2H), 2.61 (s, 3H), 2.35 (t, $J = 6.8\text{Hz}$, 2H), 2.22 (s, 3H), 2.13 (br, 6H). HRMS (ESI) (m/z): $[\text{M} + \text{H}]^+$ calcd for $\text{C}_{31}\text{H}_{37}\text{N}_8\text{O}_2$, 553.3039; found, 553.3031.


(S)-8-(1-acryloyl-3-pyrrolidinyl)-2-((4-((2-(dimethylamino)ethyl)(methyl)amino)-3-methoxyphenyl)amino)-6-phenyl-7(8H)pteridinone (Compound 41)



Dark red solids (yield 49%). mp 260.0-260.2 °C. ^1H NMR (400 MHz, $\text{DMSO}-d_6$): δ 10.09 (s, 1H), 8.88 (d, $J = 2.8\text{Hz}$, 1H), 8.131 (br, 2H), 7.49-7.48 (m, 3H), 7.34-7.25 (m, 2H), 6.81 (t, $J = 7.2\text{Hz}$, 1H), 6.76-6.49 (m, 1H), 6.23-6.05 (m, 2H), 5.75-5.65 (m, 1H), 4.25 (t, $J = 8.4\text{Hz}$, 1H), 4.06-3.82 (m, 3H), 3.77 (s, 3H), 3.74-3.67 (m, 1H), 3.05 (t, $J = 7.2\text{Hz}$, 2H), 2.94-2.79 (m, 1H), 2.68 (s, 3H), 2.36 (t, $J = 7.2\text{Hz}$, 2H), 2.13 (s, 6H). HRMS (ESI) (m/z): $[\text{M} + \text{H}]^+$ calcd for $\text{C}_{31}\text{H}_{37}\text{N}_8\text{O}_3$, 569.2989; found, 569.2988.


N-(3-(2-((3-methoxy-4-(4-methyl-1-piperazinyl)phenyl)amino)-7-oxo-6-phenyl-8(7H)pteridinyl)phenyl)acrylamide (Compound 42)

EP 16802532.8 – 8 November 2021



red solids (yield 81%). mp 264.7-265.5 °C. ¹H NMR (400 MHz, DMSO-*d*₆): δ 10.43 (s, 1H), 10.01 (s, 1H), 8.93 (s, 1H), 8.22-8.19 (m, 2H), 7.90 (d, *J* = 7.6Hz, 1H), 7.79 (s, 1H), 7.56 (t, *J* = 8.0Hz, 1H), 7.49 (t, *J* = 3.2Hz, 1H), 7.17 (d, *J* = 8.4Hz, 1H), 7.07 (br, 1H), 6.97 (br, 1H), 6.46 (dd, *J* = 16.8Hz, 10.0Hz, 1H), 6.27 (dd, *J* = 16.8Hz, 1.6Hz, 1H), 5.77 (dd, *J* = 10.0Hz, *J* = 1.6Hz, 1H), 3.56 (s, 3H), 2.83 (br, 4H), 2.41 (br, 4H), 2.21 (s, 3H). HRMS (ESI) (*m/z*): [M + H]⁺ calcd for C₃₃H₃₃N₈O₃, 589.2676; found, 589.2642.


N-(3-(6-cyclohexyl-2-((2-methoxy-4-(4-methyl-1-piperazinyl)phenyl)amino)-7-oxo-8(7H)pteridinyl)phenyl)acrylamide (Compound 43)



yellow solids (yield 82%). mp 268.9-269.4 °C. ¹H NMR (400 MHz, DMSO-*d*₆): δ 10.40 (s, 1H), 8.73 (s, 1H), 8.24 (s, 1H), 7.85 (d, *J* = 8.0Hz, 1H), 7.69 (t, *J* = 1.6Hz, 1H), 7.52 (t, *J* = 8.0Hz, 1H), 7.33 (d, *J* = 9.2Hz, 1H), 7.10-7.08 (m, 1H), 6.53 (d, *J* = 2.4Hz, 1H), 6.45 (dd, *J* = 16.8Hz, 10.0Hz, 1H), 6.26 (dd, *J* = 16.8Hz, 2.0Hz, 1H), 6.06 (br, 1H), 5.77 (dd, *J* = 10.0Hz, 2.0Hz, 1H), 3.76 (s, 3H), 3.03 (br, 4H), 2.43 (t, *J* = 4.8Hz, 4H), 2.22 (s, 3H), 3.41 (d, *J* = 12.0Hz, 2H), 1.82 (d, *J* = 12.0Hz, 2H), 1.72 (d, *J* = 12.0Hz, 2H), 1.53-1.23 (m, 6H). HRMS (ESI) (*m/z*): [M + H]⁺ calcd for C₃₃H₃₉N₈O₃, 595.3145; found, 595.3141.


(S)-8-((1-acryloyl-3-piperidinyl)methyl)-2-((2-methoxy-4-(4-methyl-1-piperazinyl)phenyl)amino)-6-phenyl-7(8H)pteridinone (Compound 44)

EP 16802532.8 – 8 November 2021



red solids (yield 49%). mp 208.3-209.3 °C. ¹H NMR (400 MHz, DMSO-*d*₆): δ 8.88 (s, 1H), 8.81 (s, 1H), 8.18-8.16 (m, 2H), 7.59-7.52 (m, 1H), 7.47 (t, *J* = 3.2Hz, 3H), 6.75 (dd, *J* = 16.4Hz, 10.0, 1H), 6.64 (s, 1H), 6.55 (dd, *J* = 17.2Hz, 8.8Hz, 1H), 6.03 (d, *J* = 16.4Hz, 1H), 5.58 (dd, *J* = 32.8Hz, 10.0Hz, 1H), 4.19-4.06 (m, 3H), 3.89-3.83 (m, 1H), 3.79 (s, 3H), 3.17 (br, 4H), 3.05-2.55 (m, 2H), 2.48 (t, *J* = 4.8Hz, 4H), 2.24 (s, 3H), 2.00 (br, 1H), 1.66 (br, 2H), 1.26-1.23 (m, 2H). HRMS (ESI) (*m/z*): [M + H]⁺ calcd for C₃₃H₃₉N₈O₃, 595.3145; found, 595.3157.


(R)-8-(1-acryloyl-3-piperidinyl)-2-((2-methoxy-4-(4-methyl-1-piperazinyl)phenyl)amino)-6-phenyl-7(8H)pteridinone (Compound 45)



red solids (yield 38%). mp 134.9-135.1 °C. ¹H NMR (400 MHz, DMSO-*d*₆): δ 9.17 (s, 1H), 8.79 (s, 1H), 8.11-8.09 (m, 2H), 7.77-7.46 (m, 3H), 7.24 (br, 1H), 6.84-6.63 (m, 2H), 6.49 (s, 1H), 6.06-6.19 (m, 1H), 5.67 (dd, *J* = 37.2Hz, 9.6Hz, 1H), 5.00-4.27 (m, 3H), 3.75 (s, 3H), 3.14 (s, 4H), 2.49 (s, 4H), 2.24 (s, 3H), 1.69 (br, 2H), 1.35 (br, 1H), 1.23 (s, 1H). HRMS (ESI) (*m/z*): [M + H]⁺ calcd for C₃₂H₃₇N₈O₃, 581.2989; found, 581.2972.


(R)-8-(1-acryloyl-3-piperidinyl)-6-cyclohexyl-2-((2-methoxy-4-(4-methyl-1-piperazinyl)phenyl)amino)-7(8H)pteridinone (Compound 46)

EP 16802532.8 – 8 November 2021



yellow solids (yield 53%). mp 94.2-94.7 °C. ¹H NMR (400 MHz, DMSO-*d*₆): δ 8.96(s, 1H), 8.65(s, 1H), 7.23 (br, 1H), 6.83-6.61(m, 2H), 6.47 (s, 1H), 6.15-6.08 (m, 1H), 5.66 (dd, *J* = 36Hz, 9.6Hz, 1H), 4.84 (br, 1H), 4.29 (br, 1H), 3.95 (br, 1H), 3.72 (s, 3H), 3.13 (br, 4H), 2.46 (br, 4H), 2.23 (s, 3H), 1.81-1.63(m, 7H), 1.44-1.23 (m, 9H). HRMS (ESI) (*m/z*): [M + H]⁺ calcd for C₃₂H₄₃N₈O₃, 587.3458; found 587.3458.


N-(3-(2-((3-methyl-4-(4-methyl-1-piperazinyl)phenyl)amino)-7-oxy-6-phenyl-8(7H)pteridiny)phenyl)acrylamide (Compound 47)



orange solids (yield 68%). mp 265.0-265.2 °C. ¹H NMR (400 MHz, DMSO-*d*₆): δ 10.43(s, 1H), 10.04(s, 1H), 8.92(s, 1H), 8.22-8.19(m, 2H), 7.89(d, *J* = 8.0Hz, 1H), 7.81(s, 1H), 7.57(t, *J* = 8.0Hz, 1H), 7.50-7.49(m, 3H), 7.25(s, 1H), 7.19(s, 1H), 7.17(s, 1H), 6.71(br, 1H), 6.46(dd, *J* = 16.8Hz, 10.4Hz, 1H), 6.26(dd, *J* = 16.8Hz, 1.6Hz, 1H), 5.77(dd, *J* = 10.4Hz, 1.6Hz, 1H), 2.7(br, 4H), 2.44(br, 4H), 2.23(s, 3H), 1.98(s, 3H). HRMS (ESI) (*m/z*): [M + H]⁺ calcd for C₃₃H₃₃N₈O₂, 573.2726; found, 573.2729.


N-(3-(2-((2-methoxy-4-(4-(4-methyl-1-piperazinyl)piperidinyl)phenyl)amino)-7-oxo-6-phenyl-8(7H)pteridiny)phenyl)acrylamide (Compound 48)

EP 16802532.8 – 8 November 2021



red solids (yield 40%). mp>300 °C. ¹H NMR (400 MHz, DMSO-*d*₆): δ 10.44(s, 1H), 8.86(s, 1H), 8.41 (s, 1H), 8.20-8.19(m, 2H), 7.90(br, 1H), 7.75(s, 1H), 7.54(t, *J* = 8.0Hz, 1H), 7.50-7.48(m, 3H), 7.33(d, *J* = 6.4Hz, 1H), 7.15(d, *J* = 8.0Hz, 1H), 6.52(br, 1H),
 5 6.47(dd, *J* = 16.8Hz, 10.0Hz, 1H), 6.27(dd, *J* = 16.8Hz, 1.6Hz, 1H), 6.04(br, 1H), 5.78(dd, *J* = 10.0Hz, 1.6Hz), 3.77(s, 3H), 3.61(d, *J* = 8.4Hz, 2H), 2.60-2.54(m, 5H), 2.39-2.30(m, 5H), 1.81(d, *J* = 11.6Hz, 2H), 1.51-1.43(m, 2H). HRMS (ESI) (*m/z*): [M + H]⁺ calcd for C₃₈H₄₂N₉O₃, 672.3411; found, 672.3407 .


8-(1-acryloyl-3-pyrrolidinyl)-2-((3-methyl-4-(4-methyl-1-piperazinyl)phenyl)amino)-6-phenyl-7(8H)pteridinone (Compound 49)



orange solids (yield 65%). ¹H NMR (400 MHz, DMSO-*d*₆): δ 10.08(s, 1H), 8.87(d, *J* = 2.8Hz, 1H), 8.13-8.11(m, 2H), 7.51-7.46(m, 5H), 6.97-6.93(m, 1H), 6.70(dd, *J* = 16.8Hz, 10.4, 1H), 6.51(dd, *J* = 16.8Hz, 10.4Hz, 1H), 6.23-6.15(m, 1H), 6.11-6.01(m, 1H),
 15 5.75-5.65(m, 1H), 4.25(t, *J* = 8.8Hz, 1H), 4.06-3.68(m, 3H), 3.52-3.45(m, 1H), 2.95-2.86(m, 1H), 2.79(br, 4H), 2.46(br, 4H), 2.23(s, 3H), 2.21(s, 3H), 2.17-2.12(m, 1H). HRMS (ESI) (*m/z*): [M + H]⁺ calcd for C₃₁H₃₅N₈O₂, 551.2883; found, 551.2883 .


(S)-8-(1-acryloyl-3-pyrrolidinyl)-6-cyclohexyl-2-((2-methoxy-4-(4-methyl-1-piperazinyl)phenyl)amino)-7(8H)pteridinone (Compound 50)

EP 16802532.8 – 8 November 2021



orange solids (yield 49%). mp 117.9-118.2 °C. ¹H NMR (400 MHz, DMSO-*d*₆): δ 8.71(s, 1H), 8.66(d, *J* = 3.6Hz, 1H), 7.39(t, *J* = 8.0Hz, 1H), 6.63-6.59(m, 0.5H), 6.57(s, 1H), 6.49-6.45(m, 0.5H), 6.41(d, *J* = 9.2Hz, 1H), 6.20-6.13(m, 1H), 5.82(br, 1H), 5.73-5.63(m, 1H), 4.11(t, *J* = 9.2Hz, 0.5H), 3.87(dd, *J* = 12.0Hz, 8.8Hz, 0.5H), 3.69-3.58(m, 2H), 3.18-3.04(m, 5H), 2.80-2.65(m, 1H), 2.45(br, 4H), 2.23(s, 3H), 2.09 (br, 0.5H), 1.99(br, 0.5H), 1.81(t, *J* = 11.2Hz, 4H), 1.70(d, *J* = 11.2Hz, 1H), 1.45-1.16(m, 6H). HRMS (ESI) (*m/z*): [M + H]⁺ calcd for C₃₁H₄₁N₈O₃, 573.3302; found, 573.3304.


8-((3-aminophenyl)-2-((2-methoxy-4-(4-methyl-1-piperazinyl)phenyl)amino)-6-phenyl-7(8H)pteridin-9(10H)-one)acrylamide (Compound 51)



orange solids (yield 84%). mp 235.5-235.7 °C. ¹H NMR (400 MHz, DMSO-*d*₆): δ 8.84 (s, 1H), 8.39 (br, 1H), 8.19-8.17 (m, 2H), 7.49-7.47 (m, 4H), 7.21 (t, *J* = 8.0Hz, 1H), 6.72 (d, *J* = 7.2Hz, 1H), 6.571-6.56 (m, 2H), 6.52 (d, *J* = 8.0Hz, 1H), 6.19 (br, 1H), 5.34 (s, 2H), 3.79 (s, 3H), 3.08 (br, 4H), 2.45 (t, *J* = 4.4Hz, 4H), 2.23 (s, 3H). HRMS (ESI) (*m/z*): [M + H]⁺ calcd for C₃₀H₃₁N₈O₂, 535.2570; found, 535.2569.


N-(3-(6-(4-fluorophenyl)-2-((2-methoxy-4-(4-methyl-1-piperazinyl)phenyl)amino)-7-oxo-8(7H)pteridin-9(10H)-yl)phenyl)acrylamide (Compound 52)

EP 16802532.8 – 8 November 2021



orange solids (yield 73%). mp 262.3-262.7 °C. ¹H NMR (400 MHz, DMSO-*d*₆): δ 10.43(s, 1H), 8.87(s, 1H), 8.44(s, 1H), 8.29(dd, *J* = 8.4Hz, 6.0Hz, 1H), 7.89(br, 1H), 7.77(s, 1H), 7.55(t, *J* = 8.0Hz, 1H), 7.36-7.31(m, 3H), 7.15(d, *J* = 8.4Hz, 1H), 6.54(s, 1H), 6.47(dd, *J* = 16.8Hz, 10.0Hz, 1H), 6.28(dd, *J* = 16.8Hz, 1.6Hz, 1H), 6.04(br, 1H), 5.78(dd, *J* = 10.0Hz, 1.6Hz, 1H), 3.78(s, 3H), 3.05(br, 4H), 2.44(t, *J* = 4.4Hz, 4H), 2.23(s, 3H). HRMS (ESI) (*m/z*): [M + H]⁺ calcd for C₃₃H₃₂N₈O₃F, 607.2581; found, 607.2589.


(S)-8-((1-acryloyl-3-piperidinyl)methyl)-2-((2-methoxy-4-(4-methyl-1-piperazinyl)phenyl)amino)-6-phenyl-7(8H)pteridinone (Compound 53)



red solids, yield 49%. mp 208.3-209.3 °C. ¹H NMR (400 MHz, DMSO-*d*₆): δ 8.88 (s, 1H), 8.81 (s, 1H), 8.18-8.16 (m, 2H), 7.59-7.52 (m, 1H), 7.47 (t, *J* = 3.2Hz, 3H), 6.75 (dd, *J* = 16.4Hz, 10.0, 1H), 6.64 (s, 1H), 6.55 (dd, *J* = 17.2Hz, 8.8Hz, 1H), 6.03 (d, *J* = 16.4Hz, 1H), 5.58 (m, 1H), 4.19-4.06 (m, 3H), 3.89-3.83 (m, 1H), 3.79 (s, 3H), 3.17 (br, 4H), 3.05-2.55 (m, 2H), 2.48 (t, *J* = 4.8Hz, 4H), 2.24 (s, 3H), 2.00 (br, 1H), 1.66 (br, 2H), 1.26-1.23 (m, 2H). HPLC purity: 95.1%, Retention time = 15.82 min. HRMS (ESI) (*m/z*): [M+H]⁺ calcd for C₃₃H₃₉N₈O₃, 595.3145; found, 595.3157.


N-(3-(2-((2-methoxy-4-(4-methyl-1-piperazinyl)phenyl)amino)-6-(4-methoxyphenyl)-7-oxo-8(7H)pteridinyl)phenyl)acrylamide (Compound 54)

EP 16802532.8 – 8 November 2021



orange solids (yield 65%). mp 296.5-297.3 °C. ¹H NMR (400 MHz, DMSO-*d*₆): δ 10.43(s, 1H), 8.84(s, 1H), 8.34(s, 1H), 8.26(d, *J* = 8.8Hz, 1H), 7.89(d, *J* = 7.2Hz, 1H), 7.74(s, 1H), 7.55(t, *J* = 8.0Hz, 1H), 7.36(d, *J* = 8.4Hz, 1H), 7.15(d, *J* = 8.4Hz, 1H), 7.05(d, *J* = 9.2Hz, 1H), 6.54(s, 1H), 6.47(dd, *J* = 16.8Hz, 10.0Hz, 1H), 6.27(dd, *J* = 16.8Hz, 1.6Hz, 1H), 6.06(br, 1H), 5.78(d, *J* = 10.0Hz, 1.6Hz, 1H), 3.84(s, 3H), 3.78(s, 3H), 3.05(br, 4H), 2.44(br, 4H), 2.23(s, 3H). HRMS (ESI) (*m/z*): [M + H]⁺ calcd for C₃₄H₃₅N₈O₄, 619.2781; found, 619.2780.


N-(3-(6-(3,5-difluorophenyl)-2-((2-methoxy-4-(4-methyl-1-piperazinyl)phenyl)amino)-7-oxo-8(7H)pteridinyloxy)phenyl)acrylamide (Compound 55)



red solids (yield 64%). mp 291.2-291.4 °C. ¹H NMR (400 MHz, DMSO-*d*₆): δ 10.43(s, 1H), 8.92(s, 1H), 8.57(s, 1H), 7.96(d, *J* = 7.2Hz, 1H), 7.87(br, 1H), 7.77(s, 1H), 7.54(t, *J* = 8.0Hz, 1H), 7.43-7.39(m, 1H), 7.33(s, 1H), 7.14(d, *J* = 7.2Hz, 1H), 6.54(s, 1H), 6.47(dd, *J* = 16.8Hz, 10.0, 1H), 6.27(dd, *J* = 16.8Hz, 1.6Hz, 1H), 6.00(br, 1H), 5.78(dd, *J* = 10.0Hz, 1.6Hz, 1H), 3.77(s, 3H), 3.05(br, 4H), 2.43(br, 4H), 2.22(s, 3H). HRMS (ESI) (*m/z*): [M + H]⁺ calcd for C₃₃H₃₁N₈O₃F₂, 625.2487; found, 625.2482.


N-(3-(6-(3,4-difluorophenyl)-2-((2-methoxy-4-(4-methyl-1-piperazinyl)phenyl)amino)-7-oxo-8(7H)pteridinyloxy)phenyl)acrylamide(Compound 56)

EP 16802532.8 – 8 November 2021



orange solids (yield 67%). Mp>300 °C. ¹H NMR (400 MHz, DMSO-*d*₆): δ 10.43(s, 1H), 8.89(s, 1H), 8.50(br, 1H), 8.28-8.23(m, 1H), 8.15(br, 1H), 7.87(s, 1H), 7.77(s, 1H), 7.61-7.52(m, 2H), 7.34(s, 1H), 7.14(d, *J* = 8.4Hz, 1H), 6.54(s, 1H), 6.47(dd, *J* = 16.8Hz, 10.0Hz, 1H), 6.27(dd, *J* = 16.8Hz, 1.6Hz, 1H), 6.02(br, 1H), 5.78(dd, *J* = 10.0Hz, 1.6Hz, 1H), 3.77(s, 3H), 3.05(br, 4H), 2.43(t, *J* = 4.4Hz, 4H), 2.22(s, 3H). HRMS (ESI) (*m/z*): [M + H]⁺ calcd for C₃₃H₃₁N₈O₃F₂, 625.2487; found, 625.2478.


N-(3-(2-((2-methoxy-4-(4-methyl-1-piperazinyl)phenyl)amino)-6-(3-methoxyphenyl)-7-oxo-8(7H)pteridinyl)phenyl)acrylamide (Compound 57)



Brown solids (yield 52%). Mp>300 °C. ¹H NMR (400 MHz, DMSO-*d*₆): δ 10.42(s, 1H), 8.88(s, 1H), 8.45(s, 1H), 7.89(s, 1H), 7.82-7.81(m, 2H), 7.76(s, 1H), 7.55(t, *J* = 8.0Hz, 1H), 7.41(t, *J* = 8.0Hz, 1H), 7.36-7.34(br, 1H), 7.16(d, *J* = 7.6Hz, 1H), 7.08(d, *J* = 8.4Hz, 1H), 6.54(s, 1H), 6.47(dd, *J* = 16.8Hz, 10.4Hz, 1H), 6.28(d, *J* = 17.2Hz, 1H), 6.04(br, 1H), 5.80(m, 1H), 3.82(s, 3H), 3.78(s, 3H), 3.05(br, 4H), 2.44(br, 4H), 2.23(s, 3H). HRMS (ESI) (*m/z*): [M + H]⁺ calcd for C₃₄H₃₅N₈O₄, 619.2781; found, 619.2780.

8-(1-acryloyl-3-piperidiny)-2-((2-methoxy-4-(4-methyl-1-piperazinyl)phenyl)amino)-6-phenyl-7(8H)pteridinone (Compound 58)

EP 16802532.8 – 8 November 2021



orange solids (yield 40%). mp ^1H NMR (400 MHz, $\text{DMSO}-d_6$): δ 9.19 (s, 1H), 8.78 (s, 1H), 8.11-8.09 (m, 2H), 7.47-7.46 (m, 3H), 7.22 (br, 1H), 6.88-6.63 (m, 2H), 6.49 (br, 1H), 6.06-6.19 (m, 1H), 5.67 (dd, $J = 36.8\text{Hz}$, 10Hz , 1H), 4.91-3.99 (m, 3H), 3.74 (s, 3H), 3.14 (br, 4H), 2.47(br, 4H), 2.24 (s, 3H), 1.69 (br, 2H), 1.35 (br, 1H), 1.23 (s, 1H). HRMS (ESI) (m/z): $[\text{M} + \text{H}]^+$ calcd for $\text{C}_{32}\text{H}_{37}\text{N}_8\text{O}_3$, 581.2989; found, 581.2992 .

Example 2

Assay on bioactivity: Activity test on molecular level (kinase)

EGFR-TK catalyses transfer of one phosphate group of adenosine triphosphate (ATP) to polypeptide substrate, poly(Glu, Tyr)_{4:1}, which is labeled with two fluorescent groups coumarin and fluorescein. Based on fluorescence energy resonance transfer (FRET) method, EGFR-TK catalyzes a reaction of ATP, resulting in two fluorescent groups approaching, exciting donator (coumarin) at 400 nM, releasing part of energy (emission wavelength at 445 nM), and transferring other part of energy to fluorescein (emission wavelength at 520 nM). Different compounds exhibited different inhibition-degrees on EGFR-TK, leading to different phosphorylation-degrees of substrate, so that the inhibition ratio of different compounds can be calculated by measuring the ratio of phosphorylation percentage of substrate catalyzed by an enzyme.

1. Experiment steps:

2.5 μL of Test Compounds, 5 μL Kinase / Peptide Substrate Mixture, 2.5 μL ATP Solution were added into a 384-well plate, and the 10 μL of reaction system was shaken for 30s and incubated at room temperature for 1 h. 5 μL of Development Solution was added and the 15 μL of reaction system was shaken for 30s, and incubated at room temperature for 1h. 5 μL of Stop Reagent was added, the reaction system in a total volume of 20 μL was shaken for 30s, and a microplate reader was used to detect fluorescence signals at excitation wavelength of 400 nm and emission wavelength of 445 nm and 520 nm, respectively. The inhibitory rates of the compounds at 7 concentration gradients were determined and the IC_{50} value for each compound was calculated by Origin 8.0 fitting curve. Positive controls were used during the experiment to confirm the feasibility of the

reaction system, and each experiment was performed in triplicate.

2. Calculation formula:

$$\text{Emission Ratio} = \frac{\text{Coumarin Emission (445 nm)}}{\text{Fluorescein Emission (520 nm)}}$$

$$\% \text{Phosphorylation} = 1 - \frac{(\text{Emission Ratio} \times F100\%) - C100\%}{(C0\% - C100\%) + \text{Emission Ratio} \times (F100\% - F0\%)}$$

$$\% \text{Inhibition} = \left(1 - \frac{\% \text{Phos sample}}{\% \text{Phos 0\%inhibition}}\right) \times 100$$

In Vitro Enzyme Activity Assay: Wild type and various mutant (T790M, L858 /
5 T790M) EGFR were purchased from Invitrogen. 10 concentration gradients from 5.1×10^{-11} mol / L to 1.0×10^{-6} mol / L were set for all compounds to be tested.

The concentration of different kinases was determined by optimization experiments and the corresponding concentrations were: EGFR (PV3872, Invitrogen) 0.287 µg/µL, EGFR-T790M (PV4803, Invitrogen) 0.174 µg/µL, EGFR-L858R/T790M (PV4879,
10 Invitrogen) 0.055 µg/µL. Compounds were diluted in DMSO for three times from 5.1×10^{-9} M to 1×10^{-4} M. 4 µL of the compound was dissolved in 96 µL of water to obtain a 4x compound solution. 40 µM ATP was dissolved in 1.33x kinase buffer and the kinase / peptide mixture contained 2x kinase, 4 µM tyrosine tetrapeptide for use. The 10 µL kinase reaction consisted of 2.5 µL compound solution, 5 µL kinase / peptide mix, 2.5 µL ATP
15 solution. 10 µL of kinase reaction included 2.5 µL of compound solution, 5 µL of kinase / peptide mixture, 2.5 µL of ATP solution. 5 µL of phosphorylated peptide solution, instead of the kinase / peptide mixture, was used as a 100% phosphorylation control. 2.5 µL of 1.33x kinase buffer, instead of ATP solution was used as a 100% inhibition control and 2.5 µL of 4% DMSO, instead of compound solution was used as a 0% inhibition control.
20 The plates were thoroughly mixed and incubated at room temperature for 1.5 hours. 5 µL of Development Solution was added into each well and then incubated for another 1 hour at room temperature, unphosphorylated peptides were cleaved during this period. Finally, 5 µL Stop Reagent was added to quench the reaction. The plates were measured with EnVision Multilabel Reader (Perkin Elmer). The experimental data were calculated using
25 GraphPad Prism version 4.0. Each experiment was repeated more than 3 times.

Example 3

Cell activity test (Inhibitory activity on cell proliferation)

Inhibition Analysis on Cell Proliferation and Growth: H1975 (non-small cell lung cancer cells, EGFR L858R / T790M), A431 (non-small cell lung cancer cells, EGFR wild
30 type) cells were obtained from the ATCC. Cell proliferation activity was assessed by MTS assay. Cells were exposed to processing conditions for 72 hours and the number of cells used in each experiment for each cell line was adjusted based on absorbance values

EP 16802532.8 – 8 November 2021

(absorbance values at 490 nm, 1.3-2.2). Six concentration gradients (0.1 nM - 10 μ M) were set for the compounds to be tested, wherein at least 6 sets of parallel controls for each concentration were used.

H1975, A431 cells were cultured in corresponding media, and cells were passaged at least twice after resuscitation and then used for experiments. Cells at Log phase were trypsinized and resuspended in culture medium. H1975 (1000 cells per well), A431 (2000 cells per well) were seeded in 96-well plates in a volume of 100 μ L; and 6 sets of parallel controls and 7 columns were set. The plate was placed in a 37°C 5% CO₂ incubator overnight. Compounds were dissolved in DMSO to a concentration of 10 μ M per liter, and then the concentration was gradually diluted to 10 μ M, 1 μ M, 0.1 μ M, 0.01 μ M, 0.001 μ M, 0.0001 μ M per liter, respectively. 2 μ L of the compound solution was added to 998 μ L of medium, and the mixture was thoroughly mixed. 100 μ L of the mixture was added to a 96-well plate. 2 μ L of DMSO, instead of compound solution was used as a 0% inhibition control. After culturing for 68 hours, 20 μ L of MTT (5 mg / mL) was added. After 4 hours, the supernatant was discarded and 150 μ L of DMSO was added. After shaking for 10 minutes, the plate was read with Synergy HT (Bio TeK) (OD490). Data were calculated using GraphPad Prism version 4.0 and IC₅₀ values were obtained using a non-linear regression model of dose response curve.

Determination of solubility in water: About 2 mg of the compound to be tested was weighed into a 1.5 mL centrifuge tube and 1 mL of PBS buffer (20 mM, pH = 6.8) was added, sonicated for 5 minutes (if the compound was completely dissolved, 2 mg of the compound was added again), and shaken overnight. The suspension was filtered through a 0.22 μ m PVDF filter, and then diluted to an appropriate concentration with methanol (v = 50%) and PBS (v = 50%), and the absorbance was measured by a UV spectrophotometer. About 3 mg of compound was accurately weighed, and methanol was added for preparing a stock solution of 1 mg / mL. The above stock solution was taken and prepared into a solution at concentrations of 0.05, 0.04, 0.03, 0.02 and 0.01 mg / mL, respectively, PBS solution (v = 50%) was added and the absorbance value was measured. Concentration c (mg / mL) as horizontal ordinate vs absorbance A as longitudinal coordinate was plotted, and a line was fitted for obtaining a standard curve (correlation coefficient greater than 0.998). The solubility of a compound in water was calculated from the standard curve.

Test results are shown in following Table 1.







Table 1. Kinase levels and selectivity, cellular level and selectivity, and partial solubility of Compound

Number of compound	Structure of compound	Inhibitory activity of EGFR on kinase (IC ₅₀ , nM)	Inhibitory activity on proliferation of cells (IC ₅₀ , μ M)	Solubility (pH 6.8, $\square$ g/mL)
--------------------	-----------------------	---	--	-------------------------------------







EP 16802532.8 – 8 November 2021

		WT	T790M/L858R	H1975	A431	
1*		250.5	129.0	12.16	13.75	1279.4
2*		>10uM	>10uM	18.25	89.61	11082
3*		>10uM	>10uM	33.67	60.62	>11352
4		4.5	3.9	0.009	0.94	1367
5		2.4	0.6	0.006	0.82	122.4
6*		105.8	14.8	0.46	1.69	51.8

EP 16802532.8 – 8 November 2021

7*		45.2	9.5	0.197	20.52	<25
8*		5.7	2.1	0.09	0.97	280.5
9*		23.4	20.8	1.04	2.81	162.7
10*		93.7	191.1	2.60	7.53	398.7
11*		2.2	7.3	0.20	8.09	48.6
12		3.5	6.2	0.006	2.52	369.4

EP 16802532.8 – 8 November 2021

13*		4.2	5.7	0.046	1.48	8.9
14		10.5	1.0	0.014	4.18	9.7
15*		2296	2250	3.60	8.13	514
16*		4704	>10000	>10	>10	956
17*		2190	191	1.42	>10	957
18*		47	1.0	0.06	1.66	208.2









EP 16802532.8 - 8 November 2021

19*		174.2	14.1	1.16	3.41	90.7
20		15.0	1.3	0.040	6.0	7.4
21*		2880	251	4.49	>10	72.2
22*		>10000	6897	>10	>10	3232
23*		1730	939	1.85	3.20	1689
24*		700	576	2.49	3.1	2918









EP 16802532.8 - 8 November 2021

25		65.2	1.5	0.016	1.75	56.67
26*		1302	559	3.23	9.5	10515
27		53.7	3.8	0.116	1.85	263.5
28*		>10000	>10000	8.38	>10	3969
29		177	791	3.09	1.6	437.5
30*		>1000	>1000	8.689	>10	> 10








EP 16802532.8 – 8 November 2021

31		522.3	9.0	2.155	>10	63.3
32*		1.3	7.1	0.033	2.753	<10
33*		33.0	1.3	0.230	3.682	24.5
34*		86.7	895.9	0.831	>10	223.1
35		20.1	17.5	0.378	0.225	63.3
36*		>1000	>1000	2.196	9.601	737
37*		26.0	4.9	0.074	3.235	<10
38		2.8	0.6	0.165	2.345	<10







EP 16802532.8 – 8 November 2021

39*		130.3	82.2	0.572	3.844	<10
40		>10000	73.1	1.052	1.377	>10
41		5346.8	38.6	1.186	1.263	>10
42		527.6	47.5	0.003	0.328	<10
43		7.7	0.6	0.112	0.433	<10
44		343.1	>1000	1.559	1.809	<10
45		>1000	>1000	1.775	4.276	<10
46		>10000	13.3	2.160	2.392	43.2

EP 16802532.8 – 8 November 2021

47		0.5	0.2	0.018	1.733	<10
48		361.6	20.8	0.031	0.640	<10
49		19.2	14.2	0.368	0.287	<10
50		2.7	86.9	1.244	1.112	719
51		16.6	250.6	5.338	2.060	<10
52		1.3	0.6	0.044	1.462	<10
53		9.6	74.2	0.479	0.593	<10

EP 16802532.8 – 8 November 2021




54		1.5	3.5	1.065	4.096	<10
55		6.6	6.1	0.262	1.299	<10
56		1.6	1.7	0.228	1.203	<10
57		2.8	1.8	0.225	1.369	<10
58						<10
AZD9291*		236	12.1	0.016	1.9	3.50

* comparative

The solubility of AZD9291 was tested at pH 7.4 (*J. Med. Chem.*, 2014, 57: 8249-8267)

Inhibition screening test on tyrosine kinase activity of Compound 53 (before and after resolution)

EP 16802532.8 – 8 November 2021

Number of compound	Structure of compound	Inhibitory IC ₅₀ means (nM) ± SD value on tyrosine kinase activity		Inhibitory IC ₅₀ means (μM) ± SD value on tumor cell growth activity	
		EGFR	EGFR/T790 M/L858R	A431	NCI-H1975
53		9.6	74.2	0.593	0.479
53-1 (R)		272.5±38.0	294.8±85.0	5.314±1.037	>10
53-2 (S)		8.0±0.8	15.2±2.3	>10	0.062±0.012

Example 4**1. H1975 cell xenograft tumor assay: Pharmacodynamic evaluation in animals**

Purpose of experiment: H1975 transplanted BALB/c nude mice were orally administered (PO) with a substance to be tested for 14 consecutive days. The body weight and tumor size of nude mice were observed and recorded. The changes of physiological activity of nude mice after administration were observed for evaluating antineoplastic efficacy.

1.1 Preparation of H1975 cell

H1975 cells were obtained from Shanghai Institute of Materia Medica. Cells were cultured in 1640 + 10% FBS (Gibco) at 37°C in a constant-temperature carbon dioxide incubator with 5% CO₂ concentration. Cells were fluid-changed and passaged for one time every 2-3 days and expanded. When the desired amount of cells was achieved, transplantation of cells was performed. Before transplantation, cell viability should be maintained above 90%, serum-free 1640 medium was used for suspending cells, and the cell concentration in suspension was 20 million / ml.

1.2 Modeling of H1975 xenograft tumor

40 male BALB/c nude mice, aged 4-5 weeks and weighing 17-23 g, were purchased from Shanghai Bikai, placed in an animal room environment for 2-3 days, and then seeded with cells. Cell suspension was subcutaneously injected at forelimb of the nude mouse rich in capillaries, and each mouse was injected with 0.1 mL of cell suspension.

Experimental conditions in animal room:

Laboratory temperature: 23 ± 3°C

EP 16802532.8 – 8 November 2021

Laboratory humidity: 30-60%

12 hours day-night rhythm

In about 2-3 weeks, xenograft tumor can be found, and mice were grouped and administered with a drug when average size of tumor reaching 300mm³.

5 1.3 Formulation of substance to be tested

Appropriate amount of a substance to be tested was weighed, suspended in 14 mL of DMSO-PEG400 saline solution (DMSO: PEG400: saline = 1:30:69) respectively, formulated according to the following table, suspended or dissolved by ultrasonic wave, filled into vials and sealed, and reserved in a 4°C refrigerator for further use.

Experiment group	Weighing (mg)	Actual concentration (mg/ml)	Dose (mg/kg)
1	0	0	0
2	17.5	1.25	5
3	35	2.5	10
4	87.5	6.25	25
5	175	12.5	50
6	350	25	100

10

1.4 Dosage and mode of administration

The mice were administered with a dosage of 0.1ml / 25g through oral gavage once a day, and the mice were observed for their status and abnormalities. The length and width of a tumor and body weight of a mouse were recorded at least twice a week. The tumor size is calculated as $V = 0.5 * \text{length} * \text{width} * \text{width}$.

15

Figure 1 shows pharmacodynamic evaluation of a series of compounds: after orally administered to nude mice of H1975 cell xenograft model for 14 days, Compound 14 exhibited superior efficacy over CO-1686.

Figure 2 shows pharmacodynamics comparison between Compound 14 and CO-1686 and AZD9291: Compound 14 (10 mg/kg) exhibited better efficacy over CO-1686 (50 mg/kg), Compound 14 (25 mg/kg) exhibited comparable efficacy to AZD9291 (25 mg/kg), both of which can effectively reduce the size of a tumor, and the mice still live and the tumor did not recur after administration was stopped for 1 month.

20

2. A431 cell xenograft tumor assay:

Purpose of experiment: A431 xenograft tumor BALB/c nude mice were orally administered (PO) with a substance to be tested for 14 consecutive days. The body weight and tumor size of nude mice were observed and recorded. The changes of physiological

25

EP 16802532.8 – 8 November 2021

activity of nude mice after administration were observed for evaluating antineoplastic efficacy.

2.1 Preparation of A431 cell

A431 cells belong to our laboratory, and in October 2015 confirmed by STR experiment. Cells were cultured in 1640 + 10% FBS (Gibco) at 37°C in a constant-temperature carbon dioxide incubator with 5% CO₂ concentration. Cells were fluid-changed and passaged for one time every 2-3 days and expanded. When the desired amount of cells was achieved, transplantation of cells was performed. Before transplantation, cell viability should be maintained above 90%, serum-free 1640 medium was used for suspending cells, and the cell concentration in suspension was 20 million/ml.

2.2 Modeling of A431 xenograft tumor

40 male BALB/c nude mice, aged 4-5 weeks and weighing 17-23 g, were purchased from Shanghai Bikai, placed in an animal room environment for 2-3 days, and then seeded with cells. Cell suspension was subcutaneously injected at forelimb of the nude mouse rich in capillaries, and each mouse was injected with 0.1 mL of cell suspension.

Experimental conditions in animal room:

Laboratory temperature: 23 ± 3°C

Laboratory humidity: 30-60%

12 hours day-night rhythm

In about 2-3 weeks, xenograft tumor can be found, and mice were grouped and administered with a drug when average size of tumor reaching 500mm³.

2.3 Formulation of substance to be tested

Appropriate amount of a substance to be tested was weighed, suspended in 14 mL of DMSO-PEG400 saline solution (DMSO: PEG400: saline = 1:30:69) respectively, formulated according to the following table, suspended or dissolved by ultrasonic wave, filled into vials and sealed, and reserved in a 4°C refrigerator for further use.

Experiment group	Weighing (mg)	Actual concentration (mg/ml)	Dose (mg/kg)
1	0	0	0
2	17.5	1.25	5
3	35	2.5	10
4	87.5	6.25	25
5	175	12.5	50
6	350	25	100

2.4 Dosage and mode of administration

EP 16802532.8 – 8 November 2021

The mice were administered with a dosage of 0.1ml / 25g through oral gavage once a day, and the mice were observed for their status and abnormalities. The length and width of a tumor and body weight of a mouse were recorded at least twice a week. The tumor size is calculated as $V = 0.5 * \text{length} * \text{width} * \text{width}$.

5 Figure 3 shows that compound **14** exhibited better selectivity on wild-type EGFR. Nude mice of EGFR wild-type A431 cell xenograft model were administered for 14 days by oral administration, compound **14** (25 mg/kg) exhibited performance comparable to the same dose of AZD9291, that is, the size of the tumor did not substantially change, and the selectivity was significantly better than afatinib (a medicament of 2nd generation).

10 **Example 4**

1. Acute Toxicity Evaluation of Compound: 10 times and higher dose of AZD9291 were selected for Acute toxicity test in ICR mice. In contrast, ZW-W-33 at 100 mg / kg, 250 mg / kg, and 500 mg / kg showed comparable toxicity to the same dose of AZD9291 with a slight decrease in body weight.

15 Figure 4 shows the toxicity assessment of Compound **14** and AZD9291.

2. Figure 5 shows the pathological analysis of tumor-bearing mice after 14 consecutive days of administration.

20 Conclusion: after long-term (14 days) administration of continuous treatment dose, no obvious pathological change was observed in all organs, and side effects were comparable to those of the same dose of positive drug AZD9291.


In addition, it should be understood that after reading the above teachings of the present invention, those skilled in the art can make various modifications or changes to the present invention.

25

EP 16802532.8 – 8 November 2021

Claims

1. A compound of formula I or a pharmaceutical acceptable salt thereof:



I

wherein

A is a divalent radical $-(CR_aR_b)_n-A'$, wherein R_a and R_b are independently selected from a group consisting of H, a C_{1-3} alkyl, a halogen, and n is an integer from 0 to 3;


m is an integer from 0 to 3;

R^1 is substituted or unsubstituted phenyl, optionally substituted C_3 - C_8 cycloalkyl;

A' is a benzene ring, a five- or six-membered nitrogen-containing heterocycle;


R^2 is independently selected from a C_2 - C_4 alkenyl substituted acylamino, substituted or unsubstituted C_2 - C_4 alkenyl substituted acyl;

R^7 is selected from a group consisting of:



R^3 , R^4 , R^5 and R^6 are independently selected from a group consisting of H, a substituted or unsubstituted C_1 - C_6 alkoxy, a substituted or unsubstituted C_1 - C_6 alkyl,


or a compound or a pharmaceutical acceptable salt thereof selected from the following group:








12.

EP 16802532.8 – 8 November 2021


2. The compound of claim 1, wherein the substituted or unsubstituted C₁-C₆ alkoxy is a C₁-C₃ alkoxy; and
the substituted or unsubstituted C₁-C₆ alkyl is a C₁-C₃ alkyl.
3. The compound or a pharmaceutical acceptable salt thereof of claim 1, selected from the following group:




EP 16802532.8 – 8 November 2021




31




38




40




41



46



52



53-2(S)

4. The compound of claim 1, wherein

R¹ is substituted or unsubstituted phenyl;

n = 0;

A' is a benzene ring, or a five-membered nitrogen-containing heterocycle;


R² is independently selected from a C₂-C₄ alkenyl substituted acylamino, or C₂-C₄

EP 16802532.8 – 8 November 2021

alkenyl substituted acyl;


m is an integer from 0 to 2;

R⁷ is selected from:



R³, R⁴, R⁵ and R⁶ are independently selected from a group consisting of H, a substituted or unsubstituted C₁-C₃ alkoxy, a substituted or unsubstituted C₁-C₃ alkyl.

5. The compound or a pharmaceutical acceptable salt thereof of claim 1, selected from the following group:



6. A pharmaceutical composition, comprising the compound or a pharmaceutical acceptable salt thereof of any one of claims 1-5 and a pharmaceutically acceptable carrier or excipient.

7. The compound of any one of claims 1-5 for use in treating or preventing EGFR-mediated diseases, or inhibiting EGFR, wherein the EGFR-mediated disease is cancer.

EP 16802532.8 – 8 November 2021

8. The compound for use of claim 7, wherein the cancer is selected from a group consisting of non-small cell lung cancer, small cell lung cancer, lung adenocarcinoma, lung squamous cell carcinoma, breast cancer, prostate cancer, glioblastoma, ovarian cancer, squamous cell carcinoma of head and neck, cervical cancer, esophageal cancer, liver cancer, kidney cancer, pancreas cancer, colon cancer, skin cancer, leukemia, lymphoma, stomach cancer, multiple myeloma and solid tumors.




Fig. 1




Fig. 2




Fig. 3
















Fig. 4

Tumor-bearing tissue	Control	Compound 14 25mg/kg	AZD9291 25mg/kg
Intestine			
	Normal tissue structure, no pathological change	Mucosal inflammatory cell focal infiltration, as indicated by the black arrow	Inflammatory cell focal infiltration seen in tissue, as indicated by the black arrow
Lung			
	Normal tissue structure	Normal tissue structure, no pathological change	Pulmonary congestion, interstitial edema
Liver			
	Normal tissue structure, no pathological change	Extensive hepatic sinusoids dilatation in the entire tissue, as indicated by the black arrow; visible mild fatty degeneration in liver tissue, with small droplets of lipid, as indicated by the yellow arrow	More severe liver injury, extensive hepatic sinusoids dilatation, as indicated by the black arrow; visible necrosis in liver tissue, as indicated by the yellow arrow
Tumor			
	Some part of nucleus shrinking in the tissue, as indicated by the yellow arrow; a slight necrosis of cells in the entire of tissue, as indicated by the black arrows	Moderate necrosis of cells in the entire tissues, as indicated by the black arrow	Moderate necrosis of cells in the entire tissues, as indicated by the black arrow










Brain			
	Normal tissue structure, no pathological change	Normal tissue structure, no pathological change	Normal tissue structure, no pathological change
Kidney			
	part of glomerular wall in the tissue thickening, as shown by the black arrow	Part of glomerular wall in the tissue thickening, as shown by the black arrow; no significant abnormality	Extensive kidney tubular lumen expansion in tissue, as shown by the black arrow; part of glomerular wall thickening, as shown by the yellow arrow; visible transparent tube
Heart			
	Moderate expansion of myocardial fiber gap in the tissue, as indicated by the black arrow; vacuoles formed in the cytoplasm of the myocardial fibers, as indicated by the yellow arrows	Moderate expansion of myocardial fiber gap in the tissue	Moderate expansion of myocardial fiber gap in the tissue, as indicated by the black arrow; vacuoles formed in the cytoplasm of the myocardial fibers, as indicated by the yellow arrows

Fig. 5