


Claims

1. A compound of formula I or a pharmaceutical acceptable salt thereof:



wherein

A is a divalent radical $-(CR_aR_b)_n-A'$ -, wherein R_a and R_b are independently selected from a group consisting of H, a C_{1-3} alkyl, a halogen, and n is an integer from 0 to 3;


m is an integer from 0 to 3;

R^1 is independently selected from a hydrogen, substituted or unsubstituted C_1-C_{10} alkyl, substituted or unsubstituted phenyl, optionally substituted C_3-C_8 cycloalkyl;

A' is a benzene ring, a five- or six-membered nitrogen-containing heterocycle;

R^2 is independently selected from a C_2-C_4 alkenyl substituted acylamino, substituted or unsubstituted C_2-C_4 alkenyl substituted acyl;

R^7 is selected from a group consisting of:



R^3 , R^4 , R^5 and R^6 are independently selected from a group consisting of H, a substituted or unsubstituted C_1-C_6 alkoxy, a substituted or unsubstituted C_1-C_6 alkyl.













2. The compound of claim 1, wherein the substituted or unsubstituted C_1-C_{10} alkyl is a substituted or unsubstituted C_1-C_6 alkyl;





the substituted or unsubstituted C_1-C_6 alkoxy is a C_1-C_3 alkoxy; and












the substituted or unsubstituted C_1-C_6 alkyl is a C_1-C_3 alkyl.


3. A compound or a pharmaceutical acceptable salt thereof selected from the

following group:









4. The compound of claim 1, wherein

R^1 is independently selected from a group consisting of a hydrogen, substituted or unsubstituted C_1 - C_6 alkyl, or substituted or unsubstituted phenyl;


$n = 0$;

A' is a benzene ring, or a five-membered nitrogen-containing heterocycle;

R^2 is independently selected from a C_2 - C_4 alkenyl substituted acylamino, or C_2 - C_4 alkenyl substituted acyl;


m is an integer from 0 to 2;


R^7 is selected from:



R^3 , R^4 , R^5 and R^6 are independently selected from a group consisting of H, a substituted or unsubstituted C_1 - C_3 alkoxy, a substituted or unsubstituted C_1 - C_3 alkyl.

5. A compound or a pharmaceutical acceptable salt thereof selected from the following group:





6. A pharmaceutical composition, comprising the compound or a pharmaceutical acceptable salt thereof of any one of claims 1-5 and a pharmaceutically acceptable carrier or excipient.

7. The compound of any one of claims 1-5 for use in treating or preventing EGFR-mediated diseases, or inhibiting EGFR, wherein the EGFR-mediated disease is cancer.

8. The compound for use of claim 7, wherein the cancer is selected from a group consisting of non-small cell lung cancer, small cell lung cancer, lung adenocarcinoma, lung squamous cell carcinoma, breast cancer, prostate cancer, glioblastoma, ovarian cancer, squamous cell carcinoma of head and neck, cervical cancer, esophageal cancer, liver cancer, kidney cancer, pancreas cancer, colon cancer, skin cancer, leukemia, lymphoma, stomach cancer, multiple myeloma and solid tumors.