

CLAIMS

1. A therapeutic agent that inhibits poly [ADP-ribose] polymerase (“anti-PARP therapy”) for use in treating cancer in a human patient, wherein the agent is niraparib, or a pharmaceutically acceptable salt thereof,
wherein the patient is characterized by previous treatment with a platinum-based therapy and an absence of a germline or a sporadic mutation in BRCA1 and/or BRCA2,
wherein the niraparib, or a pharmaceutically acceptable salt thereof, is administered as a maintenance therapy following complete or partial response to at least one platinum-based therapy, and
wherein the patient is further characterized by an absence of homologous recombination deficiency (HRD).
2. The therapeutic agent for use according to claim 1, wherein the patient has a tumor with a negative HRD status.
3. The therapeutic agent for use according to claim 1 or claim 2, wherein the niraparib, or a pharmaceutically acceptable salt thereof, is administered as a maintenance therapy following partial response to at least one platinum-based therapy.
4. The therapeutic agent for use according to any one of claims 1-3, wherein the niraparib, or a pharmaceutically acceptable salt thereof, is prepared in the form of a tosylate monohydrate.
5. The therapeutic agent for use according to any one of claims 1-4, wherein the cancer is selected from the group consisting of: ovarian cancer, fallopian tube cancer, peritoneal cancer, and breast cancer.
6. The therapeutic agent for use according to claim 5, wherein the ovarian cancer is high grade serous ovarian cancer or high grade predominantly serous histology ovarian cancer.

7. The therapeutic agent for use according to any one of claims 1-6, wherein the cancer is a recurrent cancer.
8. The therapeutic agent for use according to claim 7, wherein the cancer is selected from the group consisting of: ovarian cancer, fallopian tube cancer or primary peritoneal cancer.
9. The therapeutic agent for use according to claim 8, wherein the ovarian cancer, fallopian tube cancer or primary peritoneal cancer is platinum sensitive at the commencement of treatment.